

SAVANNAH ENVIRONMENTAL (PTY) LTD


REPORT ON:

MERAPI SOLAR PROJECT SOIL AND AGRICULTURAL POTENTIAL ASSESSMENT (FOUR PHASES)

REPORT: P276

Submitted to:

Savannah Environmental (Pty)Ltd
PO Box 148
Sunninghill
2157



VILJOEN & ASSOCIATES

DISTRIBUTION:

1 Copy - Savannah Environmental
1 Copy - Viljoen Associates
October 2012

Table of Contents

1	TERMS OF REFERENCE	1
2	INVESTIGATION OBJECTIVES	1
3	METHOD OF INVESTIGATION.....	2
	3.1 Sampling Procedures.....	2
	3.2 Inorganic Analyses.....	3
	3.3 Quality Assurance Quality Control.....	4
4	PROBLEM ANALYSES	4
	4.1 Basic Soil Forming Principles	4
	4.2 Abbreviated Legal Register for Rehabilitation.....	7
	4.3 South African Environmental Soil Legislation	10
	4.4 Objectives 1 and 2: Soil Classification and effective soil depth.....	11
	4.5 Objective 3: Agricultural potential	12
	4.6 Objective 4: Assessment of erodibility of soils and evidence of misuse	13
	4.7 Objective 5: Land Use & Land Capability	13
	4.8 Objective 6: Soil stripping utilisation guide and plan	15
	4.9 Objective 7: Overview of basic soil chemical, physical and mineralogical properties of soils.....	16
	4.10 Objective 8: Assessment of suitability of soils for rehabilitation purposes.....	17
	4.11 Objective 9: Impact assessment.....	18
5	IMPACT ASSESSMENT & RECOMMENDATIONS.....	24
6	ENVIRONMENTAL MANAGEMENT PROGRAMME	25
7	CONCLUSIONS	27
8	REFERENCES	29
9	EXPERIENCE AND EXPERTISE	29

EXECUTIVE SUMMARY

Savannah Environmental requested during May 2012 a proposal for a soil survey, land use land capability, wetland assessment for the Merapi Solar Project (four phases) near Bloemfontein in the Free State province.

The objectives of the investigation included a soil survey and mapping of study area, measurement of the effective depth of the soil(s), assessment of agriculture potential of soils, assessment of the erodibility and misuse of soils, mapping of land use & land capability, formulation of a soil stripping guide and plan, determination of chemical, mineralogical and physical properties of representative soil forms, assessment of suitability of soils for rehabilitation purposes and an impact assessment of topsoil stripping on soils with recommendations to mitigate negative impacts.

From the assessment it is conclusive the dominant soils according to the Taxonomical Soil Classification System of South Africa include Mispah, Sterkspruit and Clovelly soils. The effective depth of the soils is restricted to an average 300mm inclusive to the *Orthic A – Horizon*, however in the Clovelly soil the effective depth >300mm into the yellow brown apedalic B – Horizon.

The agricultural potential under dry land and irrigation conditions is available in **Table 3 (p13)**. The agricultural potential of the Mispah and Sterkspruit soils is considered medium to low and medium to high for the Clovelly soil under dryland (*650mm/y rainfall*) and irrigation conditions (*>10-15mm/week 33-1,500kPa plant available water*).

No evidence of soil erosion was observed on any of the soils during the investigation.

The current land use includes 235ha natural veld, 16ha ploughed land and 17ha plantation. The land capability includes 16ha arable, 235ha grazing and 17ha wilderness.

A soil stripping stockpiling strategy is given on **p16 (Table 6)**. A total area of 273ha could potentially be covered 300mm thick @ bulk density 1,275kg.m³ during rehabilitation taking into consideration a 10% loss of topsoil from the 912,000m³ due to handling, compaction etc.

The Mispah, Sterkspruit and Clovelly soils are characterised by neutral pH values (5,3 and 7,2) and low electrical conductivity values (<250mS/m). Under these conditions plant available nitrogen (15-20mg/kg), phosphorus (10-15mg/kg) and potassium (>50mg/kg) are readily available for plant uptake and sustainable plant growth. The *Orthic A-Horizon* is typically characterised by a low dense structure and texture distribution of approximately 65% sand, 20% silt and 15% clay with drainage properties in order of 10mm/h. The dominant clay mineral in the *Orthic A, Prisma-cutanic B and Red Apedalic B – Horizons* is kaolinite (*1:1 layer silicate*), with a low buffer capacity due to the low cation exchange capacity (<10cmol+/kg).

The Orthic soil horizon specified in **Section 4.4 p12** of the Mispah, Sterkspruit, Clovelly and Red Apedalic B horizon of the Clovelly soils are suitable for rehabilitation purposes.

The potential impacts and reasons/activities with proposed mitigation measures on the soil due to construction activities include:

- **Loss of topsoil:**

This is due to stripping, handling and placement of the soil associated with the pre-construction land clearing and rehabilitation and it is recommended to strip all usable soil irrespective of soil depth.

- **Change to soil's physical, chemical and biological properties:**

There is a high probability that topsoil will be lost due to wind and water erosion, which will alter the soils properties. Stockpiling and subsequent mixing of soil layers during handling will ultimately have a negative effect on altering the basic soil properties. It is suggested to implement live management and placement of topsoil where possible, improve the organic content of the soils, and maintain fertility levels through fertilisation and to curb topsoil loss as much as possible.

- **Cumulative effect of the soil:**

Alteration of the natural surface topography due to reprofiling during construction after stripping will have an accumulation effect on the soils and careful consideration should be given to minimise compaction and ensure free drainage preferential surface water pathways.

MERAPI SOLAR PROJECT SOIL AND AGRICULTURAL POTENTIAL ASSESSMENT (FOUR PHASES)

1 TERMS OF REFERENCE




Figure 1. Investigation area.

During May 2012 *Savannah Environmental (Pty) Ltd* requested a proposal for a baseline soil land use land capability wetland assessment for the Merapi Solar Project (four phases) near Bloemfontein in the Free State province. The study area is approximately 270ha (**Figure 1**).

2 INVESTIGATION OBJECTIVES

The objectives of the investigation were interpreted as:

- **Objective 1:** *Soil survey and mapping of study area.*
- **Objective 2:** *Measurement of the effective depth of the soil(s).*
- **Objective 3:** *Assessment of agriculture potential of soils.*
- **Objective 4:** *Erodibility and misuse of soils.*
- **Objective 5:** *Land use & land capability.*
- **Objective 6:** *Soil stripping guide and plan.*

- **Objective 7:** *Determination of chemical, mineralogical and physical properties of representative soil forms.*
- **Objective 8:** *Assessment of suitability of soils for rehabilitation purposes.*
- **Objective 9:** *Impact assessment of topsoil stripping on soils with recommendations to mitigate negative impacts.*

3 METHOD OF INVESTIGATION

In order to meet the objectives of the investigation, the following scope of work was conducted:

- A soil survey according to standard soil survey techniques comprising of auger holes on a flexible grid system GPS referenced (Datum 1984, degrees, minutes, seconds) to produce an electronic soil map to be available in AutoCAD, Micro Station, tif, pdf and/or jpeg.
- Soil profile studies according to the latest version of the Taxonomical Soil Classification System of South Africa.
- Representative sampling of different soil types.
- Inorganic analysis of the samples according to standard methods and techniques (**Table 1**).
- Interpretation of analytical data and field observations.
- Compilation of draft report.
- Internal review and submission of final report.

3.1 Sampling Procedures

Soil sampling carried out according to the following procedures:

- Auger holes drilled with a 75mm diameter 1,8m mechanical steel auger.
- The ground surface at the position of the auger hole cleared of loose material. If present, surface vegetation was carefully removed and the soil clinging to any roots left behind collected with the surface soil sample.
- The sampling interval in the auger holes was 150mm and consolidated to one sample per auger hole.

- The auger was advanced to the required depth and then carefully removed from the hole. The hole was covered to prevent foreign material from entering.
- Approximately 1.5kg soil sample was taken from the hole raisings and soil material removed from the auger. The samples were quartered to produce a representative sample of suitable weight, *i.e.* 500g.
- Prior to the taking of each sample both the steel auger and stainless steel trowel used to collect the soil samples were wiped clean of soil, washed with tap water, rinsed in a phosphate free detergent and finally sprayed with de-ionised water to prevent cross contamination between sampling depths.
- The soil samples were placed directly in zip-lock freezer bags, clearly labelled in indelible ink with the name of the site, auger hole number and sampling date.
- The soil samples were stored in the shade prior to being transported to an air-conditioned environment awaiting transport to the analytical laboratory.
- Chain of custody forms accompanied the soil samples to the laboratory and the samples were verified and signed for by the laboratory chemist.
- All auger hole logs were geo-referenced (GPS: datum WGS1984, decimal degrees).

3.2 Inorganic Analyses

Table 1 shows the analytical soil parameters for analyses.

TABLE 1. SOIL ANALYTICAL PARAMETERS

ELEMENT	METHOD
CHEMICAL	
Sample Preparation	Standard
pH (H ₂ O)	Standard
CEC+K+Na	NH ₄ Ac-extraction
EC+NO ₃	Saturated distilled water extract
P	Bray 1-extract
Lime Requirement	Double Buffer Titration
MINERALOGY	
Clay fraction (<0.002mm) identification	XRD-scan (6 treatments)
PHYSICAL	
Particle size distribution (3 fractions-sand+silt+clay)	Hydrometer

3.3 Quality Assurance Quality Control

The quality assurance/quality control procedure for the investigation entail a combination of the following:

- Duplicate analyses on 5% of the samples submitted.
- Carry out additional checks using standard reference materials.
- Conduct multi linear regression techniques to ensure analytical equipment are properly calibrated.
- Double check calibrated equipment with spiked standards above highest standard and confirm with 10x dilution.

4 PROBLEM ANALYSES

Section 4.1 is a brief description of basic soil forming principles to set a framework for evaluation of the draft scoping soil assessment:

4.1 Basic Soil Forming Principles




Figure 2. Typical soil profile.

According to A Glossary of Soil Science (1995), soil (**Figure 2**) can be defined as:

“the unconsolidated mineral and organic material on the immediate surface of the earth that serves as a natural medium for growth of plants, or, the unconsolidated mineral matter on the surface of the earth that has been subjected to and influenced by genetic and environmental factors of parent material, climate (including precipitation and temperature effects), macro- and micro-organisms and topography

all acting over the period of time and producing a product – soil – that differs from the material, which is derived in many physical, chemical, biological and morphological properties and characteristics”.

Soil is a thin surface covering the bedrock of most of the land area of the Earth. It is a resource that, along with water and air, provides the basis of human existence. Soil develops when rock is broken down by weathering and material is exchanged through interaction with the environment. Organic matter becomes incorporated into the soil as the result of the activity of living organisms. Soil also contains water, minerals, and gases. The soil system (**Figure 3**) is dynamic and it develops a distinct structure, often with recognizable layers or soil horizons arranged vertically through the soil profile.




Figure 3. Soil system with different layers.

Soil is essential for the development of most plants, providing physical support and nutrients. Plants are anchored in the soil by their roots. Nutrients, dissolved in soil water, are necessary for the plants' growth. Soil contains various organic matter, including dead material from plants and animals as well as animals that choose to

live in the soil. The soil is therefore a store of major nutrients such as carbon and nitrogen and plays an important role in global nutrient cycles and in regulating hydrological cycles and atmospheric systems.

Soils vary from place to place due to various conditions such as climate, rock type, topography, and the local soil-forming processes. Over time soils develop characteristics specific to their location, which relate closely to the climate and vegetation of the area. The major world biomes reflect a clear association between vegetation and soil that has developed in response to the prevailing climate. Each soil type has a distinct combination of soil horizons and associated soil properties.




Figure 4. Different stages of soil formation.

People depend on the soil for agriculture, and as such it is a valuable natural resource. Soils form continuously as the result of natural processes (**Figure 4**), and can therefore be regarded as a renewable resource. However, the soil-forming processes operate very slowly and the misuse or mismanagement of the soil may lead to damage or erosion, (**Figure 5**) or can disrupt the processes by which the soil forms.



Figure 5. Example of soil erosion (*not taken on site*).

If this happens the resource can be degraded or even lost and this is what should be prevented during topsoil stripping, stockpiling, replacement and rehabilitation. Many human activities cause damage to soils. These include bad farming techniques, overgrazing, deforestation, urbanization, construction, soil stripping, wars, contamination, pollution, and fires. The most critical result of these is soil erosion (**Figure 5**). With growing populations, the need for productive soils is increasing. Soil loss in many developing countries is a major cause for concern and will become a major issue in the future. The process of soil loss can have a detrimental effect on other systems as it produces sediment that can cause siltation of river systems and reservoirs, set off flooding downstream, and contribute to pollution and damage to estuaries, wetlands, and coral reefs. Soils need to be managed carefully in order to remain in good condition.

4.2 Abbreviated Legal Register for Rehabilitation

The following **Acts** focused on human rights, protection of the environment, accountability and financial provision should be considered with projects in South Africa:

- *Section 12 of the Minerals Act 50 of 1991.*
- *Sections 41, 42 and 43 of the Mineral & Petroleum Resources Development Act 28 of 2002, the M&PRD Regulations R527.*
- *Constitution of South Africa Act 108 of 1996.*
- *National Environmental Management Act 107 of 1998, and Amendments.*

-
- *National Water Act 36 of 1998 (Section 36), and Amendments, with specific reference to the NWA Regulations GN704 of 1999 and use of Water for Mining and Related Activities aimed at the Protection of Water Resources.*
 - *The Water Services Act 108 of 1997.*
 - *The Conservation of Agricultural Resources Act No. 43 of 1983 & Amendments (Govt. Gazette Vol. 429 No. 22166 of March 2001).*
 - *National Forest Act 84 of 1998.*
 - *Physical Planning Act of 1991.*
 - *National Environmental Management Biodiversity Act of 2003.*
 - *National Environmental Management Protected Areas Act of 2003.*
 - *National Veld and Forest Fire Act 101 of 1998.*
 - *Environment Conservation Act 73 of 1089.*
 - *Environment Conservation Amendment Act 50 of 2003.*
 - *Air Quality Act 39 of 2004.*
 - *National Heritage Resources Act 25 of 1999.*
 - *National Development Facilitation Act 67 of 1999.*
 - *National Development Facilitation Act 67 of 1995.*
 - *Promotion of Access to Information Act 2 of 2000.*
 - *National Monuments Act 28 of 1969.*
 - *Nuclear Energy Act 46 of 1999.*
 - *National Nuclear Regulator Act 47 of 1999.*
 - *Health Act 63 of 1997.*
 - *Plant Improvement Act 53 of 1976.*
 - *Occupational Health and Safety Act 85 of 1993.*

-
- *Agricultural Pests Act 36 of 1983.*
 - *Fertilisers, Farm Feeds, Agricultural remedies and Stock Remedies Act 36 of 1947.*
 - *Mine Health and Safety Act 29 of 1996.*
 - *Hazardous Substances Act 15 of 1973.*
 - *Land Survey Act 8 of 1997.*
 - *SABS 0286: 1998 Code of Practice for Mine Residue.*
 - *SABS: Water Quality.*
 - *Chamber of Mines of SA Guidelines for Environmental Protection: Engineering Design, Operation & Closure of Metalliferous, Diamond & Coal residue deposits.*
 - *Department of Mining & Energy Aide Memoir Guideline for the Peperation of EMPR'S.*
 - *Department of Mining & Energy Mineral Policy in terms of Section 12 of the Minerals Act 1995.*
 - *Department of Mining & Energy Policy on Financial Provision 1994.*
 - *Guideline on the Compilation of a Mandatory Code of Practice on Mine Residue Deposits.*
 - *Department of Water Affairs & Forestry Guideline on water & salt balances for TSF's.*
 - *Chamber of Mines Guidelines for Vegetation of Mine Residue Deposits.*
 - *Department of Water Affairs Policy and Guidelines for dealing with pollution from TFS's, and the containment and rehabilitation of abandoned TFS's, and prosecutions.*
 - *Convention of Wetlands of International Importance especially as Waterfowl Habitat RAMSAR (in force in SA from 12 Dec 1975).*
 - *International Cyanide Code.*

4.3 South African Environmental Soil Legislation

The following section outlines a summary of *South African Environmental Legislation* that needs to be considered for the proposed project with reference to management of soil:

- *The law on Conservation of Agricultural Resources (Act 43 of 1983) states that the degradation of the agricultural potential of soil is illegal.*
- *The Bill of Rights states that environmental rights exist primarily to ensure good health and well being, and secondarily to protect the environment through reasonable legislation, ensuring the prevention of the degradation of resources.*
- *The Environmental right is furthered in the National Environmental Management Act (No. 107 of 1998), which prescribes three principles, namely the precautionary principle, the “polluter pays” principle and the preventive principle.*
- *It is stated in the above-mentioned Act that the individual/group responsible for the degradation/pollution of natural resources is required to rehabilitate the polluted source.*
- *Soils and land capability are protected under the National Environmental Management Act 107 of 1998, the Environmental Conservation Act 73 of 1989, the Minerals Act 50 of 1991 and the Conservation of Agricultural Resources Act 43 of 1983.*
- *The National Veld and Forest Fire Bill of 10 July 1998 and the Fertiliser, Farm Feeds, Agricultural Remedies and Stock Remedies Act 36 of 1947 can also be applicable in some cases.*
- *The National Environmental Management Act 107 of 1998 requires that pollution and degradation of the environment be avoided, or, where it cannot be avoided be minimized and remedied.*
- *The Minerals Act of 1991, MPRDA requires an EMPR, in which the soils and land capability be described.*
- *The Conservation of Agriculture Resources Act 43 of 1983 requires the protection of land against soil erosion and the prevention of water logging and salinisation of soils by means of suitable soil conservation works to be constructed and maintained. The utilisation of marshes, water sponges and water courses are also addressed.*

Sections 4.4 to 4.11 address the investigation objectives (Section 2, p1) for the project.

4.4 Objectives 1 and 2: Soil Classification and effective soil depth




Figure 6. Soil types.

Figure 6 shows the distribution of soil types classified on the study area according to the latest version of the *South African Taxonomical Soil Classification System*.



Figure 7. Soil types: Mispah, Sterkspruit and Clovelly soils (left to right).

Figure 7 shows the diagnostic horizons of the Mispah, Sterkspruit and Clovelly soils classified according to the *South African Taxonomical Soil Classification System* summarised in **Table 2**:

TABLE 2. SOIL TYPES

SOIL TYPE	DIAGNOSTIC HORIZONS	EFFECTIVE DEPTH (MM)
Mispah	Orthic A – Horizon/Hard Rock	<300
Sterkspruit	Orthic A – Horizon/Prismacutanic B - Horizon	<300
Clovelly	Orthic A – Horizon/Yellow Brown Apedalic B – Horizon/Unspecified	>300

4.5 Objective 3: Agricultural potential

The agricultural potential was assessed using the following formula as a function of various variables:

$$YIELD (kg ha^{-1}) = R/B \times ED/A \times C \times X$$

Where:

R – Rainfall (mm)

B - Species growth characteristics factor.

ED - Effective depth of the soil.

A - Soil wetness factor for textural classes of soil above effective depth.

C - Correction factor for aeration of soil.

X - Fixed coefficient for species.

The main variables determining the soil's agricultural potential (**Table 3**) include the **effective depth** (>300mm), **clay content** (15%) and **rainfall** (650mm).

TABLE 3. AGRICULTURAL POTENTIAL OF SOILS.

SOIL TYPE	AGRICULTURAL POTENTIAL	
	DRY LAND	IRRIGATION
Mispah	Low	Medium
Sterkspruit	Low	Medium
Clovelly	Medium	High

4.6 Objective 4: Assessment of erodibility of soils and evidence of misuse

The exchangeable sodium percentage of the soils is be below 15% of the cation exchange capacity, rendering the soils free of dispersion anomalies caused by the hydration of sodium and consequent soil erosion.

4.7 Objective 5: Land Use & Land Capability



Figure 8. Land use.

Table 4 summarises the *land use* (**Figure 8**) of the area investigated:

TABLE 4. LAND USE

<u>Area</u>	<u>Land Use</u>	<u>Surface Area (ha)</u>	<u>% of Total</u>
Merapi	Natural Veld	235	89
	Plantations	17	6
	Wetlands/Dams/Pans		
	Ploughed Land	16	5
	Total	268	100




Figure 9. Land capability.

Table 5 summarises the *land capability* (**Figure 9**) of the area investigated:

TABLE 5. LAND CAPABILITY

<u>Area</u>	<u>Land Capability</u>	<u>Surface Area (ha)</u>	<u>% of Total</u>
Merapi	Arable	16	5
	Wilderness	17	6
	Grazing	235	89
	Wetlands/Dams/Pans		
	Total	268	100

4.8 Objective 6: Soil stripping utilisation guide and plan

It is recommended that all usable soil be stripped and stockpiled in advance of activities that might contaminate the soil.

The stripped soil should be stockpiled upslope of areas of disturbance or development to prevent contamination of stockpiled soils by dirty runoff or seepage. All stockpiles should also be protected by a bund wall to prevent erosion of stockpiled material and deflect surface water runoff.

Stockpiles can be used as a barrier to screen operational activities. If stockpiles are used as screens, the same preventative measures described above should be implemented to prevent loss or contamination of soil. The stockpiles should not exceed a maximum height of 6m and it is recommended that the side slopes and surface areas be vegetated in order to prevent water and wind erosion. If used to screen construction operations, the surface of the stockpile should not be used as a roadway as this will result in excessive soil compaction.

A conservative estimate of anticipated available topsoil to be stripped is summarised in **Table 6**.

TABLE 6. AVAILABLE TOPSOIL FOR REHABILITATION PURPOSES.

Soil Type & Average Effective Depth (mm)	Size (ha)	Available Volume (m ³)
Mispah (300)	21	63,000
Sterkspruit (300)	229	687,000
Clovelly (900)	18	162,000
TOTAL		912,000 @ BD: 1,275kg/m³

A total area of 273ha could potentially be covered 300mm thick @ bulk density 1,275kg.m³ during rehabilitation taking into consideration a 10% loss of topsoil from the 912,000m³ due to handling, compaction *etc.*

4.9 Objective 7: Overview of basic soil chemical, physical and mineralogical properties of soils

The Mispah, Sterkspruit and Clovelly soils are characterised by neutral pH values (5,3 and 7,2) and low electrical conductivity values (<250mS/m). Under these conditions plant available nitrogen (15-20mg/kg), phosphorus (10-15mg/kg) and potassium (>50mg/kg) are readily available for plant uptake and sustainable plant growth.

The Orthic A and Yellow Apedalic B-Horizon is typically characterised by a low dense structure and texture distribution of approximately 65% sand, 20% silt and 15% clay with drainage properties in order of 10mm/h.




Figure 10. 1:1 Clay mineral.

The dominant clay mineral in the *Orthic A – Horizon and Yellow Brown Apedalic B – Horizon* is kaolinite (*1:1 layer silicate*), with a low buffer capacity due to the low cation exchange capacity ($<10\text{cmol}^+/\text{kg}$).

4.10 Objective 8: Assessment of suitability of soils for rehabilitation purposes.

The Orthic A and Yellow Apedalic B - horizons specified in **Section 4.4 p12** of the Mispah, Sterkspruit and Clovelly soils are suitable for rehabilitation purposes.

When stockpiled soils have been replaced during rehabilitation, the soil fertility should be assessed to determine the level of fertilisation required to sustain normal plant growth. The fertility remediation requirements need to be verified at time of rehabilitation. The topsoil should be uniformly spread onto the rehabilitated areas and care should be taken to minimise compaction that would result in soil loss and poor root penetration.

When returning soil to the rehabilitation site care should be taken to place soil in a manner that will allow for levelling of soil to take place in a single pass. The soil profile should not be built up using a repeated tipping and levelling action to increase the soil depth.

Proper water control measures should be implemented to ensure a free draining rehabilitated landscape.

4.11 Objective 9: Impact assessment

The potential significance of environmental impacts identified during topsoil stripping was determined by using a ranking scale, based on the following (the terminology is from the DEAT guideline document on EIA Regulations, April 1998):

Occurrence

Probability of occurrence (how likely is it that the impact may occur?), and duration of occurrence (how long may it last?)

Severity

Magnitude (severity) of impact (will the impact be of high, moderate or low severity?), and scale/extent of impact (will the impact affect the national, regional or local environment, or only that of the site?).

In order to assess each of these factors for each impact, the following ranking scales (**Table 7**) were used:

TABLE 7. RANKING SCALES FOR IMPACT ASSESSMENT.

Probability:

5 – Definite/don't know

4 – Highly probable

3 – Medium probability

2 – Low probability

1 – Improbable

0 – None

Scale:

5 – International

4 – National

3 – Regional

Duration:

5 – Permanent

4 - Long-term (ceases with the operational life)

3 - Medium-term (5-15 years)

2 - Short-term (0-5 years)

1 – Immediate

Magnitude:

10 - Very high/don't know

8 – High

6 – Moderate

2 – Local

4 – Low

1 – Site only

2 – Minor

0 – None

Once the above factors had been ranked for each impact, the environmental significance of each was assessed using the following formula:

$$SP = (magnitude + duration + scale) \times probability$$

The maximum value is 100 significance points (SP). Environmental effects were rated as either of high, moderate or low significance on the following basis:

- More than 60 significance points indicated high environmental significance.
- Between 30 and 60 significance points indicated moderate environmental significance.
- Less than 30 significance points indicated low environmental significance.

TABLE 8. IMPACTS ON SOIL

Nature: Loss of topsoil due to stripping, handling and placement of soil associated with pre construction land clearing and rehabilitation.		
	Without Mitigation	With Mitigation
Extent	Local (1)	Local (1)
Duration	Long Term (4)	Short Term (1)
Magnitude	Moderate (6)	Low (4)
Probability	Very Probable (4)	Very Probable (4)
Significance	Moderate (44)	Low (24)
Status (positive or negative)	Negative	Negative
Reversibility	Irreversible	Irreversible
Irreplaceable loss of resources?	Irreplaceable	Irreplaceable
Can impacts be mitigated?	Yes	
Mitigation measures:		
<ul style="list-style-type: none"> Strip all usable soil, irrespective of soil depth. 		
Cumulative impact:		
<ul style="list-style-type: none"> Cumulative impact of loss of topsoil due to stripping and placement associated with pre construction land clearing and rehabilitation is considered low due to the undeveloped nature of the area but further development will increase impact. 		
Residual impact:		
<ul style="list-style-type: none"> Minor localised loss of topsoil 		

TABLE 8. IMPACTS ON SOIL/CONTINUED

Nature: Change of soil's physical, chemical and biological properties due to loss of topsoil due to erosion, stockpiling, mixing of deep and surface soils during handling, stockpiling and subsequent placement.		
	Without Mitigation	With Mitigation
Extent	Local (1)	Local (1)
Duration	Long Term (4)	Short Term (1)
Magnitude	Moderate (8)	Low (4)
Probability	Very Probable (5)	Very Probable (4)
Significance	Moderate (65)	Low (24)
Status (positive or negative)	Negative	Negative
Reversibility	Irreversible	Irreversible
Irreplaceable loss of resources?	Irreplaceable	Irreplaceable
Can impacts be mitigated?	Yes	
Mitigation measures:		
<ul style="list-style-type: none"> Implement live placement of soil where possible, improve organic status of soils, maintain fertility levels and curb topsoil loss. 		
Cumulative impact:		
<ul style="list-style-type: none"> Cumulative impact of soil's physical, chemical and biological properties due to loss of topsoil, due to erosion, stockpiling, mixing of deep surface soils during handling, stockpiling and subsequent placement is considered low due to the undeveloped nature of the area but further development will increase impact. 		
Residual impact:		
<ul style="list-style-type: none"> Minor localised degradation of topsoil's chemical, physical and biological properties. 		

TABLE 8. IMPACTS ON SOIL/CONTINUED

Nature: Change of natural surface topography due to reprofiling of surface after stripping.		
	Without Mitigation	With Mitigation
Extent	Local (1)	Local (1)
Duration	Long Term (4)	Short Term (1)
Magnitude	Moderate (8)	Low (4)
Probability	Very Probable (5)	Very Probable (4)
Significance	Moderate (65)	Low (24)
Status (positive or negative)	Negative	Negative
Reversibility	Irreversible	Irreversible
Irreplaceable loss of resources?	Irreplaceable	Irreplaceable
Can impacts be mitigated?	Yes	
Mitigation measures:		
<ul style="list-style-type: none"> Implement surface digital terrain mapping to ensure surface water control measures are implemented to ensure free draining system with minimal soil erosion. 		
Cumulative impact:		
<ul style="list-style-type: none"> Cumulative impact of the change of surface topography due to reprofiling of surface after stripping is considered low due to the undeveloped nature of the area but further development will increase impact. 		
Residual impact:		
<ul style="list-style-type: none"> Minor localised degradation of topsoil's chemical, physical and biological properties. 		

TABLE 8. IMPACTS ON SOIL/CONTINUED

Nature: Loss of land with high agricultural potential and land capability.		
	Without Mitigation	With Mitigation
Extent	Local (1)	Local (1)
Duration	Permanent (5)	Permanent (5)
Magnitude	Low (4)	Low (4)
Probability	High Probable (4)	High Probable (4)
Significance	Moderate (40)	Low (16)
Status (positive or negative)	Negative	Negative
Reversibility	Medium	Medium
Irreplaceable loss of resources?	No	No
Can impacts be mitigated?	Direct impacts cannot be mitigated but direct impacts can be minimised and avoided through adequate planning of layout.	
Mitigation measures:		
<ul style="list-style-type: none"> Loss of agricultural land is a long term loss and no mitigation measures exist. Mitigation is restricted to limitation of extent of impact to the immediate area of impact and minimisation of off-site impacts. 		
Cumulative impact:		
<ul style="list-style-type: none"> Soil erosion may arise due to altered surface water runoff. Management and erosion control measures should be implemented. 		
Residual impact:		
<ul style="list-style-type: none"> Loss of agricultural land is a long term loss, limited to the footprint of the project, which is minimal percentage surface area of the land. Agriculture can still continue on rest of unoccupied areas. 		

5 IMPACT ASSESSMENT & RECOMMENDATIONS

The results of the Impact Assessment for the proposed Merapi Solar Project (four phases) find the proposed activity will have a medium to low impact on the immediate and surrounding soil systems. Implementation and management of proposed mitigation measures will minimize loss of topsoil, prevent contamination of topsoil and stockpiled soil and prevent overall soil erosion.

Renewable energy projects contribute to clean energy generation as a sustainable resource and holds huge benefits for the local region and the country as a whole.

It is recommended that the proposed project be approved subjected to the mitigation measures stipulated in the Impact Assessment and Environmental Management Programme

6 ENVIRONMENTAL MANAGEMENT PROGRAMME

The environmental management programme summarise the key findings of the soil & agricultural mitigation measures and suggest potential management actions in order to mitigate the potential visual impacts.

TABLE 9. ENVIRONMENTAL MANAGEMENT PROGRAMME

OBJECTIVE: Mitigate the possible visual impact associated with construction phase.		
Project Component(s)	Construction site	
Potential Impact	Visual impact of general construction	
Activity/risk source	Potential impact on surrounding environmental receptors.	
Mitigation: Target/Objective	Minimal aesthetic disturbance by construction activities.	
Mitigation: Action/control	Responsibility	Timeframe
An Environmental Coordinator must manage environmental impacts in coordination with construction schedule.	Client	Pre-Construction
Contractors to sing and undertake environmental compliance.	Client	Pre-Construction
Keep disturbed areas and stockpiles to minimum to prevent soil loss.	Client/contractor	Construction
Identify suitable areas to stockpile stripped soil.	Client/contractor	Construction

TABLE 9. ENVIRONMENTAL MANAGEMENT PROGRAMME/CONTINUED

Ensure no surface runoff and seepage on site contaminate stockpiled soils and stripped areas.	Client/contractor	Construction
Minimise soil erosion through wind and water	Client/contractor	Construction
Remediate and rehabilitate disturbed areas in accordance with development plan	Client/contractor	Construction
Performance Indicator	Construction site is confined to demarcated boundaries and buffer zones. No transgression is allowed outside the set boundaries and protocol of the set specifications.	
Monitoring	Monitoring to be undertaken by a certified appointed Environmental Officer.	

7 CONCLUSIONS

- The dominant soils according to the Taxonomical Soil Classification System of South Africa are Mispah, Sterkspruit and Clovelly soils.
- The effective depth of the Mispah and Sterkspruit soils is on average 300mm restricted to the *Orthic A – Horizon*. The effective depth of the Clovelly soil is >300mm extending into the *Yellow Apedalic B-Horizon*.
- The agricultural potential under dry land and irrigation conditions is available in **Table 3 (p13)**. The agricultural potential of the Mispah and Sterkspruit soils is considered medium to low and for the Clovelly soil medium to high under dryland (*650mm/y rainfall*) and irrigation conditions (*>10-15mm/week 33-1,500kPa plant available water*).
- No evidence of soil erosion was observed on any of the soils during the investigation.
- The current land use includes 235ha natural veld, 16ha ploughed land and 17ha plantation. The land capability includes 16ha arable, 235ha grazing and 17ha wilderness.
- A soil stripping stockpiling strategy is given on **p16 (Table 6)**. A total area of 273ha could potentially be covered 300mm thick @ bulk density 1,275kg.m³ during rehabilitation taking into consideration a 10% loss of topsoil from the 912,000m³ due to handling, compaction etc.
- The Mispah, Sterkspruit and Clovelly soils are characterised by neutral pH values (*5,3 and 7,2*) and low electrical conductivity values (*<250mS/m*). Under these conditions plant available nitrogen (*15-20mg/kg*), phosphorus (*10-15mg/kg*) and potassium (*>50mg/kg*) are readily available for plant uptake and sustainable plant growth. The *Orthic A and Yellow Apedalic B--Horizon* is typically characterised by a low dense structure and texture distribution of approximately 65% sand, 20% silt and 15% clay with drainage properties in order of 10mm/h. The dominant clay mineral in the *Orthic A and Yellow Apedalic B – Horizon* is kaolinite (*1:1 layer silicate*), with a low buffer capacity due to the low cation exchange capacity (*<10cmol+/kg*).
- The Orthic A-horizon and Yellow Brown Apedalic B-Horizon specified in **Section 4.4 p12** of the Mispah, Sterkspruit and Clovelly soil types are suitable for rehabilitation purposes.
- The potential impacts and reasons/activities with proposed mitigation measures on the soil due to construction activities include:

- **Loss of topsoil:**

This is due to stripping, handling and placement of the soil associated with the pre construction land clearing and rehabilitation and it is recommended to strip all usable soil irrespective of soil depth.

- **Change to soil's physical, chemical and biological properties:**

There is a high probability that topsoil will be loss due to wind and water erosion, which will alter the soils properties. Stockpiling and subsequent mixing of soil layers during handling will ultimately have a negative effect on altering the basic soil properties. It is suggested to implement live management and placement of topsoil where possible, improve the organic content of the soils, and maintain fertility levels through fertilisation and to curb topsoil loss as much as possible.

- **Cumulative effect of the soil:**

Alteration of the natural surface topography due to reprofiling during construction after stripping will have an accumulation effect on the soils and careful consideration should be given to minimise compaction and ensure free drainage preferential surface water pathways.

- The approach for the soil assessment in the EIA will include a detailed site visit to assess the following parameters:
 - *Soil survey and mapping of study area.*
 - *Measurement of the effective depth of the soil(s).*
 - *Assessment of agriculture potential of soils.*
 - *Erodibility and misuse of soils.*
 - *Land use & land capability.*
 - *Soil stripping guide and plan.*
 - *Determination of chemical, mineralogical and physical properties of representative soil forms.*
 - *Assessment of suitability of soils for rehabilitation purposes.*
 - *Impact assessment of topsoil stripping on soils with recommendations to mitigate negative impacts.*

8 REFERENCES

VAN DER WATT H. AND VAN ROOYEN, T.H. 1990. A Glossary for Soil Science. V&R Printing Works (Pty) Ltd.

9 EXPERIENCE AND EXPERTISE

Viljoen Associates specialise in soil remediation, and have broad experience of soil surveys, geotechnical assessments, soil pollution investigations, soil remediation and rehabilitation of gold slimes dams, coal discard dams, industrial polluted areas, industrial effluent evaporation dams and footprints of gold slimes dams, principles & practise of environmental management and stabilisation of ecological sites that have been eroded naturally.

A combination of theoretical and practical soil chemistry, physics and mineralogy and 16 years professional experience of the mining and environmental industry have resulted in a sound grasp of specialist environmental remediation and rehabilitation issues.

Viljoen Associates have undertaken numerous soil specialist studies and have been a key project member of several large multi-disciplinary projects, including environmental impact assessments, mine closure planning and rehabilitation of gold tailings, coal discard dumps and industrially polluted sites.

This investigation was done on available information and subsequent interpretation of data to reveal the properties on site with the techniques described.



M.Sc. Pr. Sci. Nat. SACNASP (400131/96)