



GEOHYDROLOGY

GEOTECHNICAL

ENVIRONMENTAL

SOCIAL DEVELOPMENT



**Terrestrial Biodiversity, Plant and Animal Species Impact Assessment Report**  
**A TERRESTRIAL BIODIVERSITY IMPACT ASSESSMENT (PLANT AND ANIMAL SPECIES)**  
**FOR THE PROPOSED DEVELOPMENT OF THE MOPANE SOLAR PV 5 ON PORTION 2 OF**  
**THE FARM ROOIDRAAI 85 IQ, LOCATED WITHIN THE JB MARKS LOCAL MUNICIPALITY,**  
**DR KENNETH KAUNDA DISTRICT MUNICIPALITY, NORTH WEST PROVINCE**

November 2022

Prepared for: VOLTALIA SOUTH AFRICA (PTY) LTD

Compiled by Dr BJ Henning  
Document version 1.0 – Draft



Proudly Supporting  
TOUCHING AFRICA

Prepared by



**A TERRESTRIAL BIODIVERSITY IMPACT ASSESSMENT (INCLUDING PLANT AND ANIMAL SPECIES ASSESSMENT) FOR THE PROPOSED DEVELOPMENT OF THE MOPANE SOLAR PV 5 ON PORTION 2 OF THE FARM ROOIDRAAI 85 IQ, LOCATED WITHIN THE JB MARKS LOCAL MUNICIPALITY, DR KENNETH KAUNDA DISTRICT MUNICIPALITY, NORTH WEST PROVINCE**

November 2022

**Compiled by:**

Dr BJ Henning (*Pri Sci Nat – Ecological Science and Soil Science*)  
PhD. Plant Ecology  
M.Sc. Botany - Soil Science related

**Reviewed by:**

Mr. J.H.Botha (*Pr. Sci. Nat.- Ecological Science & Environmental Management*)  
B.Sc. Botany & Zoology, B.Sc. Hons Zoology,  
M.Sc. Geography and Environmental Management

LIMPOPO PROVINCE: 120 Marshall Street, Polokwane, 0699, PO Box 2526, Polokwane 0700,  
Tel: +27 15 291 1577 Fax: +27 15 291 1577 [www.ages-group.com](http://www.ages-group.com)

Offices: North West Eastern Cape Western Cape Limpopo Gauteng Kwazulu-Natal  
AGES Limpopo Directors: A von Well Dr. BJ Henning M Myburgh

## REPORT DISTRIBUTION LIST

<b>Name</b>	<b>Institution</b>
Ms. E. Grobler	AGES Limpopo
Mr. Joubert	VOLTALIA SOUTH AFRICA (PTY) LTD
	North West Department of Economic Development, Environment, Conservation and Tourism (NW DEDECT)
	Registered Interested and Affected Parties

## DOCUMENT HISTORY

<b>Report no</b>	<b>Date</b>	<b>Version</b>	<b>Status</b>
L22 091 E	November 2022	1.0	Draft

Although AGES Limpopo (Pty) Ltd exercises due care and diligence in rendering services and preparing documents, AGES Limpopo (Pty) Ltd accepts no liability, and the client, by receiving this document, indemnifies AGES Limpopo (Pty) Ltd and its directors, managers, agents and employees against all actions, claims, demands, losses, liabilities, costs, damages and expenses arising from or in connection with services rendered, directly or indirectly by AGES Limpopo (Pty) Ltd and by the use of the information contained in this document.

This document contains confidential and proprietary information of AGES Limpopo (Pty) Ltd and is protected by copyright in favour of AGES Limpopo (Pty) Ltd and may not be reproduced or used without the written consent of AGES Limpopo (Pty) Ltd, which has been obtained beforehand. This document is prepared exclusively for **VOLTALIA SOUTH AFRICA (PTY) LTD** and is subject to all confidentiality, copyright and trade secrets, rules, intellectual property law and practices of South Africa.

---

## Curriculum Vitae

### CURRICULUM VITAE

**B J Henning**

**PhD Plant Ecology**

#### PERSONAL DETAILS

Name: BAREND JOHANNES HENNING  
Date of Birth: 1976-09-06  
Profession/Specialization: Senior Ecologist  
Years with Firm: 6 years (previously 2006-2012 & since May 2020)  
Nationality: South African  
Years' experience: 15 years

#### QUALIFICATIONS

University attended: University of Pretoria, Pretoria (1995- 2002)  
PhD Plant Ecology, MSc (Botany), BSc (Hons.), BSc

#### COURSES

Advanced Wetland Course (UP CE, 2010)

Wetland Rehabilitation Course (UFS, 2015)

Course on wetland offsets (SANBI)

#### KEY QUALIFICATIONS AND EXPERIENCE

- Senior Ecologist / Soil Science Specialist for Ages Limpopo since September 2006 to 2012 and again since May 2020 involved in the following aspects:
  - Agricultural potential and land capability studies of soils on farms. (Reference: Mr Johan Botha, AGES Limpopo; 0152911577, Mr Herman Gildenhuys, Exigo; 0127512160;)
  - Spatial Development Frameworks.
  - Strategic Development Area Frameworks for local municipalities
  - Vegetation surveys, sensitivity, and zoning analysis of development sites, including eco-estates, mines, residential developments, shopping centres, roads, water supply and other related infrastructure etc (Reference: Mr Johan Botha, AGES Limpopo; 0152911577, Mr Herman Gildenhuys, Exigo; 0127512160;)
  - Faunal analysis and scoping reports (Reference: Mr Johan Botha, AGES Limpopo; 0152911577, Mr Herman Gildenhuys, Exigo; 0127512160)

## Terrestrial Biodiversity, Plant & Animal Species Impact Assessment Mopane Solar PV 5

- Avifauna studies related to solar plant and power line connection developments.
- Wetland delineations and functional capacity assessments (completed advanced wetland course of the Continued Education Department, University of Pretoria 2010 as well as Wetland rehabilitation course of the University of the North West).
- Wildlife Management Plans and habitat assessment for rare and endangered game species.
- GIS related functions.
- Senior Ecologist for Exigo (previously AGES Gauteng) November 2012 to November 2022. Involved in all the above mentioned aspects.
- Environmental Consultant for Envirodel Wildlife & Ecological Services cc and Dubel Integrated Environmental Services, Polokwane 2004 - 2006. Involved in the following aspects:
  - Wildlife management plans for game farms /reserves throughout the Limpopo Province
  - Environmental impact assessments (vegetation surveys and faunal scoping reports), habitat suitability analysis and report compilation.
  - Coordinating and performing grass monitoring surveys for the Limpopo Tourism and Parks Board
  - Soil potential studies.
- Environmental Consultant for Ficus – pro Environmental Services cc., Modimolle 2004 / 5. Involved mostly in fieldwork, report compilation or impact studies. Reference: Mr. R. Venter (0147173378)
- Subconsultant for AGES (Africa Geo-Environmental Services 2005-2006. Vegetation surveys and sensitivity zoning and analyses. Mr Johan Botha (0836449957)
- Eco-Agent environmental services cc, Pretoria 2002 - 2004. Involved in environmental impact studies. Prof G. J. Bredenkamp (0825767046), University of Pretoria.
- Enviroguard environmental services cc, Heidelberg 2002 - 2004. Involved in environmental impact studies. Prof L. R Brown (0825767046).
- GIS related aspects for all the above-mentioned aspects on projects

### POSITION AND DUTIES

Employed as Senior Ecological Specialist. Main duties and responsibilities include:

- Compilation of project proposals.
- Conducting specialist assessments

## Terrestrial Biodiversity, Plant & Animal Species Impact Assessment Mopane Solar PV 5

- Ecological assessments
- Soils and Land use potential studies.
- Wetland assessments.
- Wetland rehabilitation plans.
- Ecological & wetland monitoring.
- Biodiversity Action & Management Plans.
- Agricultural assessments.
- Avifauna assessments.
- Wildlife Management Plans and assessments.
- Rehabilitation Strategy & Implementation Programmes (RSIPs)
- Liaison with clients.
- GIS and map compilation.
- Project admin and management.
- Integration and interaction with the environmental consultants.
- Travelling.
- Any ad hoc duties that may be given by immediate manager.



**Declaration**

I, DR BJ Henning declare that -

- I act as the independent specialist.
- I will perform the work relating to the project in an objective manner, even if this results in views and findings that are not favourable to the project proponent.
- I declare that there are no circumstances that may compromise my objectivity in performing such work.
- I have expertise in conducting the specialist report relevant to this project, including knowledge of the National Environmental Management Act, 1998 (Act No. 107 of 1998) Gazette No. 43310 Government Notice R. 320, Plant and Animal Species Protocols, regulations and any guidelines that have relevance to the activity.
- I will comply with the Act, regulations and all other applicable legislation.
- I will consider, to the extent possible, the matters listed in Regulation 18 of the NEMA EIA Regulations.
- I have no, and will not engage in, conflicting interests in the undertaking of the activity.
- I undertake to disclose to the project proponent and the competent authority all material information in my possession that reasonably has or may have the potential of influencing - any decision to be taken with respect to the project; and - the objectivity of any report, plan or document to be prepared by myself for submission to the competent authority or project proponent.
- All the particulars furnished by me in this document are true and correct; and
- I realise that a false declaration is an offence in terms of Regulation 320 and is punishable in terms of section 24F of the Act.



---

**SIGNATURE OF SPECIALIST**  
**NOVEMBER 2022**

## Table of contents

---

<b>1</b>	<b>ASSIGNMENT .....</b>	<b>1</b>
<b>1.1</b>	<b>INFORMATION SOURCES .....</b>	<b>4</b>
<b>1.2</b>	<b>REGULATIONS GOVERNING THIS REPORT .....</b>	<b>5</b>
1.2.1	<i>National Environmental Management Act, 1998 (Act No. 107 of 1998) - Gazette No. 43310 Government Notice R. 320 .....</i>	<i>5</i>
1.2.2	<i>Conservation of Agricultural Resources Act (Act No. 43 of 1983) (CARA) .....</i>	<i>7</i>
1.2.3	<i>National Environmental Management Biodiversity Act (Act 10 Of 2004) (NEMBA) .....</i>	<i>7</i>
1.2.4	<i>The National Forest Act (Act 84 of 1998) (NFA) .....</i>	<i>7</i>
1.2.5	<i>Transvaal Nature Conservation Ordinance .....</i>	<i>7</i>
<b>1.3</b>	<b>TERMS OF REFERENCE .....</b>	<b>8</b>
1.3.1	<i>Rationale of solar plant development .....</i>	<i>8</i>
1.3.2	<i>Objectives .....</i>	<i>9</i>
1.3.3	<i>Scope .....</i>	<i>9</i>
1.3.4	<i>Limitations and assumptions .....</i>	<i>10</i>
<b>2</b>	<b>METHODS .....</b>	<b>12</b>
<b>2.1</b>	<b>VEGETATION AND PLANT SPECIES SURVEY .....</b>	<b>12</b>
2.1.1	<i>Data recorded: .....</i>	<i>12</i>
2.1.2	<i>Red data species .....</i>	<i>12</i>
2.1.3	<i>Protected trees .....</i>	<i>12</i>
2.1.4	<i>Protected plants .....</i>	<i>13</i>
2.1.5	<i>Data processing .....</i>	<i>13</i>
<b>2.2</b>	<b>FAUNA HABITATS AND ANIMALS' SPECIES SURVEY .....</b>	<b>13</b>
2.2.1	<i>Data recorded: .....</i>	<i>13</i>
2.2.2	<i>Red data species lists .....</i>	<i>14</i>
2.2.3	<i>Data processing .....</i>	<i>14</i>
<b>2.3</b>	<b>IMPACT RATING ASSESSMENT MATRIX .....</b>	<b>14</b>
<b>2.4</b>	<b>SENSITIVITY ASSESSMENT .....</b>	<b>16</b>
2.4.1	<i>Ecological function .....</i>	<i>16</i>
2.4.2	<i>Conservation importance .....</i>	<i>16</i>
2.4.3	<i>Sensitivity scale .....</i>	<i>17</i>
<b>2.5</b>	<b>EIA SCREENING TOOL .....</b>	<b>17</b>
<b>3</b>	<b>BASELINE ENVIRONMENT .....</b>	<b>18</b>
<b>3.1</b>	<b>LOCATION AND DESCRIPTION OF ACTIVITY .....</b>	<b>18</b>
<b>3.2</b>	<b>CLIMATE .....</b>	<b>23</b>
<b>3.3</b>	<b>GEOLOGY AND SOIL TYPES .....</b>	<b>23</b>
<b>3.4</b>	<b>TOPOGRAPHY, LANDUSES AND DRAINAGE .....</b>	<b>23</b>
<b>3.5</b>	<b>SENSITIVITY ANALYSIS AND CONSERVATION ANALYSIS TOOLS .....</b>	<b>24</b>
3.5.1	<i>NORTH WEST BIODIVERSITY CONSERVATION PLAN .....</i>	<i>24</i>
3.5.2	<i>PROTECTED AREAS NETWORK AND NATIONAL PROTECTED AREAS EXPANSION STRATEGY (NPAES) .....</i>	<i>26</i>
3.5.3	<i>IMPORTANT BIRD AREAS .....</i>	<i>28</i>
3.5.4	<i>NATIONALLY THREATENED ECOSYSTEMS .....</i>	<i>28</i>

3.5.5 STRATEGIC WATER SOURCE AREAS (SWSA), NATIONAL FRESHWATER ECOSYSTEM PRIORITY AREAS (NFEPA) STATUS OF RIVERS AND WETLANDS ON SITE ... 30

**4 RESULTS ..... 33**

**4.1 VEGETATION ..... 33**

4.1.1 Biome and Ecoregion ..... 33

4.1.2 Ecosystem drivers and ecological services ..... 33

4.1.3 Vegetation types ..... 34

4.1.4 Vegetation units ..... 34

**4.2 PLANT SPECIES LEVEL ASSESSMENT ..... 41**

4.2.1 Species of conservation concern ..... 41

4.2.2 Protected Plants (North West Nature Conservation Ordinance) ..... 43

4.2.3 Invasive alien species ..... 43

4.2.4 General ..... 45

**4.3 FAUNAL HABITAT AND ANIMAL SPECIES ASSESSMENT ..... 46**

4.3.1 Overview ..... 46

4.3.2 Results of desktop survey and site visits during November 2022 ..... 46

4.3.3 Fauna habitats of the project area ..... 46

4.3.4 Common fauna documented and potentially occurring on the development site ..... 46

4.3.5 Species of Conservation Concern (SCC) ..... 49

4.3.6 EIA screening tool listed species ..... 51

**5 POTENTIAL IMPACTS OF THE PROPOSED DEVELOPMENT ON THE FAUNA AND FLORA ..... 54**

**5.1 POTENTIAL IMPACTS ..... 54**

5.1.1 Direct habitat destruction ..... 54

5.1.2 Habitat fragmentation ..... 56

5.1.3 Increased Soil erosion and sedimentation ..... 57

5.1.4 Soil and water pollution ..... 57

5.1.5 Air pollution ..... 58

5.1.6 Spread and establishment of alien invasive species ..... 59

5.1.7 Negative effect of human activities and road mortalities ..... 60

**5.2 IMPACT ASSESSMENT MATRIX ..... 60**

**6 ECOLOGICAL SENSITIVITY CLASSES ..... 62**

**7 DISCUSSION ..... 64**

**8 CONCLUSION ..... 66**

**9 REFERENCES ..... 67**

**APPENDIX A. PLANT SPECIES IN QDS ..... 71**

**APPENDIX B. PLANT SPECIES FOUND ON SITE ..... 83**

**APPENDIX C. BIRD SPECIES LIST FOR QDS ..... 85**

**APPENDIX D MAMMAL SPECIES LIST ..... 89**

**APPENDIX E HERPETOFAUNA LIST ..... 90**

## List of Figures

Figure 1. Terrestrial Biodiversity Sensitivity as obtained from the EIA screening tool for the site.....	2
Figure 2. Animal Species Theme Sensitivity as obtained from the EIA screening tool for the site. ....	3
Figure 3. Plant Species Theme Sensitivity as obtained from the EIA screening tool for the site. ....	4
Figure 4. Regional location Map of the project area .....	20
Figure 5. Aerial Map indicating the proposed location of the Solar Plant and associated infrastructure .....	21
Figure 6. Topographical Map of the project area .....	22
Figure 7. North West C-Plan Map (2015) for the project area .....	25
Figure 8. Location of the project area in relation to listed protected areas. ....	27
Figure 9. Listed threatened ecosystems in proximity to the proposed development site (SANBI). ....	29
Figure 10. Location of the project area in relation to NFEPA Rivers and SWSA .....	32
Figure 11. Vegetation Unit Map of the proposed development area .....	36
Figure 12. South African red list categories indicating the categories to be used for Species of Conservation Concern .....	42
Figure 13. Sensitivity Map of the project area.....	63

## List of Tables

Table 1. Impact rating assessment weights.....	15
Table 2. Land types, geology, and dominant soil types of the proposed development site .....	23
Table 3. Botanical analysis and characteristics of <i>Schizachyrium – Trachypogon – Seriphium</i> rocky grassland.....	37
Table 4. Botanical analysis and characteristics of Open grassland with <i>Searsia pyroides</i> clumps.....	38
Table 5. Botanical analysis and characteristics of <i>Searsia pyroides</i> open woodland .....	40
Table 6. Red data and endemic species occurring in the project area of the QDS .....	42
Table 7. Declared weeds and invader plants of the study area .....	45
Table 8. Red data list of potential fauna for the study area .....	49
Table 9. Listed fauna species for the project area according to the EIA screening tool, status and habitat.....	51
Table 10. Impact assessment Matrix for the proposed development .....	61

## List of Photographs

Photograph 1. <i>Schizachyrium – Trachypogon – Seriphium</i> rocky grassland in the project area .....	38
Photograph 2. Open grassland with <i>Searsia pyroides</i> clumps in the project area .....	39
Photograph 3. <i>Searsia pyroides</i> open woodland in the project area .....	40
Photograph 4. Exotic bushclumps in the project area .....	41

## NOTATIONS AND TERMS

**Biota:** living things; plants, animals, bacteria

**Bottomland:** the lowlands along streams and rivers, on alluvial (river deposited) soil.

**Connectivity:** in this context, referring to either the upstream-downstream or lateral (between the channel and the adjacent floodplain) connectivity of a drainage line. Upstream-downstream connectivity is an important consideration for the movement of sediment as well as migratory aquatic biota. Lateral connectivity is important for the floodplain species dependent on the wetting and nutrients associated with overbank flooding.

**Endorheic:** closed drainage e.g., a pan.

**Floristic:** of flora (plants).

**Floodplain:** wetland inundated when a river overtops its banks during flood events resulting in the wetland soils being saturated for extended periods of time.

**Gley:** soil material that has developed under anaerobic conditions because of prolonged saturation with water. Grey and sometimes blue or green colours predominate but **mottles** (yellow, red, brown, and black) may be present and indicate localised areas of better aeration.

**Groundwater:** subsurface water in the zone in which permeable rocks, and often the overlying soil, are saturated under pressure equal to or greater than atmospheric.

**Horizon:** see soil horizons.

**Hydrophyte:** any plant that grows in water or on a substratum that is at least periodically deficient in oxygen because of soil saturation or flooding; plants typically found in wet habitats.

**Hydro-geomorphic:** refers to the water source and geology forms.

**Hydrology** is defined in this context as the distribution and movement of water through a wetland and its soils.

**Geomorphology** is defined in this context as the distribution and retention patterns of sediment within the wetland.

**Infilling:** dumping of soil or solid waste onto the wetland surface. Infilling generally has a very high and permanent impact on wetland functioning and is like drainage in that the upper soil layers are rendered less wet, usually so much so that the area no longer functions as a wetland.

**Mottles:** soils with variegated colour patterns are described as being mottled, with the "background colour" referred to as the matrix and the spots or blotches of colour referred to as mottles.

**Organic soil material:** soil material with a high abundance of un-decomposed plant material and humus.

**Palustrine (wetland):** all non-tidal wetlands dominated by persistent emergent plants (e.g., reeds) emergent mosses or lichens, or shrubs or trees (see Cowardin *et al.*, 1979).

**Perched water table:** the upper limit of a zone of saturation in soil, separated by a relatively impermeable unsaturated zone from the main body of groundwater.

**Permanently wet soil:** soil which is flooded or waterlogged to the soil surface throughout the year, in most years.

**Riparian:** the area of land adjacent to a stream or river that is influenced by stream-induced or related processes. Riparian areas which are saturated or flooded for prolonged periods would be considered wetlands and could be described as **riparian wetlands**. Some riparian areas are not wetlands (e.g., an area where alluvium is periodically deposited by a stream during floods, but which is well drained).

**Roughness coefficient:** an index of the roughness of a surface; a reflection of the frictional resistance offered by the surface to water flow.

**Runoff:** total water yield from a catchment including surface and subsurface flow.

**Seasonally wet soil:** soil which is flooded or waterlogged to the soil surface for extended periods (>1 month) during the wet season but is predominantly dry during the dry season.

**Sedges:** grass-like plants belonging to the family *Cyperaceae*, sometimes referred to as nutgrasses. Papyrus is a member of this family.

**Soil drainage classes:** describe the soil moisture conditions as determined by the capacity of the soil and the site for removing excess water. The classes range from very well drained, where excess water is removed very quickly, to very poorly drained, where excess water is removed very slowly. Wetlands include all soils in the very poorly drained and poorly drained classes, and some soils in the somewhat poorly drained class. These three classes are roughly equivalent to the permanent, seasonal and temporary classes.

**Soil horizons:** layers of soil that have uniform characteristics and have developed through pedogenic processes; they are bound by air, hard rock or other horizons (i.e., soil material that has different characteristics).

**Soil profile:** the vertically sectioned sample through the soil mantle, usually consisting of two or three horizons (Soil Classification Working Group, 1991).

**Soil saturation:** the soil is considered saturated if the water table or **capillary fringe** reaches the soil surface (Soil Survey Staff, 1992).

**Temporarily wet soil:** the soil close to the soil surface (i.e., within 50 cm) is wet for periods > 2 weeks during the wet season in most years. However, it is seldom flooded or saturated at the surface for longer than a month.

**Terrain unit classes:** areas of the land surface with homogenous form and slope. Terrain may be seen as being made up of all or some of the following units: crest (1), scarp (2), mid-slope (3), foot-slope (4) and valley bottom (5).

**Transpiration:** the transfer of water from plants into the atmosphere as water vapour

**Unchanneled valley bottom:** linear fluvial, net depositional valley bottom surfaces which do not have a channel. The valley floor is a depositional environment composed of fluvial or colluvial deposited sediment. These systems tend to be found in the upper catchment areas.

**Vegetation** is defined in this context as the vegetation structural and compositional state.

**Water regime:** when and for how long the soil is flooded or saturated.

**Water Quality** largely self-explanatory and reflecting the changes in quality because of changes in land use or as a direct result of activities within the wetland itself that could lead to changes in the quality of the water flowing through and within the wetland.

**Waterlogged:** soil or land saturated with water long enough for anaerobic conditions to develop.

**Wetland:** land which is transitional between terrestrial and aquatic systems where the water table is usually at or near the surface, or the land is periodically covered with shallow water, and which under normal circumstances supports or would support vegetation typically adapted to life in saturated soil.

**Wetland catchment:** the area up-slope of the wetland from which water flows into the wetland and including the wetland itself.

**Wetland delineation:** The determination and marking of the boundary of a wetland on a map.

**LIST OF ABBREVIATIONS**

<b>Abbreviation</b>	<b>Description</b>
ARC	Agricultural Research Council
C-Plan	Conservation Plan
CSIR	Council for Scientific and Industrial Research
DAFF	Department of Agriculture, Forestry and Fisheries
DEA	Department of Environmental Affairs
DME	Department of Minerals and Energy Affairs
DWS	Department of Water and Sanitation
EAP	Environmental Assessment Practitioner
EIA	Environmental Impact Assessment
EIS	Ecological Importance and Sensitivity
EMPR	Environmental Management Programme Report
ENPAT	Environmental Potential Atlas
GIS	Geographic Information Systems
GPS	Geographical Positioning System
HGM	Hydro-Geomorphic
HFI	Hydrological Function and Importance
IHI	Index of Habitat Integrity
IUCN	World Conservation Union
MAE	Mean Annual Evaporation
MAMSL	Meter Above Mean Sea Level
MAP	Mean Annual Precipitation
MAR	Mean Annual Runoff
NEMA	National Environmental Management Act
PES	Present Ecological State
PESC	Present Ecological Status Class
PQ4	Priority Quaternary Catchment
QDS	Quarter Degree Square
SADC	Southern African Development Community
SANBI	South African National Biodiversity Institute
WMA	Water Management Area
WHO	World Health Organisation



## 1 ASSIGNMENT

AGES Limpopo (Pty) Ltd was appointed by **VOLTALIA SOUTH AFRICA (PTY) LTD** to conduct a terrestrial biodiversity, plant species and animal species impact assessment for the proposed development of a solar plant named as follows:

- Mopane Solar PV 5.

The project site includes the establishment of a renewable energy generation facilities (Photovoltaic Power Plants) with associated infrastructure and structures, and power lines on Portion 2 of the farm Roodraai 85 IQ, located within the JB Marks Local Municipality, Dr Kenneth Kaunda District Municipality, North West Province. The project site is located  $\pm 7$  km northwest of Welverdiend along the border between Gauteng and the North West Province. The Eskom Carmel Main Transmission Substation (MTS) is located 16.4 km South-East of project sites.

The Species Environmental Impact Assessments Guideline has been developed in support of the Terrestrial Biodiversity, Plant and Animal Species protocols that were gazetted 30th October 2020 (Government Notice number 1150). This guideline provides details for implementing relevant species protocols and is available for use to plant and animal specialists, environmental assessment practitioners and Competent Authorities.

According to the national web-based environmental screening tool in terms of National Environmental Management Act (NEMA), 1998 (Act No. 107 of 1998), the site has the following sensitivities:

- Terrestrial Biodiversity: Very High Sensitivity (Figure 1).
- Animal Species Theme: Medium or Low Sensitivity (Figure 2).
- Plant Species Theme: Medium or Low Sensitivity (Figure 3).

A pre-screening site visit was therefore conducted to determine if the assessment was accurate and if the studies recommended should be conducted. After the site visit the following was concluded:

- The site has a HIGH Sensitivity from a terrestrial biodiversity perspective due to the presence of indigenous grassland.
- The site has a Medium Sensitivity from an Animal Species Theme Perspective due to the presence of natural fauna habitats.
- The site has a Medium Sensitivity from a Plant Species Theme Perspective due to the presence of indigenous grassland.

After the assessment, it was concluded that a detailed terrestrial biodiversity, plant species theme and animal species theme assessment should be conducted.

This report will include a detailed impact assessment of the proposed development site on the biodiversity of the site. This assessment is essential as it will contribute to meeting the


requirements of the National Environmental Management Act (NEMA), 1998 (Act No. 107 of 1998) in compliance with Gazette No. 43310 Government Notice R320.

The activities pertinent to this application are reflected below:


- **Activity 15 - The clearance of an area of 20 hectares or more of indigenous vegetation.**

“indigenous vegetation” refers to vegetation consisting of indigenous plant species occurring naturally in an area, regardless of the level of alien infestation and where the topsoil has not been lawfully disturbed during the preceding ten years.

The assignment is interpreted as follows: Compile a terrestrial biodiversity assessment on the flora (vegetation units), fauna and general ecology of the site and determine the potential impacts of the proposed development on the fauna and flora of the area as well as any impacts on the wetlands and proposed mitigation measures. The study will be done according to guidelines and criteria set by North West Department of Economic Development, Environment, Conservation and Tourism (NW DEDECT) and the regulations recently gazetted for biodiversity studies as well as animal and plant species protocols.



**Figure 1. Terrestrial Biodiversity Sensitivity as obtained from the EIA screening tool for the site.**



**Figure 2. Animal Species Theme Sensitivity as obtained from the EIA screening tool for the site.**




Figure 3. Plant Species Theme Sensitivity as obtained from the EIA screening tool for the site.

**1.1 INFORMATION SOURCES**

- All relevant topographical maps, aerial photographs and information (previous studies and environmental databases) related to the ecological components in the study area.
- Requirements regarding the fauna and flora survey as regulated by the newest terrestrial biodiversity, plant species theme and animal species theme protocols (National Environmental Management Act No. 107 of 1998 - Gazette No. 43310 Government Notice R. 320).

- Requirements regarding the fauna and flora survey as requested by North West Department of Rural, Environment and Agricultural Development (NW DREAD).
- Legislation pertaining to the fauna and flora study as relevant.
- Red data species list from the South African National Biodiversity Institute (SANBI), including the species data for the terrestrial biodiversity and the red listed species potentially occurring on site was obtained from the EIA screening tool prior to the site visit.
- Information on plant and animal species recorded for the various Quarter Degree Squares was extracted from the SABIF/SIBIS database hosted by SANBI and the faunal databases hosted by the Animal Demography Unit (ADU). This includes is a considerably larger area than the study area, but this is necessary to ensure a conservative approach as well as counter the fact that the site itself has not been well sampled in the past.
- Vegetation types and their conservation status were extracted from the South African National Vegetation Map (Mucina and Rutherford 2006) as well as the National List of Threatened Ecosystems (2011), where relevant.
- Critical Biodiversity Areas were obtained from the various coverages produced by the North West C-Plan.

## **1.2 REGULATIONS GOVERNING THIS REPORT**

### **1.2.1 National Environmental Management Act, 1998 (Act No. 107 of 1998) - Gazette No. 43310 Government Notice R. 320**

This report was prepared in terms of the National Environmental Management Act, 1998 (Act No. 107 of 1998) Gazette No. 43310 Government Notice R. 320. Specialist reports includes a list of requirements to be included in a specialist report for a Terrestrial Biodiversity, Plant Species and Animal Species Assessment

1. A specialist report or a report prepared in terms of these regulations must contain:
  - a. Details of
    - i. The specialist who prepared the report; and
    - ii. The expertise of that specialist to compile a specialist report, including a curriculum vitae.
  - b. A declaration that the specialist is independent in a form as may be specified by the competent authority.
  - c. An indication of the scope of, and purpose for which, the report was prepared.
  - d. The date and season of the site investigation and the relevance of the season to the outcome of the assessment.

- e. A description of the methodology adopted in preparing the report or carrying out the specialized process.
- f. The specific identified sensitivity of the site related to the activity and its associated structures and infrastructure.
- g. An identification of any areas to be avoided, including buffers.
- h. A map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers.
- i. A description of any assumptions made and any uncertainties or gaps in knowledge.
- j. A description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives, on the environment.
- k. Any mitigation measures for inclusion in the EMPr.
- l. Any conditions for inclusion in the environmental authorisation.
- m. Any monitoring requirements for inclusion in the EMPr or environmental authorisation.
- n. A reasoned opinion –
  - i. As to whether the proposed activity or portions thereof should be authorised and
  - ii. If the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr and where applicable, the closure plan.
- o. A description of any consultation process that was undertaken while preparing the specialist report.
- p. A summary and copies of any comments received during any consultation process and where applicable all responses thereto; and
- q. Any other information requested by the competent authority.

This Act also embraces all three fields of environmental concern namely: resource conservation and exploitation; pollution control and waste management; and land-use planning and development. The environmental management principles include the duty of care for wetlands and special attention is given to management and planning procedures.

### **1.2.2 Conservation of Agricultural Resources Act (Act No. 43 of 1983) (CARA)**

This Act regulates the utilization and protection of wetlands, soil conservation and all matters relating thereto; control and prevention of veld fires, control of weeds and invader plants, the prevention of water pollution resulting from farming practices and losses in biodiversity.

### **1.2.3 National Environmental Management Biodiversity Act (Act 10 of 2004) (NEMBA)**

The following aspects of the NEMBA (2004) are important to consider in the compilation of an ecological report. It:

- Lists ecosystems that are threatened or in need of national protection.
- Links to Integrated Environmental Management processes.
- Must be considered in EMPs and IDPs.
- The Minister may make regulations to reduce the threats to listed ecosystems.

### **1.2.4 The National Forest Act (Act 84 of 1998) (NFA)**

In terms of section 15(1) of the National Forests Act, 1998, no person may cut, disturb, damage, or destroy any protected tree; or possess, collect, remove, transport, export, purchase, sell, donate or in any other manner acquire or dispose of any protected tree or any product derived from a protected tree, except under a licence or exemption granted by the Minister of Agriculture, Forestry and Fisheries.

### **1.2.5 Transvaal Nature Conservation Ordinance**

This Act deals with the following:

- To provide for the sustainable utilisation and protection of biodiversity within the North West Province.
- To provide for professional hunting.
- To provide for the preservation of caves and cave formations.
- To provide for the establishment of zoos and similar institutions.
- To provide for the appointment of nature conservators.
- To provide for the issuing of permits and other authorisations.
- To provide for offences and penalties for contravention of the Act.
- To implement the provisions of the Ordinance and to provide for matters connected therewith.

### 1.3 TERMS OF REFERENCE

#### 1.3.1 Rationale of solar plant development

South Africa currently relies principally on fossil fuels (coal and oil) for the generation of electricity. At the present date, Eskom generates approximately 90% of the electricity used in South Africa. However, South Africa has a largely unexploited potential in renewable energy resources such as solar, wind, biomass, and hydro to produce electricity as opposed to other energy types (liquid fuel or coal). South Africa's electricity supply still heavily relies upon coal power plants, whereas the current number of renewable energy power plants is still limited. In the last few years, the demand for electricity in South Africa has been growing at a rate approximately 3% per annum.

These factors, if coupled with the rapid advancement in community development, have determined the growing consciousness of the significance of environmental impacts, climate change and the need for sustainable development. The use of renewable energy technologies is a sustainable way in which to meet future energy requirements.

The development of clean, green, and renewable energy has been qualified as a priority by the Government of South Africa with a target for 2013 of 10,000 GWh, as planned in the Integrated Resource Plan 1 (IRP1) and with the Kyoto Protocol. Subsequently the Department of Energy of South Africa (DoE) decided to undertake a detailed process to determine South Africa's 20-year electricity plan, called Integrated Resources Plan 2010-2030 (IRP 2010). The IRP1 (2009) and IRP 2010 (2011, updated in March 2014 and in October 2019) outline the Government's vision, policy, and strategy in matter of the use of energy resources and the current status of energy policies in South Africa.

The purpose of the proposed Solar Photovoltaic Plant is to add new capacity for the generation of renewable electric energy to the national electricity supply in compliance with the updated IRP 2019 in order to meet the "sustainable growth" of the North West Province. The use of solar radiation for power generation is considered as a non-consumptive use and a renewable natural resource which does not produce greenhouse gas emissions. The generation of renewable energy will contribute to the growth of South Africa's electricity market, which has been primarily dominated up to this date by coal-based power generation. With specific reference to photovoltaic energy, and the proposed project, it is important to consider that South Africa has one of the highest levels of solar radiation in the world.

The proposed solar park will assist the Eskom grid to meet the high energy demand related to the industrial activities conducted in the Carletonville area. The purpose of the proposed Mopane Solar PV 5 is to add new capacity for the generation of electrical energy to the national electricity supply, in compliance with the Minister's Determinations and to meet "electricity consumptions growth" of the North West Province.



### 1.3.2 Objectives

1. The primary aim of this project is to investigate options for enhancing and / or maintaining biodiversity to mitigate the impact of the development and related infrastructure with the overall objective of preventing further loss of biodiversity. The product would be a tool for promoting and lobbying for the recognition of the importance of species habitat and habitat conservation. Options available to maintain the current level of floral diversity include:
  - a. Protection of native vegetation restored elsewhere in return for unavoidable clearing.
  - b. Minimisation of habitat fragmentation.
  - c. Minimisation of any threats to the native flora and fauna and their habitats during the construction and operational phases of the developments and.
  - d. Rehabilitation to establish plant communities / landscaping that will provide future habitat values.
2. To produce a clear and agreed species and habitat priorities for conservation actions. This includes the following:
  - i. Determine the ecological impacts and actions the developments will have on the biodiversity on a species and habitat level.
  - ii. Conduct a risk analyses of the impacts identified to determine the significance of the impacts on the fauna and flora of the study area.
  - iii. Protection and enhancement of vegetation habitats of high conservation value.
  - iv. Retention of a substantial amount of native vegetation/habitat of adequate size and configuration to promote conservation of existing flora communities.
  - v. Retention and/or creation of vegetation links, wildlife corridors and vegetation buffers wherever possible, subject to appropriate bush fire risk management;
  - vi. The protection of water quality in the locality so as not to threaten native aquatic flora that rely on the watercourse for survival.
3. Provide recommendations on the ecological mitigation measures to be implemented by the developer and the way forward.

### 1.3.3 Scope

1. Conduct a field study to determine the state of the vegetation on site:
  - i. After studying the aerial photograph determine the previous state of the vegetation compared to the current state of the vegetation on site.
  - ii. Conduct a site visit and list the plant species (trees, shrubs, grasses, succulents and other herbaceous species of special interest) present for plant communities still present after construction.
  - iii. Identify potential red data plant species, possible encroacher species, medicinal plants of value and exotic plant species.

2. Determine the ecological impact the development will have on the fauna and flora of the site and conduct an impact rating assessment.
3. Fauna scoping
  - a. List the potential fauna (mammal species, red data birds, reptiles, amphibians, invertebrates) present linked to the specific potential habitats that occur as identified in the vegetation survey.
  - b. Analyse the data and identify potential red data fauna species, as well as other endemic or protected species of importance.
  - c. Indicate species mitigation measures and management measures to be implemented to prevent any negative impacts on the fauna of the area.
4. General
  - a. Identify and describe ecologically sensitive areas. Create a sensitivity map to indicate specific sensitive areas based on various environmental parameters like natural vegetation in a good condition, rockiness, slopes, floodlines etc.
  - b. Identify problem areas in need of special treatment or management, e.g., bush encroachment, erosion, degraded areas, reclamation areas.
  - c. Make recommendations, impact ratings and risk assessments for each specific impact.

#### **1.3.4 Limitations and assumptions**

- Maintaining due cognisance of the integrity and accuracy of the ecological survey, it should be stated that the ecological resources identified during the study do not necessarily represent all the ecological resources present on the property.
- To obtain a comprehensive understanding of the dynamics of communities and the status of endemic, rare or threatened species in an area, ecological studies should ideally be replicated over several seasons and over a few years. However, due to project time constraints such long-term studies are not feasible.
- Most threatened plant species are extremely seasonal and only flower during specific periods of the year,
- Most threatened faunal species are extremely secretive and difficult to survey even during thorough field surveys conducted over several seasons.
- The detailed surveys focused on the proposed development footprint of the solar park. Although surveys were conducted in other areas of the site during the pre-screening and siting exercise, these areas were identified as sensitive and unsuitable for the development, and therefore no further surveys in these areas were considered necessary.

Thus, even though it might be assumed that survey findings are representative of the ecosystem of the site for the development activities, it should be stated that the possibility exists that individual plants species might have been missed due to the nature of the terrain and size of the study area. Therefore, maintaining due cognisance of the integrity and accuracy of the ecological survey, it should be stated that the ecological resources identified during the study do not necessarily represent all the ecological resources present on the property.

## 2 METHODS

### 2.1 VEGETATION AND PLANT SPECIES SURVEY

Two basic methods were used during the vegetation survey:

- Line transects were walked on the site surveyed to record the plant species present. Rare and threatened plant species and any botanically sensitive sites or habitats were searched for in the various vegetation units.
- The Braun-Blanquet survey technique to describe plant communities as ecological units was also used for this study. It allows for the mapping of vegetation and the comparison of the data with similar studies in the area.

**The site surveys were conducted on 8 and 9 November 2022. The relevance of the season (summer months) had NO impact on the outcome of the assessment.** The vegetation was in a moderate to good condition and most species could be identified, although some species might have been missed due to timing of the flowering season.

The field work was conducted during November 2022 and the timing of the seasonal survey was considered as sufficient due to adequate early rains that fell in the area during October and Early November 2022.

The seasonal survey in November 2022 is considered as sufficient to identify fauna habitats, vegetation units as well as potential red listed flora and fauna. Red listed flora have varying flowering times and early season was considered as a suitable time to conduct the survey and identify red listed and protected flora associated with the grassland vegetation types.

#### 2.1.1 Data recorded:

Plant names used in this report are in accordance with Arnold & De Wet (1993), except for a few newly revised species. A list of all plant species present, including trees, shrubs, grasses, forbs, geophytes, and succulents were compiled. All identifiable plant species were listed. Notes were additionally made of any other features that might have an ecological influence as well as potential fauna habitat that might occur.

#### 2.1.2 Red data species

A species list of the red data species previously recorded in the vicinity of the development was obtained from the EIA screening tool as well as the South African Biodiversity Institute (SANBI), South Africa as classified by the IUCN red data list categories.

#### 2.1.3 Protected trees

A species list of the protected tree species was obtained from the Department of Forestry. These trees are listed by the NFA (Act 84 of 1998) as protected.

#### 2.1.4 Protected plants

A list of protected and specially protected plants was obtained from the Transvaal Nature Conservation Ordinance.

#### 2.1.5 Data processing

A classification of vegetation data was done to identify, describe and map vegetation types. The descriptions of the vegetation units include the tree, shrub, and herbaceous layers.

Conservation priority of each vegetation unit was assessed by evaluating the plant species composition in terms of the present knowledge of the vegetation of the North West Province, as well as the vegetation type.

The following four conservation priority categories were used for each vegetation unit:

- High: Ecologically sensitive and valuable land with high species richness that should be conserved, and no development allowed.
- Medium: Land that should be conserved but on which low impact development could be considered with the provision of mitigation measures.
- Medium-low: Land that has some conservation value but on which development could be considered with limited impact on the vegetation / ecosystem. It is recommended that certain sections of the vegetation be maintained.
- Low: Land that has little conservation value and that could be considered for developed with little to no impact on the vegetation / ecosystem.

## 2.2 FAUNA HABITATS AND ANIMALS' SPECIES SURVEY

The fauna survey was conducted as follows:

- A site survey was done to identify potential habitats after identifying the vegetation units. Fauna observed on site or any specific indication of species was noted as confirmed in the species lists.
- A scoping survey was then conducted by comparing the habitat types identified with the preferred habitats of species occurring in the area.
- A survey was thereafter conducted to document species occurring in the habitats on site.

#### 2.2.1 Data recorded:

A list of all species of fauna and their status as observed on the site or that could potentially occur on the site. Notes were made of any specific sensitive or specialized habitats that occur on the site.

### 2.2.2 Red data species lists

A species list of the red data species of the different faunal classes was obtained from the following references:

- EIA screening tool as relevant for the project area.
- Red Data Book of the Mammals of South Africa (Friedman & Daly, 2004)
- The Atlas of the Southern African Birds - digital data on quarter degree grid data (Avian Demography Unit, University of Cape Town)
- Atlas and red data book of the frogs of South Africa, Lesotho, and Swaziland (Minter et al. 2004)
- South African Red Data Book – Reptiles and Amphibians. National Scientific Programmes Report no. 151.

### 2.2.3 Data processing

A comparison of the habitats (vegetation units) occurring on the property was made to the preferred habitats of the faunal species. In addition to species observed on the site, lists of the potential mammal, bird, reptile, amphibian, and insect species were compiled and mitigating measures recommended if needed.

## 2.3 IMPACT RATING ASSESSMENT MATRIX

An impact can be defined as any change in the physical-chemical, biological, cultural and/or socio-economic environmental system that can be attributed to human activities related to alternatives under study for meeting a project need.

The significance of the impacts will be determined through a synthesis of the criteria below (Plomp, 2004):

**Probability.** This describes the likelihood of the impact occurring:

- **Improbable:** The possibility of the impact occurring is very low, due to the circumstances, design, or experience.
- **Probable:** There is a probability that the impact will occur to the extent that provision must be made, therefore.
- **Highly Probable:** It is most likely that the impact will occur at some stage of the development.
- **Definite:** The impact will take place regardless of any prevention plans, and there can only be relied on mitigation actions or contingency plans to contain the effect.

**Duration.** The lifetime of the impact

- **Short term:** The impact will either disappear with mitigation or will be mitigated through natural processes in a time span shorter than any of the phases.

- **Medium term:** The impact will last up to the end of the phases, where after it will be negated.
- **Long term:** The impact will last for the entire operational phase of the project but will be mitigated by direct human action or by natural processes thereafter.
- **Permanent:** Impact that will be non-transitory. Mitigation either by man or natural processes will not occur in such a way or in such a time span that the impact can be considered transient.

**Scale.** The physical and spatial size of the impact

- **Local:** The impacted area extends only as far as the activity, e.g., footprint.
- **Site:** The impact could affect the whole, or a measurable portion of the above-mentioned properties.
- **Regional:** The impact could affect the area including the neighbouring areas.

**Magnitude/ Severity.** Does the impact destroy the environment or alter its function?

- **Low:** The impact alters the affected environment in such a way that natural processes are not affected.
- **Medium:** The affected environment is altered, but functions and processes continue in a modified way.
- **High:** Function or process of the affected environment is disturbed to the extent where it temporarily or permanently ceases.

**Significance.** This is an indication of the importance of the impact in terms of both physical extent and time scale, and therefore indicates the level of mitigation required.

- **Negligible:** The impact is non-existent or unsubstantial and is of no or little importance to any stakeholder and can be ignored.
- **Low:** The impact is limited in extent, has low to medium intensity; whatever its probability of occurrence is, the impact will not have a material effect on the decision and is likely to require management intervention with increased costs.
- **Moderate:** The impact is of importance to one or more stakeholders, and its intensity will be medium or high; therefore, the impact may materially affect the decision, and management intervention will be required.
- **High:** The impact could render development options controversial or the project unacceptable if it cannot be reduced to acceptable levels; and/or the cost of management intervention will be a significant factor in mitigation.

The following weights will be assigned to each attribute (Table 1):

**Table 1. Impact rating assessment weights**

Aspect	Description	Weight
<b>Probability</b>	Improbable	1
	Probable	2
	Highly Probable	4
<b>Duration</b>	Definite	5
	Short term	1
	Medium term	3
	Long term	4
<b>Scale</b>	Permanent	5
	Local	1
	Site	2
	Regional	3
<b>Magnitude/Severity</b>	Low	2
	Medium	6
	High	8
<b>Significance</b>	<b>Sum (Duration, Scale, Magnitude) x Probability</b>	
	Negligible	<20
	Low	<40
	Moderate	<60
	High	>60

The significance of each activity will be rated without mitigation measures and with mitigation measures for the development.

The mitigation effect of each impact will be indicated without and with mitigation measures as follows:

- Can be reversed.
- Can be avoided, managed or mitigated.
- May cause irreplaceable loss of resources.

## 2.4 SENSITIVITY ASSESSMENT

The ecological sensitivity of any piece of land is based on its inherent ecosystem service and overall preservation of biodiversity.

### 2.4.1 Ecological function

The ecological function relates to the degree of ecological connectivity between systems within a landscape matrix. Therefore, systems with a high degree of landscape connectivity amongst one another are perceived to be more sensitive and will be those contributing to ecosystem service (e.g., wetlands) or overall preservation of biodiversity.

### 2.4.2 Conservation importance

Conservation importance relates to species diversity, endemism (unique species or unique



processes) and the high occurrence of threatened and protected species or ecosystems protected by legislation.

#### 2.4.3 Sensitivity scale

- High – sensitive ecosystem with either low inherent resistance or low resilience towards disturbance factors or highly dynamic systems considered being important for the maintenance of ecosystem integrity. Most of these systems represent ecosystems with high connectivity with other important ecological systems or with high species diversity and usually provide suitable habitat for a few threatened or rare species. These areas should be protected.
- Medium – These are slightly modified systems which occur along gradients of disturbances of low-medium intensity with some degree of connectivity with other ecological systems or ecosystems with intermediate levels of species diversity but may include potential ephemeral habitat for threatened species.
- Low – Degraded and highly disturbed / transformed systems with little ecological function and which are generally very poor in species diversity.

### 2.5 EIA SCREENING TOOL

The significance of a site or natural feature may only become apparent when it is evaluated in terms of a broader biodiversity context. Put differently, local impacts on biodiversity may seem unimportant, but can become highly significant when interpreted beyond the immediate boundaries of a site. Even if a locality has a history of disturbance such as alien infestation, cultivation, or recurrent fires, and it does not host any plant or animal species of special concern, it may nevertheless be significant for biodiversity conservation when viewed from a landscape or even national perspective.

**According to the national web-based environmental screening tool in terms of section 24(5)(h) of the NEMA, 1998 (Act No 107 of 1998) and regulation 16(1)(b)(v) of the EIA regulations, 2014, as amended, the following listed fauna and flora species occur in the project area.**

#### Fauna:

- *Tyto capensis*:
  - Status: Vulnerable
  - Sensitivity: Medium
- *Eupodotis senegalensis*:
  - Status: Vulnerable
  - Sensitivity: Medium

#### Flora:

- *Sensitive species 1261:*
  - Status: Vulnerable
  - Sensitivity: Medium
- *Sensitive species 1147:*
  - Status: Endangered
  - Sensitivity: Medium

### 3 BASELINE ENVIRONMENT

#### 3.1 LOCATION AND DESCRIPTION OF ACTIVITY

In view of the growing electricity demand and to use renewable energy resources, VOLTALIA SOUTH AFRICA (PTY) LTD is assessing the feasibility of energy generation facilities, consisting of the construction, operation and maintenance of Photovoltaic (PV) Power Plants with a maximum generation capacity up to 120 MW, at the point of connection.

The project site is on Portion 2 of the farm Rooidraai 85 IQ, located within the JB Marks Local Municipality, Dr Kenneth Kaunda District Municipality, North West Province.

The project site is located  $\pm 7$  km north west of Welverdiend along the border between Gauteng and the North West Province (Figure 4). The Eskom Carmel Main Transmission Substation (MTS) is located 16.4 km South-East of project sites.

The development is located 5.5km north of R501 with access from both the D859 (Preferred) and R501 (alternative).

The developed area (footprint) required for the proposed Mopane Solar PV 5 will be up to 182 hectares. The final size and location of the project footprint will be assessed following the outcomes of the Public Participation Process and of the recommendations and conclusions of the Specialist Studies to be conducted during the Environmental Impact Assessment (EIA) process.

The proposed development (the Photovoltaic (PV) Power Plants and connection infrastructure) consists of the installation of the following equipment:

- Photovoltaic modules (mono-crystalline, poly-crystalline, or bi-facial modules)
- Mounting systems for the PV arrays (single-axis horizontal trackers or fixed structures) and related foundations
- Internal cabling and string boxes
- DC/AC inverters
- Medium voltage stations, hosting LV/MV power transformers
- Medium voltage receiving station(s)

- Workshops & warehouses
- One on-site high-voltage substation and one high-voltage busbar with metering and protection devices
- One on-site high-voltage substation with high-voltage power transformers, stepping up the voltage to 400kV/132kV and one high-voltage busbar with metering and protection devices
- One on-site switching station, with one high-voltage busbar with metering and protection devices
- Battery Energy Storage Systems (BESS), with a Maximum Export Capacity up to 120 MW and a 5-hour storage capacity up to 1250 MWh, with a footprint up to 10 ha, next to the on-site high-voltage substation, within the PV plant footprint / fenced areas
- Electrical system and UPS (Uninterruptible Power Supply) devices
- Lighting system
- Grounding system
- Internal roads
- Fencing of the site and alarm and video-surveillance system
- Water access point, water supply pipelines, water treatment facilities
- Small scale patented wastewater treatment system

During the construction phase, the site may be provided with additional activities which will be removed at the end of construction.

- Water access point, water supply pipelines, water treatment facilities
- Prefabricated buildings
- Workshops & warehouses

The aerial map of the site (including the footprints) is indicated in Figure 5, while the topographical map of the proposed development is presented in Figure 6.




Figure 4. Regional location Map of the project area




Figure 5. Aerial Map indicating the proposed location of the Solar Plant and associated infrastructure




Figure 6. Topographical Map of the project area

### 3.2 CLIMATE

Climate in the broad sense is a major determinant of the geographical distribution of species and vegetation types. However, on a smaller scale, the microclimate, which is greatly influenced by local topography, is also important. Within areas, the local conditions of temperature, light, humidity and moisture vary greatly, and it is these factors which play an important role in the production and survival of plants (Tainton, 1981). The climate for the region can be described as warm-temperate. In terrestrial environments, limitations related to water availability are always important to plants and plant communities.

The spatial and temporal distribution of rainfall is very complex and has great effects on the productivity, distribution and life forms of the major terrestrial biomes (Barbour et al. 1987). The study area is situated within the summer and autumn rainfall region with very dry winters and frequent frost that occurs during the colder winter months. The spatial and temporal distribution of rainfall is very complex and has great effects on the productivity, distribution and life forms of the major terrestrial biomes (Barbour et al. 1987).

The climate for the region is warm-temperate, summer rainfall region, with overall mean annual precipitation of 593mm. Severe, frequent frost occurs, although summer temperatures are high. The mean annual temperature for the area is 16.1°C, and the mean annual frost days is 37 days. Mean Annual Potential Evaporation is 2407mm, with Mean Annual Soil Moisture Stress of 78%.

### 3.3 GEOLOGY AND SOIL TYPES

Geology is directly related to soil types and plant communities that may occur in a specific area (Van Rooyen & Theron, 1996). A Land type unit is a unique combination of soil pattern, terrain and macroclimate, the classification of which is used to determine the potential agricultural value of soils in an area. The land type unit represented within the study area include the Fa14 land type (Land Type Survey Staff, 1987) (ENPAT, 2001). The land type, geology and associated soil types is presented in Table 2 below as classified by the Environmental Potential Atlas, South Africa (ENPAT, 2000).

**Table 2. Land types, geology, and dominant soil types of the proposed development site**

Landtype	Soils	Geology
Fa14	Glenrosa and/or Mispah forms (other soils may occur), lime rare or absent in the entire landscape	Dolomite and chert of the Chuniespoort Group; chert gravels are abundant on middle and footslopes including valley bottoms.

Soils associated with the site vary between slightly deeper, loamy red apedal soils, to shallow rocky soils.

### 3.4 TOPOGRAPHY, LANDUSES AND DRAINAGE

The study area lies completely within the Upper Vaal Water Management Area (WMA) and entirely within the Highveld ecoregion (Kleynhans et al., 2005).

The topography is characterised by slightly undulating plains. The topography of the site can be described as generally favourable, when considering that most of the area consists of slopes of less than 1:5. The site is located at an altitude of 1460 meters above mean sea level (AMSL).

Most properties situated within a 500m radius are being used for livestock grazing and crop cultivation. The proposed development land is used for livestock farming at present. The natural vegetation of the varies from intact to planted pastures.

The site is located within the C23G quaternary catchment and is situated in the Upper Vaal Water Management Area. Drainage occurs as sheet-wash into the drainage channels and wetlands on site that eventually drains into the major river namely the Mooi River and Mooiriviersloop River that occurs to the west and south of the site respectively.

### **3.5 SENSITIVITY ANALYSIS AND CONSERVATION ANALYSIS TOOLS**

There are several assessments for South Africa as a whole, as well as on provincial levels that allow for detailed conservation planning as well as meeting biodiversity targets for the country's variety of ecosystems. These guides are essential to consult for development projects and will form an important part of the sensitivity analysis. Areas earmarked for conservation in the future, or that are essential to meet biodiversity and conservation targets should not be developed and have a high sensitivity as they are necessary for overall functioning. In addition, sensitivity analysis in the field based in much finer scale data can be used to ground truth the larger scale assessments and put it into a more localised context.

#### **3.5.1 NORTH WEST BIODIVERSITY CONSERVATION PLAN**

The purpose of the North West Biodiversity Conservation Plan is to develop the spatial component of a bioregional plan (i.e., map of Critical Biodiversity Areas (CBA) and associated land-use guidelines). The North West Conservation Plan categories for the developments are presented in Figure 7. The following can be concluded regarding developments:

- Most of the proposed development footprints represent CBA1 areas although most of these areas should rather be classified as ESA1 or ESA2 areas. The management objective for this area is to maintain ecosystem functionality and connectivity allowing for limited loss of biodiversity pattern.



# Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5




Figure 7. North West C-Plan Map (2015) for the project area

### 3.5.2 PROTECTED AREAS NETWORK AND NATIONAL PROTECTED AREAS EXPANSION STRATEGY (NPAES)

Officially protected areas, either Provincially or Nationally that occur close to a project site could have consequences as far as impacts on these areas are concerned. For the proposed development and associated infrastructure no protected areas occur in proximity, with the closest being the Abe Bailey Provincial Nature Reserve that occurs to the east of the project area (Figure 8).

The NPAES are areas designated for future incorporation into existing protected areas (both National and informal protected areas). These areas are large, mostly intact areas required to meet biodiversity targets, and suitable for protection. They may not necessarily be proclaimed as protected areas in the future and are a broad scale planning tool allowing for better development and conservation planning. No NPAES occur within the project area, although the Vaal Grasslands NPAES occur to the east of the project area (Figure 8).

Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5




Figure 8. Location of the project area in relation to listed protected areas.

### 3.5.3 IMPORTANT BIRD AREAS

An Important Bird Area (IBA) is an area recognized as being globally important habitat for the conservation of bird populations. Currently there are about 10,000 IBAs worldwide. At present, South Africa has 124 IBA's, covering over 14 million hectares of habitat for our threatened, endemic and congregatory birds. Yet only million hectares of the total land surface covered by our IBA's legally protected. The BirdLife SA IBA programme continues a programme of stewardship which will ultimately achieve formal protection (Birdlife, 2013). The project area is not located within or close to any IBA.

### 3.5.4 NATIONALLY THREATENED ECOSYSTEMS

The list of national Threatened Ecosystems has been gazetted (NEM:BA: National list of ecosystems that are threatened and in need of protection) and result in several implications in terms of development within these areas. Four basic principles were established for the identification of threatened ecosystems. These include:

- The approach must be explicit and repeatable.
- The approach must be target driven and systematic, especially for threatened ecosystems.
- The approach must follow the same logic as the IUCN approach to listing threatened species, whereby a few criteria are developed, and an ecosystem is listed based on its highest-ranking criterion: and
- The identification of ecosystems to be listed must be based on scientifically credible, practical, and simple criteria, which must translate into spatially explicit identification of ecosystems.

Areas were delineated based on as fine a scale as possible and are defined by one of several assessments: These areas are essential for conservation of the country's ecosystems as well as meeting conservation targets. The project area is not located within any listed threatened ecosystem, although the Vaal-Vet Sandy Grasslands and Eastern Temperate Freshwater Wetlands Listed Threatened Ecosystems occur in close proximity to the project area (Figure 9).

The indigenous grassland vegetation units on the proposed development site is not considered as Critical Habitat in line with IFC Performance Standard PS6. The Carletonville Dolomite Grassland vegetation type has a Least Concern Conservation Status (Sanbi, 2016) with almost 25% of it being transformed for cultivation, urban sprawl or by mining activities. Small extent is conserved in statutory and in at least six private conservation areas, with the conservation target being 24%.

# Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5




Figure 9. Listed threatened ecosystems in proximity to the proposed development site (SANBI).

**3.5.5 STRATEGIC WATER SOURCE AREAS (SWSA), NATIONAL FRESHWATER ECOSYSTEM PRIORITY AREAS (NFEPA) STATUS OF RIVERS AND WETLANDS ON SITE**

NFEPA maps provide strategic spatial priorities for conserving South Africa's freshwater ecosystems and supporting sustainable use of water resources. These strategic spatial priorities are known as Freshwater Ecosystem Priority Areas, or 'FEPAs'. NFEPA maps were developed using the principles of systematic biodiversity planning, also known as systematic conservation planning (Margules and Pressey 2000). Systematic biodiversity planning is a well-established field of science in which South Africa is considered a world leader (Balmford 2003). The NFEPA maps and supporting information form part of a comprehensive approach to sustainable and equitable development of South Africa's scarce water resources. For integrated water resources planning, NFEPA provides guidance on how many rivers, wetlands and estuaries, and which ones, should remain in a natural or near-natural condition to support the water resource protection goals of the National Water Act (Act 36 of 1998). NFEPA products are therefore directly applicable to the National Water Act, feeding into Catchment Management Strategies, water resource classification, reserve determination, and the setting and monitoring of resource quality objectives. NFEPA products are also directly relevant to the National Environmental Management: Biodiversity Act (Act 10 of 2004), informing both the listing of threatened freshwater ecosystems and the process of bioregional planning provided for by this Act. NFEPA products support the implementation of the National Environmental Management: Protected Areas Act (Act 57 of 2003) by informing the expansion of the protected area network. The project area is located close to the listed NFEPA rivers, named Mooi River and Mooiriviersloop River, although these rivers will not be impacted on by the development footprints. The rivers also represent a NFEPA wetlands as indicated in Figure 10.

Strategic Water Source Areas (SWsAs) are now defined as areas of land that either:

- Supply a disproportionate (i.e., relatively large) quantity of mean annual surface water runoff in relation to their size and so are considered nationally important; or
- Have high groundwater recharge and where the groundwater forms a nationally important resource; or
- Areas that meet both criteria (a) and (b).

They include transboundary Water Source Areas that extend into Lesotho and Swaziland. All surface water SWsAs are in high rainfall areas where baseflow is at least 11 25 mm/a, which is evidence of a strong link between groundwater and surface water in the SWsAs. The aquifers sustain baseflow, contribute to runoff and, especially, contribute to dry season flows. Sustained river flows are important as they support people and communities who depend directly on rivers for their water, especially during the dry season and droughts.

The 2018 national and transboundary surface-water SWsAs cover about 124 075 km<sup>2</sup> (10% of the region) and provide a MAR of 24 954 million m<sup>3</sup> (50% of the total). The greatest volume of

MAR is generated by the Southern Drakensberg (9% of national and transboundary MAR), followed by the Eastern Cape, Northern Drakensberg and Maloti Drakensberg, and the Boland. The Boland has the highest MAR per unit area (3588 m<sup>3</sup>/ha/year), followed by Table Mountain, the Northern Drakensberg and the Mpumalanga Drakensberg.

Seven of these SWSAs are transboundary areas because Lesotho and Swaziland include portions of important SWSAs for South Africa. The portions of the SWSAs that fall within Lesotho (Eastern Cape, and the Southern, Northern and Maloti Drakensberg) cover 18 570 km<sup>2</sup> and generate a MAR of about 3522 million m<sup>3</sup>. This MAR sustains the Orange and Caledon Rivers and supplies water to Gauteng via the Lesotho Highlands water supply system. In the case of Swaziland, the portions of the SWSAs falling in this country (Ekangala Drakensberg, Mbabane Hills, Upper Usutu) total 9376 km<sup>2</sup> and produce a MAR of about 2053 million m<sup>3</sup>. In total, the SWSAs in these two countries produce about 11% of the total MAR, which is a substantial contribution that needs to be protected.

The project area is located within the Far West Carst Groundwater SWSA.

# Terrestrial Biodiversity, Plant & Animal Species Impact Assessment Mopane Solar PV 5




Figure 10. Location of the project area in relation to NFEPA Rivers and SWSA



## **4 RESULTS**

### **4.1 VEGETATION**

#### **4.1.1 Biome and Ecoregion**

The development site lies within the Grassland Biome which is found chiefly on the high central plateau of South Africa. Grasslands are dominated by a single layer of grasses. The amount of cover depends on rainfall and the degree of grazing. Trees are absent except in a few localised habitats. Geophytes are often abundant. Frost, fire and grazing maintain the grass dominance and prevent the establishment of trees (Low & Rebelo, 1996).

The Highveld Ecoregion draws its name from the high interior plateau known as the Highveld, and the expansive cover of species-rich communities of grasses. The ecoregion is bordered by the Drakensberg in the east, the arid Karoo and Kalahari in the west, and the low-lying bushveld to the north. The Highveld Plateau is flat with elevations varying from 1,400 m to 1,800 m. The flat topography means that the landscape is traversed by many meandering rivers, with the grassland community historically playing an important role in natural water purification of the westward flowing rivers that originate on the Drakensberg escarpment (Davies and Day 1998). The functioning of this ecosystem has been disrupted in many areas by water transfer projects that have been built to supply greater Johannesburg with water (Davies and Day 1998).

The Highveld Grassland Ecoregion has further suffered extensive degradation. Because it is one of the best areas for farming in South Africa, large tracts of land have already been converted to agriculture, mainly for corn production. Urban expansion, fire, and overgrazing have led to increased fragmentation, as has coal mining and afforestation for stands of exotic trees, especially by species of Eucalyptus (Low and Rebelo, 1998; Cowling et al. 1997). Over several hundred years, particularly around towns, planted wattle (*Acacia mearnsii*) has become invasive, and is prone to rapid expansion upriver watersheds. In the future, expanded surface activity associated with mining below the grassland may become a greater concern as companies develop new technology to make deep mining of coal more profitable (Mallett 1999).

#### **4.1.2 Ecosystem drivers and ecological services**

Fire and grazing are two of the most important ecological drivers in grassland. Any land-use change that results in reduced ability to manage fire or grazing in the remaining natural areas will have significant implications for grassland biodiversity. Invasive alien species and soil erosion are two of the most pervasive management issues affecting all grassland ecosystems and are key indicators that the limits of acceptable change have been exceeded.

The Highveld also plays an important role in natural water purification, as the peat formed here has been shown to filter out 90 percent of the harmful chemicals in herbicides. Peat is also useful in absorbing various other pollutants, as a source of fuel, in horticulture, and for medicinal purposes. In South Africa, where clean water resources are already particularly valuable, this natural filter is being extracted from the Highveld at an unprecedented rate. Approximately 60

percent of locally extracted peat is used to grow mushrooms, while the remaining 40 percent comprises "environmentally friendly" potting soil and compost. Peat has an extremely slow regeneration rate, increasing between 0.7 mm to 1.2 mm per year depending on environmental conditions (Dada 1999). Given its slow formation process, it is unlikely this resource will recover from the damage caused by its rapid removal. Hence, the Highveld's role as a natural filtration element for scarce water resources could be in danger. The preservation of this resource is imperative and could be fulfilled by moderating or halting the use of peat for gardening purposes.

#### **4.1.3 Vegetation types**

The most recent classification of the area by Mucina & Rutherford (2006) shows that the site is classified as Carletonville Dolomite Grassland.

Carletonville Dolomite Grasslands (Gh15) are predominantly found in the North West Province, in the regions around Potchefstroom, Ventersdorp and Carletonville. Vegetation and Landscape Features Carletonville Dolomite Grasslands occur on slightly undulating plains which are typically intersected by rocky chert ridges. They are species rich and according to Mucina and Rutherford (2006), dominated by many plant species.

**Important Plant Taxa Grasses:** *Aristida congesta*, *Brachiaria serrata*, *Cynodon dactylon*, *Digitaria tricholaenoides*, *Diheteropogon amplexens*, *Eragrostis chloromelas*, *E. racemosa*, *Heteropogon contortus*, *Loudetia simplex*, *Schizachyrium sanguineum*, *Setaria sphacelata*, *Themeda triandra* and *Alloteropsis semialata*.

**Herbs:** *Acalypha angusta*, *Barleria macrostegia*, *Chamaecrista mimosoides*, *Chamaesyce inaequilatera*, *Crabbea angustifolia*, *Dianthus mooiensis*, *Dicoma anomala*, *Helichrysum caespitium*, *H. miconiifolium*, *H. nudifolium*, *Ipomoea ommaneyi*, *Kyphocarpa angustifolia* and *Senecio coronatus*. **Shrubs:** *Anthospermum rigidum*, *Indigofera comosa*, *Pygmaeothamnus zeyheri*, *Englerophytum magalismontana*, *Tylosema esculentum* and *Ziziphus zeyheriana*.

This vegetation type is described as Least Concern. Almost 25% of it has been transformed for cultivation, urban sprawl or by mining activities. Small extent is conserved in statutory and in at least six private conservation areas, with the conservation target being 24%.

#### **4.1.4 Vegetation units**

The proposed development site occurs on a slightly undulating landscape. The importance to survey the area to have a better understanding of the ecosystem and the potential impact of the solar development on the natural environment was identified as a key factor, and subsequently the footprint areas was completely surveyed. The site forms part of a larger farm used for livestock farming. The vegetation units on the site vary according to soil characteristics, topography, and land-use. Vegetation units were identified on the footprint development sites and can be divided into 4 distinct vegetation units according to soil types and topography.

The vegetation communities identified on the proposed development site are classified as

physiographic physiognomic units, where physiognomic refers to the outer appearance of the vegetation, and physiographic refers to the position of the plant communities in the landscape. The physiographic-physiognomic units will be referred to as vegetation units in the following sections. These vegetation units are divided in terms of the land-use, plant species composition, topographical and soil differences that had the most definitive influence on the vegetation units. Each unit is described in terms of its characteristics and detailed descriptions of vegetation units are included in the following section. A species list for the site is included in Appendix B, while a plant species list for the quarter degree grid square (QDS) is included in Appendix A. Photographs of each unit is included in the next section to illustrate the grass layer, woody structure, and substrate (soil, geology etc.). The following vegetation units were identified during the survey.

1. *Schizachyrium – Trachypogon – Seriphium* rocky grassland.
2. Open grassland with *Searsia pyroides* clumps.
3. *Searsia pyroides* open woodland
4. Exotic bushclumps.

The vegetation units for the solar development are presented in Figure 11:




Figure 11. Vegetation Unit Map of the proposed development area

**4.1.4.1 Schizachyrium – Trachypogon – Seriphium rocky grassland**

This vegetation unit comprises an isolated section of the project area on undulating terrain. The soil is shallow and red soils of the Glenrosa and Hutton soil forms derived from quartzite. There are little to no trees present with the grasses having the highest cover. The grass layer is dominated by species such as *Schizachyrium sanguineum*, *Trachypogon spicatus*, *Tristachya leucothrix*, and *Elionorus*, while the dwarf shrub *Seriphium plumosum* is also prominent indicating that the area was probably overgrazed in the past. The state of the vegetation is indicated in photograph 1, while the characteristics of the variations of this vegetation unit are summarized in Table 3.

**Table 3. Botanical analysis and characteristics of *Schizachyrium – Trachypogon – Seriphium* rocky grassland.**

Vegetation unit characteristics	
<b>State of the vegetation:</b>	Natural grassland in a slightly degraded state
<b>Need for rehabilitation</b>	Low
<b>Conservation priority</b>	Medium
<b>Soils &amp; Geology</b>	Red-yellow apedal sandy soils of the Hutton / Glenrosa soils derived from quartzite
<b>Density of woody layer</b>	Trees: <1% (avg. height: 3-6m) Shrubs:<1% (avg. height: 1-2m)
<b>Density of herbaceous layer</b>	Grasses: 70-80% (avg. height: 0.8-1.2m) Forbs: <1% (avg. height: 0.8m)
<b>Sensitivity</b>	Medium
<b>Red data species</b>	None observed
<b>Protected species</b>	<i>Boophane distichya</i>

The following specific recommendations for the vegetation unit regarding the proposed development should be adhered to:

- The vegetation unit is classified as having a medium sensitivity due to the due to the widespread status of this vegetation unit within the larger project area.
- The eradication of protected plant species *Boophane distycha* would need a permit from local authorities in the North West Province.
- The development of the solar development is considered suitable in this area.



Photograph 1. *Schizachyrium – Trachypogon – Seriphium* rocky grassland in the project area

#### 4.1.4.2 Open grassland with *Searsia pyroides* clumps

A large section of the proposed development footprint forms medium tall grassland with scattered bushclumps of the woody species *Searsia pyroides* on red- apedal soils of the Hutton soil form. The grass layer is well developed and dominated by species such as *Elionorus muticus*, *Hyparrhenia hirta* and *Schizachyrium sanguineum*. The state of the vegetation is indicated in photograph 2, while the characteristics of the variations of this vegetation unit are summarized in Table 4.

Table 4. Botanical analysis and characteristics of Open grassland with *Searsia pyroides* clumps

Vegetation unit characteristics	
State of the vegetation:	Natural grassland in a slightly degraded state
Need for rehabilitation	Low
Conservation priority	Medium
Soils & Geology	Red-yellow apedal sandy soils of the Clovelly / Hutton soils
Density of woody layer	Trees: 1-2% (avg. height: 3-6m) Shrubs:5-10% (avg. height: 1-2m)
Density of herbaceous layer	Grasses: 70-80% (avg. height: 0.8-1.2m) Forbs: <1% (avg. height: 0.8m)

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Vegetation unit characteristics	
Sensitivity	Medium
Red data species	None observed
Protected species	<i>Boophane disticha</i>

The following specific recommendations for the vegetation unit regarding the proposed development should be adhered to:

- The vegetation unit is classified as having a Medium sensitivity due its widespread occurrence in the Grassland Biome.
- The eradication of protected plant species *Boophane* would need a permit from local authorities in the North West.
- The development of the solar development is considered suitable in this area.



Photograph 2. Open grassland with *Searsia pyroides* clumps in the project area

### 4.1.4.3 *Searsia pyroides* open woodland

This vegetation unit occurs on red apedal soils in the south-western section of the site and represent secondary old fields. The woody layer forms an open woodland mostly dominated by species such as *Searsia pyroides*, *Grewia flava*, *Diospyros lycioides* and *Vachellia karroo*, while the grass layer is dominated by species such as *Elionorus muticus*, *Aristida congesta* and *Schizachyrium sanguineum*. The state of the vegetation is indicated in photograph 3, while the characteristics of the variations of this vegetation unit are summarized in Table 5.

Table 5. Botanical analysis and characteristics of *Searsia pyroides* open woodland

Vegetation unit characteristics	
State of the vegetation:	Natural grassland in a slightly degraded state
Need for rehabilitation	Low
Conservation priority	Medium
Soils & Geology	Deep red apedal soils of the Hutton soil form derived from quartzite
Density of woody layer	Trees: 1-2% (avg. height: 3-6m) Shrubs:10% (avg. height: 1-2m)
Density of herbaceous layer	Grasses: 70-80% (avg. height: 0.8-1.2m) Forbs: <1% (avg. height: 0.8m)
Sensitivity	Medium
Red data species	None observed
Protected species	None observed

The following specific recommendations for the vegetation unit regarding the proposed development should be adhered to:

- The vegetation unit is classified as having a Medium sensitivity due its widespread occurrence in the Grassland Biome.
- The development of the solar development is considered suitable in this area.



Photograph 3. *Searsia pyroides* open woodland in the project area

#### 4.1.4.4 Exotic bushclumps

The exotic bushclumps represent homogenous stands of *Eucalyptus camaldulensis* red apedal



## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

soils of the Hutton soil form (Photograph 4). Exotic weeds and pioneer grasses often colonize the areas beneath the exotic bushclumps. No detailed survey was considered for this area due to the completely modified state of the vegetation.




**Photograph 4. Exotic bushclumps in the project area**

### 4.2 PLANT SPECIES LEVEL ASSESSMENT

South Africa has been recognized as having remarkable plant diversity with high levels of endemism. The major threats to plants in the study area are urban expansion, non-sustainable harvesting, collecting, overgrazing/browsing, mining and agriculture. The objective of this section was to compile a list of plant species for which there is conservation concern. This included threatened, rare, declining, protected, and endemic species.

#### 4.2.1 Species of conservation concern

Species of conservation concern are species that have a high conservation importance in terms of preserving South Africa's high floristic diversity and include not only threatened species, but also those classified in the categories Extinct in the Wild (EW), Regionally Extinct (RE), Near Threatened (NT), Critically Rare, Rare, Declining and Data Deficient – Insufficient Information (DDD). It should also be noted that not all species listed as protected are threatened or vice versa. A list of SCC plant species previously recorded in the study area in which the proposed development is planned was obtained from the Plants of Southern Africa (POSA) database of SANBI. Figure 12 indicates the classification system used by Sanbi for SCC:



**Figure 12. South African red list categories indicating the categories to be used for Species of Conservation Concern**

Habitat degradation is one of the main reasons for plant species becoming extinct in a particular area. Threatened species are also seen as indicators of the overall health of an ecosystem (Hilton-Taylor, 1996).

A list of red data plant species previously recorded in the grid square in which the proposed development is planned was obtained from SANBI as indicated in Table 6.

**Table 6. Red data and endemic species occurring in the project area of the QDS**

Species	Threat status	Confirmed presence on site
<i>Myrothamnus flabellifolius</i>	DATA DEFICIENT	NO
<i>Acalypha caperonioides</i>	DATA DEFICIENT	NO
<i>Habenaria mossii</i>	ENDANGERED	NO
<i>Boophane distycha</i>	Declining	YES

Only the red data species *Boophane distycha* listed above was observed during the surveys. The species can be relocated from its current conditions if needed through a rescue and relocation programme should the development activities impact on populations.

Ecological monitoring should however still be implemented during the construction phase and specific sensitive habitats (riparian) needs to be avoided to ensure that any potential red data species potentially missed during the field surveys are preserved and not potentially impacted on.

The EIA screening tool highlight the following red listed flora.

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

### 4.2.1.1 Sensitive species 1261

A relatively widespread (EOO 13 374 km<sup>2</sup>), but very rare species that has lost a large proportion of its habitat to agriculture, urban expansion and mining. It is known from fewer than 10 locations and continue to decline due to ongoing habitat loss and degradation. Habitat includes sandy loam soils in thornveld and *Themeda*-grassland

**Probability of occurrence on site:** LOW due to the absence of suitable habitat on the proposed development footprint.

**Probability of impact during vegetation clearance:** LOW, no suitable habitat observed on site and population of the species was documented.

### 4.2.1.2 Sensitive species 1147

Surveys of remaining habitat within Gauteng Province revealed that there are only about 230 mature individuals. These occur as six scattered subpopulations, the largest of which only has 70-80 mature individuals, but there are generally fewer than 40 mature individuals per subpopulation. There is a continuing decline due to the rapid urban expansion.

Occurs in Open grassland on dolomite or in black, sandy soil. Threats include invasive alien species (direct effects), habitat loss and habitat degradation.

**Probability of occurrence on site:** Moderate due to the presence of suitable habitat on the proposed development footprint.

**Probability of impact during vegetation clearance:** LOW, no population of the species was documented, although monitoring should be implemented during the construction phase of the development.

### 4.2.2 Protected Plants (North West Nature Conservation Ordinance)

Plant species are also protected in the North West Province according to the North West Nature Conservation Ordinance. According to this ordinance, no person may pick, import, export, transport, possess, cultivate, or trade in a specimen of a specially protected or protected plant species. The Appendices to the ordinance provide an extensive list of species that are protected, comprising a significant component of the flora expected to occur on site. Communication with Provincial authorities indicates that a permit is required for all these species if they are expected to be affected by the proposed project.

After a detailed survey was conducted during November 2022, the listed species *Boophane disticha* confirmed for the site. No eradication should be allowed without a permit.

### 4.2.3 Invasive alien species

Invasive alien plants pose a direct threat not only to South Africa's biological diversity, but also to water security, the ecological functioning of natural systems and the productive use of land. They intensify the impact of fires and floods and increase soil erosion. Of the estimated 9000

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

plants introduced to this country, 198 are currently classified as being invasive. It is estimated that these plants cover about 10% of the country and the problem is growing at an exponential rate.

The Alien and Invasive Species Regulations (GNR 599 of 2014) are stipulated as part of the National Environmental Management: Biodiversity Act (10/2004). The regulation listed a total of 559 alien species as invasive and further 560 species are listed as prohibited and may not be introduced into South Africa. Below is a brief explanation of the four categories of Invasive Alien Plants as per the regulation.

- Category 1a: Invasive species requiring compulsory control. Remove and destroy. Any specimens of Category 1a listed species need, by law, to be eradicated from the environment. No permits will be issued.
- Category 1b: Invasive species requiring compulsory control as part of an invasive species control programme. Remove and destroy. These plants are deemed to have such a high invasive potential that infestations can qualify to be placed under a government sponsored invasive species management programme. No permits will be issued.
- Category 2: Invasive species regulated by area. A demarcation permit is required to import, possess, grow, breed, move, sell, buy, or accept as a gift any plants listed as Category 2 plants. No permits will be issued for Category 2 plants to exist in riparian zones.
- Category 3: Invasive species regulated by activity. An individual plant permit is required to undertake any of the following restricted activities (import, possess, grow, breed, move, sell, buy, or accept as a gift) involving a Category 3 species. No permits will be issued for Cat 3 plants to exist in riparian zones.

The fight against invasive alien plants is spearheaded by the Working for Water (WfW) programme, launched in 1995 and administered through the DWA. This programme works in partnership with local communities, to whom it provides jobs, and with Government departments including the Departments of Environmental Affairs and Tourism, Agriculture, and Trade and Industry, provincial departments of agriculture, conservation and environment, research foundations and private companies.

WfW currently runs over 300 projects in all nine of South Africa's provinces. Scientists and field workers use a range of methods to control invasive alien plants. These include:

- Mechanical methods - felling, removing, or burning invading alien plants.
- Chemical methods - using environmentally safe herbicides.
- Biological control - using species-specific insects and diseases from the alien plant's country of origin. To date 76 bio-control agents have been released in South Africa against 40 weed species.

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

- Integrated control - combinations of the above three approaches. Often an integrated approach is required to prevent enormous impacts.

Vehicles often transport many seeds, and some may be of invader species, which may become established along the roads through the area, especially where the area is disturbed. The construction phase of the development will almost certainly carry the greatest risk of alien invasive species being imported to the site, and the high levels of habitat disturbance also provide the greatest opportunities for such species to establish themselves, since most indigenous species are less tolerant of disturbance. The biggest risk is that invasive alien species such as the seeds of noxious plants may be carried onto the site along with materials that have been stockpiled elsewhere at already invaded sites.

Continued movement of personnel and vehicles on and off the site, as well as occasional delivery of materials required for maintenance, will result in a risk of importation of alien species throughout the life of the project. The following alien invasive and exotic plant species were recorded on site during the surveys as stipulated in the Alien and Invasive Species Regulations (GNR 599 of 2014) (Table 7):

**Table 7. Declared weeds and invader plants of the study area.**

Species	Category
<i>Argemone ochroleuca</i>	1b
<i>Datura stramonium</i>	1b
<i>Eucalyptus camaldulensis</i>	1b
<i>Verbena brasiliensis</i>	1b
<i>Xanthium strumarium</i>	1b

According to the amended regulations (No. R280) of March 2001 of the Conservation of Agricultural Resources Act 1983 (Act no. 43 of 1983), it is the legal duty of the land user/landowner to control invasive alien plants occurring on the land under their control. The State has the right to clear invasive plants at the landowner's expense if the landowner refuses to remove invasive plants.

### 4.2.4 General

An important aspect relating to the proposed development should be to protect and manage the biodiversity (structure and species composition) of the vegetation types which are represented on the proposed development site. Vegetation removal should be kept to a minimum during the construction phase of the development and only vegetation on the footprint areas should be removed. Mitigation measures and monitoring should however be implemented should the development be approved.

### 4.3 FAUNAL HABITAT AND ANIMAL SPECIES ASSESSMENT

#### 4.3.1 Overview

A healthy environment is inhabited by animals that vary from micro-organisms to the birds and mammals. The species composition and diversity are often parameters taken into consideration when determining the state of the environment. A comprehensive survey of all animals is a time-consuming task that will take a long time and several specialists to conduct. The alternative approach to such a study is to do a desktop study from existing databases and conduct a site visit to verify the habitat requirements and condition of the habitat. If any rare or endangered species are discovered in the desktop study that will be negatively influenced by the proposed development, specialist surveys will be conducted.

#### 4.3.2 Results of desktop survey and site visits during November 2022

A survey was conducted during November 2022 to identify specific fauna habitats, and to compare these habitats with habitat preferences of the different fauna groups (birds, mammals, reptiles, amphibians) occurring in the quarter degree grid.

The number of mammal species supported by a plant community depends on several factors like the primary production, seasonal availability of resources, floral heterogeneity, diversity of plant structure, nature of the substratum and previous history (Delany, 1982). Each mammal species has a particular niche, which can be regarded as the sum of all ecological requirements of a species namely food, space, shelter, and physical conditions. Mills & Hes (1997) stated that the distribution and abundance of animal species does not rigorously follow that of plant communities or biomes. Instead, mammal species seem to have certain preferences for a specific habitat type (Skinner & Smithers, 1990). Several authors have shown this preference of mammals to certain habitats through analysis (Beardall et al. 1984; Ben-Shahar, 1991; Dekker et al. 1996). The area represents a diverse vegetation structure and height class. A detailed species list for the fauna of the area is included in Appendix C, D and E.

#### 4.3.3 Fauna habitats of the project area

Three major fauna habitats were observed in the area namely:

- Grassland.
- Open woodland / exotic bushclumps
- Open water habitats / wetlands.

#### 4.3.4 Common fauna documented and potentially occurring on the development site

##### 4.3.4.1 Mammals

Much of the large and medium-sized mammal fauna that previously occurred on the project site is now locally extinct or occurs in small, fragmented populations in reserves. Most of the habitat

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

types on the respective study sites are fragmented. Therefore, the expected mammalian richness on these areas is considered low, although slightly higher richness values are expected from the more intact grassland, woodland and wetland habitats.

The Highveld Ecoregion contains a higher number of mammals, although only the orange mouse (*Mus orangiae*) is restricted to the ecoregion, and the rough-haired golden mole (*Chrysofalax villosa*) is near-endemic. The ecoregion also supports populations of several large mammal species, some of which are rare in southern Africa (Stuart and Stuart 1995). Among these are the brown hyena (*Hyaena brunnea*), African civet (*Civettictis civetta*), leopard (*Panthera pardus*), pangolin (*Manis temminckii*), honey badger (*Mellivora capensis*), striped weasel (*Poecilogale albinucha*), aardwolf (*Proteles cristatus*), oribi (*Ourebia ourebi*), and mountain zebra (*Equus zebra hartmannae*).

Predators that still roam freely in the area include larger predators such as brown hyena, while smaller predators such as caracal, serval and honey badger are common throughout the larger area. Antelope species such as duiker and steenbok will roam freely through the area and are not restricted by game fences. Smaller mammal species such as honey badgers and serval can become habituated to anthropogenic influences, while other species such as brown hyena will rather move away from the construction activities and will seldom use the area.

The connectivity<sup>1</sup> of the project site to the remainder of the larger area is Moderate due to other surrounding areas representing natural grassland and wetlands. Of significance is the role of the wetlands and indigenous grasslands as zoogeographical dispersal corridor.

Most mammal species are highly mobile and will move away during construction of the solar development. The most important corridors that need to be preserved for free-roaming mammal species in the area include the wetlands and indigenous grasslands.

### 4.3.4.2 Birds (avifauna)

Bird species richness is relatively high within the Highveld Ecoregion (Harrison et al. 1997). However, Botha's lark (*Spizocorys fringillaris*) is the only bird species strictly endemic to the ecoregion, where it inhabits heavily grazed grassland. An additional six species of birds are near-endemics including whitewinged flufftail (*Sarothrura ayresii*), blue korhaan (*Eupodotis caerulescens*), southern whitebellied korhaan (*Eupodotis cafra*), Rudd's lark (*Heteromirafra ruddi*), melodious lark (*Mirafra cheniana*), buff-streaked chat (*Oenanthe bifasciata*), and yellow-breasted pipit (*Hemimacronyx chloris*) (Harrison et al. 1997).

Many grassland birds, several of which are endemic to southern Africa, show a clear preference for sour over sweet and mixed grassland, and some of these are essentially absent from the last two grassland types, e.g. Bald Ibis, Redwing Francolin, Blackwinged Plover, Rudd's Lark, Botha's Lark, Blue Swallow, Buffstreaked Chat, Palecrowned Cisticola and Yellowbreasted

---

<sup>1</sup> **Connectivity (habitat connectivity)** - Allowing for the conservation or maintenance of continuous or connected habitats, so as to preserve movements and exchanges associated with the habitat.

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Pipit. Examples of grassland species preferring sweet and mixed grasslands appear fewer but include Melodious Lark and South African Cliff Swallow. The extensive human pressures on the grassland biome have severe conservation implications for its avifauna: many of the globally threatened species present on the mainland of South Africa, Lesotho and Swaziland have major strongholds in the grassland biome and five of these (Bald Ibis, Whitewinged Flufftail, Rudd's and Botha's larks, and Yellowbreasted Pipit) are entirely restricted to this biome in the region.

The grassland occurs throughout the project area. Bird species such as crowned plovers, crested guineafowls, francolin species as well as the birds of prey the smaller bird species attract utilize these areas. Although this microhabitat is in a degraded state, the area is a popular habitat for bird species, especially as foraging area, while species such as crowned plover and other smaller non-passerine birds also breed on the ground in this area.

More than 250 bird species have been recorded in the project area and surroundings. Globally threatened species include Secretarybird. Congregatory birds are Egyptian Goose, Western Cattle Egret, Spur-winged Goose, South African Shelduck, Cape Shoveler and African Spoonbill.

According to Birdlife South Africa, the study area falls outside of any Important Bird Areas (IBA), identified within South Africa ([www.birdlife.org.za](http://www.birdlife.org.za)). The conservation status of many of the bird species that are dependent on wetlands reflects the critical status of wetland nationally, with many having already been destroyed. In the study area, man-made dams represent wetland areas.

### 4.3.4.3 Herpetofauna (Reptiles and Amphibians)

Twenty-nine amphibians occur within the ecoregion but none are endemic (Passmore and Carruthers 1995). Breeding habitat of frogs and toads can be found mostly in the permanent wet zone of the wetlands and dams in the larger area. Amphibian species potentially occurring in the larger area include Common River Frog, Natal Sand Frog, Gutteral Toad, Raucous Toad and Bubbling Kassina. These species are non-threatened and widespread, and as such the development will not have any impact on amphibian conservation within the region.

Relatively few reptile species occur within the Highveld Ecoregion, mainly due to its cool climate. However, the ecoregion supports some of Africa's most characteristic reptile species, including Nile crocodile (*Crocodylus niloticus*), African rock-python (*Python sebae*), water monitor (*Varanus niloticus*) and veld monitor (*Varanus exanthematicus albigularis*). There are also two strict endemic reptiles: giant girdled lizard (*Cordylus giganteus*), and *Agama distanti* (Branch 1998). Several additional reptile species are near-endemics, including Drakensberg rock gecko (*Afroendura niravia*), giant spinytail lizard (*Cordylus giganteus*), and Breyer's whiptail (*Tetradactylus breyeri*) (Branch 1998).

In the presence of dead termitaria, the small geckos listed are probably found on the site. A few terrestrial lizards (Yellow-throated Plated Lizard, Variegated Skink), typical for Highveld



## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Grassveld, are expected to be present. A variety of smaller snake species characteristic for Highveld Grassveld will be present (Common Wolf Snake, Brown House Snake), although some might be dependent on by the presence of dead termitaria. The only venomous snakes, which has been reported as being present and common, is as expected, the Rinkhals, Mozambique spitting cobra, snouted cobra and the Puffadder for this QDS. All the reptile species are common and widespread, and as such the development will not have any impact on reptile conservation within the region. The sungazer lizard occurs in some of the grassland areas, while the southern spiny agama and the striped harlequin snake may occur in small numbers in suitable habitat.

### 4.3.5 Species of Conservation Concern (SCC)

According to the existing databases and field survey the following number of fauna species included in the IUCN red data lists can potentially be found in the study area (Table 8):

**Table 8. Red data list of potential fauna for the study area**

English Name	Conservation Status	Probability of occurrence on site
<b>BIRDS</b>		
Pallid Harrier	Near Threatened	Moderate
Greater Flamingo	Near Threatened	Low (wetlands)
Grass Owl	Vulnerable	Moderate
Secretarybird	Vulnerable	Moderate
Caoe Vulture	Endangered	Moderate
Macoa Duck	Near Threatened	Low (wetlands)
<b>MAMMALS</b>		
Serval	Near Threatened (2016)	Moderate
African Clawless Otter	Near Threatened (2016)	Low (wetlands)

The following impacts might occur during the development phase on the fauna populations of the area:

- Destruction/permanent loss of individuals of rare, endangered, endemic and/or protected species through habitat loss or fragmentation.
- Disturbance of remnant terrestrial wild mammal, avian, amphibian and insect fauna would probably occur through physical habitat destruction, noise, traffic, and movement of people.
- Potential increase in feral animals and impact on indigenous fauna e.g., cats, rats.
- Illegal hunting or disturbance.

The following management measures are proposed regarding the conservation of these and other fauna which might occur on the property:

- The development would not have a significant impact on the above-mentioned red data fauna since adequate and natural habitat/vegetation would be available on the

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

peripheral grassland habitats surrounding the development site. The most probable habitat to find any of the red data species in the study area would be in the more natural areas of the grassland and wetlands where little or no disturbances from humans or livestock occur at a regular interval. Fauna will therefore rather move away from the area and utilize adjacent, more natural areas. The importance to preserve the wetland habitat to the south of the development footprint should still be considered a high priority though.

- The removal of vegetation should be confined to the footprints of the proposed development site. This will be on small sections in relation to the total available surrounding habitat for fauna. Development also will not influence the natural feeding and movement patterns of the existing fauna in the area.
- If one considers the habitat descriptions of the red data species, most of them are not directly threatened by habitat loss. The impact of development on the red data species would therefore be less than predicted.
- The protection of different habitat types in the area will be important to ensure the survival of the different animals due to each species' individual needs and requirements. Sufficient natural corridor sections should be protected around the proposed development footprints to allow fauna to move freely between the different vegetation units on the property. The drainage channels and sections of natural vegetation will be preserved as corridors in the area and mitigation measures should be implemented to ensure that the habitats are protected.
- The taller (>3m) indigenous trees within this area also provide resting/perching sites for larger birds like birds of prey, arboreal reptiles and mammals that might occur/pass through the area and should preferably be preserved. These larger trees should be protected as far as possible and be incorporated into the proposed development. The removal of large dead trees is also not advised as these trees also provide smaller habitats for the mentioned bat species as well as rodents. The grass layer on the other hand also provides a valuable food source (insects, reptiles, small mammals that occur in/on the grass layer) for fauna.
- A monitoring programme needs to be implemented by a specialist if any rare species are confirmed on the property.

The following practical recommendations with regards to the fauna of the area apply with regards to the construction of the proposed development:

- Where trenches pose a risk to animal safety during construction, they should be adequately cordoned off to prevent animals falling in and getting trapped and/or injured. This could be prevented by the constant excavating and backfilling of trenches during the construction.
- No animals may be poached. Many animals are protected by law and poaching, or

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

other interference could result in a fine or jail term.

- Do not feed any wild animals on site.
- Poisons for the control of problem animals should rather be avoided since the wrong use thereof can have disastrous consequences for the raptors occurring in the area. The use of poisons for the control of rats, mice or other vermin should only be used after approval from an ecologist.
- Walkways and roads should be designed without vertical pavements to allow for the movement of small mammals.
- Waste bins and foodstuffs should be made scavenger proof.
- Monitoring of the environmental aspects is recommended for the future phases of the proposed development should the authorities approve the application. The monitoring phase would ensure that negative impacts on the fauna and flora of the area are limited to a minimum during the construction phase.

### 4.3.6 EIA screening tool listed species

Table 9 indicate the listed species for the project area according to the EIA screening tool:

**Table 9. Listed fauna species for the project area according to the EIA screening tool, status and habitat.**

Species	Status	Habitat
<i>Tyto capensis</i>	Vulnerable	Habitat includes moist grassland and open savanna up to an elevation of 3200m. May also be found in dry grassland and at higher altitudes. In each case, habitat is normally characterised by long dense grass.
<i>Eupodotis senegalensis</i>	Vulnerable	It is widespread in sub-Saharan Africa in grassland and open woodland habitats.

#### 4.3.6.1 *Tyto capensis*

Habitat includes moist grassland and open savanna up to an elevation of 3200m. May also be found in dry grassland and at higher altitudes. In each case, habitat is normally characterised by long dense grass.

The African Grass-Owl *Tyto capensis* has experienced a reduction in regional population size and satisfies the population-size criterion for Vulnerable (an observed, estimated, inferred or suspected population size reduction of 30% over the last 10 years or three generations, whichever is the longer, where the reduction or its causes may not have ceased OR may not be understood OR may not be reversible). In addition, the population of less than 10 000 mature individuals is projected to decline by at least 10% within the next three generations. For these reasons, the species is listed as regionally Vulnerable.

The species is habitat-specific, and only a proportion of its distribution represents suitable

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

ecological conditions. A basic model of African Grass Owl habitat, generated for Gauteng using high-resolution land-cover data, showed that a maximum of 25% of rural Gauteng potentially comprised suitable habitat for the species.

The species is believed to have undergone a reduction in population size of greater than 30% in the past three generations and is projected to decline by at least 10% within the next three generations. However, in the absence of more detailed historical population data, the rate of population decline is difficult to estimate accurately.

The primary threat to the African Grass Owl in the region is loss of habitat. Between 1994 and 2005, the combined footprint of urbanisation, afforestation, mining and cultivation in the three provinces that comprise the remaining core of the African Grass Owl's range in South Africa, i.e. KwaZulu-Natal, Mpumalanga and Gauteng, increased by an estimated 8.5%. The extent of potential African Grass Owl habitat affected remains to be assessed, but loss of wetland and associated grassland habitat is expected to exceed 20% in the next three generations. In support of this somewhat speculative suggestion, in Mpumalanga alone development applications between 2005 and 2010 covered 72% of the province. Applications were primarily for prospecting and mining for coal and covered 90% of areas in the province regarded as having high importance for groundwater re-charge and 80% of those areas with high importance for water run-off, thus increasing the probability that African Grass Owl habitat will be affected.

Fire and grazing are important tools for the management of grassland and wetland habitats, but regular heavy grazing pressure and too frequent burning prevent the development of rank grassland habitat required by African Grass Owls (Brooke 1984, Jansen et al. 1999). Their habit of nesting on the ground makes eggs and chicks vulnerable to fire and trampling by livestock (Tarboton and Erasmus 1998, Whittington-Jones 2010). Wetland drainage schemes and incompatible farming practices may explain the apparent absence of this species from Lesotho, much of Eastern Cape (Brooke 1984) and more recently from large areas of its former range in the rest of South Africa. In a slightly more positive vein, given that it is nocturnal and roosts in tall, dense grass during the day, under-recording is a problem. The species frequently hunts along road verges (Ansara 2004), where it is likely preying on rodents attracted to grain spilled by passing trucks. Consequently, African Grass Owls are frequently killed by vehicles at night, and are well-represented in museum collections. Collisions with vehicles are a significant cause of direct mortality: 27% of the 554 owl carcasses recovered from portions of two roads in Gauteng between October 2001 to September 2003 were those of African Grass Owls (Ansara 2004). Entanglement with barbed-wire fences is another potentially significant, but poorly documented cause of mortality.

**Probability of occurrence on site: MODERATE** due to the presence of suitable habitat on the proposed development footprint, although no population of the species occur on site.

**Probability of impact during vegetation clearance: MODERATE, no populations documented although some habitat considered suitable.**

### 4.3.6.2 *Eupodotis senegalensis*

Habitat includes grassland and open woodland habitats. The regional population of White-bellied Korhaan *Eupodotis senegalensis* satisfies the population size-reduction criterion for regionally Vulnerable (=30% decline over the past three generations where the reduction or its causes may not have ceased and may not be reversible, based on a decline in AoO, EoO and/or quality of habitat). It is believed that this trend will continue for the next three generations.

The global population is suspected to be decreasing due to habitat destruction, but not at a rate sufficient to qualify the species as globally Vulnerable (BirdLife International 2014b). Assuming the population estimates provided above are accurate, the decrease in the regional population over the past three generations (10.3 years) would be c. 73%, hence an assessment of regionally Vulnerable. Generation length was calculated based on extrapolated mean age at first breeding and extrapolated maximum longevity in the wild (BirdLife International 2014b). Confidence in this is low.

The main threats is a familiar list of problems also facing other bustard species. The relative severity and potential impact of these threats have not been quantified or even assessed, but of particular concern are habitat loss and degradation due to agriculture, afforestation (invasive alien vegetation and timber plantations), overgrazing, urban development, unsuitable burning practices, and other habitat modifications as a result of growing human populations (Moreira 2004, Allan 2005i). Clancey (1972) mentioned fires as a threat to chicks, but this is unlikely to constitute a major threat during the breeding season in the summer rainfall area. The loss of habitat to bush encroachment poses a threat to White-bellied Korhaans, although the species appears to be adaptable to low levels of woody cover in grasslands. Apart from habitat loss, the threat of subsistence hunting and poaching, due to high human densities, also needs to be considered. Collisions with power-lines do not seem to pose as serious a threat as it does to larger bustard species (Shaw 2009), with only a single record of a male killed in this fashion on record (Allan 2005i).

**Probability of occurrence on site: MODERATE** due to the presence of suitable habitat on the proposed development footprint, although no population of the species occur on site.

**Probability of impact during vegetation clearance: MODERATE, no populations documented although some habitat considered suitable.**

### 5 POTENTIAL IMPACTS OF THE PROPOSED DEVELOPMENT ON THE FAUNA AND FLORA

An environmental impact is defined as a change in the environment, be it the physical/chemical, biological, cultural and or socio-economic environment. Any impact can be related to certain aspects of human activities in this environment and this impact can be either positive or negative. It could also affect the environment directly or indirectly and the effect of it can be cumulative. There are three major categories of impacts on biodiversity namely:

- Impacts on habitat resulting in loss, degradation and / or fragmentation.
- Direct impacts on fauna and flora and species, for example plants and animals that are endemic / threatened / special to a habitat will not be able to survive if that habitat is destroyed or altered by the development.
- Impact on natural environmental processes and ecosystem functioning. This can lead to an accumulated effect on both habitat and species.

This biodiversity assessment focused on the description of ecosystem- and species-related biodiversity. It can be expected that if ecosystem diversity is managed effectively, species and genetic diversity should also be protected. Emphasis was therefore placed on the ecosystem diversity (landscape/habitat types) within the proposed development area, with reference to biota observed and expected to utilise these landscapes or habitat types.

#### 5.1 POTENTIAL IMPACTS

##### 5.1.1 Direct habitat destruction

###### 5.1.1.1 *Description of impact:*

The construction phase of the development and associated infrastructure will result in loss of and damage to natural habitats if the vegetation is cleared for the development of the solar plant. Rehabilitation of some areas would be possible but there is likely to be long-term damage in large areas. Most habitat destruction will be caused during the construction phase. Vegetation communities are likely to be impacted on a small spatial scale in comparison to the extent of the vegetation communities' total area in the region.

The impact of the habitat destruction will be on the flora and fauna of the study area in the following ways:

- The construction will lead to the loss of individual plants such as grasses, forbs, trees, and shrubs that will be cleared on the footprint area. This will mostly occur during the construction phase.
- Loss of threatened, near-threatened and endemic taxa: The anticipated loss of some of the natural habitats that support endemic species will result in the local displacement of endemic listed flora.

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

- Due to habitat loss and construction activities animals will migrate from the construction area and animal numbers will decrease.
- Loss of threatened, “near-threatened” and conservation important taxa: The anticipated loss of the natural woodland will result in the local displacement of some fauna species. In some cases, isolated populations of threatened fauna might be removed from the area, although no such populations or knowledge thereof was found in the study area. This impact could also take place because of hunting and snaring of animals in natural areas not used for the mine or its infrastructure.
- Changes in the community structure: It is expected that the faunal species composition will shift, due to an anticipated loss in habitat surface area. In addition, it is predicted that more generalist species (and a loss of functional guilds) will dominate the study area. Attempts to rehabilitate will attract taxa with unspecialized and generalist life-histories. It is predicted that such taxa will persist for many years before conditions become suitable for succession to progress.

### 5.1.1.2 *Mitigation measures:*

- The removal of indigenous plants should be kept to a minimum necessary. Trim, rather than fell of woody species along the edges of the development site where possible. The clearing and damage of plant growth in the riparian and wetland areas should be restricted to the actual road crossing where possible, and not into the sensitive adjacent areas. Where protected plants such as geophytes will need to be cleared or pruned, permits should be obtained from the relevant authority.
- Peripheral impacts around the development footprint sites on the surrounding vegetation of the area should be avoided and a monitoring programme should be implemented to ensure the impacts are kept to a minimum, while the rehabilitation of the site should be prioritized after construction has been completed.
- During construction, sensitive habitats must be avoided by construction vehicles and equipment, wherever possible, to reduce potential impacts. Only necessary damage must be caused and, for example, unnecessary driving around in the veld or bulldozing natural habitat must not take place.
- An avifauna specialist should be consulted to conduct a specialist study for the project area and monitoring of the potential impact of the solar plant in the future.
- All development activities should be restricted to specific recommended areas. The Environment Control Officer (ECO) should control these areas. Storage of equipment, fuel and other materials should be limited to demarcated areas. Layouts should be adapted to fit natural patterns rather than imposing rigid geometries. The entire development footprint should be clearly demarcated prior to initial site clearance and prevent construction personnel from leaving the demarcated area. This would only be applicable to the construction phase of the proposed development.

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

- The ECO should advise the construction team in all relevant matters to ensure minimum destruction and damage to the environment. The ECO should enforce any measures that he/she deem necessary. Regular environmental training should be provided to construction workers to ensure the protection of the habitat, fauna and flora and their sensitivity to conservation.
- Where holes for poles pose a risk to animal safety, they should be adequately cordoned off to prevent animals falling in and getting trapped and/or injured. This could be prevented by the constant excavating and backfilling during planting of the poles along the lines.
- Poisons for the control of problem animals should rather be avoided since the wrong use thereof can have disastrous consequences for the raptors occurring in the area. The use of poisons for the control of rats, mice or other vermin should only be used after approval from an ecologist.
- Limit pesticide use to non-persistent, immobile pesticides and apply in accordance with label and application permit directions and stipulations for terrestrial and aquatic applications.
- Monitoring should be implemented during the construction phase of the development to ensure that minimal impact is caused to the fauna and flora of the area.
- A detailed wetland assessment should be conducted to determine the exact edges of potential wetlands and drainage channels.

### 5.1.2 Habitat fragmentation

#### 5.1.2.1 *Description of impact:*

The construction of the development and associated infrastructure will result in natural movement patterns being disrupted for a limited period and, to a varying degree depending on how different species react to these barriers will result in the fragmentation of natural populations, although the impact will be minimal and restricted to the construction phase.

#### 5.1.2.2 *Mitigation measures:*

- Use existing facilities (e.g., impacted areas) to the extent possible to minimize the amount of new disturbance.
- Ensure protection of important resources by establishing protective buffers to exclude unintentional disturbance. All possible efforts must be made to ensure as little disturbance as possible to the sensitive features such as surrounding woodland and riparian woodland outside the project area during construction.
- During construction, sensitive habitats must be avoided by construction vehicles and equipment, wherever possible, to reduce potential impacts. Only necessary damage must be caused and, for example, unnecessary driving around in the veld or bulldozing natural habitat must not take place.



## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

- Construction activities must remain within defined construction areas. No construction / disturbance will occur outside these areas.

### 5.1.3 Increased Soil erosion and sedimentation

#### 5.1.3.1 *Description of impact:*

The construction activities associated with the development may result in widespread soil disturbance and is usually associated with accelerated soil erosion. Soil erosion promotes a variety of terrestrial ecological changes associated with disturbed areas, including the establishment of alien invasive plant species, altered plant community species composition and loss of habitat for indigenous flora.

#### 5.1.3.2 *Mitigation measures:*

The following mitigation measures should be implemented to prevent erosion during construction:

- The project should be divided into as many phases as possible, to ensure that the exposed areas prone to erosion are minimal at any specific time.
- Cover disturbed soils as completely as possible, using vegetation or other materials.
- Minimize the amount of land disturbance and develop and implement stringent erosion and dust control practices.
- Protect sloping areas and drainage channel banks that are susceptible to erosion and ensure that there is no undue soil erosion resultant from activities within and adjacent to the construction camp and Work Areas.
- Repair all erosion damage as soon as possible to allow for sufficient rehabilitation growth.
- Gravel roads to the construction sites must be well drained to limit soil erosion.
- Control the flow of runoff to move the water safely off the site without destructive gully formation.
- Protect all areas susceptible to erosion and ensure that there is no undue soil erosion resultant from activities within and adjacent to the construction camp and Work Areas.

### 5.1.4 Soil and water pollution

#### 5.1.4.1 *Description of impact:*

Construction work for the proposed development will always carry a risk of soil and water pollution, with large construction vehicles contributing substantially due to oil and fuel spillages. If not promptly dealt with, spillages or accumulation of waste matter can contaminate the soil

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

and surface or ground water, leading to potential medium/long-term impacts on fauna and flora. During the constructional phase heavy machinery and vehicles would be the main contributors to potential pollution problems.

### 5.1.4.2 *Mitigation measures:*

- Any excess or waste material or chemicals should be removed from the site and discarded in an environmentally friendly way. The ECO should enforce this rule rigorously.
- Spill kits should be on-hand to deal with spills immediately.
- All vehicles should be inspected for oil and fuel leaks on a regular basis. Vehicle maintenance yards on site should make provision for drip trays that will be used to capture any spills. Drip trays should be emptied into a holding tank and returned to the supplier.

### 5.1.5 **Air pollution**

#### 5.1.5.1 *Description of impact:*

The environmental impacts of wind-borne dust, gases and particulates from the construction activities associated with the proposed development are primarily related to human health and ecosystem damage. The proposed development will typically comprise the following sources and associated air quality pollutants:

- Materials handling operations (truck loading & unloading, tipping, stockpiling).
- Vehicle entrainment on paved and unpaved roads.
- Windblown dust-fugitive emissions.

One of the primary impacts on the biophysical environment is linked to emission of dusts and fumes from both the transportation system. Dust pollution will impact the most severe during the construction phase. Construction vehicles and equipment are the major contributors to the impact on air quality. Dust is generated during site clearance for the construction of infrastructure. Diesel exhaust gasses and other hydrocarbon emissions all add to the deterioration in air quality during this phase. Vehicles travelling at high speeds on dirt roads significantly aggravate the problem.

Although the potential for severe fugitive dust impacts is greatest within 100 m of dust-generating activities, there is still the potential for dust to affect vegetation up to five kilometres or more downwind from the source. Dust deposited on the ground may cause changes in soil chemistry (chemical effects) and may over the long-term result in changes in plant chemistry, species composition and community structure. Sensitivities to dust deposition of the various plant species present in the area are not known. It is therefore difficult to predict which species may be susceptible.

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Poor air quality results in deterioration of visibility and aesthetic landscape quality of the region, particularly in winter due to atmospheric inversions.

### 5.1.5.2 *Mitigation measures:*

- A speed limit should be enforced on dirt roads (preferably 30-40km/h).
- Implement standard dust control measures, including periodic spraying (frequency will depend on many factors including weather conditions, soil composition and traffic intensity and must thus be adapted on an on-going basis) of construction areas and access roads, and ensure that these are continuously monitored to ensure effective implementation.

### 5.1.6 **Spread and establishment of alien invasive species**

#### 5.1.6.1 *Description of impact:*

Continued movement of vehicles on and off the site during the construction phase will result in a risk of importation of alien species. Vehicles often transport many seeds, and some may be of invader species, which may become established along the access road, especially where the area is disturbed. The construction carries by far the greatest risk of alien invasive species being imported to the site, and the high levels of habitat disturbance also provide the greatest opportunities for such species to establish themselves, since most indigenous species are less tolerant of disturbance. The biggest risk is that seeds of noxious plants may be carried onto the site along with materials that have been stockpiled elsewhere at already invaded sites.

#### 5.1.6.2 *Mitigation measures:*

- Control involves killing the plants present, killing the seedlings which emerge, and establishing and managing an alternative plant cover to limit re-growth and re-invasion. Weeds and invader plants will be controlled in the manner prescribed for that category by the CARA or in terms of Working for Water guidelines. The control of these species should even begin prior to the construction phase considering that small populations of these species was observed during the field surveys.
- Institute strict control over materials brought onto site, which should be inspected for seeds of noxious plants and steps taken to eradicate these before transport to the site. Routinely fumigate or spray all materials with appropriate low-residual herbicides prior to transport to or in a quarantine area on site. The contractor is responsible for the control of weeds and invader plants within the construction site for the duration of the construction phase. Alien invasive tree species listed by the CARA regulations should be eradicated.
- Rehabilitate disturbed areas as quickly as possible to reduce the area where invasive species would be at a strong advantage and most easily able to establish.
- Institute a monitoring programme to detect alien invasive species early, before they

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

become established and, in the case of weeds, before the release of seeds. Once detected, an eradication/control programme should be implemented to ensure that the species' do not spread to surrounding natural ecosystems.

### 5.1.7 Negative effect of human activities and road mortalities

#### 5.1.7.1 *Description of impact:*

An increase in human activity on the site and surrounding areas is anticipated. The risk of snaring, killing, and hunting of certain faunal species is increased. If staff compounds are erected for construction workers, the risk of pollution because of litter and inadequate sanitation and the introduction of invasive fauna and flora are increased. The presence of many construction workers or regular workers during the construction phase on site over a protracted period will result in a greatly increased risk of uncontrolled fires arising from cooking fires, improperly disposed cigarettes etc.

Large numbers of fauna are also killed daily on roads. They are either being crushed under the tyres of vehicles in the case of crawling species, or by colliding with the vehicle itself in the case of avifauna or flying invertebrates. The impact is intensified at night, especially for flying insects, as result of their attraction to the lights of vehicles.

#### 5.1.7.2 *Mitigation measures:*

- No staff should be accommodated on the site. If practical, construction workers should stay in one of the nearby villages and transported daily to the site.
- The ECO should regularly inspect the site, including storage facilities and compounds and eradicate any invasive or exotic plants and animals.
- Maintain proper firebreaks around entire development footprint.
- Educate construction workers regarding risks and correct disposal of cigarettes.
- More fauna is normally killed the faster vehicles travel. A speed limit should be enforced (preferably 40 km/hour). It can be considered to install speed bumps in sections where the speed limit tends to be disobeyed. (Speed limits will also lessen the probability of road accidents and their negative consequences).
- Travelling at night should be avoided or limited as much as possible.

## 5.2 IMPACT ASSESSMENT MATRIX

Table 10 indicate the impacts described above and specific ratings of significance the development impact will potentially have on the ecological components of the study area.

Table 10. Impact assessment Matrix for the proposed development

Nr	Activity	Impact	Without or With Mitigation	Nature (Negative or Positive Impact)	Probability		Duration		Scale		Magnitude/ Severity		Significance		Mitigation Measures	Mitigation Effect
					Magnitude	Score	Magnitude	Score	Magnitude	Score	Magnitude	Score	Score	Magnitude		
<b>Terrestrial Biodiversity Impact Assessment</b>																
<b>Construction Phase</b>																
1	Clearing of vegetation for construction of infrastructure, access roads etc.	Habitat destruction & Fragmentation	WOM	Negative	Definite	5	Permanent	5	Local	1	Medium	8	70	High	Refer to Sections 5.1.1.2 and 5.1.2.2	May cause irreplaceable loss of resources
			WM	Negative	Definite	5	Permanent	5	Local	1	Low	6	60	Moderate		
2	Topsoil & subsoil stripping, exposure of soils to wind and rain during construction causing erosion and sedimentation in wetlands	Soil erosion and sedimentation	WOM	Negative	Definite	5	Permanent	5	Regional	3	High	8	80	High	Refer to section 5.1.3.2	Can be reversed
			WM	Negative	Highly Probable	4	Medium term	3	Site	2	Medium	6	44	Moderate		
3	Exposure of soils to rainfall and wind during construction	Dust pollution	WOM	Negative	Definite	5	Medium term	3	Site	2	Medium	6	55	Moderate	Refer to section 5.1.4.2	Can be reversed
			WM	Negative	Highly Probable	5	Medium term	3	Site		Low	2	25	Low		
4	Heavy machinery and vehicle movement on site	Spillages of harmful substances	WOM	Negative	Highly Probable	4	Long term	4	Regional	3	Medium	6	52	Moderate	Refer to section 5.1.5.2	Can be avoided, managed, or mitigated
			WM	Negative	Probable	2	Long term	4	Site	2	Low	2	16	Negligible		
5	Continued movement of personnel and vehicles on and off the site during the construction phase, as well as occasional delivery of materials required for maintenance	Spreading of alien invasive species	WOM	Negative	Highly Probable	4	Permanent	5	Site	2	Medium	6	52	Moderate	Refer to section 5.1.6.2	Can be reversed
			WM	Negative	Probable	2	Medium term	3	Site	2	Low	2	14	Negligible		
6	Construction of infrastructure, access roads etc.	Negative effect of human activities on fauna and flora	WOM	Negative	Highly Probable	4	Medium term	3	Site	2	Medium	6	44	Moderate	Refer to section 5.1.7.2	Can be avoided, managed, or mitigated
			WM	Negative	Probable	2	Medium term	3	Site	2	Low	2	14	Negligible		
7	Continued movement of vehicles on and off the site during the construction phase, as well as occasional delivery of materials required for maintenance	Road mortalities of fauna	WOM	Negative	Highly Probable	4	Medium term	3	Site	2	Medium	6	44	Moderate	Refer to section 5.1.8.2	Can be avoided, managed, or mitigated
			WM	Negative	Highly Probable	4	Medium term	3	Site	2	Low	2	28	Low		

### 6 ECOLOGICAL SENSITIVITY CLASSES

Following the ecological surveys, the classification of the study area into different sensitivity classes and development zones was based on information collected at various levels on different environmental characteristics. Factors which determined sensitivity classes were as follows:

- Presence, density and potential impact of development on rare, endemic and protected plant species.
- Conservation status of vegetation units.
- Soil types, soil depth and soil clay content.
- Previous land-use.
- State of the vegetation in general as indicated by indicator species.

Below included is the sensitivity map for the proposed solar development, (Figure 13). Only criteria applicable to the specific vegetation units were used to determine the sensitivity of the specific unit.

Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5




Figure 13. Sensitivity Map of the project area

### 7 DISCUSSION

Following the investigation and potential ecological impact of the proposed solar development on the biodiversity (including plant and animal species theme) of the area, some conclusions can be made:

All aspects of the environment, especially living organisms, are vulnerable to disturbance of their habitat. The proposed development activities will modify the vegetation and faunal habitats of the development site to a certain extent varying according to the habitats on the site, although in general the vegetation on site where the development footprint is planned are classified as pristine to slightly degraded.

**Most sensitive sections:** It is evident from the distribution of biodiversity, presence of threatened species and sites of scientific interest, that the proposed development has the potential for negative impact on the flora and faunal of the study area. This is particularly true of the sensitive vegetation associated with the riverine and wetland ecosystems and the project area.

**Most sensitive habitats:** Many threatened species are grassland specialists, linked to these habitats either for breeding, feeding or shelter. Major impacts on wetland areas adjacent to the site should be avoided wherever possible during construction. Where unavoidable impacts will occur on grassland and wetland zones, strict mitigation measures and legislation should be implemented (licence for eradication of protected plants, IWUL application etc.).

**Monitoring of threatened species:** Many endemic and protected species have been recorded in region. The EMP for the development should highlight the conservation status of these species and note that steps must be undertaken in conjunction with conservation authorities to protect or translocate any populations encountered during project actions. Ecological monitoring is recommended for the construction phase of the development considering the presence of protected trees and potential red data fauna on areas surrounding the site.

The importance of rehabilitation and implementation of mitigation processes to prevent negative impacts on the environment during and after the construction phase of the solar development should be considered a high priority. The proposed site for the development varies from being in a pristine to slightly degraded state.

A sensitivity analyses was conducted to identify the most suitable site for the development. From this investigation and ecological surveys, the following main observations was made:

- Most of the natural grassland and woodland have a Medium Sensitivity and development can be supported in the area provided certain mitigation measures are implemented. Where the clearance of the vegetation would cause protected



## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

plants or other fauna to be removed, permits should be obtained from the relevant authorities.

- The exotic bushclumps have a low sensitivity and unlimited development can be supported in this area.

The protected plant species *Boophane disticha* occur on the site and specific mitigation measures (permit applications, avoidance, relocation) should be implemented to avoid negative impacts on the species.

Some potential rare fauna may also occur in the area, and specific mitigation measures need to be implemented to ensure that the impact of the development on the species' habitat will be low. Specific mitigation relating to red data fauna includes the following:

- Disturbances in close vicinity of the development (periphery) should be limited to the smallest possible area to protect species habitat.
- Corridors are important to allow fauna to move freely between the areas of disturbance.

The indigenous grassland vegetation units on the proposed development site is not considered as Critical Habitat in line with IFC Performance Standard PS6.

Several potential impacts were identified and assessed. A few of these were assessed as having potentially medium or high significance, including the following:

- Destruction or disturbance to sensitive ecosystems leading to reduction in the overall extent of a particular habitat.
- Increased soil erosion.
- Impairment of the movement and/or migration of animal species resulting in genetic and/or ecological impacts.
- Destruction/permanent loss of individuals of rare, endangered, endemic and/or protected species.
- Soil and water pollution through spillages.
- Establishment and spread of declared weeds and alien invader plants.
- Impacts of human activities on fauna and flora of the area during construction.
- Air pollution through dusts and fumes from construction vehicles (construction).

Mitigation measures are provided that would reduce these impacts from a higher to a lower significance. Furthermore, the proposed layout plan of the development should be consistent with the sensitivity map and recommendations stipulated in this report, and the impact on the sensitive habitats on site should be kept to a minimum.

### 8 CONCLUSION

All aspects of the environment, especially living organisms, are vulnerable to disturbance of their habitat. If we can bring about a more integrated approach to living within our ecosystems, we are much more likely to save the fundamental structure of biodiversity. Positive contributions can be made even on a small scale such as within the proposed solar development. All stakeholders, such as business, government and environmental groups need to be involved to the impacts associated with the development from causing a significant loss.

The proposed development should allow corridors of indigenous grassland and wetlands on areas outside the development footprint to be preserved. Where sensitive areas of natural vegetation cannot be avoided, a few mitigation measures have been recommended to minimise and/or offset impacts (licence application for eradication of protected species.). Negative impacts can be minimised by strict enforcement and compliance with an Environmental Management Plan which considers the recommendations for managing impacts detailed above.

**Provided that the proposed development and layout plans is consistent with the sensitivity map and take all the mitigation measures into consideration stipulated in this report, the planned development can be supported.**

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

### 9 REFERENCES

- Acocks, J.P.H. 1988. Veld types of South Africa, 3rd ed. Memoirs of the Botanical Survey of South Africa. 57: 1–146.
- Barbour, M.G., J.H. Burk, and W.D. Pitts. 1987. Terrestrial Plant Ecology. Second Edition. Benjamin/Cummings Publishing, Menlo Park, CA.
- BOTHMA, J. DU. P. 1996. Game Ranch Management. Van Schaick, Pretoria.
- Bredenkamp, G.J. & Brown, L.R. 2001. Vegetation – A reliable ecological basis for environmental planning. Urban Greenfile Nov-Dec 2001: 38-39.
- BRADY, N. C. & WEIL, R. R. 1996. The Nature and properties of Soils. Prentice Hall, New Jersey.
- Branch, B. (1998). Field guide to snakes and other reptiles of Southern Africa. Struik Publishers. Cape Town.
- Briza publications. 2001. Problem plants of South Africa. Pretoria.
- CHECHI, F. & ROBERTS, L. 2005. Interpreting and using mortality data in humanitarian emergencies: A primer for non-epidemiologists. Humanitarian practice Network at ODI.
- CONSERVATION OF AGRICULTURAL RESOURCES ACT, 1983. (ACT No. 43 OF 1983)
- Convention on Biological Diversity. Signed 1993 and ratified 2 November 1995.
- Cowling, W. E. 2005. Tourism- A Catalyst for Attitudinal Changes in Aitutaki, Cook Islands University of Waikato, Hamilton, New Zealand
- DEAT, 1998. Guideline Document on the EIA Regulations implementation of sections 21, 22 and 26 of the Environment Act, Government Printer, Pretoria.
- DEAT, 2002. Impact Significance, Integrated Environmental Management, Information Series 5, Department of Environmental Affairs and Tourism, Pretoria
- DWAF. 2003. A practical field procedure for identification and delineation of wetlands and riparian areas. Department of Water Affairs and Forestry, Pretoria.
- Enpat, 2000. Environmental Potential Atlas. Department of Environmental Affairs and Tourism, Pretoria.
- Fabian, A & Germishuizen, G. 1997. Wildflowers of Northern South Africa. Fernwood Press.
- Friedman, Y & Daly, B. 2004. Red Data Book of the Mammals of South Africa: A Conservation Assessment: CBSG Southern Africa, Conservation Breeding Specialist Group (SSC/IUCN), Endangered Wildlife Trust. South Africa.
- Germishuizen, G. and Clarke, B. (2003). Illustrated Guide to the Wildflowers of Northern South Africa. Briza Publications, Pretoria
- GERTENBACH, W. P. D. 1983. Landscapes of the Kruger National Park. Koedoe 26: 9-121.
- GOLDING, J. (Ed.) 2002. Southern African Plant Red Data Lists. Southern African Botanical Diversity Network report no. 14. National Botanical Institute. pp. 237.
- HILTON-TAYLOR, C. 1996a. Red Data list of southern African plants. Strelitzia 4: 1 - 117.

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

HILTON-TAYLOR, C. 1996b. Red Data list of southern African plants. 1. corrections and additions. *Bothalia* 26: 177 - 182.

HILTON-TAYLOR, C. 1997. Red Data list of southern African plants. 2. corrections and additions. *Bothalia* 27: 195 - 209.

IFC. Performance Standard 6 Biodiversity Conservation and Sustainable Natural Resource Management

Kent, LE. 1980. Stratigraphy of South Africa. Part 1: Lithostratigraphy of the Republic of South Africa, South West Africa/Namibia and the Republics of Bophuthatswana, Transkei, and Venda. Pretoria: Department of Mineral and Energy Affairs, Handbook 8.

KOTZE, D. C., MARNEWECK, G. C., BATCHELOR, A. L., LINDLEY, D. S. & COLLINS, N. B. 2005. Wet-ecoServices: A technique for rapidly assessing ecosystem services supplied by wetlands. South Africa National Biodiversity Institute, Pretoria.

Land type Survey Staff, 1987. Land types of the maps. Mem. Agric. Nat. Resour. S. Afr. no. 8.

LEE, K. E. & WOOD, T. G. 1971. Termites and Soils. Academic Press, London.

LOW, A. B. & REBELO, A. G. 1996. Vegetation of South Africa, Lesotho, and Swaziland. Dept. Environmental Affairs and Tourism, Pretoria.

MacKay, H. 1998: Towards a Classification System for Water Resources in South Africa. Institute for Water Quality Studies. Internal Report. Department of Water Affairs and Forestry, Pretoria, South Africa.

MACVICAR, C. N. 1991. Soil Classification: A Taxonomic system for South Africa. Department of Agriculture, Pretoria.

MCLEESE, R.L. AND WHITESIDE, E.P. 1977. Ecological effects of highway construction upon Michigan woodlots and wetlands: soil relationships. *Journal of Environmental Quality*. v6 n4, 476-471.

Manning, J. (2003). *Photographic Guide to the Wildflowers of South Africa*. Briza Publications. Pretoria.

Minter, L.R., Burger, M., Harrison, J.A., Braack, H.H., Bishop, P.J. and Kloepfer, D. (2004). *Atlas and Red Data Book of the Frogs of South Africa, Lesotho, and Swaziland*. Smithsonian Institute, Washington, DC.

Mucina, L., Bredenkamp, G.J., Hoare, D.B. & McDonald, D.J. 2000. A National vegetation database for South Africa. *South Africa Journal of Science* 96:497-498.

Mueller-Dombois, D. & Ellenberg, H. 1974. *Aims and methods of vegetation ecology*. Wiley, New York.

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

- Mucina, L & Rutherford, M. C. 2006. The vegetation of South Africa, Lesotho, and Swaziland. *Strelitzia* 19, SANBI, Pretoria.
- NATIONAL FOREST ACT, 1998 (Act No. 84 of 1998). Government Gazette No. 29062, Notice 897, 8 September 2006)
- NATIONAL WATER ACT, 1998. Act No 36 of 1998.
- Onderstall, J. (1996). *Wildflower Guide for Mpumalanga and Northern Province*. DynamicAd. Nelspruit.
- Palgrave, M.C. (2002). *Trees of Southern Africa*. Struik Publishers. Cape Town.
- Pooley, E. 1998. *A field guide to wildflowers of Kwazulu Natal and the Eastern Region*. Natal Flora Publications Trust.
- SANBI & DEAT. 2009. *Threatened Ecosystems in South Africa: Descriptions and Maps*. DRAFT for Comment. South African National Biodiversity Institute, Pretoria, South Africa.
- Sinclair, A. R. E. & A. E. Byrom. 2006. Understanding ecosystem dynamics for conservation of biota. *Journal of Animal Ecology*, 75: 64–79
- Smithers, R.H.N. (1983). *Soogdiere van die Suider-Afrikaanse Substreek*. Universiteit van Pretoria. Pretoria
- Tainton, N. M. (ed.), 1981. *Veld and Pasture Management in South Africa*. Shuter and Shooter, Pietermaritzburg, 481pp.
- The National Environmental Management Biodiversity Act, 2004. (Act 10 Of 2004). Government Gazette RSA Vol. 467, 26436, Cape Town, June 2004.
- The National Environmental Management Biodiversity Act, 2004. (Act 10 Of 2004). Draft. List of Threatened Ecosystems. Government Gazette RSA Vol. 1477, 32689, Cape Town, 6 Nov 2009.
- The Natural Scientific Professions Act (Act 27 of 2003)
- THOMPSON H (2006) *Water Law: A Practical Approach to Resource Management and the Provision of Services*. Juta, Cape Town.
- Van Der Merwe, C. R. 1952. Soil Groups and subgroups of South Africa. *Science Bulletin* 356.
- VAN WYK, B-E. & GERICKE, N. 2000. *People's Plants: A Guide to useful plants of southern Africa*. Briza publications, Pretoria.
- Van Wyk, B & Malan, S. 1988. *Field Guide to the wildflowers of the Highveld*. Struik Publishers.
- Van Wyk, B. & Van Wyk, P. 1997. *Field Guide to Trees of Southern Africa*. Struik Publishers. Cape Town.
- Van Wyk, B.E., Van Oudtshoorn, B. & Gericke, N. 1997. *Medicinal plants of South Africa*. Briza, Pretoria.
- Van Oudtshoorn, F. (1991) *Gids tot grasse van Suid Afrika*. Briza Publikasies. Pretoria.
- WERGER, M.J.A. 1978. *Biogeography and Ecology of Southern Africa*. Monographie Biologicae vol. 31. Junk, The Hague.
- Westhoff, V. & Van der Maarel, E. 1978. The Braun-Blanquet approach. In: Whittaker, R.H. (ed.)

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Classification of plant communities. W. Junk, The Hague.

WHITE, F. 1983. The vegetation of Africa: a descriptive memoir to accompany the UNESCO/AETFAT/UNSO vegetation map of Africa. UNESCO, Paris, France.

WINTER, C. 1988. A conceptual framework for assessing cumulative impacts on the hydrology of nontidal wetlands. Environmental Management. v12, n5, 605-620. APPENDIX A. PLANT SPECIES LISTs FOR QDS

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

### APPENDIX A. PLANT SPECIES IN QDS

Family	Species	IUCN	Ecology
Acanthaceae	<i>Crabbea hirsuta</i>	LC	Indigenous
Convolvulaceae	<i>Ipomoea bathycolpos</i>	LC	Indigenous; Endemic
Scrophulariaceae	<i>Jamesbrittenia aurantiaca</i>	LC	Indigenous
Apocynaceae	<i>Gomphocarpus fruticosus</i>	LC	Indigenous
Cyperaceae	<i>Kyllinga alba</i>	LC	Indigenous
Poaceae	<i>Sporobolus fimbriatus</i>	LC	Indigenous
Polygalaceae	<i>Polygala transvaalensis</i>	LC	Indigenous
Ricciaceae	<i>Riccia okahandjana</i>		Indigenous
Apocynaceae	<i>Raphionacme hirsuta</i>	LC	Indigenous
Convolvulaceae	<i>Ipomoea obscura</i>	LC	Indigenous
Scrophulariaceae	<i>Limosella longiflora</i>	LC	Indigenous
Asteraceae	<i>Lactuca serriola</i>		Not indigenous; Naturalised
Araliaceae	<i>Cussonia spicata</i>	LC	Indigenous
Boraginaceae	<i>Ehretia rigida</i>	LC	Indigenous
Rhamnaceae	<i>Ziziphus mucronata</i>	LC	Indigenous
Thymelaeaceae	<i>Lasiosiphon canoargenteus</i>	LC	Indigenous; Endemic
Poaceae	<i>Sporobolus discosporus</i>	LC	Indigenous
Hyacinthaceae	<i>Ledebouria burkei</i>	LC	Indigenous
Poaceae	<i>Eragrostis superba</i>	LC	Indigenous
Fabaceae	<i>Indigofera cryptantha</i>	LC	Indigenous
Asteraceae	<i>Senecio oxyriifolius</i>	LC	Indigenous
Ricciaceae	<i>Riccia atropurpurea</i>		Indigenous
Malvaceae	<i>Hibiscus calyphyllus</i>	LC	Indigenous
Poaceae	<i>Urochloa panicoides</i>	LC	Indigenous
Fabaceae	<i>Indigofera hiliaris</i>	LC	Indigenous
Malvaceae	<i>Triumfetta sonderi</i>	LC	Indigenous; Endemic
Thymelaeaceae	<i>Lasiosiphon capitatus</i>	LC	Indigenous
Pteridaceae	<i>Cheilanthes viridis</i>	LC	Indigenous
Amaranthaceae	<i>Guilleminea densa</i>		Not indigenous; Naturalised; Invasive
Campanulaceae	<i>Wahlenbergia denticulata</i>	LC	Indigenous; Endemic
Rubiaceae	<i>Vangueria pygmaea</i>	LC	Indigenous
Asteraceae	<i>Dicoma sp.</i>		
Asteraceae	<i>Senecio affinis</i>	LC	Indigenous
Apocynaceae	<i>Riocreuxia polyantha</i>	LC	Indigenous
Malvaceae	<i>Hermannia sp.</i>		
Proteaceae	<i>Protea caffra</i>	LC	Indigenous
Fabaceae	<i>Eriosema burkei</i>	LC	Indigenous
Rubiaceae	<i>Anthospermum hispidulum</i>	LC	Indigenous
Ophioglossaceae	<i>Ophioglossum polyphyllum</i>	LC	Indigenous
Verbenaceae	<i>Chascanum pinnatifidum</i>	LC	Indigenous
Poaceae	<i>Eragrostis tef</i>	NE	Not indigenous; Naturalised
Poaceae	<i>Elionurus muticus</i>	LC	Indigenous
Caryophyllaceae	<i>Silene burchellii</i>		Indigenous

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Dipsacaceae	<i>Cephalaria pungens</i>	LC	Indigenous
Asteraceae	<i>Senecio venosus</i>	LC	Indigenous
Orobanchaceae	<i>Striga elegans</i>	LC	Indigenous
Asteraceae	<i>Xanthium strumarium</i>		Not indigenous; Naturalised; Invasive
Limeaceae	<i>Limeum viscosum</i>	NE	Indigenous
Orchidaceae	<i>Habenaria mossii</i>	EN	Indigenous; Endemic
Asteraceae	<i>Helichrysum nudifolium</i>	LC	Indigenous
Hyacinthaceae	<i>Albuca virens</i>	LC	Indigenous
Euphorbiaceae	<i>Euphorbia sp.</i>		
Rhamnaceae	<i>Ziziphus zeyheriana</i>	LC	Indigenous
Malvaceae	<i>Sida dregei</i>	LC	Indigenous
Asteraceae	<i>Galinsoga parviflora</i>		Not indigenous; Naturalised; Invasive
Poaceae	<i>Aristida stipitata</i>	LC	Indigenous
Asteraceae	<i>Osteospermum scariosum</i>	NE	Indigenous
Asphodelaceae	<i>Trachyandra saltii</i>	LC	Indigenous
Celastraceae	<i>Gymnosporia buxifolia</i>	LC	Indigenous
Poaceae	<i>Digitaria tricholaenoides</i>	LC	Indigenous
Amaryllidaceae	<i>Scadoxus puniceus</i>	LC	Indigenous
Poaceae	<i>Panicum schinzii</i>	LC	Indigenous
Hyacinthaceae	<i>Ledebouria marginata</i>	LC	Indigenous
Poaceae	<i>Eragrostis obtusa</i>	LC	Indigenous
Asteraceae	<i>Osteospermum muricatum</i>	LC	Indigenous
Rubiaceae	<i>Pentanisia angustifolia</i>	LC	Indigenous
Oleaceae	<i>Olea europaea</i>		Indigenous
Gentianaceae	<i>Exochaenium grande</i>	LC	Indigenous
Rosaceae	<i>Agrimonia bracteata</i>	LC	Indigenous
Poaceae	<i>Cymbopogon caesius</i>	LC	Indigenous
Poaceae	<i>Eustachys paspaloides</i>	LC	Indigenous
Hypoxidaceae	<i>Hypoxis interjecta</i>	LC	Indigenous; Endemic
Asteraceae	<i>Erigeron bonariensis</i>		Not indigenous; Naturalised; Invasive
Poaceae	<i>Setaria pumila</i>	LC	Indigenous
Iridaceae	<i>Gladiolus elliotii</i>	LC	Indigenous
Poaceae	<i>Eragrostis sclerantha</i>	LC	Indigenous
Convolvulaceae	<i>Falkia oblonga</i>	LC	Indigenous
Malvaceae	<i>Sida chrysantha</i>	LC	Indigenous
Poaceae	<i>Eragrostis cilianensis</i>	LC	Indigenous
Ebenaceae	<i>Euclea crispa</i>	LC	Indigenous
Apocynaceae	<i>Asclepias sp.</i>		
Asteraceae	<i>Chrysocoma ciliata</i>	LC	Indigenous
Apocynaceae	<i>Pentarrhinum insipidum</i>	LC	Indigenous
Poaceae	<i>Setaria sphacelata</i>	LC	Indigenous
Poaceae	<i>Bromus catharticus</i>	NE	Not indigenous; Naturalised; Invasive
Poaceae	<i>Bewsia biflora</i>	LC	Indigenous
Poaceae	<i>Chloris virgata</i>	LC	Indigenous



## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Cucurbitaceae	<i>Cucumis hirsutus</i>	LC	Indigenous
Amaranthaceae	<i>Amaranthus deflexus</i>		Not indigenous; Naturalised
Acanthaceae	<i>Blepharis stainbankiae</i>	LC	Indigenous; Endemic
Fabaceae	<i>Indigostrum burkeanum</i>	LC	Indigenous
Lamiaceae	<i>Salvia sp.</i>		
Poaceae	<i>Digitaria velutina</i>	LC	Indigenous
Apocynaceae	<i>Asclepias eminens</i>	LC	Indigenous
Malvaceae	<i>Grewia occidentalis</i>	LC	Indigenous
Fabaceae	<i>Erythrina zeyheri</i>	LC	Indigenous
Rubiaceae	<i>Pygmaeothamnus zeyheri</i>	LC	Indigenous
Poaceae	<i>Paspalum dilatatum</i>	NE	Not indigenous; Naturalised; Invasive
Asteraceae	<i>Artemisia afra</i>	LC	Indigenous
Hyacinthaceae	<i>Dipcadi viride</i>	LC	Indigenous
Anacardiaceae	<i>Searsia discolor</i>	LC	Indigenous
Amaranthaceae	<i>Achyranthes aspera</i>		Not indigenous; Naturalised
Poaceae	<i>Brachiaria serrata</i>	LC	Indigenous
Asteraceae	<i>Xanthium spinosum</i>		Not indigenous; Naturalised; Invasive
Acanthaceae	<i>Barleria macrostegia</i>	LC	Indigenous
Cucurbitaceae	<i>Kedrostis africana</i>	LC	Indigenous
Poaceae	<i>Tristachya leucothrix</i>	LC	Indigenous
Cyperaceae	<i>Carex cognata</i>	LC	Indigenous
Lamiaceae	<i>Leonotis martinicensis</i>	LC	Indigenous
Caryophyllaceae	<i>Pollichia campestris</i>	LC	Indigenous
Poaceae	<i>Melinis nerviglumis</i>	LC	Indigenous
Amaranthaceae	<i>Cyphocarpa angustifolia</i>	LC	Indigenous
Anacardiaceae	<i>Searsia pyroides</i>	LC	Indigenous
Rubiaceae	<i>Kohautia amatymbica</i>	LC	Indigenous
Asteraceae	<i>Berkheya radula</i>	LC	Indigenous
Amaranthaceae	<i>Cyathula uncinulata</i>	LC	Indigenous
Poaceae	<i>Diheteropogon amplexans</i>	LC	Indigenous
Iridaceae	<i>Freesia grandiflora</i>	LC	Indigenous
Verbenaceae	<i>Verbena bonariensis</i>		Not indigenous; Naturalised; Invasive
Thymelaeaceae	<i>Lasiosiphon sericocephalus</i>	LC	Indigenous
Asteraceae	<i>Senecio inornatus</i>	LC	Indigenous
Poaceae	<i>Themeda triandra</i>	LC	Indigenous
Asteraceae	<i>Helichrysum paronychioides</i>	LC	Indigenous
Onagraceae	<i>Oenothera jamesii</i>		Not indigenous; Naturalised; Invasive
Apocynaceae	<i>Raphionacme velutina</i>	LC	Indigenous
Poaceae	<i>Sporobolus stapfianus</i>	LC	Indigenous
Asteraceae	<i>Cineraria albicans</i>	LC	Indigenous
Asteraceae	<i>Launaea rarifolia</i>	LC	Indigenous
Acanthaceae	<i>Blepharis innocua</i>	LC	Indigenous; Endemic
Polygonaceae	<i>Persicaria lapathifolia</i>		Not indigenous; Naturalised; Invasive
Amaryllidaceae	<i>Crinum sp.</i>		

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Malvaceae	<i>Hibiscus microcarpus</i>	LC	Indigenous
Poaceae	<i>Digitaria eriantha</i>	LC	Indigenous
Fabaceae	<i>Sphenostylis angustifolia</i>	LC	Indigenous
Asteraceae	<i>Brachylaena sp.</i>		
Geraniaceae	<i>Pelargonium luridum</i>	LC	Indigenous
Achariaceae	<i>Kiggelaria africana</i>	LC	Indigenous
Asteraceae	<i>Tragopogon dubius</i>		Not indigenous; Naturalised
Urticaceae	<i>Didymodoxa caffra</i>	LC	Indigenous
Lamiaceae	<i>Salvia stenophylla</i>		Indigenous
Polygalaceae	<i>Polygala uncinata</i>	LC	Indigenous
Cyperaceae	<i>Cyperus esculentus</i>	LC	Indigenous
Apiaceae	<i>Deverra burchellii</i>	LC	Indigenous
Cyperaceae	<i>Schoenoplectus tabernaemontani</i>		Not indigenous; Naturalised
Papaveraceae	<i>Papaver aculeatum</i>	LC	Indigenous
Asphodelaceae	<i>Aloe transvaalensis</i>		Indigenous
Asteraceae	<i>Helichrysum cerastioides</i>	LC	Indigenous
Fabaceae	<i>Melolobium microphyllum</i>	LC	Indigenous
Asparagaceae	<i>Asparagus laricinus</i>	LC	Indigenous
Asteraceae	<i>Helichrysum lepidissimum</i>	LC	Indigenous
Asteraceae	<i>Bidens pilosa</i>		Not indigenous; Naturalised
Commelinaceae	<i>Commelina africana</i>	LC	Indigenous
Asteraceae	<i>Helichrysum callicomum</i>	LC	Indigenous
Anacardiaceae	<i>Searsia magalismontana</i>	LC	Indigenous
Lamiaceae	<i>Ocimum obovatum</i>	NE	Indigenous
Malvaceae	<i>Pavonia burchellii</i>	LC	Indigenous
Asteraceae	<i>Polydora angustifolia</i>	LC	Indigenous
Anacampserotaceae	<i>Anacampseros subnuda</i>	LC	Indigenous
Poaceae	<i>Eragrostis plana</i>	LC	Indigenous
Poaceae	<i>Alloteropsis semialata</i>	LC	Indigenous
Anacardiaceae	<i>Searsia pyroides</i>	LC	Indigenous
Apocynaceae	<i>Orbea lutea</i>	LC	Indigenous
Poaceae	<i>Andropogon schirensis</i>	LC	Indigenous
Asteraceae	<i>Dicoma macrocephala</i>	LC	Indigenous
Asteraceae	<i>Seriphium plumosum</i>		Indigenous
Lamiaceae	<i>Plectranthus ramosior</i>	LC	Indigenous; Endemic
Poaceae	<i>Paspalum distichum</i>	LC	Not indigenous; Naturalised; Invasive
Apocynaceae	<i>Ceropegia chlorantha</i>		Indigenous
Cyperaceae	<i>Cyperus congestus</i>	LC	Indigenous
Papaveraceae	<i>Argemone ochroleuca</i>		Not indigenous; Naturalised; Invasive
Poaceae	<i>Trisetopsis imberbis</i>		Indigenous
Asteraceae	<i>Tolpis capensis</i>	LC	Indigenous
Poaceae	<i>Hyparrhenia anamesa</i>	LC	Indigenous
Meliaceae	<i>Melia azedarach</i>	NE	Not indigenous; Naturalised; Invasive
Malvaceae	<i>Hibiscus trionum</i>		Not indigenous; Naturalised

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Amaranthaceae	<i>Einadia nutans</i>		Not indigenous; Naturalised
Polygonaceae	<i>Oxygonum dregeanum</i>	NE	Indigenous
Poaceae	<i>Phragmites australis</i>	LC	Indigenous
Rubiaceae	<i>Kohautia caespitosa</i>	LC	Indigenous
Solanaceae	<i>Solanum lichtensteinii</i>	LC	Indigenous
Poaceae	<i>Aristida canescens</i>	LC	Indigenous
Rhamnaceae	<i>Helinus integrifolius</i>	LC	Indigenous
Plantaginaceae	<i>Plantago major</i>		Not indigenous; Naturalised
Malvaceae	<i>Grewia flava</i>	LC	Indigenous
Asteraceae	<i>Sonchus dregeanus</i>	LC	Indigenous
Cucurbitaceae	<i>Cucumis heptadactylus</i>	LC	Indigenous; Endemic
Fabaceae	<i>Eriosema cordatum</i>	LC	Indigenous
Poaceae	<i>Echinochloa colona</i>	LC	Indigenous
Lobeliaceae	<i>Lobelia sonderiana</i>	LC	Indigenous
Convolvulaceae	<i>Convolvulus multifidus</i>	LC	Indigenous; Endemic
Cyperaceae	<i>Cladium mariscus</i>	LC	Indigenous
Asteraceae	<i>Conyza podocephala</i>		Indigenous
Plantaginaceae	<i>Plantago lanceolata</i>	LC	Indigenous
Commelinaceae	<i>Commelina africana</i>	LC	Indigenous
Fabaceae	<i>Pearsonia uniflora</i>	LC	Indigenous
Talinaceae	<i>Talinum caffrum</i>	LC	Indigenous
Anacardiaceae	<i>Searsia rigida</i>	LC	Indigenous; Endemic
Agavaceae	<i>Chlorophytum transvaalense</i>	LC	Indigenous
Malvaceae	<i>Hermannia tomentosa</i>	LC	Indigenous
Crassulaceae	<i>Crassula lanceolata</i>	LC	Indigenous
Salicaceae	<i>Salix babylonica</i>		Not indigenous; Naturalised
Poaceae	<i>Chloris pycnothrix</i>	LC	Indigenous
Asteraceae	<i>Gazania krebsiana</i>	LC	Indigenous
Celastraceae	<i>Gymnosporia polyacantha</i>	LC	Indigenous; Endemic
Rubiaceae	<i>Richardia brasiliensis</i>	NE	Not indigenous; Naturalised
Cyperaceae	<i>Cyperus margaritaceus</i>	LC	Indigenous
Acanthaceae	<i>Crabbea angustifolia</i>	LC	Indigenous; Endemic
Malvaceae	<i>Hermannia cordata</i>	LC	Indigenous; Endemic
Agavaceae	<i>Chlorophytum cooperi</i>	LC	Indigenous
Apocynaceae	<i>Araujia sericifera</i>		Not indigenous; Naturalised; Invasive
Convolvulaceae	<i>Convolvulus sagittatus</i>	LC	Indigenous
Cyperaceae	<i>Coleochloa setifera</i>	LC	Indigenous
Apocynaceae	<i>Asclepias adscendens</i>	LC	Indigenous
Verbenaceae	<i>Lantana rugosa</i>	LC	Indigenous
Oleaceae	<i>Menodora africana</i>	LC	Indigenous
Campanulaceae	<i>Wahlenbergia undulata</i>	LC	Indigenous
Asteraceae	<i>Acanthospermum glabratum</i>		Not indigenous; Naturalised
Asteraceae	<i>Nidorella hottentotica</i>	LC	Indigenous
Amaranthaceae	<i>Achyranthes aspera</i>		Indigenous

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Apocynaceae	<i>Orthanthera jasminiflora</i>	LC	Indigenous
Scrophulariaceae	<i>Manulea paniculata</i>	LC	Indigenous
Apocynaceae	<i>Asclepias fallax</i>	LC	Indigenous; Endemic
Asteraceae	<i>Gerbera piloselloides</i>	LC	Indigenous
Phytolaccaceae	<i>Phytolacca octandra</i>		Not indigenous; Naturalised; Invasive
Acanthaceae	<i>Blepharis angusta</i>	LC	Indigenous; Endemic
Fabaceae	<i>Elephantorrhiza elephantina</i>	LC	Indigenous
Polygonaceae	<i>Polygonum aviculare</i>		Not indigenous; Naturalised
Brassicaceae	<i>Lepidium africanum</i>	LC	Indigenous
Fabaceae	<i>Vachellia karroo</i>	LC	Indigenous
Amaranthaceae	<i>Alternanthera pungens</i>		Not indigenous; Naturalised
Amaranthaceae	<i>Amaranthus thunbergii</i>	LC	Indigenous
Crassulaceae	<i>Crassula setulosa</i>	NE	Indigenous
Fabaceae	<i>Rhynchosia pedunculata</i>		Indigenous; Endemic
Acanthaceae	<i>Blepharis squarrosa</i>	LC	Indigenous; Endemic
Poaceae	<i>Eragrostis chloromelas</i>	LC	Indigenous
Boraginaceae	<i>Cynoglossum lanceolatum</i>	LC	Indigenous
Acanthaceae	<i>Justicia anagalloides</i>	LC	Indigenous
Commelinaceae	<i>Commelina livingstonii</i>	LC	Indigenous
Hyacinthaceae	<i>Eucomis autumnalis</i>	NE	Indigenous
Poaceae	<i>Andropogon eucomus</i>	LC	Indigenous
Amaranthaceae	<i>Gomphrena celosioides</i>		Not indigenous; Naturalised
Rubiaceae	<i>Kohautia cynanchica</i>	LC	Indigenous
Acanthaceae	<i>Crabbea acaulis</i>	LC	Indigenous
Lythraceae	<i>Ammannia involucrata</i>		Indigenous
Santalaceae	<i>Thesium resedoides</i>	LC	Indigenous
Limeaceae	<i>Limeum viscosum</i>	NE	Indigenous
Amaranthaceae	<i>Chenopodium album</i>		Not indigenous; Naturalised; Invasive
Orobanchaceae	<i>Graderia subintegra</i>	LC	Indigenous
Convolvulaceae	<i>Ipomoea crassipes</i>	LC	Indigenous
Fabaceae	<i>Chamaecrista comosa</i>	LC	Indigenous
Pteridaceae	<i>Pteris vittata</i>	LC	Indigenous
Asteraceae	<i>Nolletia rarifolia</i>	LC	Indigenous; Endemic
Hypoxidaceae	<i>Hypoxis acuminata</i>	LC	Indigenous
Asphodelaceae	<i>Bulbine abyssinica</i>	LC	Indigenous
Fabaceae	<i>Melilotus albus</i>	NE	Not indigenous; Naturalised; Invasive
Asteraceae	<i>Senecio hieracioides</i>	LC	Indigenous
Iridaceae	<i>Gladiolus papilio</i>	LC	Indigenous
Rutaceae	<i>Zanthoxylum capense</i>	LC	Indigenous
Verbenaceae	<i>Chascanum adenostachyum</i>	LC	Indigenous
Poaceae	<i>Brachiaria deflexa</i>	LC	Indigenous
Poaceae	<i>Eragrostis patentipilosa</i>	LC	Indigenous
Caryophyllaceae	<i>Dianthus mooiensis</i>	NE	Indigenous; Endemic
Poaceae	<i>Melinis repens</i>	LC	Indigenous

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Malvaceae	<i>Corchorus aspleniifolius</i>	LC	Indigenous
Phyllanthaceae	<i>Phyllanthus parvulus</i>	LC	Indigenous
Santalaceae	<i>Thesium multiramulosum</i>	LC	Indigenous
Cucurbitaceae	<i>Coccinia sessilifolia</i>	LC	Indigenous
Geraniaceae	<i>Monsonia burkeana</i>	LC	Indigenous
Asteraceae	<i>Senecio burchellii</i>	LC	Indigenous; Endemic
Poaceae	<i>Aristida aequiglumis</i>	LC	Indigenous
Poaceae	<i>Heteropogon contortus</i>	LC	Indigenous
Cyperaceae	<i>Bulbostylis oritrephes</i>	LC	Indigenous
Poaceae	<i>Eragrostis trichophora</i>	LC	Indigenous
Apocynaceae	<i>Cryptolepis oblongifolia</i>	LC	Indigenous
Asteraceae	<i>Hilliardiella elaeagnoides</i>		Indigenous
Poaceae	<i>Sporobolus congoensis</i>	LC	Indigenous
Poaceae	<i>Hyparrhenia dregeana</i>	LC	Indigenous
Apocynaceae	<i>Ceropegia circinata</i>		Indigenous
Asparagaceae	<i>Asparagus asparagoides</i>	LC	Indigenous
Asteraceae	<i>Dicoma anomala</i>	LC	Indigenous
Poaceae	<i>Enneapogon scoparius</i>	LC	Indigenous
Cucurbitaceae	<i>Acanthosicyos naudinianus</i>	LC	Indigenous
Poaceae	<i>Triraphis andropogonoides</i>	LC	Indigenous
Exorhizaceae	<i>Exorhiza pustulosa</i>		Indigenous
Cyperaceae	<i>Schoenoplectus brachyceras</i>	LC	Indigenous
Agavaceae	<i>Chlorophytum angulicaule</i>	LC	Indigenous
Poaceae	<i>Oropetium capense</i>	LC	Indigenous
Poaceae	<i>Aristida congesta</i>	LC	Indigenous
Apocynaceae	<i>Asclepias meyeriana</i>	LC	Indigenous
Iridaceae	<i>Moraea pallida</i>	LC	Indigenous
Asteraceae	<i>Cirsium vulgare</i>		Not indigenous; Naturalised; Invasive
Peraceae	<i>Clusia pulchella</i>	LC	Indigenous
Phrymaceae	<i>Mimulus gracilis</i>	LC	Indigenous
Scrophulariaceae	<i>Zaluzianskya elongata</i>	LC	Indigenous
Poaceae	<i>Eragrostis capensis</i>	LC	Indigenous
Euphorbiaceae	<i>Euphorbia clavarioides</i>	LC	Indigenous
Ebenaceae	<i>Diospyros lycioides</i>	LC	Indigenous
Phyllanthaceae	<i>Phyllanthus incurvus</i>	LC	Indigenous
Apocynaceae	<i>Pachycarpus schinzianus</i>	LC	Indigenous
Araceae	<i>Zantedeschia sp.</i>		
Fabaceae	<i>Leobordea hirsuta</i>	LC	Indigenous; Endemic
Asteraceae	<i>Cineraria aspera</i>	LC	Indigenous
Solanaceae	<i>Datura stramonium</i>		Not indigenous; Naturalised; Invasive
Asteraceae	<i>Senecio coronatus</i>	LC	Indigenous
Poaceae	<i>Lolium multiflorum</i>	NE	Not indigenous; Naturalised; Invasive
Asteraceae	<i>Ursinia nana</i>	LC	Indigenous
Rubiaceae	<i>Anthospermum rigidum</i>	LC	Indigenous

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Asteraceae	<i>Taraxacum brunneum</i>		Not indigenous; Naturalised
Poaceae	<i>Leersia hexandra</i>	LC	Indigenous
Poaceae	<i>Pogonarthria squarrosa</i>	LC	Indigenous
Ruscaceae	<i>Eriospermum cooperi</i>	LC	Indigenous
Pteridaceae	<i>Cheilanthes hirta</i>	LC	Indigenous
Fabaceae	<i>Indigofera hedyantha</i>	LC	Indigenous
Brassicaceae	<i>Sisymbrium turczaninowii</i>	LC	Indigenous
Fabaceae	<i>Pearsonia cajanifolia</i>	LC	Indigenous; Endemic
Poaceae	<i>Setaria sphacelata</i>	LC	Indigenous
Zygophyllaceae	<i>Tribulus terrestris</i>	LC	Indigenous
Amaranthaceae	<i>Amaranthus hybridus</i>		Not indigenous; Naturalised
Scrophulariaceae	<i>Selago densiflora</i>	LC	Indigenous
Araliaceae	<i>Cussonia paniculata</i>	LC	Indigenous
Alliaceae	<i>Tulbaghia sp.</i>		
Asteraceae	<i>Helichrysum setosum</i>	LC	Indigenous
Apocynaceae	<i>Aspidoglossum biflorum</i>	LC	Indigenous
Scrophulariaceae	<i>Chaenostoma leve</i>	LC	Indigenous
Aspleniaceae	<i>Asplenium aethiopicum</i>	LC	Indigenous
Boraginaceae	<i>Lappula heteracantha</i>		Not indigenous; Naturalised
Apocynaceae	<i>Ceropegia rehmannii</i>		Indigenous
Polygalaceae	<i>Polygala gracilentia</i>	LC	Indigenous
Fabaceae	<i>Lotononis laxa</i>	LC	Indigenous
Asteraceae	<i>Bidens bipinnata</i>		Not indigenous; Naturalised
Asteraceae	<i>Cosmos bipinnatus</i>		Not indigenous; Naturalised
Polygalaceae	<i>Polygala hottentotta</i>	LC	Indigenous
Poaceae	<i>Monocymbium ceresiforme</i>	LC	Indigenous
Pittosporaceae	<i>Pittosporum viridiflorum</i>	LC	Indigenous
Lamiaceae	<i>Teucrium trifidum</i>	LC	Indigenous
Iridaceae	<i>Tritonia nelsonii</i>	LC	Indigenous
Poaceae	<i>Perotis patens</i>	LC	Indigenous
Fabaceae	<i>Vigna vexillata</i>	LC	Indigenous
Amaryllidaceae	<i>Ammocharis coranica</i>	LC	Indigenous
Amaranthaceae	<i>Dysphania ambrosioides</i>		Not indigenous; Naturalised; Invasive
Cyperaceae	<i>Afroscleroides dioeca</i>		Indigenous
Anacardiaceae	<i>Searsia rigida</i>	LC	Indigenous; Endemic
Scrophulariaceae	<i>Jamesbrittenia atropurpurea</i>	LC	Indigenous
Poaceae	<i>Paspalum sp.</i>		
Cyperaceae	<i>Cyperus capensis</i>	LC	Indigenous; Endemic
Fabaceae	<i>Tephrosia semiglabra</i>	LC	Indigenous
Poaceae	<i>Agrostis lachnantha</i>	LC	Indigenous
Apocynaceae	<i>Ceropegia rendallii</i>	LC	Indigenous
Solanaceae	<i>Solanum nigrum</i>		Not indigenous; Naturalised
Lamiaceae	<i>Rotheca hirsuta</i>	LC	Indigenous
Cucurbitaceae	<i>Momordica balsamina</i>	LC	Indigenous

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Fabaceae	<i>Tephrosia elongata</i>	LC	Indigenous
Lamiaceae	<i>Acrotome hispida</i>	LC	Indigenous
Convolvulaceae	<i>Ipomoea oblongata</i>	LC	Indigenous
Typhaceae	<i>Typha capensis</i>	LC	Indigenous
Asteraceae	<i>Denekia capensis</i>	LC	Indigenous
Asteraceae	<i>Nidorella resedifolia</i>	LC	Indigenous
Cyperaceae	<i>Scirpoides burkei</i>	LC	Indigenous
Hyacinthaceae	<i>Ledebouria cooperi</i>	LC	Indigenous
Brassicaceae	<i>Diplotaxis muralis</i>		Not indigenous; Naturalised; Invasive
Asteraceae	<i>Tarchonanthus camphoratus</i>	LC	Indigenous
Cyperaceae	<i>Abildgaardia ovata</i>	LC	Indigenous
Hypoxidaceae	<i>Hypoxis argentea</i>	LC	Indigenous
Fabaceae	<i>Tylosema esculentum</i>	LC	Indigenous
Euphorbiaceae	<i>Euphorbia spartaria</i>	LC	Indigenous
Hypoxidaceae	<i>Hypoxis iridifolia</i>	LC	Indigenous
Commelinaceae	<i>Commelina benghalensis</i>	LC	Indigenous
Aizoaceae	<i>Delosperma sp.</i>		
Asphodelaceae	<i>Bulbine narcissifolia</i>	LC	Indigenous
Aspleniaceae	<i>Asplenium cordatum</i>	LC	Indigenous
Santalaceae	<i>Thesium utile</i>	LC	Indigenous
Brassicaceae	<i>Raphanus raphanistrum</i>		Not indigenous; Naturalised; Invasive
Asteraceae	<i>Erigeron canadensis</i>		Not indigenous; Naturalised; Invasive
Asteraceae	<i>Symphotrichum squamatum</i>		Not indigenous; Naturalised; Invasive
Euphorbiaceae	<i>Euphorbia striata</i>	LC	Indigenous
Asphodelaceae	<i>Aloe subspicata</i>		Indigenous
Amaranthaceae	<i>Dysphania multifida</i>		Not indigenous; Naturalised; Invasive
Fabaceae	<i>Indigofera confusa</i>	LC	Indigenous
Poaceae	<i>Eragrostis curvula</i>	LC	Indigenous
Poaceae	<i>Eleusine coracana</i>	LC	Indigenous
Fabaceae	<i>Dichilus gracilis</i>	LC	Indigenous
Fabaceae	<i>Vigna unguiculata</i>	LC	Indigenous
Cyperaceae	<i>Cyperus semitrifidus</i>	LC	Indigenous
Asteraceae	<i>Berkheya zeyheri</i>	LC	Indigenous
Poaceae	<i>Ehrharta erecta</i>	LC	Indigenous
Fabaceae	<i>Chamaecrista biensis</i>	LC	Indigenous
Asteraceae	<i>Zinnia peruviana</i>		Not indigenous; Naturalised; Invasive
Dipsacaceae	<i>Scabiosa columbaria</i>	LC	Indigenous
Commelinaceae	<i>Cyanotis speciosa</i>	LC	Indigenous
Hypoxidaceae	<i>Hypoxis rigidula</i>	LC	Indigenous
Boraginaceae	<i>Lithospermum cinereum</i>	LC	Indigenous
Cannabaceae	<i>Celtis africana</i>	LC	Indigenous
Cucurbitaceae	<i>Cucumis zeyheri</i>	LC	Indigenous
Crassulaceae	<i>Crassula capitella</i>	LC	Indigenous
Poaceae	<i>Aristida diffusa</i>	LC	Indigenous

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Scrophulariaceae	<i>Buddleja saligna</i>	LC	Indigenous
Asteraceae	<i>Pseudognaphalium oligandrum</i>	LC	Indigenous
Menispermaceae	<i>Antizoma angustifolia</i>	LC	Indigenous
Amaryllidaceae	<i>Boophone disticha</i>	LC	Indigenous
Verbenaceae	<i>Lippia scaberrima</i>	LC	Indigenous
Poaceae	<i>Microchloa caffra</i>	LC	Indigenous
Poaceae	<i>Loudetia simplex</i>	LC	Indigenous
Ricciaceae	<i>Riccia albolimbata</i>		Indigenous
Cyperaceae	<i>Cyperus longus</i>	NE	Indigenous
Poaceae	<i>Tristachya rehmannii</i>	LC	Indigenous
Euphorbiaceae	<i>Acalypha angustata</i>	LC	Indigenous
Fabaceae	<i>Tephrosia capensis</i>	LC	Indigenous
Malvaceae	<i>Hermannia lancifolia</i>	LC	Indigenous; Endemic
Convolvulaceae	<i>Ipomoea purpurea</i>		Not indigenous; Naturalised; Invasive
Poaceae	<i>Tragus berteronianus</i>	LC	Indigenous
Poaceae	<i>Trichoneura grandiglumis</i>	LC	Indigenous
Fabaceae	<i>Medicago sativa</i>	NE	Not indigenous; Cultivated; Naturalised; Invasive
Amaryllidaceae	<i>Nerine laticoma</i>	LC	Indigenous
Solanaceae	<i>Solanum pseudocapsicum</i>		Not indigenous; Naturalised; Invasive
Asteraceae	<i>Pseudopegolettia tenella</i>		Indigenous
Poaceae	<i>Eragrostis racemosa</i>	LC	Indigenous
Ranunculaceae	<i>Clematis brachiata</i>	LC	Indigenous
Amaryllidaceae	<i>Haemanthus montanus</i>	LC	Indigenous
Fabaceae	<i>Dichilus lebeckioides</i>	LC	Indigenous
Anacampserotaceae	<i>Anacampseros filamentosa</i>		Indigenous; Endemic
Asteraceae	<i>Schkuhria pinnata</i>		Not indigenous; Naturalised
Poaceae	<i>Setaria nigrirostris</i>	LC	Indigenous
Asteraceae	<i>Lopholaena coriifolia</i>	LC	Indigenous
Convolvulaceae	<i>Ipomoea ommanneyi</i>	LC	Indigenous
Solanaceae	<i>Withania somnifera</i>	LC	Indigenous
Cyperaceae	<i>Carex glomerabilis</i>	LC	Indigenous
Fabaceae	<i>Leobordea divaricata</i>	LC	Indigenous
Poaceae	<i>Setaria verticillata</i>	LC	Indigenous
Iridaceae	<i>Gladiolus crassifolius</i>	LC	Indigenous
Polygonaceae	<i>Rumex crispus</i>		Not indigenous; Naturalised; Invasive
Chrysobalanaceae	<i>Parinari capensis</i>	LC	Indigenous
Asteraceae	<i>Tagetes minuta</i>		Not indigenous; Naturalised; Invasive
Asteraceae	<i>Helichrysum caespitium</i>	LC	Indigenous
Poaceae	<i>Eragrostis gummiflua</i>	LC	Indigenous
Myrothamnaceae	<i>Myrothamnus flabellifolius</i>	DD	Indigenous
Asteraceae	<i>Senecio erubescens</i>	NE	Indigenous
Poaceae	<i>Schizachyrium sanguineum</i>	LC	Indigenous
Agavaceae	<i>Chlorophytum trichophlebium</i>	LC	Indigenous; Endemic
Apocynaceae	<i>Aspidoglossum glabrescens</i>	LC	Indigenous; Endemic



## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Malvaceae	<i>Malva parviflora</i>		Not indigenous; Naturalised
Brassicaceae	<i>Erucastrum austroafricanum</i>	LC	Indigenous
Poaceae	<i>Panicum natalense</i>	LC	Indigenous
Solanaceae	<i>Solanum sisymbriifolium</i>		Not indigenous; Naturalised; Invasive
Orchidaceae	<i>Bonatea antennifera</i>	LC	Indigenous
Apiaceae	<i>Heteromorpha arborescens</i>	LC	Indigenous
Lamiaceae	<i>Syncolostemon canescens</i>	LC	Indigenous
Asteraceae	<i>Dicoma anomala</i>	LC	Indigenous
Fabaceae	<i>Indigofera oxytropis</i>	LC	Indigenous
Juncaceae	<i>Juncus exsertus</i>	LC	Indigenous
Anacardiaceae	<i>Searsia pyroides</i>	LC	Indigenous
Portulacaceae	<i>Portulaca quadrifida</i>	LC	Indigenous
Lobeliaceae	<i>Cyphia persicifolia</i>	LC	Indigenous; Endemic
Poaceae	<i>Trachypogon spicatus</i>	LC	Indigenous
Acanthaceae	<i>Dyschoriste costata</i>	LC	Indigenous; Endemic
Cleomaceae	<i>Cleome maculata</i>	LC	Indigenous
Onagraceae	<i>Oenothera rosea</i>		Not indigenous; Naturalised; Invasive
Rubiaceae	<i>Rubia horrida</i>	LC	Indigenous
Dipsacaceae	<i>Cephalaria zeyheriana</i>	LC	Indigenous
Amaranthaceae	<i>Aerva leucura</i>	LC	Indigenous
Oxalidaceae	<i>Oxalis corniculata</i>		Not indigenous; Naturalised; Invasive
Malvaceae	<i>Hermannia depressa</i>	LC	Indigenous
Apocynaceae	<i>Asclepias fulva</i>	LC	Indigenous
Poaceae	<i>Eragrostis biflora</i>	LC	Indigenous
Poaceae	<i>Panicum repens</i>	LC	Indigenous
Asteraceae	<i>Sonchus oleraceus</i>		Not indigenous; Naturalised; Invasive
Poaceae	<i>Cynodon hirsutus</i>	LC	Indigenous
Iridaceae	<i>Gladiolus antholyzoides</i>	LC	Indigenous; Endemic
Apocynaceae	<i>Asclepias brevipes</i>	LC	Indigenous; Endemic
Agavaceae	<i>Chlorophytum bowkeri</i>	LC	Indigenous
Fabaceae	<i>Zornia milneana</i>	LC	Indigenous
Santalaceae	<i>Thesium magalismsontanum</i>	LC	Indigenous
Santalaceae	<i>Thesium transvaalense</i>	LC	Indigenous; Endemic
Scrophulariaceae	<i>Nemesia fruticans</i>	LC	Indigenous
Asteraceae	<i>Helichrysum nudifolium</i>	LC	Indigenous
Poaceae	<i>Cynodon dactylon</i>	LC	Indigenous
Asteraceae	<i>Helichrysum rugulosum</i>	LC	Indigenous
Apiaceae	<i>Berula repanda</i>	LC	Indigenous
Cyperaceae	<i>Bulbostylis burchellii</i>	LC	Indigenous
Asphodelaceae	<i>Bulbine capitata</i>	LC	Indigenous
Asteraceae	<i>Helichrysum chionosphaerum</i>	LC	Indigenous
Pteridaceae	<i>Pellaea calomelanos</i>	LC	Indigenous
Asteraceae	<i>Helichrysum dregeanum</i>	LC	Indigenous
Amaranthaceae	<i>Dysphania carinata</i>		Not indigenous; Naturalised; Invasive

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Family	Species	IUCN	Ecology
Thymelaeaceae	<i>Lasiosiphon kraussianus</i>		Indigenous
Orobanchaceae	<i>Striga asiatica</i>	LC	Indigenous
Asteraceae	<i>Geigeria burkei</i>	NE	Indigenous
Geraniaceae	<i>Monsonia angustifolia</i>	LC	Indigenous
Fabaceae	<i>Leobordea mucronata</i>		Indigenous
Verbenaceae	<i>Priva meyeri</i>	LC	Indigenous
Hyacinthaceae	<i>Ledebouria luteola</i>	LC	Indigenous
Onagraceae	<i>Oenothera tetraptera</i>		Not indigenous; Naturalised; Invasive
Solanaceae	<i>Datura ferox</i>		Not indigenous; Naturalised; Invasive
Euphorbiaceae	<i>Acalypha caperonioides</i>	DD	Indigenous

**APPENDIX B. PLANT SPECIES FOUND ON SITE**

<b>Woody species</b>
<i>Eucalyptus camaldulensis</i>
<i>Grewia flava</i>
<i>Vachellia karroo</i>
<i>Ziziphus mucronata</i>
<i>Diospyros lycioides</i>
<i>Searsia lancea</i>
<i>Searsia pyroides</i>
<b>Grass species</b>
<i>Aristida congesta</i>
<i>Aristida junciformes</i>
<i>Brachiaria nigropedata</i>
<i>Brachiaria serata</i>
<i>Cymbopogon pospischilli</i>
<i>Cynodon dactylon</i>
<i>Dichanthium annulatum</i>
<i>Digitaria eriantha</i>
<i>Diplachne fusca</i>
<i>Eragrostis bicolor</i>
<i>Eragrostis biflora</i>
<i>Eragrostis curvula</i>
<i>Eragrostis gummiflua</i>
<i>Eragrostis lehmanniana</i>
<i>Eragrostis plana</i>
<i>Heteropogon contortus</i>
<i>Hyparrhenia hirta</i>
<i>Hyparrhenia tamba</i>
<i>Melinis repens</i>
<i>Panicum natalense</i>
<i>Eragrostis racemosa</i>
<i>Setaria sphacelata</i>
<i>Sporobolus africanus</i>
<i>Themeda triandra</i>
<i>Trachypogon spicatus</i>
<i>Trichoneura grandiglumis</i>
<i>Triraphis andropogonooides</i>
<i>Urochloa mosambicensis</i>
<i>Urochloa panicoides</i>
<b>Dwarf shrubs, Forbs, geophytes &amp; succulents</b>
<i>Acalypha angusta</i>
<i>Altenanthera pungens</i>

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

<i>Amaranthus spinosa</i>
<i>Anthospermum rigidum</i>
<i>Argemone ochroleuca</i>
<i>Asparagus larycinus</i>
<i>Asparagus suaveolens</i>
<i>Athrixia elata</i>
<i>Barleria macrostegia</i>
<i>Berkheya onopordifolia</i>
<i>Berkheya purpurea</i>
<i>Berkheya rigida</i>
<i>Berkheya speciosa</i>
<i>Bidens bipinnata</i>
<i>Bidens pilosa</i>
<i>Boophane disticha</i>
<i>Chamaecrista mimosoides</i>
<i>Chamaesyce inaequilatera</i>
<i>Clematis brachiata</i>
<i>Conyza albida</i>
<i>Conyza bonariensis</i>
<i>Crabbea angustifolia</i>
<i>Cyperus obtusiflorus</i>
<i>Cyperus sexangularis</i>
<i>Datura stramonium</i>
<i>Dianthus mooiensis</i>
<i>Dicoma anomala</i>
<i>Felicia muricata</i>
<i>Helichrysum caespitium</i>
<i>Helichrysum miconiifolium</i>
<i>Helichrysum nudifolium</i>
<i>Hermbstaedtia linearis</i>
<i>Hypoxis rigidula</i>
<i>Indigofera comosa</i>
<i>Indigofera daleioides</i>
<i>Ipomoea ommaneyi</i>
<i>Kyling alba</i>
<i>Kyphocarpa angustifolia</i>
<i>Nidorella anomala</i>
<i>Opuntia ficus indica</i>
<i>Oxalis spp.</i>
<i>Pentzia incana</i>
<i>Persicaria serrulata</i>
<i>Pygmaeothamnus zeyheri</i>
<i>Senecio coronatus</i>
<i>Senecio inornatus</i>

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

<i>Solanum incanum</i>
<i>Stoebe vulgaris</i>
<i>Tagetes minuta</i>
<i>Tylosema esculentum</i>
<i>Tylosema fassoglense</i>
<i>Typha capensis</i>
<i>Vernonia oligocephala</i>
<i>Wahlenbergia caledonica</i>
<i>Xanthium strumarium</i>
<i>Zinnia peruviana</i>
<i>Ziziphus zeyheriana.</i>

### APPENDIX C. BIRD SPECIES LIST FOR QDS

Common_group	Common_species	Genus	Species
	Bokmakierie	<i>Telophorus</i>	<i>zeylonus</i>
	Neddicky	<i>Cisticola</i>	<i>fulvicapilla</i>
	Quailfinch	<i>Ortygospiza</i>	<i>atricollis</i>
	Ruff	<i>Calidris</i>	<i>pugnax</i>
	Secretarybird	<i>Sagittarius</i>	<i>serpentarius</i>
Barbet	Acacia Pied	<i>Tricholaema</i>	<i>leucomelas</i>
Barbet	Crested	<i>Trachyphonus</i>	<i>vallantii</i>
Bee-eater	Blue-cheeked	<i>Merops</i>	<i>persicus</i>
Bee-eater	European	<i>Merops</i>	<i>apiaster</i>
Bee-eater	Little	<i>Merops</i>	<i>pusillus</i>
Bishop	Southern Red	<i>Euplectes</i>	<i>orix</i>
Bishop	Yellow-crowned	<i>Euplectes</i>	<i>afer</i>
Bittern	Little	<i>Ixobrychus</i>	<i>minutus</i>
Bulbul	African Red-eyed	<i>Pycnonotus</i>	<i>nigricans</i>
Bulbul	Dark-capped	<i>Pycnonotus</i>	<i>tricolor</i>
Buzzard	Common	<i>Buteo</i>	<i>buteo</i>
Buzzard	Jackal	<i>Buteo</i>	<i>rufofuscus</i>
Canary	Black-throated	<i>Crithagra</i>	<i>atrogularis</i>
Canary	Yellow	<i>Crithagra</i>	<i>flaviventris</i>
Canary	Yellow-fronted	<i>Crithagra</i>	<i>mozambica</i>
Chat	Ant-eating	<i>Myrmecocichla</i>	<i>formicivora</i>
Cisticola	Cloud	<i>Cisticola</i>	<i>textrix</i>
Cisticola	Desert	<i>Cisticola</i>	<i>aridulus</i>
Cisticola	Levaillant's	<i>Cisticola</i>	<i>tinniens</i>
Cisticola	Wing-snapping	<i>Cisticola</i>	<i>ayresii</i>
Cisticola	Zitting	<i>Cisticola</i>	<i>juncidis</i>
Coot	Red-knobbed	<i>Fulica</i>	<i>cristata</i>
Cormorant	Reed	<i>Microcarbo</i>	<i>africanus</i>
Cormorant	White-breasted	<i>Phalacrocorax</i>	<i>lucidus</i>
Crake	Black	<i>Zapornia</i>	<i>flavirostra</i>
Crow	Pied	<i>Corvus</i>	<i>albus</i>

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Common_group	Common_species	Genus	Species
Cuckoo	Diederik	<i>Chrysococcyx</i>	<i>caprius</i>
Darter	African	<i>Anhinga</i>	<i>rufa</i>
Dove	Cape Turtle	<i>Streptopelia</i>	<i>capicola</i>
Dove	Laughing	<i>Spilopelia</i>	<i>senegalensis</i>
Dove	Namaqua	<i>Oena</i>	<i>capensis</i>
Dove	Red-eyed	<i>Streptopelia</i>	<i>semitorquata</i>
Duck	African Black	<i>Anas</i>	<i>sparsa</i>
Duck	White-faced Whistling	<i>Dendrocygna</i>	<i>viduata</i>
Duck	Yellow-billed	<i>Anas</i>	<i>undulata</i>
Eagle	African Fish	<i>Haliaeetus</i>	<i>vocifer</i>
Eagle	Black-chested Snake	<i>Circaetus</i>	<i>pectoralis</i>
Eagle	Brown Snake	<i>Circaetus</i>	<i>cinereus</i>
Egret	Great	<i>Ardea</i>	<i>alba</i>
Egret	Little	<i>Egretta</i>	<i>garzetta</i>
Egret	Western Cattle	<i>Bubulcus</i>	<i>ibis</i>
Falcon	Peregrine	<i>Falco</i>	<i>peregrinus</i>
Fiscal	Southern	<i>Lanius</i>	<i>collaris</i>
Flamingo	Greater	<i>Phoenicopterus</i>	<i>roseus</i>
Flycatcher	Fiscal	<i>Melaenornis</i>	<i>silens</i>
Francolin	Coqui	<i>Peliperdix</i>	<i>coqui</i>
Francolin	Orange River	<i>Scleroptila</i>	<i>gutturalis</i>
Goose	Egyptian	<i>Alopochen</i>	<i>aegyptiaca</i>
Goose	Spur-winged	<i>Plectropterus</i>	<i>gambensis</i>
Goshawk	Pale Chanting	<i>Melierax</i>	<i>canorus</i>
Grassbird	Cape	<i>Sphenoecacus</i>	<i>afer</i>
Grebe	Great Crested	<i>Podiceps</i>	<i>cristatus</i>
Grebe	Little	<i>Tachybaptus</i>	<i>ruficollis</i>
Greenshank	Common	<i>Tringa</i>	<i>nebularia</i>
Guineafowl	Helmeted	<i>Numida</i>	<i>meleagris</i>
Gull	Grey-headed	<i>Chroicocephalus</i>	<i>cirrocephalus</i>
Harrier	African Marsh	<i>Circus</i>	<i>ranivorus</i>
Harrier	Pallid	<i>Circus</i>	<i>macrourus</i>
Heron	Black	<i>Egretta</i>	<i>ardesiaca</i>
Heron	Black-crowned Night	<i>Nycticorax</i>	<i>nycticorax</i>
Heron	Black-headed	<i>Ardea</i>	<i>melanocephala</i>
Heron	Grey	<i>Ardea</i>	<i>cinerea</i>
Heron	Purple	<i>Ardea</i>	<i>purpurea</i>
Heron	Squacco	<i>Ardeola</i>	<i>ralloides</i>
Hoopoe	African	<i>Upupa</i>	<i>africana</i>
Ibis	African Sacred	<i>Threskiornis</i>	<i>aethiopicus</i>
Ibis	Glossy	<i>Plegadis</i>	<i>falcinellus</i>
Ibis	Hadada	<i>Bostrychia</i>	<i>hagedash</i>
Kestrel	Greater	<i>Falco</i>	<i>rupicoloides</i>
Kingfisher	Pied	<i>Ceryle</i>	<i>rudis</i>

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Common_group	Common_species	Genus	Species
Kite	Black-winged	<i>Elanus</i>	<i>caeruleus</i>
Korhaan	Northern Black	<i>Afrotis</i>	<i>afraoides</i>
Lapwing	African Wattled	<i>Vanellus</i>	<i>senegallus</i>
Lapwing	Blacksmith	<i>Vanellus</i>	<i>armatus</i>
Lapwing	Crowned	<i>Vanellus</i>	<i>coronatus</i>
Lark	Eastern Clapper	<i>Mirafra</i>	<i>fasciolata</i>
Lark	Melodious	<i>Mirafra</i>	<i>cheniana</i>
Lark	Red-capped	<i>Calandrella</i>	<i>cinerea</i>
Lark	Rufous-naped	<i>Mirafra</i>	<i>africana</i>
Lark	Sabota	<i>Calendulauda</i>	<i>sabota</i>
Lark	Spike-heeled	<i>Chersomanes</i>	<i>albofasciata</i>
Longclaw	Cape	<i>Macronyx</i>	<i>capensis</i>
Martin	Banded	<i>Riparia</i>	<i>cincta</i>
Martin	Brown-throated	<i>Riparia</i>	<i>paludicola</i>
Moorhen	Common	<i>Gallinula</i>	<i>chloropus</i>
Mousebird	Red-faced	<i>Urocolius</i>	<i>indicus</i>
Mousebird	White-backed	<i>Colius</i>	<i>colius</i>
Myna	Common	<i>Acridotheres</i>	<i>tristis</i>
Ostrich	Common	<i>Struthio</i>	<i>camelus</i>
Owl	Marsh	<i>Asio</i>	<i>capensis</i>
Pigeon	Speckled	<i>Columba</i>	<i>guinea</i>
Pipit	African	<i>Anthus</i>	<i>cinnamomeus</i>
Pipit	Buffy	<i>Anthus</i>	<i>vaalensis</i>
Plover	Three-banded	<i>Charadrius</i>	<i>tricoloris</i>
Pratincole	Black-winged	<i>Glareola</i>	<i>nordmanni</i>
Prinia	Black-chested	<i>Prinia</i>	<i>flavicans</i>
Quelea	Red-billed	<i>Quelea</i>	<i>quelea</i>
Rail	African	<i>Rallus</i>	<i>caerulescens</i>
Robin-Chat	Cape	<i>Cossypha</i>	<i>caffra</i>
Sandpiper	Wood	<i>Tringa</i>	<i>glareola</i>
Scrub Robin	Kalahari	<i>Cercotrichas</i>	<i>paena</i>
Shelduck	South African	<i>Tadorna</i>	<i>cana</i>
Shrike	Lesser Grey	<i>Lanius</i>	<i>minor</i>
Shrike	Red-backed	<i>Lanius</i>	<i>collurio</i>
Sparrow	Cape	<i>Passer</i>	<i>melanurus</i>
Sparrow	House	<i>Passer</i>	<i>domesticus</i>
Sparrow	Southern Grey-headed	<i>Passer</i>	<i>diffusus</i>
Sparrow-Weaver	White-browed	<i>Plocepasser</i>	<i>mahali</i>
Spoonbill	African	<i>Platalea</i>	<i>alba</i>
Spurfowl	Swainson's	<i>Pternistis</i>	<i>swainsonii</i>
Starling	Cape	<i>Lamprotornis</i>	<i>nitens</i>
Starling	Pied	<i>Lamprotornis</i>	<i>bicolor</i>
Starling	Wattled	<i>Creatophora</i>	<i>cinerea</i>
Stilt	Black-winged	<i>Himantopus</i>	<i>himantopus</i>

## Terrestrial Biodiversity Impact Assessment Mopane Solar PV 5

Common_group	Common_species	Genus	Species
Stonechat	African	<i>Saxicola</i>	<i>torquatus</i>
Swallow	Barn	<i>Hirundo</i>	<i>rustica</i>
Swallow	Greater Striped	<i>Cecropis</i>	<i>cucullata</i>
Swallow	Lesser Striped	<i>Cecropis</i>	<i>abyssinica</i>
Swallow	South African Cliff	<i>Petrochelidon</i>	<i>spilodera</i>
Swallow	White-throated	<i>Hirundo</i>	<i>albigularis</i>
Swamphen	African	<i>Porphyrio</i>	<i>madagascariensis</i>
Swift	African Palm	<i>Cypsiurus</i>	<i>parvus</i>
Swift	Little	<i>Apus</i>	<i>affinis</i>
Swift	White-rumped	<i>Apus</i>	<i>caffer</i>
Teal	Blue-billed	<i>Spatula</i>	<i>hottentota</i>
Teal	Red-billed	<i>Anas</i>	<i>erythrorhyncha</i>
Tern	Whiskered	<i>Chlidonias</i>	<i>hybrida</i>
Thick-knee	Spotted	<i>Burhinus</i>	<i>capensis</i>
Vulture	Cape	<i>Gyps</i>	<i>coprotheres</i>
Wagtail	African Pied	<i>Motacilla</i>	<i>aguimp</i>
Wagtail	Cape	<i>Motacilla</i>	<i>capensis</i>
Warbler	African Reed	<i>Acrocephalus</i>	<i>baeticatus</i>
Warbler	Chestnut-vented	<i>Curruca</i>	<i>subcoerulea</i>
Warbler	Lesser Swamp	<i>Acrocephalus</i>	<i>gracilirostris</i>
Warbler	Little Rush	<i>Bradypterus</i>	<i>baboecala</i>
Waxbill	Common	<i>Estrilda</i>	<i>astrild</i>
Weaver	Scaly-feathered	<i>Sporopipes</i>	<i>squamifrons</i>
Weaver	Southern Masked	<i>Ploceus</i>	<i>velatus</i>
Weaver	Thick-billed	<i>Amblyospiza</i>	<i>albifrons</i>
Wheatear	Capped	<i>Oenanthe</i>	<i>pileata</i>
Wheatear	Mountain	<i>Myrmecocichla</i>	<i>monticola</i>
White-eye	Cape	<i>Zosterops</i>	<i>virens</i>
Whydah	Pin-tailed	<i>Vidua</i>	<i>macroura</i>
Widowbird	Long-tailed	<i>Euplectes</i>	<i>progne</i>
Widowbird	Red-collared	<i>Euplectes</i>	<i>ardens</i>
Widowbird	White-winged	<i>Euplectes</i>	<i>albonotatus</i>
Wryneck	Red-throated	<i>Jynx</i>	<i>ruficollis</i>



APPENDIX D MAMMAL SPECIES LIST

Family	Scientific name	Common name	Red list
Bovidae	<i>Antidorcas marsupialis</i>	Springbok	Least Concern (2016)
Bovidae	<i>Connochaetes gnou</i>	Black Wildebeest	Least Concern (2016)
Bovidae	<i>Connochaetes taurinus</i>	Blue Wildebeest	Least Concern (ver 3.1, 2017)
Bovidae	<i>Damaliscus pygargus phillipsi</i>	Blesbok	Least Concern (2016)
Bovidae	<i>Sylvicapra grimmia</i>	Bush Duiker	Least Concern (2016)
Bovidae	<i>Tragelaphus strepsiceros</i>	Greater Kudu	Least Concern (2016)
Canidae	<i>Canis mesomelas</i>	Black-backed Jackal	Least Concern (2016)
Cercopithecidae	<i>Chlorocebus pygerythrus</i>	Vervet Monkey	Least Concern (2016)
Felidae	<i>Leptailurus serval</i>	Serval	Near Threatened (2016)
Herpestidae	<i>Atilax paludinosus</i>	Marsh Mongoose	Least Concern (2016)
Herpestidae	<i>Cynictis penicillata</i>	Yellow Mongoose	Least Concern (2016)
Hyaenidae	<i>Proteles cristata</i>	Aardwolf	Least Concern (2016)
Hystricidae	<i>Hystrix africaeaustralis</i>	Cape Porcupine	Least Concern
Leporidae	<i>Lepus capensis</i>	Cape Hare	Least Concern
Leporidae	<i>Lepus saxatilis</i>	Scrub Hare	Least Concern
Macroscelididae	<i>Elephantulus myurus</i>	Eastern Rock Elephant Shrew	Least Concern (2016)
Muridae	<i>Aethomys ineptus</i>	Tete Veld Aethomys	Least Concern (2016)
Muridae	<i>Gerbilliscus leucogaster</i>	Bushveld Gerbil	Least Concern (2016)
Mustelidae	<i>Aonyx capensis</i>	African Clawless Otter	Near Threatened (2016)
Rhinolophidae	<i>Rhinolophus clivosus</i>	Geoffroy's Horseshoe Bat	Least Concern (2016)
Sciuridae	<i>Paraxerus cepapi</i>	Smith's Bush Squirrel	Least Concern (2016)
Sciuridae	<i>Xerus inauris</i>	South African Ground Squirrel	Least Concern
Suidae	<i>Phacochoerus africanus</i>	Common Warthog	Least Concern (2016)
Viveridae	<i>Genetta maculata</i>	Common Large-spotted Genet	Least Concern
Viverridae	<i>Genetta genetta</i>	Common Genet	Least Concern (2016)
Viverridae	<i>Genetta tigrina</i>	Cape Genet (Cape Large-spotted Genet)	Least Concern (2016)

## APPENDIX E HERPETOFAUNA LIST

### REPTILES

Family	Scientific name	Common name	Red list
Agamidae	<i>Agama aculeata distanti</i>	Distant's Ground Agama	Least Concern (SARCA 2014)
Agamidae	<i>Agama atra</i>	Southern Rock Agama	Least Concern (SARCA 2014)
Colubridae	<i>Dasypeltis scabra</i>	Rhombic Egg-eater	Least Concern (SARCA 2014)
Cordylidae	<i>Cordylus vittifer</i>	Common Girdled Lizard	Least Concern (SARCA 2014)
Elapidae	<i>Hemachatus haemachatus</i>	Rinkhals	Least Concern (SARCA 2014)
Gekkonidae	<i>Lygodactylus capensis</i>	Common Dwarf Gecko	Least Concern (SARCA 2014)
Gekkonidae	<i>Pachydactylus capensis</i>	Cape Gecko	Least Concern (SARCA 2014)
Gerrhosauridae	<i>Gerrhosaurus flavigularis</i>	Yellow-throated Plated Lizard	Least Concern (SARCA 2014)
Lamprophiidae	<i>Aparallactus capensis</i>	Black-headed Centipede-eater	Least Concern (SARCA 2014)
Lamprophiidae	<i>Lycophidion capense capense</i>	Cape Wolf Snake	Least Concern (SARCA 2014)
Lamprophiidae	<i>Psammophis brevirostris</i>	Short-snouted Grass Snake	Least Concern (SARCA 2014)
Lamprophiidae	<i>Psammophylax rhombeatus</i>	Spotted Grass Snake	Least Concern (SARCA 2014)
Lamprophiidae	<i>Pseudaspis cana</i>	Mole Snake	Least Concern (SARCA 2014)
Pelomedusidae	<i>Pelomedusa galeata</i>	South African Marsh Terrapin	Not evaluated
Scincidae	<i>Trachylepis capensis</i>	Cape Skink	Least Concern (SARCA 2014)
Scincidae	<i>Trachylepis varia sensu lato</i>	Common Variable Skink Complex	Least Concern (SARCA 2014)
Typhlopidae	<i>Rhinotyphlops lalandei</i>	Delalande's Beaked Blind Snake	Least Concern (SARCA 2014)
Viperidae	<i>Causus rhombeatus</i>	Rhombic Night Adder	Least Concern (SARCA 2014)

### AMPHIBIANS

Family	Scientific name	Common name	Red list
Bufonidae	<i>Schismaderma carens</i>	Red Toad	Least Concern
Bufonidae	<i>Sclerophrys gutturalis</i>	Guttural Toad	Least Concern (IUCN, 2016)
Hyperoliidae	<i>Kassina senegalensis</i>	Bubbling Kassina	Least Concern
Phrynobatrachidae	<i>Phrynobatrachus natalensis</i>	Snoring Puddle Frog	Least Concern (IUCN, 2013)
Pipidae	<i>Xenopus laevis</i>	Common Platanna	Least Concern
Pyxicephalidae	<i>Amietia delalandii</i>	Delalande's River Frog	Least Concern (2017)
Pyxicephalidae	<i>Cacosternum boettgeri</i>	Common Caco	Least Concern (2013)