

PALAEONTOLOGICAL HERITAGE REPORT: COMBINED DESKTOP & FIELD-BASED ASSESSMENT

PROPOSED SOVENTIX SOLAR PV PROJECT ON VARIOUS FARMS, NEAR HANOVER, ENTHANJENI MUNICIPALITY, PIXLEY KA SEME DISTRICT, NORTHERN CAPE

Dr John E. Almond
Natura Viva cc
PO Box 12410 Mill Street
CAPE TOWN 8010, RSA

November 2017

EXECUTIVE SUMMARY

The proposed Soventix 225 MW solar PV plant situated between Hanover and De Aar, Enthanjeni Municipality, Pixley Ka Seme District, Northern Cape, is underlain at depth by Late Permian continental sediments of the Lower Beaufort Group (Adelaide Subgroup), Karoo Supergroup. So far very few vertebrate or other fossils have been recorded in this region, in part because of the thick mantle of alluvial sediments covering the potentially-fossiliferous bedrocks. A palaeontological field survey of the project area confirmed that bedrock exposures are very limited indeed here and the Karoo sediments have been intruded and baked by several dolerite sills.

No palaeontological No-Go areas or highly-sensitive fossil sites have been identified within the development footprint of the Soventrix solar PV project. Apart from questionable fossil invertebrate burrows, the only fossils recorded comprise locally common blocks of well-preserved petrified wood that have been reworked from the Beaufort Group bedrocks into overlying, semi-consolidated alluvial sediments (Fig. 30). The fossils found are all assigned a low Proposed Field Rating (IIIC) and are not considered to require specialist mitigation. Similar derived fossils are likely to occur widely within a zone of older calcretised alluvium along the banks of the Brakrivier and its major tributaries. This palaeontologically more sensitive zone (outlined in blue in Fig. 30 herein) lies within the protected riverine buffer zone and outside the proposed solar PV footprint.

The ECO responsible for the construction phase of the project should be aware of the potential for important new fossil finds – such as vertebrate bones and teeth, or petrified logs - and the necessity to conserve them for possible professional mitigation (See, for example, Macrae 1999 for a well-illustrated popular account of Karoo fossils). The ECO should monitor all substantial excavations into sedimentary rocks for fossil remains on an on-going basis during the construction phase. Recommended mitigation of chance fossil finds involves safeguarding of the fossils (preferably *in situ*) by the responsible ECO and reporting of finds to SAHRA for the Northern Cape (Contact details: SAHRA, 111 Harrington Street, Cape Town. PO Box 4637, Cape Town 8000, South Africa. Phone: +27 (0)21 462 4502. Fax: +27 (0)21 462 4509. Web: www.sahra.org.za). Where appropriate, judicious sampling and recording of fossil material and associated geological data by a qualified palaeontologist, appointed by the developer, may be required by the relevant heritage

regulatory authorities. Any fossil material collected should be curated within an approved repository (museum / university fossil collection) by a qualified palaeontologist.

There are no fatal flaws in the proposed alternative energy project from a palaeontological heritage viewpoint. Cumulative impacts of fossil heritage in the context of several proposed or authorised alternative energy developments in the region (especially around De Aar) are assessed as low, given their comparatively small footprint compared with the outcrop area of the rock units concerned. There are no objections to authorisation of the proposed solar development, provided that the recommended mitigation measures are incorporated into the EMPr for this project and fully implemented.

1. PROJECT DESCRIPTION & BRIEF

The company Soventix South Africa (Pty) Ltd is proposing to develop a 225 MW solar PV plant on various farms located between Hanover and De Aar, Enthanjeni Municipality, Pixley Ka Seme District, Northern Cape (Fig. 1). The land portions concerned include the Remainder of Goedehoop 26C, Portion 6 of Leuwefontein 27C, the Remainder of and Portion 1 of Rietfontein 39C, the Remainder of and Portion 1 of Kwanselaars Hoek 40C, Portion 4 of Taaiboschfontein 41C and Portion 1 of Kafferspoort 56C. The land is currently used for sheep farming.

The footprint of the proposed solar plant is c. 520 ha. This area will include three interconnected 75 MW solar PV plants (170 ha each) with associated infrastructure including a substation that will link into the ESKOM overhead 400 kV powerlines that run through the project area (Fig. 2). Existing roads will be used for main access, which may need to be enlarged to facilitate access to the site during construction. Three PV plant sites have been identified as preferred in consultation with the EAP, Client and Landowner (Fig. 2). The 225 MW plant is considered the first phase and, depending on available capacity on the ESKOM powerlines, additional phase may be implemented in future; these will undergo independent authorisation processes.

The SAHRA Archaeology, Palaeontology and Meteorites (APM) Unit has requested that a Palaeontological Impact Assessment (PIA) inclusive a field assessment be conducted for the proposed development as part of the EIA process (Interim Comment of 1 September 2017. Case ID: 10210).

The present combined desktop and field-based palaeontological heritage study of the Soventrix solar PV plant study area contributes to the EIA and heritage aspects of the Environmental Management Programme (EMPr). The EIA and EMPr process for the project is being co-ordinated by Ecoleges Environmental Consultants, Machadodorp (Contact details: Mr Justin Bowers. Ecoleges Environmental Consultants, Generaal Street, Machadodorp, 1170. PO Box 516 Machadodorp, 1170. Mobile: 082 451 5608. E-mail: juston@ecoleges.co.za).



Figure 1: Google Earth satellite image showing the location of the proposed Soventix solar PV plant some 30 km northwest of Hanover (H), Northern Cape (Scale bar = 10 km.). Three interconnected solar plant sites (1,2,3) make up the integrated PV project (See Fig. 2).




Figure 2: Google Earth satellite image of the project area for the proposed Soventix 225 MW solar PV plant near Hanover, Northern Cape (orange polygon). The three component 75 MW solar plant sites (1, 2, 3) are shown in yellow. Scale bar = 1 km. Arrow points N.

2. APPROACH TO THE PALAEOLOGICAL HERITAGE STUDY

The approach to this palaeontological heritage study is briefly as follows. Fossil bearing rock units occurring within the broader study area are determined from geological maps and satellite images. Known fossil heritage in each rock unit is inventoried from scientific literature, previous assessments of the broader study region, and the author's field experience and palaeontological database. Based on this data as well as field examination of representative exposures of all major sedimentary rock units present, the impact significance of the proposed development is assessed with recommendations for any further studies or mitigation.

In preparing a palaeontological desktop study the potentially fossiliferous rock units (groups, formations *etc*) represented within the study area are determined from geological maps and satellite images. The known fossil heritage within each rock unit is inventoried from the published scientific literature, previous palaeontological impact studies in the same region, and the author's field experience (Almond & Pether 2008). Consultation with professional colleagues as well as examination of institutional fossil collections may play a role here, or later following field assessment during the compilation of the final report. This data is then used to assess the palaeontological sensitivity of each rock unit to development. The likely impact of the proposed development on local fossil heritage is then determined on the basis of (1) the palaeontological sensitivity of the rock units concerned and (2) the nature and scale of the development itself, most significantly the extent of fresh bedrock excavation envisaged. When rock units of moderate to high palaeontological sensitivity are present within the development footprint, a Phase 1 field assessment study by a professional palaeontologist is usually warranted to identify any palaeontological hotspots and make specific recommendations for any monitoring or mitigation required before or during the construction phase of the development.

On the basis of the desktop and Phase 1 field assessment studies, the likely impact of the proposed development on local fossil heritage and any need for specialist mitigation are determined. Adverse palaeontological impacts normally occur during the construction rather than the operational or decommissioning phase. Phase 2 mitigation by a professional palaeontologist – normally involving the recording and sampling of fossil material and associated geological information (*e.g.* sedimentological data) may be required (a) in the pre-construction phase where important fossils are already exposed at or near the land surface and / or (b) during the construction phase when fresh fossiliferous bedrock has been exposed by excavations. To carry out mitigation, the palaeontologist involved will need to apply for palaeontological collection permits from the relevant heritage management authorities, *i.e.* SAHRA for the Northern Cape (Contact details: SAHRA, 111 Harrington Street, Cape Town. P.O. Box 4637, Cape Town 8000, South Africa. Phone: +27 (0)21 462 4502. Fax: +27 (0)21 462 4509. Web: www.sahra.org.za). It should be emphasized that, *providing appropriate mitigation is carried out*, the majority of developments involving bedrock excavation can make a *positive* contribution to our understanding of local palaeontological heritage.

GPS data for geological and fossil localities mentioned in the text and figure legends are provided separately in the Appendix to this report.

2.1. Information sources

The information used in this scoping palaeontological heritage study was based on the following:

1. A short project description, maps and kmz files kindly provided by Ecoleges Environmental Consulting;
2. A review of the relevant satellite images, topographical maps and scientific literature, including published geological maps and accompanying sheet explanations, as well as several previous desktop and field-based palaeontological assessment studies in the broader Hanover – De Aar study region (e.g. Almond 2010a-b, 2011, 2012a-c, 2013a-d, Millstead 2014, Cedar Tower Services 2015, Groenewald 2012).
3. The author's previous field experience with the formations concerned and their palaeontological heritage (Almond & Pether 2008);
4. A one-day palaeontological field assessment in October 2017 by the author and one assistant.

2.2. Assumptions & limitations

The accuracy and reliability of palaeontological specialist studies as components of heritage impact assessments are generally limited by the following constraints:

1. Inadequate database for fossil heritage for much of the RSA, given the large size of the country and the small number of professional palaeontologists carrying out fieldwork here. Most development study areas have never been surveyed by a palaeontologist.
2. Variable accuracy of geological maps which underpin these desktop studies. For large areas of terrain these maps are largely based on aerial photographs alone, without ground-truthing. The maps generally depict only significant ("mappable") bedrock units as well as major areas of superficial "drift" deposits (alluvium, colluvium) but for most regions give little or no idea of the level of bedrock outcrop, depth of superficial cover (soil *etc*), degree of bedrock weathering or levels of small-scale tectonic deformation, such as cleavage. All of these factors may have a major influence on the impact significance of a given development on fossil heritage and can only be reliably assessed in the field.
3. Inadequate sheet explanations for geological maps, with little or no attention paid to palaeontological issues in many cases, including poor locality information.
4. The extensive relevant palaeontological "grey literature" - in the form of unpublished university theses, impact studies and other reports (e.g. of commercial mining companies) - that is not readily available for desktop studies.
5. Absence of a comprehensive computerized database of fossil collections in major RSA institutions which can be consulted for impact studies. A Karoo fossil vertebrate database is now accessible for impact study work.

In the case of palaeontological desktop studies without supporting Phase 1 field assessments these limitations may variously lead to either:

- (a) *underestimation* of the palaeontological significance of a given study area due to ignorance of significant recorded or unrecorded fossils preserved there, or

(b) *overestimation* of the palaeontological sensitivity of a study area, for example when originally rich fossil assemblages inferred from geological maps have in fact been destroyed by tectonism or weathering, or are buried beneath a thick mantle of unfossiliferous “drift” (soil, alluvium *etc*).

Since most areas of the RSA have not been studied palaeontologically, a palaeontological desktop study usually entails *inferring* the presence of buried fossil heritage within the study area from relevant fossil data collected from similar or the same rock units elsewhere, sometimes at localities far away. Where substantial exposures of bedrocks or potentially fossiliferous superficial sediments are present in the study area, the reliability of a palaeontological impact assessment may be significantly enhanced through field assessment by a professional palaeontologist.

In the case of the present study area near Hanover in the Northern Cape preservation of potentially fossiliferous bedrocks is favoured by the semi-arid climate and sparse vegetation but bedrock exposure is very limited – especially within the flatter-lying PV plant development areas - by extensive superficial deposits, especially alluvium, sandy soils and scree. However, it is considered that sufficient bedrock and cover sediment exposures were examined during the course of this study to assess the broader palaeontological heritage sensitivity of the study area (See Appendix). Comparatively few academic palaeontological studies or field-based fossil heritage impact studies have been carried out in the region, so any new data from impact studies here are of scientific interest.

2.3. Legislative context for palaeontological assessment studies

The proposed alternative energy project is located in an area that is underlain by potentially fossiliferous sedimentary rocks of Palaeozoic, Mesozoic and younger, Late Tertiary or Quaternary, age (Sections 3 and 4). The construction phase of the proposed development will entail substantial excavations into the superficial sediment cover and locally into the underlying bedrock as well. These may include, for example, excavations for the PV panel footings, internal access roads, underground cables, transmission line pylon footings, electrical substation, operations and services workshop area/office building and construction site camp. All these developments may adversely affect potential fossil heritage within the study area by destroying, disturbing or permanently sealing-in fossils at or beneath the surface of the ground that are then no longer available for scientific research or other public good. The operational and decommissioning phases of the wind energy facility are unlikely to involve further adverse impacts on local palaeontological heritage, however.

The present combined desktop and field-based palaeontological heritage study contributes to the Heritage Assessment for the Soventrix 225 MW solar PV plant project and falls under the South African Heritage Resources Act (Act No. 25 of 1999). It will also inform the Environmental Management Programme for this project.

The various categories of heritage resources recognised as part of the National Estate in Section 3 of the National Heritage Resources Act include, among others:

- geological sites of scientific or cultural importance;
- palaeontological sites;
- palaeontological objects and material, meteorites and rare geological specimens.

According to Section 35 of the National Heritage Resources Act, dealing with archaeology, palaeontology and meteorites:

(1) The protection of archaeological and palaeontological sites and material and meteorites is the responsibility of a provincial heritage resources authority.

(2) All archaeological objects, palaeontological material and meteorites are the property of the State.

(3) Any person who discovers archaeological or palaeontological objects or material or a meteorite in the course of development or agricultural activity must immediately report the find to the responsible heritage resources authority, or to the nearest local authority offices or museum, which must immediately notify such heritage resources authority.

(4) No person may, without a permit issued by the responsible heritage resources authority—

(a) destroy, damage, excavate, alter, deface or otherwise disturb any archaeological or palaeontological site or any meteorite;

(b) destroy, damage, excavate, remove from its original position, collect or own any archaeological or palaeontological material or object or any meteorite;

(c) trade in, sell for private gain, export or attempt to export from the Republic any category of archaeological or palaeontological material or object, or any meteorite; or

(d) bring onto or use at an archaeological or palaeontological site any excavation equipment or any equipment which assist in the detection or recovery of metals or archaeological and palaeontological material or objects, or use such equipment for the recovery of meteorites.

(5) When the responsible heritage resources authority has reasonable cause to believe that any activity or development which will destroy, damage or alter any archaeological or palaeontological site is under way, and where no application for a permit has been submitted and no heritage resources management procedure in terms of section 38 has been followed, it may—

(a) serve on the owner or occupier of the site or on the person undertaking such development an order for the development to cease immediately for such period as is specified in the order;

(b) carry out an investigation for the purpose of obtaining information on whether or not an archaeological or palaeontological site exists and whether mitigation is necessary;

(c) if mitigation is deemed by the heritage resources authority to be necessary, assist the person on whom the order has been served under paragraph (a) to apply for a permit as required in subsection (4); and

(d) recover the costs of such investigation from the owner or occupier of the land on which it is believed an archaeological or palaeontological site is located or from the person proposing to undertake the development if no application for a permit is received within two weeks of the order being served.

Minimum standards for the palaeontological component of heritage impact assessment reports (PIAs) have been published by SAHRA (2013).

3. GEOLOGICAL CONTEXT

The study area for the proposed Soventix 225 MW solar PV plant is situated some 30 km NW of Hanover and 35 km SE of De Aar in the Great Karoo region of the Northern Cape (Fig. 1). The terrain is semi-arid, topographically-subdued Karoo with rocky dolerite ridges and *koppies* (especially towards the western margins of the area) and extensive alluvial *vlaktes* associated with the NW-flowing Brakrivier and its numerous tributary streams (Figs.3 to 6). Most of the landscape lies between 1300-1380 m amsl. Judging from place names on the 1: 50 000 maps (3024CC, CD) there are numerous fountains in the area, probably related to local dolerite intrusions. Due to the pervasive cover by superficial sediments (alluvium, soils, colluvium), levels of Karoo sedimentary bedrock exposure in the region are very low indeed, and mainly confined to small borrow pits, road and railway cuttings, farm dams and the more deeply-incised drainage courses.

The geology of the study area is outlined on the 1: 250 000 geology sheet 3024 Colesberg (Le Roux 1993) (Fig. 7). The area is underlain by Late Permian sedimentary rocks of the Karoo Supergroup that are intruded by Early Jurassic dolerites. According to the 1: 250 000 geological map the area is largely underlain at depth by Permian continental sediments of the **Adelaide Subgroup (Lower Beaufort Group)** (Pa) that in this region are extensively intruded by dolerite dykes of the **Karoo Dolerite Suite** (Jd). The great majority of the Beaufort Group outcrop area is obscured by superficial sediments of probable Pleistocene to Holocene age, as well as by abundant karroid shrub and grassy vegetation (Figs. 4 to 6). These superficial sediments include silty to sandy soils and alluvium related to the broader Brakrivier drainage system, doleritic colluvium (scree, hillwash slope deposits), and downwasted surface gravels dominated by resistant clasts of hornfels, sandstone and dolerite that are modified by sheet wash processes. The dolerite intrusions weather out at surface as low rocky ridges and *koppies* that show up in rusty-brown colours in satellite images (Fig. 2). They have baked (thermally metamorphosed) the adjacent Karoo Supergroup mudrocks to hornfels, and any sandstones to quartzites. Dolerite colluvial rubble extends well beyond the intrusions themselves to blanket adjacent slopes and *vlaktes*.



Figure 3: Flat grassy *bossieveld* terrain on Goedehoop 26 showing lack of bedrock exposure here.



Figure 4: Eroded silty to sandy *vlaktes* on Kwanselaarshoek 40. The Karoo Supergroup bedrocks in such areas are deeply buried beneath thick alluvial deposits.



Figure 5: Incised course of the Brakrivier on Rietfontain 39 with good sections through Late Caenozoic alluvial deposits along the steep banks. Note lack of bedrock exposure in the shallow river bed here.



Figure 6: Typical bouldery *koppies* of Karoo dolerite in the western portion of the study area (Farm 56).

3.1. Adelaide Subgroup

The Adelaide Subgroup (Pa) (Lower Beaufort Group, Karoo Supergroup) was deposited by large-scale meandering river systems flowing northwards from the youthful Cape Fold Belt across the extensive floodplains of the ancient Karoo Basin (Smith 1980, Rubidge 1995, Johnson *et al.* 2006). The sediments mainly comprise fine-grained overbank mudrocks with subordinate lenticular channel sandstones. These last commonly have a basal conglomeratic lag of rolled mudflake pellets and calcrete nodules, the latter reflecting the prevailing semi-arid climates in Middle to Late Permian times. Small, often transient playa lakes were also present on the floodplain. In the Colesberg 1: 250 000 sheet area the Lower Beaufort succession consists largely of blocky-weathering, blue-grey and subordinate reddish floodplain mudrocks, showing occasional mudcracks. There are also subordinate siltstones, fine-grained, lenticular, current cross-bedded channel sandstones, flat-laminated crevasse-splay sandstones, and occasional playa lake deposits (Le Roux 1993). Carbonate lenses and concretions, including ferruginous *koffieklip*, as well as calcrete nodules (pedogenic limestones) and silicified gypsum rosettes (“desert roses”) are common.

The precise stratigraphic assignment of the Lower Beaufort Group sediments in the study area near Hanover is unresolved on the 1: 250 000 geology map (Le Roux 1993). According to the most recent fossil biozonation map of the Beaufort Group (Van der Walt *et al.* 2010) (Fig. 24) the sediments here are assigned to the *Tropidostoma* Assemblage Zone that characterises the mudrock-dominated **Hoedemaker Member** of the Teekloof Formation west of longitude 24° East, as well as the Middleton Formation to the east (Rubidge 1995). De Aar is situated on the (arbitrary) cut-off line between these two stratigraphic schemes.

The geology of the Hoedemaker Member, which is up to 240 m thick, is outlined by Smith (1980, 1993a, b) and later by Smith and Keyser (1995) as well as Cole and Smith (2008). The Hoedemaker succession is dominated by greenish-grey to purple-brown overbank mudrocks, with occasional single-storey sheet sandstones. Palaeosol (ancient soil) horizons characterized by calcrete nodules and rhizocretions (root casts) are common, as are also lacustrine (transient to long-lived playa lake) sediments deposited in depressions on the Late Permian floodplain. These last are associated with limestone crusts, gypsum crystals (“desert roses”) as well as a range of fine-scale sedimentary features such as wave rippled sandstones, falling water marks, mudcracks, and trace fossils (Stear 1978, Smith 1980, 1986, 1993a).

Only very limited exposures of the Adelaide Subgroup bedrocks are seen within the present study area. Prominent-weathering, pale brown channel sandstones crop out on hillslopes on Goedehoop 26 (Fig. 10) but are largely obscured by doleritic colluvium. Small rafts or xenoliths of sandstone can be seen enclosed within the dolerite itself, while exposures of streaky, foliated sandstone reflect remobilization and baking of the Karoo Supergroup country rocks by hot dolerite magma (Figs. 15 & 16). The overbank mudrocks in the vicinity of the intrusions have been thermally metamorphosed to form dark grey hornfels (often brown-patinated) which dominates surface gravels locally (Fig. 21). Small exposures of well-jointed, thin sandstone beds (probably crevasse splays) and grey-green hackly-weathering mudrocks occur along more deeply-incised tributaries of the Brakrivier (Figs. 8 & 9). Good exposures of the mudrock facies is also seen in borrow pits near the railway line where they contain well-developed pale grey to pinkish-grey calcrete nodules and lenses reflecting palaeosols (Fig. 12). Some of the calcrete

nodules show internal “septarian” cracking, probably caused by diagenetic shrinkage (Fig. 13). Occasional resistant-weathering patches or lenses of dark brown ferruginous carbonate (*koffieklip*) are encountered in the field (Fig. 11) and may be a target for plant fossil recording.

3.2. Karoo Dolerites

The Karoo Dolerite Suite (Jd) is an extensive network of basic igneous bodies (dykes, sills) that were intruded into sediments of the Main Karoo Basin in the Early Jurassic Period, about 183 million years ago (Duncan & Marsh 2006). These dolerites form part of the Karoo Igneous Province of Southern Africa that developed in response to crustal doming and stretching preceding the break-up of Gondwana. Hard cappings of blocky, reddish-brown to rusty-weathering dolerite are a very typical feature of the flat-topped *koppies* in the Great Karoo region. As seen from geological maps (Fig. 7), extensive dolerite intrusion of the Lower Beaufort Group rocks is observed in the Hanover region. The country rocks adjacent to the intrusions have often been extensively baked or thermally metamorphosed. Mudrocks are altered to flinty hornfels (“lydianite” of some authors), while sandstones are metamorphosed to resistant-weathering, siliceous quartzites that may show complex deformation of the original bedding (Figs. 15 & 16). The Karoo rocks within the thermal aureole of the dolerite intrusions are also often chemically altered; they tend to be silicified, more brittle and contain numerous irregular *vugs* (cavities) lined or infilled with secondary minerals. Blocky, well-jointed outcrops of Karoo dolerite are seen on the slopes and summits of low *koppies* and ridges or *rante* in various parts of the study area, especially towards the western margins (Fig. 14).

3.3. Kimberlite intrusions

Numerous **kimberlite pipes and dykes** of Jurassic to Cretaceous age intrude the Karoo Supergroup rocks between Hanover and De Aar, including several examples to the north and east of the present study area (black diamonds in Fig. 7). They are variously assigned to the Victoria West and Group II Provinces (Skinner & Truswell 2006) and do not contain diamonds. According to Le Roux (1993) the ultramafic kimberlite pipe rocks in the Colesberg sheet area are highly weathered with no obvious surface expression. They can usually be located only on the basis of characteristic mineral assemblages (garnet, phlogopite mica) found in ant heaps, termite mounds and prospecting holes. Kimberlite rocks are unfossiliferous, although rich Cretaceous to Paleocene fossil assemblages may be found in associated crater lake facies (not present here).

3.4. Late Cenozoic superficial deposits

Quaternary to Recent superficial deposits cover all but the steepest slopes of the dolerite *koppies* and *rante* as well as most of the plains at their feet, including dry river courses such as the Brakrivier in the study region (Figs. 3 to 5). Various types of superficial deposits of geologically young, Late Cenozoic (Miocene / Pliocene to Recent) age (< 5 Ma) occur throughout the Great Karoo region (Prinsloo 1989, Le Roux 1993, with more extensive discussion in Holmes & Marker 1995, Cole *et al.* 2004, Partridge *et al.* 2006). They include pedocretes (*e.g.* calcretes), colluvial slope deposits (dolerite, sandstone and hornfels scree *etc.*), sandy, gravelly and bouldery river alluvium, as well as spring and pan sediments.

These colluvial and alluvial deposits may be extensively calcretised (*i.e.* cemented with soil limestone), especially in the neighbourhood of dolerite intrusions.

The dominant superficial sediments in the present study area comprise thick (up to several m), fine-grained alluvial deposits of various ages associated with the Brakrivier and its tributaries (Figs. 17 to 19). These deposits are well-exposed along the more deeply-incised stream banks as well as adjoining zones of gully erosion (*dongas*). The younger (probably Holocene) alluvium is pale brown, silty to sandy and unconsolidated (without calcrete), and contains thin lenses and horizons of gravels as well as dispersed gravel clasts. These last include locally abundant hornfels stone artefacts that tend to down-waste in eroded areas onto the underlying firmer alluvium. The distinctive older alluvial deposits are orange-brown in hue, sandy and partially consolidated with numerous creamy calcrete veins. They contain sparse to locally abundant gravels, including reworked petrified wood (Section 4) and are probably Pleistocene in age. Fine-grained alluvial soils cover much of the lower-lying parts of the study area (pale areas in Fig. 2). Bare soil patches expose surface gravels of sheet-washed hornfels (sometimes flaked), dolerite, quartzite and calcrete but – in contrast with previous studies near De Aar – no petrified wood was observed in this setting (Fig. 20).




Figure 7: Geological map of the Soventix solar PV plant project area between De Aar and Hanover, Northern Cape (yellow polygon) (Map abstracted from 1: 250 000 geology sheet 3024 Colesberg, Council for Geoscience, Pretoria). The following main rock units are mapped within the study areas: green (Pa) = Adelaide Subgroup (Lower Beaufort Group); pink (Jd) = intrusive dykes and sills of the Karoo Dolerite Suite; white = Pleistocene to Recent superficial deposits

(alluvium, colluvium etc); small black diamond symbols = Kimberlite pipe; blue lines – kimberlite dykes (k).



Figure 8: Rare exposures of Adelaide Subgroup sandstones (foreground) and mudrocks (background) in the bed and banks of the Brakrivier on Rietfontain 39 (Loc. 011).



Figure 9: Thin crevasse splay sandstones and grey-green overbank mudrocks exposed at the locality shown above, Rietfontain 39 (Loc. 011) (Hammer = 27 cm).



Figure 10: Exposure of pale brown channel sandstones of the Adelaide Subgroup on Goodehoop 26 (Loc. 004) (Hammer = 27 cm).



Figure 11: Extensive lens of dark brown ferruginous carbonate (*koffieklip*) exposed in the *vlaktes* on Goedehoop 26 (Loc. 007).



Figure 12: Pale pinkish-grey calcrete nodules within a palaeosol horizon, borrow pit exposure of Adelaide Subgroup overbank mudrocks on Goedehoop 26 (Loc. 001) (Hammer = 27 cm). Such ancient soil units are a primary focus for fossil recording.



Figure 13: Ovoid grey calcrete nodules showing septarian cracking, surface gravels in bare patch on Kwanselaarshoek 40 (Loc. 010) (Scale in cm).



Figure 14: Well-jointed rusty-brown *koppies* reflecting surface expression of a major dyke of resistant-weathering , Kafferspoort 56.



Figure 15: Xenolith of Beaufort Group quartzite enclosed within a dolerite intrusion, Goodehoop 26 (Loc. 003) (Hammer = 27 cm).



Figure 16: Foliated and contorted quartzite in the vicinity of a dolerite intrusion, Goodehoop 26 (Loc. 005) (Hammer = 27 cm).



Figure 17: Vertical section through calcrete-veined, semi-consolidated older alluvial deposits underlying brownish unconsolidated younger alluvium exposed along the banks of the Brakrivier, Rietfontain 39 (Loc. 018) (Hammer = 27 cm).



Figure 18: Good gullied exposures of older (probably Pleistocene) consolidated , orange-brown alluvial deposits showing calcrete, Kafferspoort 56 (Loc. 13) (Hammer = 27 cm).



Figure 19: Riverbank exposure of older, more orange-brown and younger, brownish alluvium, Kafferspoort 56 (Loc 13) (Hammer = 27 cm).



Figure 20: Typical bare patch with sparse, fine surface gravels of hornfels, dolerite, quartzite and calcrete, Kafferspoort 56 (Loc. 015). Such patches are a search target for downwasted blocks of petrified wood.



Figure 21: Downwasted colluvial gravels of brownish-patinated hornfels (with several flaked stone artefacts), Goodehoop 26 (Loc. 009) (Hammer = 27 cm).



Figure 22: Well-developed pedocrete of cream-coloured calcrete within soils overlying dolerite, Goodehoop 26 (Loc. 008) (Hammer = 27 cm).

4. PALAEOONTOLOGICAL HERITAGE

4.1. Fossils within the Lower Beaufort Group

A chronological series of mappable fossil biozones or assemblage zones (AZ), defined mainly on their characteristic tetrapod faunas, has been established for the Main Karoo Basin of South Africa (Rubidge 1995). Maps showing the distribution of the Beaufort assemblage zones within the Main Karoo Basin have been provided by Kitching (1977), Keyser and Smith (1979), Rubidge (1995) and Van der Walt *et al.* (2010) (Fig. 24). These sources establish that the study area *probably* lies within the ***Tropidostoma* Assemblage Zone** (AZ) that characterizes the Hoedemaker Member of the Teekloof Formation (Smith & Keyser, 1995).

The following major categories of fossils might be expected within *Tropidostoma* AZ sediments in the study area (Kitching 1977, Keyser & Smith 1977-78, Anderson & Anderson 1985, Smith & Keyser 1995, MacRae 1999, Cole *et al.* 2004, Smith *et al.* 2012):

- isolated petrified bones as well as rare articulated skeletons of **terrestrial vertebrates (tetrapods)** such as true **reptiles** (notably large herbivorous pareiasaurs) and **therapsids** or “mammal-like reptiles” (e.g. diverse herbivorous dicynodonts, flesh-eating gorgonopsians, and insectivorous therocephalians) (Fig. 23);

- aquatic vertebrates such as large **temnospondyl amphibians** (*Rhinesuchus* spp., usually disarticulated), and **palaeoniscoid bony fish** (*Atherstonia*, *Namaichthys*, often represented by scattered scales rather than intact fish);
- freshwater **bivalves** (e.g. *Palaeomutela*);
- **trace fossils** such as worm, arthropod and tetrapod burrows and trackways, coprolites (fossil droppings), fish swimming trails;
- **vascular plant remains** including leaves, twigs, roots and petrified woods ("*Dadoxylon*") of the *Glossopteris* Flora (usually sparse, fragmentary), especially glossopterid trees and arthropytes (horsetails).

According to Smith and Keyser (1995) the tetrapod fauna of the *Tropidostoma* Assemblage Zone is dominated by the small burrowing dicynodont *Diictodon* that constitutes some 40% of the fossil remains recorded here. There are several genera of toothed dicynodonts (e.g. *Emydops*, *Priesterodon*) as well as medium-sized forms like *Rachiocephalus*, *Tropidostoma* and *Endothiodon*. Carnivores are represented by medium-sized gorgonopsians (e.g. *Lycaenops*, *Gorgonops*) as well as smaller, insectivorous therocephalians such as *Ictidosuchoides* and the early cynodont *Procynosuchus*. Among the large (2.3 to 3 m long), lumbering pareiasaur reptiles the genus *Pareiasaurus* replaces the more primitive *Bradysaurus* seen in older Beaufort Group assemblages.

As far as the biostratigraphically important tetrapod remains are concerned, the best fossil material within the Hoedemaker Member succession is generally found within overbank mudrocks, whereas fossils preserved within channel sandstones tend to be fragmentary and water-worn (Rubidge 1995, Smith 1993b). Many vertebrate fossils are found in association with ancient soils (palaeosol horizons) that can usually be recognised by bedding-parallel concentrations of calcrete nodules. Smith and Keyser (1995) report that in the *Tropidostoma* Assemblage Zone / Hoedemaker Member most tetrapod fossils comprise isolated disarticulated skulls and post-cranial bones, although well-articulated skeletons of the small dicynodont *Diictodon* are locally common, associated with burrows (See also Smith 1993b for a benchmark study of the taphonomy of vertebrate remains in the Hoedemaker Member).




Figure 23: Skulls of two important medium-sized dicynodont therapsids that characterise the *Tropidostoma* Assemblage Zone: A – *Endothiodon*; B – *Tropidostoma* (From Smith & Keyser 1995).

Due to the generally very poor exposure of Lower Beaufort Group (Adelaide Subgroup) bedrocks in the region between De Aar and Hanover, there have been very few identifiable vertebrate or other fossil finds here (See fossil site map from Nicolas 2007 in Fig. 25). Fragmentary skeletal remains of small-bodied therapsids, mainly dicynodonts, as well as of the small tortoise-like reptile *Eunotosaurus* have been recorded from the Lower Beaufort Group near De Aar by Almond (2012a, 2012b) but these belong to a slightly older horizon within the Lower Beaufort Group. Associated fossils include scrappy plant remains – mainly sphenophyte ferns and well-preserved silicified wood – as well as low-diversity trace fossil assemblages.

As a consequence of their proximity to numerous dolerite intrusions, the Beaufort Group sediments in the study area have been thermally metamorphosed or “baked” (*i.e.* recrystallised, impregnated with secondary minerals). Embedded fossil material of phosphatic composition, such as bones and teeth, is frequently altered by baking – bones may become blackened, for example - and can be very difficult to extract from the hard matrix by mechanical preparation (Smith & Keyser 1995). Thermal metamorphism by dolerite intrusions therefore tends to reduce the palaeontological heritage potential of Beaufort Group sediments.

No new vertebrate fossils were recorded during the present site visit. *Possible* small-scale invertebrate burrows with a curious rugose surface texture were seen within *koffieklip* exposures on Goedehoop 26 (Fig. 26). They are associated with mudflake intraclast breccias and their biogenic nature is equivocal. The only Permian body fossil remains found comprise locally common blocks of well-preserved silicified

wood showing clear development of seasonal growth banding (*cf* Bamford 1999, 2004) (Figs. 27 to 29). This fossil wood was not found embedded *in situ* within the Lower Beaufort Group bedrocks however, but reworked into the overlying Late Caenozoic alluvial deposits. Recorded fossil wood sites (See Appendix) associated with older alluvial deposits of the Brakrivier are shown on the satellite image in Fig. 30. The fossil wood specimens encountered are likely to be of widespread occurrence and not of high scientific or conservation value. They are therefore assigned a provisional field rating of IIC Local Resource. It is likely that reworked petrified wood occurs widely within the older consolidated alluvium along the Brakrivier and its major tributaries. These older alluvial deposits are therefore considered to be palaeontologically sensitive. They are likely to be protected in large part within the buffer zone for riverine areas and the best exposures lie outside the proposed solar plant development areas (elongate blue area marked in Fig. 30).




Figure 24: Extract from the latest Karoo fossil biozonation map (Van der Walt *et al.* 2010). The present study area near Hanover (yellow triangle) *probably* lies within the *Tropidostoma* Assemblage Zone of Late Permian age, indicated in brown, although this has not been confirmed through palaeontological field data.




Figure 25: Map of Beaufort Group vertebrate fossil localities showing the lack of fossil finds in the vicinity of the present study area (yellow triangle) between De Aar (DA) and Hanover (H) (Map abstracted from Nicolas 2007). Pink – N. Cape. Dark blue – Eastern Cape.

4.2. Fossils within Pleistocene to Recent alluvium

The Pleistocene to Recent superficial deposits in the Karoo have been comparatively neglected in palaeontological terms for the most part. However, they may occasionally contain important fossil biotas, notably the bones, teeth and horn cores of mammals (e.g. Skead 1980, Klein 1984, MacRae 1999, Partridge & Scott 2000). These may include ancient human remains of considerable palaeoanthropological significance (e.g. Grine *et al.*, 2007). Other late Caenozoic fossil biotas from these superficial deposits include non-marine molluscs (bivalves, gastropods), ostrich egg shells, trace fossils (e.g. calcretised termitaria, coprolites, rhizoliths), and plant remains such as peats or palynomorphs (pollens) in fine-grained, organic-rich alluvial horizons. Quaternary alluvial sediments may contain reworked Stone Age artifacts that are useful for constraining their maximum age.

As noted in the previous section, the older, partially-consolidated alluvial deposits along the Brakrivier and its major tributaries contain locally common blocks of petrified fossil wood that have been reworked from the Permian sedimentary bedrocks. No fossil mammalian or invertebrate remains were seen during the site visit.



Figure 26: Unidentified burrow-like structures (c. 1 cm across) with a rugose surface texture and possible mudrock infill, *koffieklip* lens on Goodehoop 26 (Loc. 007).



Figure 27: Small cherty blocks of reworked silicified wood eroded out of older alluvial deposits on Kafferspoort 56 (Loc. 014) (Scale in cm and mm).



Figure 28: Small blocks of silicified wood from older alluvial deposits along the Brakrivier on Rietfontain 39 (Loc. 017) (Scale in cm and mm).




Figure 29. Larger angular block of silicified wood showing clear seasonal growth-banding from calcretised older alluvium on Rietfontain 39 (Loc. 020) (Scale in cm and mm).




Figure 30: Google Earth satellite image of the Soventix solar plant project area showing numbered new fossil localities in yellow. 007 – questionable invertebrate burrows in *koffieklip*. 014, 017, 020 – locally common blocks of reworked Permian petrified wood eroding out of consolidated older alluvium along the Brakrivier (approximately indicated by the blue area). Note absence of fossil sites within three 75 MW solar PV development areas (red polygons). See Appendix for locality details & brief descriptions. Scale bar = 5 km.

5. EVALUATION OF IMPACTS ON PALAEOLOGICAL HERITAGE

The Soventix solar PV plant study area is located in a region of the Great Karoo that is underlain by potentially-fossiliferous sedimentary rocks of Permian and younger, Pleistocene to Holocene age (Sections 3 & 4). The construction phase of the proposed solar energy facility will entail substantial excavations into the superficial sediment cover and locally into the underlying bedrock as well. These include, for example, surface clearance and excavations for the PV panel footings, laydown areas, internal access roads, underground cables, transmission line pylon footings, electrical substation, operations and services workshop area/office building and construction camp. All these developments may adversely affect potential fossil heritage within the study area by destroying, disturbing or permanently sealing-in fossils preserved at or beneath the surface of the ground that are then no longer available for scientific research or other public good.

The inferred impact of the proposed PV solar plant on local fossil heritage resources – including all three of the component 75 MW solar PV plants - is briefly evaluated here in Table 1. This assessment applies only to the *construction phase* of the development since further significant impacts on fossil heritage during the planning, operational and decommissioning phases of the facilities are not anticipated. Confidence levels in this assessment are *medium*, given (1) the extensive palaeontological literature on the Karoo bedrocks concerned weighed against (2) very low levels of bedrock exposure within the study area and (3) the unpredictable distribution of well-preserved fossils in the subsurface.

As motivated in Table 1, the impact significance of the proposed development in terms of palaeontological heritage is assessed as *medium (negative)*. It should be noted that, should the recommended mitigation measures for the construction phase of the WEF development, as outlined in Section 6 of this report, be consistently followed-though, the impact significance would remain *medium (negative)* but would entail both positive and negative impacts. Residual negative impacts from inevitable loss of some valuable fossil heritage would be partially offset by an improved palaeontological database for the study region as a direct result of appropriate mitigation. This is a *positive* outcome because any new, well-recorded and suitably-curated fossil material from this palaeontologically little-known region would constitute a useful addition to our scientific understanding of Karoo Basin fossil heritage.

There are no fatal flaws in the proposed solar PV project from a palaeontological heritage viewpoint and no objects to authorisation of the development, provided that the recommended mitigation measures are fully implemented.

Table 1: Evaluation of anticipated impacts on fossil heritage resources of the proposed Soventix solar PV project near Hanover (Construction Phase)

CRITERIA	CATEGORY	COMMENTS
Extent	Low (2)	Limited to development footprint which is small.
Magnitude	Low (2)	Highly significant fossil material (e.g. vertebrate remains) is – at most - sparsely distributed at or near surface within the study area with no known sites within the development footprint, a significant portion of which is underlain by unfossiliferous dolerite. Fossils recorded occur outside the footprint and are of common types (e.g. petrified wood) with Provisional Field Rating of IIC Local Resource.
Duration	High (4)	Permanent.
Mitigatory Potential	Moderate (3)	Avoidance of riverine alluvial areas (already incorporated into proposed layout) + reporting of chance fossil finds to SAHRA. Residual impacts unavoidable.
Acceptability	Low (2)	Provided that mitigation measures are implemented
Degree of certainty / Probability of impact occurring	Low (2)	Fossils of some sort occur widely, so <i>some</i> level impact is unavoidable. However, most fossils are of low conservation value. The probability of loss of <i>well-preserved, scientifically important fossils</i> is rated as low.
Status	Negative	Loss of fossils preserved at or near surface during construction. Offset by new fossils reported as chance finds (positive impact).
Significance	Medium (24)	Specialist monitoring or mitigation measures therefore not proposed for this project.

5.2. Cumulative impact assessment

Previous palaeontological assessments (PIAs) for several proposed or authorized alternative energy projects within a 30 km radius of the Soventix solar PV project area have been briefly reviewed (e.g. Almond 2010a-b, 2011, 2012a-c, 2013a-d, Millstead 2014, Cedar Tower Services 2015, Groenewald 2012). It is noted that in several cases heritage assessments for alternative energy projects have been submitted, and even approved, without a PIA, or with only a desktop study. Several of the PIA studies referenced include field-based assessments involving Lower Beaufort Group bedrocks in the De Aar region, but in most cases these are for older bedrocks (e.g. Ecca Group, *Pristerognathus* Assemblage Zone) – and fossil assemblages – than those represented in the Soventrix study area.

In the author's opinion:

- Palaeontological impact significances inferred for these projects that range from low to medium to unassessed reflecting different assessment approaches rather than contrasting palaeontological sensitivities and impact levels;
- Meaningful cumulative impact assessments require comprehensive data on *all* major developments within a region, not just those involving alternative energy, as well as an understanding of the extent to which recommended mitigation measures are followed through;
- Trying to assess cumulative impacts on fossil assemblages from different stratigraphic units (for example, Late Permian fossils from the Adelaide Subgroup / Lower Beaufort Group *versus* Middle Permian assemblages from the Ecca Group) has limited value.

Given the comparatively small combined footprint of the alternative energy projects under consideration compared with the very extensive outcrop areas of the Lower Beaufort Group and Late Caenozoic alluvial sediments in the Great Karoo, the cumulative impact significance of the Soventrix solar PV proposal is assessed as LOW. There are therefore no objections on palaeontological grounds to approval of this last project.

6. RECOMMENDATIONS FOR MONITORING AND MITIGATION

Proposed monitoring and mitigation measures for the Soventix solar PV plant, to be incorporated into the Environmental Management Programme for the development, are given in Table 2.

No palaeontological No-Go areas or fossil sites requiring specialist mitigation have been identified within the Soventix PV solar development footprint near Hanover; fossil sites shown in Fig. 30 are rated as of low sensitivity (Proposed Field Rating IIIC). It is recommended that the older consolidated fluvial deposits along the Brakrivier (blue area outlined in Fig. 30) be avoided during construction since they do contain fossil wood. This area lies within the riverine buffer zone and outside the proposed solar PV plant footprint.

The ECO responsible for the construction phase of the project should be aware of the potential for important fossil finds and the necessity to conserve them for possible

professional mitigation (See, for example, Macrae 1999 for a well-illustrated popular account of Karoo fossils). The ECO should monitor all substantial excavations into sedimentary rocks for fossil remains on an on-going basis during the construction phase.

Recommended mitigation of chance fossil finds during the construction phase of the solar PV plant and associated grid connection involves safeguarding of the fossils (preferably *in situ*) by the responsible ECO and reporting of finds to SAHRA for the Northern Cape (Contact details: SAHRA, 111 Harrington Street, Cape Town. PO Box 4637, Cape Town 8000, South Africa. Phone: +27 (0)21 462 4502. Fax: +27 (0)21 462 4509. Web: www.sahra.org.za). Where appropriate, judicious sampling and recording of fossil material and associated geological data by a qualified palaeontologist, appointed by the developer, may be required by the relevant heritage regulatory authorities. Any fossil material collected should be curated within an approved repository (museum / university fossil collection) by a qualified palaeontologist. These recommendations should be included within the Environmental Management Programme for the proposed alternative energy project.

7. ACKNOWLEDGEMENTS

Mr Justin Bowers of Ecoleges Environmental Consultants, Machadodorp is thanked for commissioning this study and for providing the relevant background information. As always, the logistical support and effective assistance of Ms Madelon Tusenius in the field is very much appreciated.

Table 2: Proposed monitoring and mitigation measures for incorporation into the EMP for the Soventix solar PV project

No.	Potential Impacts	Desired Outcomes	Targets & Indicators	Management Actions & Mitigation Measures	Responsibility	Timeframe / Frequency	Monitoring
FOSSIL HERITAGE RESOURCES							
	Disturbance, destruction or damage to fossils preserved at or below surface through surface clearance and excavations during construction phase .	Avoidance of palaeontologically sensitive areas (riverine alluvium). Reporting of chance fossil finds to SAHRA.	Older (orange-brown) consolidated alluvial deposits along major water courses (e.g. Brakrivier) – see blue area in Fig. 30.	Ongoing monitoring for chance fossil finds within development footprint during construction phase. Substantial fossils (vertebrate bones, teeth, large blocks of petrified wood) to be safeguarded, preferably in situ, and reported to SAHRA for recording and sampling by professional palaeontologist.	ECO Developer to appoint palaeontologist following significant new fossil finds	Ongoing during construction phase.	Compliance to be verified by ECO.

8. REFERENCES

ABEL, O. 1935. Vorzeitliche Lebensspuren. xv+ 644 pp. Gustav Fischer, Jena.

ALMOND, J.E. 1998. Non-marine trace fossils from the western outcrop area of the Permian Ecca Group, southern Africa. Tercera Reunión Argentina de Icnología, Mar del Plata, 1998, Abstracts p. 3.

ALMOND, J.E. 2008a. Fossil record of the Loeriesfontein sheet area (1: 250 000 geological sheet 3018). Unpublished report for the Council for Geoscience, Pretoria, 32 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2008b. Palaeozoic fossil record of the Clanwilliam sheet area (1: 250 000 geological sheet 3218). Unpublished report for the Council for Geoscience, Pretoria, 49 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2009. Contributions to the palaeontology and stratigraphy of the Alexander Bay sheet area (1: 250 000 geological sheet 2816), 117 pp. Unpublished technical report prepared for the Council for Geoscience by Natura Viva cc, Cape Town.

ALMOND, J.E. 2010. Eskom Gamma-Omega 765kV transmission line: Phase 2 palaeontological impact assessment. Sector 1, Tanqua Karoo to Omega Substation (Western and Northern Cape Provinces), 95 pp + appendix. Natura Viva cc, Cape Town.

ALMOND, J.E. 2010a. Proposed windfarm at Maanhaarberg near De Aar, Northern Cape Province. Palaeontological impact assessment: desktop study, 21 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2010b. Proposed photovoltaic power generation facility at De Aar, Northern Cape Province. Palaeontological impact assessment: desktop study, 17 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2011. Proposed Mainstream Solar Park at De Aar, Northern Cape Province. Palaeontological impact assessment: desktop study, 17 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2012a. Two wind energy facilities on the Eastern Plateau near De Aar, Northern Cape Province, proposed by Mulilo Renewable Energy (Pty) Ltd. Palaeontological specialist study: combined desktop and field-based assessments, 55 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2012b. Proposed Mulilo Renewable Energy PV2, PV3 and PV4 photovoltaic energy facilities on Farms Paarde Valley, Badenhorst Dam and Annex Du Plessis Dam near De Aar, Northern Cape Province. Palaeontological specialist study: combined desktop and field-based assessments, 45 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2012c. Proposed solar power generation facilities on the remaining extent of the farm Vetlaagte No. 4, De Aar, Northern Cape Province. Palaeontological specialist study: combined desktop and field-based assessments, 32 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2013a. Proposed photovoltaic (solar) energy facilities on Badenhorst Dam Farm near De Aar, Northern Cape. Palaeontological specialist study: combined desktop and field-based assessments, 55 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2013b. Proposed photovoltaic (solar) energy facilities on Du Plessis Dam Farm near De Aar, Northern Cape. Palaeontological specialist study: combined desktop and field-based assessments, 44 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2013c. Proposed Renosterberg PV Solar Facility near De Aar, Northern Cape Province. Palaeontological specialist assessment: combined desktop and field study, 63 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2013d. Proposed 16 mtpa expansion of Transnet's existing manganese ore export railway line & associated infrastructure between Hotazel and the Port of Ngqura, Northern & Eastern Cape. Part 3: Kimberley to De Aar, Northern Cape. Palaeontological specialist assessment: combined field-based and desktop study, 65 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. & PETHER, J. 2008. Palaeontological heritage of the Northern Cape. Interim SAHRA technical report, 124 pp. Natura Viva cc., Cape Town.

ANDERSON, A.M. 1974. Arthropod trackways and other trace fossils from the Early Permian lower Karoo Beds of South Africa. Unpublished PhD thesis, University of Witwatersrand, Johannesburg, 172 pp.

ANDERSON, A.M. 1975. Turbidites and arthropod trackways in the Dwyka glacial deposits (Early Permian) of southern Africa. Transactions of the Geological Society of South Africa 78: 265-273.

ANDERSON, A.M. 1976. Fish trails from the Early Permian of South Africa. Palaeontology 19: 397-409, pl. 54.

ANDERSON, A.M. 1981. The *Umfolozia* arthropod trackways in the Permian Dwyka and Ecca Groups of South Africa. Journal of Paleontology 55: 84-108, pls. 1-4.

ANDERSON, A.M. & MCLACHLAN, I.R. 1976. The plant record in the Dwyka and Ecca Series (Permian) of the south-western half of the Great Karoo Basin, South Africa. Palaeontologia africana 19: 31-42.

ANDERSON, J.M. & ANDERSON, H.M. 1985. Palaeoflora of southern Africa. Prodomus of South African megafloras, Devonian to Lower Cretaceous, 423 pp. Botanical Research Institute, Pretoria & Balkema, Rotterdam.

BAMFORD, M. 1999. Permo-Triassic fossil woods from the South African Karoo Basin. Palaeontologia africana 35, 25-40.

BAMFORD, M.K. 2004. Diversity of woody vegetation of Gondwanan southern Africa. Gondwana Research 7, 153-164.

BENDER, P.A. 2004. Late Permian actinopterygian (palaeoniscid) fishes from the Beaufort Group, South Africa: biostratigraphic and biogeographic implications. Bulletin 135, 84pp. Council for Geoscience, Pretoria.

BENDER, P.A., RUBIDGE, B.S., GARDINER, B.S., LOOCK, J.C. & BREMNER, A.T. 1991. The stratigraphic range of the palaeoniscoid fish *Namaichthys digitata* in rocks of the Karoo sequence and its palaeoenvironmental significance. South African Journal of Science 87: 468-469.

BENDER, P.A. & BRINK, J.S. 1992. A preliminary report on new large mammal fossil finds from the Cornelia-Uitzoek site. South African Journal of Science 88: 512-515.

BENTON, M.J. 2003. When life nearly died. The greatest mass extinction of them all, 336 pp. Thames & Hudson, London.

BOUSMAN, C.B. *et al.* 1988. Palaeoenvironmental implications of Late Pleistocene and Holocene valley fills in Blydefontein Basin, Noupoot, C.P., South Africa. *Palaeoecology of Africa* 19: 43-67.

BAMFORD, M. 1999. Permo-Triassic fossil woods from the South African Karoo Basin. *Palaeontologia africana* 35, 25-40.

BAMFORD, M.K. 2004. Diversity of woody vegetation of Gondwanan southern Africa. *Gondwana Research* 7, 153-164.

BRADDY, S.J. & BRIGGS, D.E.G. 2002. New Lower Permian nonmarine arthropod trace fossils from New Mexico and South Africa. *Journal of Paleontology* 76: 546-557.

BRINK, J.S. 1987. The archaeozoology of Florisbad, Orange Free State. *Memoirs van die Nasionale Museum* 24, 151 pp.

BRINK, J.S. *et al.* 1995. A new find of *Megalotragus priscus* (Alcephalini, Bovidae) from the Central Karoo, South Africa. *Palaeontologia africana* 32: 17-22.

BUATOIS, L. & MANGANO, M.G. 2004. Animal-substrate interactions in freshwater environments: applications of ichnology in facies and sequence stratigraphic analysis of fluvio-lacustrine successions. In: McIlroy, D. (Ed.) *The application of ichnology to palaeoenvironmental and stratigraphic analysis*. Geological Society, London, Special Publications 228, pp 311-333.

BUATOIS, L.A. & MANGANO, M.G. 2007. Invertebrate ichnology of continental freshwater environments. In: Miller, W. III (Ed.) *Trace fossils: concepts, problems, prospects*, pp. 285-323. Elsevier, Amsterdam.

CEDAR TOWER SERVICES 2015, Heritage screener: Solar energy facility east of De Aar, Northern Cape, South Africa (Solaris Power Ltd), 28 pp.

CHURCHILL, S.E. *et al.* 2000. Erfkroon: a new Florisian fossil locality from fluvial contexts in the western Free State, South Africa. *South African Journal of Science* 96: 161-163.

CLUVER, M.A. 1978. Fossil reptiles of the South African Karoo. 54pp. South African Museum, Cape Town.

COLE, D.I., SMITH, R.M.H. & WICKENS, H. DE V. 1990. Basin-plain to fluvio-lacustrine deposits in the Permian Ecca and Lower Beaufort Groups of the Karoo Sequence. *Guidebook Geocongress '90*, Geological Society of South Africa, PO2, 1-83.

COLE, D.I. & VORSTER, C.J. 1999. The metallogeny of the Sutherland area, 41 pp. Council for Geoscience, Pretoria.

COLE, D.I., NEVELING, J., HATTINGH, J., CHEVALLIER, L.P., REDDERING, J.S.V. & BENDER, P.A. 2004. The geology of the Middelburg area. *Explanation to 1: 250 000 geology Sheet 3124 Middelburg*, 44 pp. Council for Geoscience, Pretoria.

COLE, D. & SMITH, R. 2008. Fluvial architecture of the Late Permian Beaufort Group deposits, S.W. Karoo Basin: point bars, crevasse splays, palaeosols, vertebrate fossils and uranium. *Field Excursion FT02 guidebook*, AAPG International Conference, Cape Town October 2008, 110 pp.

- COOKE, H.B.S. 1974. The fossil mammals of Cornelia, O.F.S., South Africa. In: Butzer, K.W., Clark, J.D. & Cooke, H.B.S. (Eds.) The geology, archaeology and fossil mammals of the Cornelia Beds, O.F.S. *Memoirs of the National Museum, Bloemfontein* 9: 63-84.
- COOPER, M.R. & KENSLEY, B. 1984. Endemic South American Permian bivalve molluscs from the Ecca of South Africa. *Journal of Paleontology* 58: 1360-1363.
- DE BEER, C.H., GRESSE, P.G., THERON, J.N. & ALMOND, J.E. 2002. The geology of the Calvinia area. Explanation to 1: 250 000 geology Sheet 3118 Calvinia. 92 pp. Council for Geoscience, Pretoria.
- COLE, D.I., NEVELING, J., HATTINGH, J., CHEVALLIER, L.P., REDDERING, J.S.V. & BENDER, P.A. 2004. The geology of the Middelburg area. Explanation to 1: 250 000 geological sheet 3124 Middelburg, 43 pp. Council for Geoscience, Pretoria.
- DUNCAN, A.R. & MARSH, J.S. 2006. The Karoo Igneous Province. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) *The geology of South Africa*, pp. 501-520. Geological Society of South Africa, Marshalltown.
- GROENEWALD, G. 2012. Palaeontological desktop assessment of the proposed development of the Eastern Plateau and Maanhaarberg 132 kV power lines at De Aar in the Northern Cape, 8 pp. Metsi Metseng Geological & Environmental Services, Bethlehem.
- HOLMES, P.J. & MARKER, M.E. 1995. Evidence for environmental change from Holocene valley fills from three central Karoo upland sites. *South African Journal of Science* 91: 617-620.
- JOHNSON, M.R. 1966. The stratigraphy of the Cape and Karoo Systems in the Eastern Cape Province. Unpublished MSc Thesis, Rhodes University, Grahamstown.
- JOHNSON, M.R. 1976. Stratigraphy and sedimentology of the Cape and Karoo sequences in the Eastern Cape Province. Unpublished PhD thesis, Rhodes University, Grahamstown, xiv + 335 pp, 1pl.
- JOHNSON, M.R., VAN VUUREN, C.J., VISSER, J.N.J., COLE, D.I., De V. WICKENS, H., CHRISTIE, A.D.M., ROBERTS, D.L. & BRANDL, G. 2006. Sedimentary rocks of the Karoo Supergroup. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) *The geology of South Africa*, pp. 461-499. Geological Society of South Africa, Marshalltown.
- KEYSER, A.W. & SMITH, R.M.H. 1977-78. Vertebrate biozonation of the Beaufort Group with special reference to the Western Karoo Basin. *Annals of the Geological Survey of South Africa* 12: 1-36.
- KITCHING, J.W. 1977. The distribution of the Karoo vertebrate fauna, with special reference to certain genera and the bearing of this distribution on the zoning of the Beaufort beds. *Memoirs of the Bernard Price Institute for Palaeontological Research, University of the Witwatersrand, No. 1*, 133 pp (incl. 15 pls).
- KLEIN, R.G. 1984. The large mammals of southern Africa: Late Pliocene to Recent. In: Klein, R.G. (Ed.) *Southern African prehistory and paleoenvironments*, pp 107-146. Balkema, Rotterdam.
- LE ROUX, F.G. 1993. Die geologie van die gebied Colesberg. Explanation to 1: 250 000 geology Sheet 3024, 12 pp. Council for Geoscience, Pretoria.
- LE ROUX, F.G. & KEYSER, A.W. 1988. Die geologie van die gebied Victoria-Wes. Explanation to 1: 250 000 geology Sheet 3122, 31 pp. Council for Geoscience, Pretoria.

- LUCAS, D.G. 2009. Global Middle Permian reptile mass extinction: the dinocephalian extinction event. *Geological Society of America Abstracts with Programs* 41, No. 7, p. 360.
- MACRAE, C. 1999. Life etched in stone. *Fossils of South Africa*, 305 pp. The Geological Society of South Africa, Johannesburg.
- MEADOWS, M.E. & WATKEYS, M.K. 1999. Palaeoenvironments. In: Dean, W.R.J. & Milton, S.J. (Eds.) *The karoo. Ecological patterns and processes*, pp. 27-41. Cambridge University Press, Cambridge.
- MILLSTEED, B.D. 2014. Full palaeontological heritage impact assessment report on a portion of a proposed win energy generation facility (the Castle Project), this being on the eastern extent of the farm Knapdaar 8 near de Aar, Northern Cape Province, 45 pp.
- NEL, L. 1977. Die geologie van die gebied suid van Hopetown. Unpublished PhD thesis, University of the Free State, 171 pp.
- NICOLAS, M.V. 2007. Tetrapod diversity through the Permo-Triassic Beaufort Group (Karoo Supergroup) of South Africa. Unpublished PhD thesis, University of Witwatersrand, Johannesburg.
- NICOLAS, M. & RUBIDGE, B.S. 2010. Changes in Permo-Triassic terrestrial tetrapod ecological representation in the Beaufort Group (Karoo Supergroup) of South Africa. *Lethaia* 43, 45-59.
- PARTRIDGE, T.C. & SCOTT, L. 2000. Lakes and pans. In: Partridge, T.C. & Maud, R.R. (Eds.) *The Cenozoic of southern Africa*, pp.145-161. Oxford University Press, Oxford.
- PARTRIDGE, T.C., BOTHA, G.A. & HADDON, I.G. 2006. Cenozoic deposits of the interior. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) *The geology of South Africa*, pp. 585-604. Geological Society of South Africa, Marshalltown.
- PRINSLOO, M.C. 1989. Die geologie van die gebied Britstown. Explanation to 1: 250 000 geology Sheet 3022 Britstown, 40 pp. Council for Geoscience, Pretoria.
- RETALLACK, G.J., METZGER, C.A., GREAVER, T., HOPE JAHREN, A., SMITH, R.M.H. & SHELDON, N.D. 2006. Middle – Late Permian mass extinction on land. *GSA Bulletin* 118, 1398-1411.
- RUBIDGE, B.S. (Ed.) 1995. Biostratigraphy of the Beaufort Group (Karoo Supergroup). South African Committee for Biostratigraphy, Biostratigraphic Series No. 1., 46 pp. Council for Geoscience, Pretoria.
- RUBIDGE, B.S. 2005. Re-uniting lost continents – fossil reptiles from the ancient Karoo and their wanderlust. 27th Du Toit Memorial Lecture. *South African Journal of Geology* 108, 135-172.
- RUBIDGE, B.S., HANCOX, P.J. & CATUNEANU, O. 2000. Sequence analysis of the Ecca-Beaufort contact in the southern Karoo of South Africa. *South African Journal of Geology* 103, 81-96.
- RUBIDGE, B.S., ERWIN, D.H., RAMEZANI, J., BOWRING, S.A. & DE KLERK, W.J. 2010. The first radiometric dates for the Beaufort Group, Karoo Supergroup of South Africa. Proceedings of the 16th conference of the Palaeontological Society of Southern Africa, Howick, August 5-8, 2010, pp. 82-83.
- RUST, I.C., SHONE, R.W. & SIEBRITS, L.B. 1991. Carnarvon Formasie: golf-oorheesde sedimentasie in 'n vlak Karoosee. *South African Journal of Science* 87, 198-202.

- RYAN, P.J. 1967. Stratigraphic and palaeocurrent analysis of the Ecca Series and lowermost Beaufort Beds in the Karoo Basin of South Africa. Unpublished PhD thesis, University of the Witwatersrand, Johannesburg, 210 pp.
- SAHRA 2013. Minimum standards: palaeontological component of heritage impact assessment reports, 15 pp. South African Heritage Resources Agency, Cape Town.
- SCOTT, L. 2000. Pollen. In: Partridge, T.C. & Maud, R.R. (Eds.) The Cenozoic of southern Africa, pp.339-35. Oxford University Press, Oxford.
- SEILACHER, A. 2007. Trace fossil analysis, xiii + 226pp. Springer Verlag, Berlin.
- SIEBRITS, L.B. 1987. Die sedimentology van die Formasie Carnarvon in die omgewing van Carnarvon. Unpublished MSc thesis, University of Port Elizabeth, v + 92 pp.
- SIEBRITS, L.B. 1989. Die geologie van die gebied Sakrivier. Explanation of 1: 250 000 geology sheet 3020, 19 pp. Council for Geoscience, Pretoria.
- SKEAD, C.J. 1980. Historical mammal incidence in the Cape Province. Volume 1: The Western and Northern Cape. 903pp. Department of Nature and Environmental Conservation, Cape Town.
- SKINNER, E.M.W. & TRUSWELL, J.F. 2006. Kimberlites. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) The geology of South Africa, pp. 651-659. Geological Society of South Africa, Marshalltown.
- SMITH, A.M. & ZAWADA, P.K. 1988. The Ecca-Beaufort transition zone near Philipstown, Cape Province: a marine shelf sequence. South African Journal of Geology 91, 75-82.
- SMITH, A.M. & ZAWADA, P.K. 1989. Permian storm current-produced offshore bars from an ancient shelf sequence: northwestern Karoo basin, Republic of South Africa. Journal of African Earth Sciences 9, 363-370.
- SMITH, A.B. 1999. Hunters and herders in the Karoo landscape. Chapter 15 in Dean, W.R.J. & Milton, S.J. (Eds.) The Karoo; ecological patterns and processes, pp. 243-256. Cambridge University Press, Cambridge.
- SMITH, R.M.H. 1980. The lithology, sedimentology and taphonomy of flood-plain deposits of the Lower Beaufort (Adelaide Subgroup) strata near Beaufort West. Transactions of the Geological Society of South Africa 83, 399-413.
- SMITH, R.M.H. 1993. Sedimentology and ichnology of floodplain paleosurfaces in the Beaufort Group (Late Permian), Karoo Sequence, South Africa. Palaios 8, 339-357.
- SMITH, R.M.H. & KEYSER, A.W. 1995. Biostratigraphy of the *Tapinocephalus* Assemblage Zone. Pp. 8-12 in Rubidge, B.S. (ed.) Biostratigraphy of the Beaufort Group (Karoo Supergroup). South African Committee for Stratigraphy, Biostratigraphic Series No. 1. Council for Geoscience, Pretoria.
- SMITH, R.M.H. & ALMOND, J.E. 1998. Late Permian continental trace assemblages from the Lower Beaufort Group (Karoo Supergroup), South Africa. Abstracts, Tercera Reunión Argentina de Icnología, Mar del Plata, 1998, p. 29.
- SMITH, R., RUBIDGE, B. & VAN DER WALT, M. 2012. Therapsid biodiversity patterns and paleoenvironments of the Karoo Basin, South Africa. Chapter 2 pp. 30-62 in Chinsamy-

- Turan, A. (Ed.) Forerunners of mammals. Radiation, histology, biology. xv + 330 pp. Indiana University Press, Bloomington & Indianapolis.
- STEAR, W.M. 1978. Sedimentary structures related to fluctuating hydrodynamic conditions in flood plain deposits of the Beaufort Group near Beaufort West, Cape. Transactions of the Geological Society of South Africa 81, 393-399.
- THERON, J.N. 1983. Die geologie van die gebied Sutherland. Explanation of 1: 250 000 geological Sheet 3220, 29 pp. Council for Geoscience, Pretoria.
- VAN DER WALT M., DAY M., RUBIDGE B., COOPER A.K. & NETTERBERG I. 2010. A new GIS-based biozone map of the Beaufort Group (Karoo Supergroup), South Africa. *Palaeontologia Africana*, 45, 5pp.
- VAN DIJK, D.E., CHANNING, A. & VAN DEN HEEVER, J.A. 2002. Permian trace fossils attributed to tetrapods (Tierberg Formation, Karoo Basin, South Africa). *Palaeontologia africana* 38: 49-56.
- VILJOEN, J.H.A. 1989. Die geologie van die gebied Williston. Explanation to geology sheet 3120 Williston, 30 pp. Council for Geoscience, Pretoria.
- VILJOEN, J.H.A. 1992. Lithostratigraphy of the Collingham Formation (Ecca Group), including the Zoute Kloof, Buffels River and Wilgehout River Members and the Matjiesfontein Chert Bed. South African Committee for Stratigraphy, Lithostratigraphic Series No. 22, 10 pp.
- VILJOEN, J.H.A. 1994. Sedimentology of the Collingham Formation, Karoo Supergroup. *South African Journal of Geology* 97: 167-183.
- VILJOEN, J.H.A. 2005. Tierberg Formation. SA Committee for Stratigraphy, Catalogue of South African Lithostratigraphic Units 8: 37-40.
- VISSER, J.N.J. & LOOCK, J.C. 1974. The nature of the Ecca-Beaufort transition in the western and central Orange Free State. Transactions of the Geological Society of South Africa 77, 371-372.
- WELLS, L.H. & COOKE, H.B.S. 1942. The associated fauna and culture of Vlakkraal thermal springs, O.F.S.; III, the faunal remains. Transactions of the Royal Society of South Africa 29: 214-232.
- WERNER, M. 2006. The stratigraphy, sedimentology and age of the Late Palaeozoic Mesosaurus Inland Sea, SW-Gondwana: new implications from studies on sediments and altered pyroclastic layers of the Dwyka and Ecca Group (lower Karoo Supergroup) in southern Namibia. Dr rer. nat. thesis, University of Würzburg, 428 pp, 167 figs, 1 table.
- WICKENS, H. DE V. 1980. Verslag oor kartering in die Calvinia gebied. Unpublished report, Council for Geoscience, Pretoria, 19 pp.
- WICKENS, H. DE V. 1984. Die stratigraphie en sedimentologie van die Group Ecca wes van Sutherland. Unpublished MSc thesis, University of Port Elizabeth, viii + 86 pp.
- WICKENS, H. DE V. 1992. Submarine fans of the Permian Ecca Group in the SW Karoo Basin, their origin and reflection on the tectonic evolution of the basin and its source areas. In: De Wit, M.J. & Ransome, I.G.D. (Eds.) Inversion tectonics of the Cape Fold Belt, Karoo and Cretaceous Basins of southern Africa, pp. 117-126. Balkema, Rotterdam.
- WICKENS, H. DE V. 1994. Submarine fans of the Ecca Group. Unpublished PhD thesis, University of Port Elizabeth. 350 pp.

WICKENS, H. DE V. 1996. Die stratigraphie en sedimentologie van die Ecca Groep wes van Sutherland. Council for Geosciences, Pretoria Bulletin 107, 49pp.

ZAWADA, P.K. 1992. The geology of the Koffiefontein area. Explanation of 1: 250 000 geology sheet 2924, 30 pp. Council for Geoscience, Pretoria.

9. QUALIFICATIONS & EXPERIENCE OF THE AUTHOR

Dr John Almond has an Honours Degree in Natural Sciences (Zoology) as well as a PhD in Palaeontology from the University of Cambridge, UK. He has been awarded post-doctoral research fellowships at Cambridge University and in Germany, and has carried out palaeontological research in Europe, North America, the Middle East as well as North and South Africa. For eight years he was a scientific officer (palaeontologist) for the Geological Survey / Council for Geoscience in the RSA. His current palaeontological research focuses on fossil record of the Precambrian - Cambrian boundary and the Cape Supergroup of South Africa. He has recently written palaeontological reviews for several 1: 250 000 geological maps published by the Council for Geoscience and has contributed educational material on fossils and evolution for new school textbooks in the RSA.

Since 2002 Dr Almond has also carried out palaeontological impact assessments for developments and conservation areas in the Western, Eastern and Northern Cape, Mpumalanga, Free State, Limpopo, Northwest and Kwazulu-Natal under the aegis of his Cape Town-based company *Natura Viva* cc. He has been a long-standing member of the Archaeology, Palaeontology and Meteorites Committee for Heritage Western Cape (HWC) and an advisor on palaeontological conservation and management issues for the Palaeontological Society of South Africa (PSSA), HWC and SAHRA. He is currently compiling technical reports on the provincial palaeontological heritage of Western, Northern and Eastern Cape for SAHRA and HWC. Dr Almond is an accredited member of PSSA and APHP (Association of Professional Heritage Practitioners – Western Cape).

Declaration of Independence

I, John E. Almond, declare that I am an independent consultant and have no business, financial, personal or other interest in the proposed development project, application or appeal in respect of which I was appointed other than fair remuneration for work performed in connection with the activity, application or appeal. There are no circumstances that compromise the objectivity of my performing such work.



**Dr John E. Almond,
Palaeontologist,
Natura Viva cc**

APPENDIX: GPS LOCALITY DATA

All GPS readings were taken in the field using a hand-held Garmin GPSmap 60CSx

001	S30° 50' 29.1" E24° 18' 47.7"	Goedehoop 26. Extensive shallow borrow pit excavated into grey-green and subordinate purple-brown mudrocks of Adelaide Subgroup. Well-indurated (possibly baked). Well-developed pedogenic calcrete horizons with rusty-brown to pinkish concretions, some showing septarian cracking. Fine-grained, grey-green sandstones / wackes, ripple cross-laminated to massive. Bedrocks near-surface with calcrete veins and overlain by orange-brown sandy soils and surface gravels (dolerite, sandstone, wacke, hornfels, occasional sparry calcite). Bare soil patches in region with fine sheetwash gravels of hornfels (some flaked but patinated, rounded).
002	S30° 50' 37.1" E24° 19' 22.5"	Goedehoop 26. Carpet of surface gravels on low hillslopes, dominated by patinated hornfels (occasionally flaked), locally also quartzite. Baked, well-jointed greyish quartzite in vicinity of dolerite intrusion.
003	S30° 50' 24.6" E24° 20' 13.6"	Goedehoop 26. Major ridge-forming dolerite intrusion with scree of coarse dolerite rubble on flanks. Rafts (xenoliths) of baked quartzite incorporated into dolerite magma.
004	S30° 50' 23.9" E24° 20' 17.0"	Goedehoop 26. Pale brown, medium-grained, medium-bedded channel sandstone with sparse, large (several dm diam.) oblate ferruginous carbonate concretions (<i>koffieklip</i>).
005	S30° 50' 21.2" E24° 20' 09.6"	Goedehoop 26. Ridge of baked, convolute-laminated sandstone with a wood-like, streaky and foliated fabric, a consequence of remobilisation and metamorphism of Adelaide Subgroup country rocks during dolerite intrusion.
006	S30° 49' 36.3" E24° 21' 48.0"	Goedehoop 26. Bare patch of soils and sheetwash gravels (no petrified wood seen). Occasional flaked artefacts of hornfels and quartzite.
007	S30° 50' 17.8" E24° 21' 14.7"	Goedehoop 26. Extensive horizon of dark brown, sandy, ferruginous carbonate <i>koffieklip</i> with lenses of greenish mudflakes associated with problematic rounded to elongate, speckled, coarse-grained inclusions, some perhaps with a mudrock core. Possibly biogenic (e.g. burrows) but are currently unidentified. Provisional Field rating IIC Local Resource.
008	S30° 51' 04.9" E24° 20' 01.2"	Goedehoop 26. Area with doleritic surface rubble as well as local development of pale creamy, nodular calcrete hardpan beneath orange-brown sandy soil.
009	S30° 50' 38.6" E24° 19' 25.4"	Goedehoop 26. Surface carpeted with patinated, angular hornfels surface gravels (many flaked).
010	S30° 52' 38.9" E24° 19' 20.3"	Kwanselaarshoek 40. Area of low relief, with extensive exposure of pale brown silty alluvium, bare patches with sheetwash surface gravels (calcrete, dolerite, hornfels, palaeocalcrete nodules with septarian cracking).
011	S30° 53' 18.5" E24° 17' 39.7"	Rietfontain 39. Stream bank exposures of Adelaide Subgroup well-jointed, thin crevasse splay sandstones and hackly-weathering, grey-green overbank mudrocks, locally intruded, baked, deformed and disrupted by dolerite dyke.
012	S30° 52' 56.9" E24° 15' 48.4"	Kafferspoort 56. Extensive riverbank sections through pale brown silty modern alluvium (Holocene) with thin fine gravel lenses. Orange-brown, incipiently calcretised older alluvial deposits (probably Pleistocene) exposed away from the river banks by erosion gullies.
013	S30° 52' 59.1" E24° 15' 45.5"	Kafferspoort 56. Good gullied exposures of older (probably Pleistocene) consolidated, orange-brown alluvial deposits.
014	S30° 52' 58.8" E24° 15' 45.1"	Kafferspoort 56. Occasional to locally common small, angular blocks of fossil wood, showing good seasonal growth banding, eroding out of older calcretised orange-brown alluvium (occasional embedded blocks seen). Provisional Field rating IIC Local Resource. Abundant dark grey hornfels artefacts (patinated and fresh) at surface, probably downwasted from younger alluvial deposits.
015	S30° 52' 47.5" E24° 14' 48.0"	Kafferspoort 56. Bare soil patch with surface gravels of hornfels, dolerite.
016	S30° 52' 49.1" E24° 15' 04.7"	Kafferspoort 56. Good examples of blocky (ruiniform) dolerite weathering among koppies towards western edge of study area. Flatter grassy areas between koppies with sandy orange soils.

017	S30° 52' 42.3" E24° 15' 45.8"	Rietfontain 39. Occasional to locally common weathered-out small blocks of cherty fossil wood (up to 6 cm diam.) showing well-developed seasonal growth lines, overlying eroded older calcretised alluvium. Provisional Field rating IIIC Local Resource.
018	S30° 52' 40.2" E24° 15' 47.5"	Rietfontain 39. Good shallow Brakrivier riverine bank exposures of calcretised, well-consolidated older alluvial deposits underlying brownish, uncalcretised younger alluvial deposits.
019	S30° 52' 39.0" E24° 15' 47.7"	Rietfontain 39. Local concentration of flaked fresh, unpatinated hornfels artefacts overlying younger alluvial deposits. LSA upper grind stone at surface in this area.
020	S30° 52' 39.4" E24° 15' 47.8"	Rietfontain 39. Concentration of fossil wood blocks up to c. 10 cm across downwasted from older alluvium. Some blocks show marked colour banding due to seasonal growth pattern. Provisional Field rating IIIC Local Resource.
021	S30° 52' 26.2" E24° 16' 00.3"	Rietfontain 39. Banks of Brakrivier incised into younger alluvial deposits.
022	S30° 50' 58.0" E24° 16' 35.8"	Rietfontain 39. Banks of Brakrivier incised into younger alluvial deposits.