

BASIC ASSESSMENT FOR THE GUNSTFONTEIN BATTERY ENERGY STORAGE SYSTEM  
(BESS), NORTHERN CAPE PROVINCE:

**FAUNA, AVIFAUNA & FLORA BASIC ASSESSMENT SPECIALIST REPORT**



**PRODUCED FOR SAVANNAH ENVIRONMENTAL (Pty) Ltd**

**BY**



**[Simon.Todd@3foxes.co.za](mailto:Simon.Todd@3foxes.co.za)**

**September 2020**

**EXECUTIVE SUMMARY**

Gunstfontein Wind Farm (Pty) Ltd would like to provide for the installation of a Battery Energy Storage System (BESS) at the authorised Gunstfontein Wind Energy Facility (WEF) in the Northern Cape. The BESS is proposed to be located near to (within 500m of) the facility substation, and will be up to 5ha in total extent. Overhead or underground MV cabling (33kV or less) will connect the BESS to the substation. An area of ~500m around the boundary of the authorised WEF substation was assessed, to allow for the optimization of the placement of the BESS anywhere within the assessment region. Savannah Environmental is conducting the required Basic Assessment (BA) process for the Gunstfontein BESS and has appointed 3Foxes Biodiversity Solutions to provide a specialist terrestrial biodiversity (fauna and flora) impact assessment study of the proposed BESS.

A field assessment as well as a review of the available ecological information for the area was conducted. The vegetation within the Gunstfontein BESS 500m assessment region consists entirely of Roggeveld Shale Renosterveld which is considered to represent a moderately sensitive vegetation type due to its low total extent and relatively high abundance of plant Species of Conservation Concern (SCC). A low impact on plant SCC is expected to occur as a result of habitat loss associated with the development, but with the appropriate mitigation (pre-construction walk-through), this is highly unlikely to compromise the local populations of any species. In terms of fauna, there are few species of conservation concern that are likely to be present or abundant at the site and the primary impact of the development on fauna would be minor habitat loss for the more common resident species. As such, no high long-term post-mitigation impacts on fauna are expected to occur. Although several avifaunal species of concern are confirmed present in the area and likely use the BESS location for foraging at least on occasion, the extent of the development is sufficiently low such that it would not generate significant habitat loss for any species of concern. Furthermore, there are no nesting sites or other important habitats within the affected area, with the result that overall impact on avifauna would be low. Consequently, the impacts of the development on fauna and flora are considered acceptable and would be of low significance after mitigation.

Although direct impacts on fauna and flora are considered acceptable with mitigation, the BESS 500m assessment region falls within an area that has been classified as CBA 1. As these are areas that have been identified as being of significance for biodiversity maintenance and ecological processes, development in these areas is generally not preferred. However, the footprint of the development would be less than 5ha and would also be within close proximity to the approved Gunstfontein WEF substation, with the result that the additional extent of disturbance and habitat loss would be small. As a result, the low overall footprint of the BESS would be very unlikely to compromise the ecological functioning of the affected CBAs in any way.

Cumulative impacts within the broader study area are of potential concern due to the proliferation of WEF energy development in the wider Roggeveld area. The contribution of the BESS would however be very minor and is not considered to represent a significant contributor to cumulative impact in the area. Cumulative impacts associated with the development of the BESS are therefore considered acceptable.

*Ecological Impact Statement*

There are no impacts associated with the establishment of Gunstfontein BESS that cannot be mitigated to a low significance. Although cumulative impacts in the area are a concern due to the high density of wind energy developments in the area, the contribution of the Gunstfontein BESS would be low and is not considered to be of significance. As such, there are no fatal flaws or high post-mitigation impacts that should prevent the development from proceeding. Based on the location of the BESS as provided for this assessment and the sensitivity determined on site, the Gunstfontein BESS can be supported from a terrestrial and avifaunal ecology point of view at the current location, as well as any other region within the 500m assessment zone provided no infrastructure is placed within regions of high or very high sensitivity.

**CONTENTS**

Executive Summary .....2

Contents .....4

List of Figures .....6

Compliance with GNR 320 OF 20 MARCH 2020.....7

Short CV/Summary of Expertise – Simon Todd .....9

Specialist Declaration ..... 10

1 INTRODUCTION ..... 11

Scope of Study ..... 11

1.1 Assessment Approach & Philosophy ..... 13

1.2 Relevant Aspects of the Development ..... 15

2 METHODOLOGY ..... 16

2.1 Data Sourcing and Review ..... 16

2.2 Site Visit ..... 18

2.3 Sensitivity Mapping & Assessment ..... 18

2.4 Sampling Limitations and Assumptions ..... 19

3 DESCRIPTION OF THE AFFECTED ENVIRONMENT- BASELINE ..... 20

3.1 Broad-Scale Vegetation Description ..... 20

3.2 Fine-Scale Vegetation Description ..... 21

3.3 Listed and Protected Plant Species ..... 25

3.4 Faunal Communities ..... 26

3.5 Critical Biodiversity Areas & Broad-Scale Processes ..... 31

3.6 Current Baseline & Cumulative Impact ..... 34

3.7 Site Sensitivity Assessment ..... 34

4 IDENTIFICATION & NATURE OF IMPACTS ..... 35

4.1 Identification of Impacts ..... 36

5 ASSESSMENT OF IMPACTS ..... 37

5.1 Planning & Construction Phase ..... 37

5.2 Operational Phase Impacts ..... 41

5.3 Decommissioning Phase ..... 44

5.4 Cumulative Impacts ..... 46

6 CONCLUSION & RECOMMENDATIONS ..... 47

7 Activities for Inclusion into the EMPr ..... 49

8 REFERENCES ..... 53

Appendix 1. Listed Plant Species ..... 54

Appendix 2. List of Mammals ..... 56  
Appendix 3. List of Reptiles. .... 60  
Appendix 4. List of Amphibians ..... 62

**LIST OF FIGURES**

Figure 1. Map showing the location of the BESS 500m assessment area with the preliminary location of the BESS footprint adjacent to the substation within the Gunstfontein WEF area. .... 16

Figure 2. Broad-scale overview of the vegetation in and around the Gunstfontein substation and BESS study area. The vegetation map is an extract of the national vegetation map (Mucina & Rutherford 2006 & 2018 update) and also includes drainage lines delineated by the NFEPA assessment (Nel et al. 2011)..... 20

Figure 3. Ecological habitats observed within the Gunstfontein BESS 500m assessment region (entire frame) and which are described in detail below. .... 22

Figure 4. The typical Roggeveld Shale Renosterveld on shallow soils within the BESS 500m study region. This habitat is considered relatively low sensitivity and is considered acceptable for the location of the BESS..... 23

Figure 5. Roggeveld Shale Renosterveld on deeper soils within the BESS study area, with taller shrubs dominated by *Euryops lateriflorus*. .... 24

Figure 6. There are several areas of exposed bedrock within the BESS study area. Although there is very little vegetation associated with these areas and they are not considered important from a botanical point of view, they were observed to be a relatively important local habitat for reptiles and other fauna which prefer rocky habitats..... 24

Figure 7. Common reptiles observed at the site include, from top left, the Variegated Skink, Common Sand Lizard, Ground Agama and Karoo Girdled Lizard..... 28

Figure 8. Extract of the Northern Cape Critical Biodiversity Areas map for the broader study area around the Gunstfontein BESS, showing that the BESS occurs within an area classified as CBA 1..... 32

Figure 9. Sensitivity map for the Gunstfontein BESS 500m assessment region. .... 35

**COMPLIANCE WITH GNR 320 OF 20 MARCH 2020.**

| <b>Requirements of a Terrestrial Biodiversity Specialist Assessment as per GNR 320 of 20 March, 2020.</b>   | <b>Addressed in the Specialist Report</b> |
|---|---|
| 2.1. The assessment must be prepared by a specialist registered with the South African Council for Natural Scientific Professionals (SACNASP) with expertise in the field of terrestrial biodiversity.  | Page 9                                    |
| 2.2. The assessment must be undertaken on the preferred site and within the proposed development footprint.   | Chapter 1                                 |
| 2.3. The assessment must provide a baseline description of the site which includes, as a minimum, the following aspects:  | Section 3.5                               |
| 2.3.1. a description of the ecological drivers or processes of the system and how the proposed development will impact these;   | Section 3.5                               |
| 2.3.2. ecological functioning and ecological processes (e.g. fire, migration, pollination, etc.) that operate within the preferred site;  | Section 3.5                               |
| 2.3.3. the ecological corridors that the proposed development would impede including migration and movement of flora and fauna;   | Section 3.5                               |
| 2.3.4. the description of any significant terrestrial landscape features (including rare or important flora- faunal associations, presence of strategic water source areas (SWSAs) or freshwater ecosystem priority area (FEPA) sub catchments; | Chapter 3                                 |
| 2.3.5. a description of terrestrial biodiversity and ecosystems on the preferred site, including:   | Chapter 3                                 |
| (a) main vegetation types;  |   |
| (b) threatened ecosystems, including listed ecosystems as well as locally important habitat types identified;   | Chapter 3                                 |
| (c) ecological connectivity, habitat fragmentation, ecological processes and fine - scale habitats; and   | Chapter 3                                 |
| (d) species, distribution, important habitats (e.g. feeding grounds, nesting sites, etc.) and movement patterns identified;   | Section 3                                 |
| 2.3.6. the assessment must identify any alternative development footprints within the preferred site which would be of a low" sensitivity as identified by the screening tool and verified through the site sensitivity verification; and       | Section 3.7                               |
| 2.3.7. the assessment must be based on the results of a site inspection undertaken on the preferred site and must identify:   | Section 2.2                               |
| 2.3.7.1. terrestrial critical biodiversity areas (CBAs), including:   | Section 3.5                               |
| (a) the reasons why an area has been identified as a CBA;   |   |
| (b) an indication of whether or not the proposed development is consistent with maintaining the CBA in a natural or near natural state or in achieving the goal of rehabilitation;  | Section 3.5                               |
| (c) the impact on species composition and structure of vegetation with an indication of the extent of clearing activities in proportion to the remaining extent of the ecosystem type(s);   | Section 3.5                               |
| (d) the impact on ecosystem threat status;  | Section 3.5                               |
| (e) the impact on explicit subtypes in the vegetation;  | Section 3.2                               |
| (f) the impact on overall species and ecosystem diversity of the site; and  | Chapter 3                                 |
| (g) the impact on any changes to threat status of populations of species of conservation concern in the CBA;  | Section 3.5                               |
| 2.3.7.2. terrestrial ecological support areas (ESAs), including:  | N/A – CBA only, refer Section 3.5         |
| (a) the impact on the ecological processes that operate within or across the site;  |   |
| (b) the extent the proposed development will impact on the functionality of the ESA; and  | N/A – CBA only, refer Section 3.5         |
| (c) loss of ecological connectivity (on site, and in relation to the broader landscape) due to the degradation and severing of ecological corridors or introducing barriers that impede migration and movement of flora and fauna;              | N/A – CBA only, refer Section 3.5         |
| 2.3.7.3. protected areas as defined by the National Environmental Management: Protected Areas Act, 2004 including-  | Section 3.5, Table 1 onwards              |

|  |   |
|--|---|
| (a) an opinion on whether the proposed development aligns with the objectives or purpose of the protected area and the zoning as per the protected area management plan;                                 |   |
| 2.3.7.4. priority areas for protected area expansion, including-<br>(a) the way in which in which the proposed development will compromise or contribute to the expansion of the protected area network; | Section 3.5 and 4.1                                     |
| 2.3.7.5. SWSAs including:<br>(a) the impact(s) on the terrestrial habitat of a SWSA; and   | N/A – please refer section 3.5                          |
| (b) the impacts of the proposed development on the SWSA water quality and quantity (e.g. describing potential increased runoff leading to increased sediment load in water courses);                     | N/A – please refer section 3.5                          |
| 2.3.7.6. FEPA sub catchments, including-<br>(a) the impacts of the proposed development on habitat condition and species in the FEPA sub catchment;  | N/A – please refer section 3.5                          |
| 2.3.7.7. indigenous forests, including:  | N/A – no indigenous forests within 500m assessment zone |
| (a) impact on the ecological integrity of the forest; and  | N/A – no indigenous forests within 500m assessment zone |
| (b) percentage of natural or near natural indigenous forest area lost and a statement on the implications in relation to the remaining areas   | N/A – no indigenous forests within 500m assessment zone |
| 2.4. The findings of the assessment must be written up in a Terrestrial Biodiversity Specialist Assessment Report.   | This report   |



**SHORT CV/SUMMARY OF EXPERTISE – SIMON TODD**

|  |  |  |
|--|--|--|
|  <p><b>3Foxes Biodiversity Solutions</b><br/><b>ECOLOGICAL SPECIALIST SERVICES</b><br/>Assessment/Management/Research</p> | <p>Simon Todd Pr.Sci.Nat<br/>Director &amp; Principle Scientist<br/>C: 082 3326502<br/>O: 021 782 0377<br/>Simon.Todd@3foxes.co.za<br/>60 Forrest Way<br/>Glencairn<br/>7975</p> | <p>Ecological Solutions for<br/>People &amp; the Environment</p> |
|--|--|--|

Simon Todd is Director and principal scientist at 3Foxes Biodiversity Solutions and has over 20 years of experience in biodiversity measurement, management and assessment. He has provided specialist ecological input on more than 200 different developments distributed widely across the country. This includes input on the Wind and WEF SEA (REDZ) as well as the Eskom Grid Infrastructure (EGI) SEA and Karoo Shale Gas SEA. He is on the National Vegetation Map Committee as representative of the Nama and Succulent Karoo Biomes. Simon Todd is a recognised ecological expert and is a past chairman and current deputy chair of the Arid-Zone Ecology Forum. He is registered with the South African Council for Natural Scientific Professions (No. 400425/11).

A selection of recent work is as follows:

**Strategic Environmental Assessments**

- Co-Author. Chapter 7 – Biodiversity & Ecosystems - Shale Gas SEA. CSIR 2016.
- Co-Author. Chapter 1 – Scenarios and Activities – Shale Gas SEA. CSIR 2016.
- Co-Author – Ecological Chapter – Wind and Solar SEA. CSIR 2014.
- Co-Author – Ecological Chapter – Eskom Grid Infrastructure SEA. CSIR 2015.
- Contributor – Ecological & Conservation components to SKA SEA. CSIR 2017.


**Recent Specialist Ecological Studies in the Vicinity of the Current Site**

- Esizayo Wind Energy Facility, Roggeveld. WSP 2017.
- Maralla East & Maralla West WEFS, Roggeveld. WSP. 2017.
- Gunstfontein Wind Energy Facility, Sutherland. Savannah Environmental. 2016.
- Brandvalley Wind Energy Facility, Roggeveld. EOH. 2016.
- Kareebosch Wind Energy Facility, Roggeveld. Savannah Environmental 2015.
- Roggeveld Wind Energy Facility. 2013.
- Komsberg East & Komsberg West WEFS. Arcus Consulting. 2016

**SPECIALIST DECLARATION**

I, ..Simon Todd....., as the appointed independent specialist, in terms of the 2014 EIA Regulations, hereby declare that I:

- 
- I act as the independent specialist in this application;
- I perform the work relating to the application in an objective manner, even if this results in views and findings that are not favourable to the applicant;
- regard the information contained in this report as it relates to my specialist input/study to be true and correct, and do not have and will not have any financial interest in the undertaking of the activity, other than remuneration for work performed in terms of the NEMA, the Environmental Impact Assessment Regulations, 2014 and any specific environmental management Act;
- I declare that there are no circumstances that may compromise my objectivity in performing such work;
- I have expertise in conducting the specialist report relevant to this application, including knowledge of the Act, Regulations and any guidelines that have relevance to the proposed activity;
- I will comply with the Act, Regulations and all other applicable legislation;
- I have no, and will not engage in, conflicting interests in the undertaking of the activity;
- I have no vested interest in the proposed activity proceeding;
- I undertake to disclose to the applicant and the competent authority all material information in my possession that reasonably has or may have the potential of influencing - any decision to be taken with respect to the application by the competent authority; and - the objectivity of any report, plan or document to be prepared by myself for submission to the competent authority;
- I have ensured that information containing all relevant facts in respect of the specialist input/study was distributed or made available to interested and affected parties and the public and that participation by interested and affected parties was facilitated in such a manner that all interested and affected parties were provided with a reasonable opportunity to participate and to provide comments on the specialist input/study;
- I have ensured that the comments of all interested and affected parties on the specialist input/study were considered, recorded and submitted to the competent authority in respect of the application;
- all the particulars furnished by me in this specialist input/study are true and correct; and
- I realise that a false declaration is an offence in terms of regulation 48 and is punishable in terms of section 24F of the Act.

Signature of the specialist:  \_\_\_\_\_

Name of Specialist: \_\_\_\_ Simon Todd \_\_\_\_\_

Date: \_\_\_\_ 30 September 2020 \_\_\_\_\_

# **1 INTRODUCTION**

---

Gunstfontein Wind Farm (Pty) Ltd would like to provide for the installation of a Battery Energy Storage System (BESS) at the authorised Gunstfontein WEF in the Northern Cape. The BESS is proposed to be located near to (within 500m of) the facility substation, and will be approximately 5a in total extent. Overhead or underground MV cabling (33kV or less) will connect the BESS to the substation. An area of ~500m around the boundary of the WEF substation was assessed, to allow for the optimization of the placement of the BESS. A Basic Assessment process is therefore required for Environmental Authorisation, requiring a Terrestrial Biodiversity Specialist Assessment as per GNR 320 of 20 March 2020.

The purpose of the Gunstfontein Terrestrial Biodiversity Specialist Assessment is to describe and detail the ecological features of the proposed BESS site, provide an assessment of the ecological sensitivity of the affected area, and identify the likely impacts associated with the development of the proposed BESS infrastructure. A field assessment and desktop review of the available ecological information for the proposed BESS site and adjacent areas was conducted in order to identify and characterise the ecological features of the affected area. Impacts are assessed for the pre-construction, construction, operation, and decommissioning phases of the development. A variety of avoidance and mitigation measures associated with each identified impact are recommended to reduce the likely impact of the development, which should be included in the Environmental Management Programme (EMPr) for the development. The full scope of study is detailed below.

## ***SCOPE OF STUDY***

The scope of the study includes the following activities, as it relates to the minimum requirements of GNR 320 of 20 March 2020 (please refer page 6 and 7 for a detailing of the contents of this report against the protocol requirements).

- a description of the environment that may be affected by the activity and the manner in which the environment may be affected by the proposed project.
- a description and evaluation of environmental issues and potential impacts (incl. using direct, indirect and cumulative impacts) that have been identified.
- a statement regarding the potential significance of the identified issues based on the evaluation of the issues/impacts.
- an indication of the methodology used in determining the significance of potential environmental impacts.
- an assessment of the significance of direct, indirect, and cumulative impacts in terms of the following criteria:
  - the nature of the impact, which shall include a description of what causes the effect, what will be affected, and how it will be affected

- the extent of the impact, indicating whether the impact will be local (limited to the immediate area or site of development), regional, national or international
  - the duration of the impact, indicating whether the lifetime of the impact will be of a short-term duration (0-5 years), medium-term (5- 15 years), long-term (> 15 years, where the impact will cease after the operational life of the activity), or permanent
  - the probability of the impact, describing the likelihood of the impact actually occurring, indicated as improbable (low likelihood) probable (distinct possibility), highly probable (most likely), or definite (impact will occur regardless of any preventable measures)
  - the severity/beneficial scale indicating whether the impact will be very severe/beneficial (a permanent change which cannot be mitigated/permanent and significant benefit with no real alternative to achieving this benefit), severe/beneficial (long-term impact that could be mitigated/long-term benefit), moderately severe/beneficial (medium- to long-term impact that could be mitigated/ medium- to long-term benefit), slight, or have no effect
  - the significance which shall be determined through a synthesis of the characteristics described above and can be assessed as low medium or high
  - the status which will be described as either positive, negative or neutral
  - the degree to which the impact can be reversed
  - the degree to which the impact may cause irreplaceable loss of resources
  - the degree to which the impact can be mitigated
- a description and comparative assessment of all alternatives (where applicable)
  - recommendations regarding practical mitigation measures for potentially significant impacts, for inclusion in the EMPr.
  - an indication of the extent to which the issue could be addressed by the adoption of mitigation measures.
  - a description of any assumptions, uncertainties and gaps in knowledge.
  - an environmental impact statement which contains:
    - a summary of the key findings of the environmental impact assessment;
    - an assessment of the positive and negative implications of the proposed activity;
    - a comparative assessment of the positive and negative implications of identified alternatives.

**General Considerations:**

- Disclose any gaps in information or assumptions made.
- Identify recommendations for mitigatory measures to minimise impacts.
- Outline additional management guidelines.

- Provide monitoring requirements, mitigation measures and recommendations in a table format as input into the Environmental Management Programme (EMPr) for faunal related issues.

A description of the potential impacts of the development and recommended mitigation measures are to be provided, which will be separated into the following project phases:

- Pre-construction;
- Construction;
- Operation Phase; and
- Decommissioning.

### **1.1 ASSESSMENT APPROACH & PHILOSOPHY**

This assessment is conducted according to the 2014 EIA Regulations (Government Notice Regulation 326) in terms of the National Environmental Management Act (Act 107 of 1998) as amended (NEMA), as well as best-practice guidelines and principles for biodiversity assessment as outlined by Brownlie (2005) and De Villiers et al. (2005). This includes adherence to the following broad principles:

- That a precautionary and risk-averse approach be adopted towards projects which may result in substantial detrimental impacts on biodiversity and ecosystems, especially the irreversible loss of habitat and ecological functioning in threatened ecosystems or designated sensitive areas: i.e. Critical Biodiversity Areas (as identified by systematic conservation plans, Biodiversity Sector Plans or Bioregional Plans) and Freshwater Ecosystem Priority Areas.
- Demonstrate how the proponent intends on complying with the principles contained in section 2 of the National Environmental Management Act, 1998 (Act No. 107 of 1998), as amended (NEMA), which, amongst other things, indicates that environmental management should:
  - In order of priority aim to: avoid, minimise or remedy disturbance of ecosystems and loss of biodiversity;
  - Avoid degradation of the environment;
  - Avoid jeopardising ecosystem integrity;
  - Pursue the best practicable environmental option by means of integrated environmental management;
  - Protect the environment as the people's common heritage;
  - Control and minimise environmental damage; and
  - Pay specific attention to management and planning procedures pertaining to sensitive, vulnerable, highly dynamic or stressed ecosystems.

These principles serve as guidelines for all decision-making concerning matters that may affect the environment. As such, it is incumbent upon the proponent to show how proposed activities would comply with these principles and thereby contribute towards the achievement of sustainable development as defined by the NEMA.

In order to adhere to the above principles and best-practice guidelines, the following approach forms the basis for the study approach and assessment philosophy:

The study will include data searches, desktop studies, site walkovers / field survey of the property and baseline data collection, describing the broad ecological characteristics of the site and its surrounds in terms of any mapped spatial components of ecological processes and/or patchiness, patch size, relative isolation of patches, connectivity, corridors, disturbance regimes, ecotones, buffering, viability, etc.

In terms of **pattern**, the following will be identified or described:

***Community and ecosystem level***

- The main vegetation type, its aerial extent and interaction with neighbouring types, soils or topography.
- Threatened or vulnerable ecosystems (*cf. SA vegetation map/National Spatial Biodiversity Assessment, fine-scale systematic conservation plans, etc.*).

***Species level***

- Red Data Book (RDB) species (giving location if possible using GPS).
- The viability of an estimated population size of the RDB species that are present (include the degree of confidence in prediction based on availability of information and specialist knowledge, i.e. High=70-100% confident, Medium 40-70% confident, Low 0-40% confident).
- The likelihood of other RDB species, or species of conservation concern, occurring in the vicinity (include degree of confidence).

***Fauna***

- Describe and assess the terrestrial fauna present in the area that will be affected by the proposed development.
- Conduct a faunal assessment that can be integrated into the ecological study.
- Describe the existing impacts of current land use as they affect the fauna.
- Clarify species of special concern (SSC) and that are known to be:
  - endemic to the region;
  - that are considered to be of conservational concern;
  - that are in commercial trade (CITES listed species); or
  - are of cultural significance.

- Provide monitoring requirements as input into the EMPr for faunal related issues.

**Other pattern issues**


- Any significant landscape features or rare or important vegetation associations such as seasonal wetlands, alluvium, seeps, quartz patches or salt marshes in the vicinity.
- The extent of alien plant cover of the site, and whether the infestation is the result of prior soil disturbance such as ploughing or quarrying (alien cover resulting from disturbance is generally more difficult to restore than infestation of undisturbed sites).
- The condition of the site in terms of current or previous land uses.

In terms of **process**, the following will be identified or described:

- The key ecological “drivers” of ecosystems on the site and in the vicinity, such as fire.
- Any mapped spatial component of an ecological process that may occur within the grid connection corridor or within its vicinity (i.e. *corridors* such as watercourses, upland-lowland gradients, migration routes, coastal linkages or inland-trending dunes, and *vegetation boundaries* such as edaphic interfaces, upland-lowland interfaces or biome boundaries).
- Any possible changes in key processes, e.g. increased fire frequency or drainage/artificial recharge of aquatic systems.
- Furthermore, any further studies that may be required during or after the BA process will be outlined.
- All relevant legislation, permits and standards that would apply to the development will be identified.
- The opportunities and constraints for development will be described and shown graphically on an aerial photograph, satellite image or map delineated at an appropriate level of spatial accuracy.

**1.2 RELEVANT ASPECTS OF THE DEVELOPMENT**

The BESS would be located near to (within 500m of) the approved Gunstfontein Wind Farm substation, and would be approximately 5ha in total extent. MV cabling (33kV or less) will connect the BESS to the adjacent substation. The MV cabling may be overhead or below-ground.



**Figure 1.** Map showing the location of the BESS 500m assessment area with the preliminary location of the BESS footprint adjacent to the substation within the Gunstfontein WEF area.

## 2 METHODOLOGY

### 2.1 DATA SOURCING AND REVIEW

Data sources from the literature consulted and used where necessary in the study includes the following:

#### Vegetation:

- Vegetation types and their conservation status were extracted from the South African National Vegetation Map (Mucina & Rutherford 2006 and 2018 update) as well as the National List of Threatened Ecosystems (2011), where relevant.
- Information on plant species recorded for the broad area around the site was extracted from the SANBI POSA database hosted by SANBI. The species list was derived from a considerably larger area than the study site, but this is necessary to ensure a conservative approach as well as counter the fact that the site itself or the immediate area has not been well sampled in the past.



- The IUCN conservation status of the species in the list was also extracted from the database and is based on the Threatened Species Programme, Red List of South African Plants (2020).

#### *Ecosystem*

- Critical Biodiversity Areas (CBAs) were extracted from the Northern Cape Critical Biodiversity Areas Map (Oosthuysen & Holness 2016 (latest update as available on BGIS)).
- Freshwater and wetland information was extracted from the National Freshwater Ecosystem Priority Areas assessment (NFEPA) (Nel et al. 2011).
- Important catchments and protected areas expansion areas were extracted from the Northern Cape Protected Areas Expansion Strategy 2016 (NC-PAES).

#### *Fauna*

- Lists of mammals, reptiles and amphibians which are likely to occur at the site were derived based on distribution records from the literature and Animal Demography Unit (ADU) Virtual Museum spatial database (<http://vmus.adu.org.za/>).
- Literature consulted includes Branch (1988) and Alexander and Marais (2007) for reptiles, Du Preez and Carruthers (2009) for amphibians, Friedmann and Daly (2004) and Skinner and Chimimba (2005) for mammals.
- The faunal species lists provided are based on species which are known to occur in the broad geographical area, as well as a preliminary assessment of the availability and quality of suitable habitat at the site.
- The conservation status of mammals is based on the IUCN Red List Categories (EWT/SANBI 2016), while reptiles are based on the South African Reptile Conservation Assessment (Bates et al. 2013) and amphibians on Minter et al. (2004) as well as the IUCN (2020).
- The Southern African Bird Atlas Project 1 (SABAP1; Harrison *et al.*, 1997), which obtained bird distribution data between 1987 and 1992, was consulted to determine the bird species likely to occur within the broader project site. The relevant quarter-degree grid cell (QDGC) that covers the broader area is 3220DC (17 cards, 98 species). More recent bird distribution data were also obtained from the second bird atlas project, which has been on-going since its inception in 2007 (SABAP 2; <http://sabap2.adu.org.za/>). SABAP2 employs a finer resolution using the pentad scale (5' latitude x 5' longitude), of which the following two pentads were used, 3230\_2035 (21 cards, 94 species).
- The Important Bird Areas of South Africa (IBA; Marnewick *et al.*, 2015) was consulted to determine the location of the nearest IBAs to the project site.

- The conservation status, endemism and biology of all species considered likely to occur within the broader project site were determined from Hockey *et al.* (2005) and Taylor *et al.* (2015).

## **2.2 SITE VISIT**

The site was visited and sampled on the 6<sup>th</sup> of August 2020. During the site visit, all plant and animal species observed on walked transects through the veld within the study area were recorded. As the Komsberg/Roggeveld area is known to have a high abundance of plant species of conservation concern, a list of these was generated prior to the site visit and used for targeted searching of the site for these species. Active searches for reptiles and amphibians were also conducted within habitats likely to harbour or be important for such species. The presence of sensitive habitats such as wetlands or pans and unique edaphic environments such as rocky outcrops were confirmed in the field if present and recorded on a GPS and mapped onto satellite imagery of the site and included in the sensitivity map where necessary.

## **2.3 SENSITIVITY MAPPING & ASSESSMENT**

An ecological sensitivity map of the development area was produced by integrating data collected during the site survey with the available ecological and biodiversity information available in the literature and various spatial databases with mapping based on the satellite imagery and personal knowledge of the area. This includes delineating different habitat units identified on the satellite imagery and assigning likely sensitivity values to the units based on their ecological properties, conservation value and the potential presence of species of conservation concern. The ecological sensitivity of the different units identified in the mapping procedure was rated according to the following scale:

- **Low** – Areas of natural or transformed habitat with a low sensitivity where there is likely to be a negligible impact on ecological processes and terrestrial biodiversity. Most types of development can proceed within these areas with little ecological impact.
- **Medium**- Areas of natural or previously transformed land where the impacts are likely to be largely local and the risk of secondary impact such as erosion low. These areas usually comprise the bulk of habitats within an area. Development within these areas can proceed with relatively little ecological impact provided that appropriate mitigation measures are taken.
- **High** – Areas of natural or transformed land where a high impact is anticipated due to the high biodiversity value, sensitivity or important ecological role of the area. These areas may contain or be important habitat for faunal species or provide important ecological services such as water flow regulation or forage provision. Development

within these areas is undesirable and should only proceed with caution as it may not be possible to mitigate all impacts appropriately.

- **Very High** – Critical and unique habitats that serve as habitat for rare/endangered species or perform critical ecological roles. These areas are essentially no-go areas from a developmental perspective and should be avoided.

#### **2.4 SAMPLING LIMITATIONS AND ASSUMPTIONS**

This study is based on a field assessment as well as a desktop review of the available information. The major potential limitation associated with the sampling approach is the narrow temporal window of sampling. Ideally, a site should be visited several times during different seasons to ensure that the full complement of plant and animal species present are captured. However, this is rarely possible due to time and cost constraints and therefore, the representivity of the species sampled at the time of the site visit should be critically evaluated.


The site visit for the current study took place in spring, near the optimal time for such a visit. As such, the abundance of geophytes, annuals and forbs was high and the presence of species of conservation concern at the site could be well documented. It is not likely that additional site visits and field assessment would significantly alter the results of the study as the current baseline is adequate to describe the site at an appropriate level of detail. The timing and duration of the site visit, is therefore not seen as a significant limitation for the current study and is not considered to be a limiting factor which might compromise the results in any way. In addition to the current site visit, the Gunstfontein Wind Farm area has been sampled by the consultant several times in the past as part of the EIA for the Wind Farm and its grid connection infrastructure. This information was used to inform the current study, as appropriate.

The lists of amphibians, reptiles, mammals and birds for the site are based on those observed at the site as well as those likely to occur in the area based on their distribution and habitat preferences. Several site visits have been conducted during various seasons to the broader area and information on fauna observed in the area is included where relevant. This represents a sufficiently conservative and cautious approach which takes the study limitations into account.

### 3 DESCRIPTION OF THE AFFECTED ENVIRONMENT- BASELINE

#### 3.1 BROAD-SCALE VEGETATION DESCRIPTION


According to the national vegetation map (Mucina & Rutherford 2006/2018), there are two within the affected area Roggeveld Shale Renosterveld and Tanqua Escarpment Shrubland (Figure 2). The BESS 500m assessment region is however restricted entirely to the Roggeveld Shale Renosterveld vegetation type. Roggeveld Shale Renosterveld occurs in the Northern and Western Cape and occupies the majority of the Roggeveld from the Western edge of the Great Escarpment mostly above the Tanqua Basin, reaching as far east as the higher-lying areas of the Teekloof Pass south of Fraserburg along the northwest summit plateaus of the Nuweveldberge. It occupies undulating, slightly sloping plateau landscapes, with low hills and broad shallow valleys supporting mainly moderately tall shrublands dominated by renosterbos with a rich geophytic flora in the wetter and rocky habitats. It occurs mostly on mudrocks and sandstones of the Adelaide Subgroup. The land types present are mostly Fc and Da. Mucina & Rutherford (2006) list 12 endemic species for this vegetation type, which is a large number given that the total extent of the vegetation type is only 2917 km<sup>2</sup>.



**Figure 2.** Broad-scale overview of the vegetation in and around the Gunstfontein substation and BESS study area. The vegetation map is an extract of the national vegetation map (Mucina & Rutherford 2006 & 2018 update) and also includes drainage lines delineated by the NFEPA assessment (Nel et al. 2011).

### **3.2 FINE-SCALE VEGETATION DESCRIPTION**

The area around the approved substation site for the Gunstfontein WEF consists of typical Roggeveld Shale Renosterveld. There are however several different habitats present within the 500m assessment region around the substation site which are illustrated and described below (Figure 3).



**Figure 3.** Ecological habitats observed within the Gunstfontein BESS 500m assessment region (entire frame) and which are described in detail below.


The majority of the BESS area consists of Roggeveld Shale Renosterveld typical of the Sutherland Plateau area. The soils are fine-textured but sandy soils and are generally quite shallow with several areas of exposed bedrock within the BESS area. There are also a few areas of deeper soils which can be recognized by their somewhat taller vegetation. Typical and dominant species observed within the BESS study area include *Euryops lateriflorus*, *Dimorphotheca cuneata*, *Selago saxatilis*, *Rosenia oppositifolia*, *Pteronia tricephala*, *Pentzia punctata*, *Euryops annae*, *Dicerotheramnus rhinocerotis*, *Ehrharta calycina*, *Ehrharta eburnea* (NT), *Senecio erosus*, *Romulea tortuosa* subsp. *tortuosa*, *Asparagus capensis*, *Euryops multifidus*, *Poa bulbosa*, *Oxalis obtusa*, *Berkheya spinosa*, *Chrysocoma ciliata*, *Romulea atrandra* var. *atrandra*, *Colchicum coloratum* subsp. *burchellii*, *Othonna auriculifolia*, *Diospyros austro-africana*, *Oxalis melanosticta* var. *melanosticta* and *Oxalis pocockiae*.



**Figure 4.** The typical Roggeveld Shale Renosterveld on shallow soils within the BESS 500m study region. This habitat is considered relatively low sensitivity and is considered acceptable for the location of the BESS.



**Figure 5.** Roggeveld Shale Renosterveld on deeper soils within the BESS study area, with taller shrubs dominated by *Euryops lateriflorus*.



**Figure 6.** There are several areas of exposed bedrock within the BESS study area. Although there is very little vegetation associated with these areas and they are not considered important from a botanical point of view, they were observed to be a relatively important local habitat for reptiles and other fauna which prefer rocky habitats.



### 3.3 LISTED AND PROTECTED PLANT SPECIES

It is important to note that the site falls within the Komsberg Centre of Diversity and Endemism and as such is an area with a known high abundance of species of concern and endemism. A list of species of conservation concern recorded from the wider area is provided in Annex 1. Species of concern observed at the site during the field assessment includes *Eriocephalus grandiflorus* (Rare) and *Ehrharta eburnea* (NT) which are both quite widespread species that have healthy populations outside of the affected area. Although it is possible that the development would generate some impact on these species, this would be minor as it is highly unlikely that the local populations would be compromised in any way by the development. Overall, the abundance of plant SCC within the site is low and the impact of the development on SCC would be acceptable and low.

In terms of the provincial legislation the following species and genera are protected and would require specific consideration during the pre-construction walk-through of the BESS footprint. The example species provided are to illustrate the typical species present and is not intended as an exhaustive list.

#### **Schedule 1** (Specially Protected Species):

- All species of the genus *Pelargonium* (Family: Geraniaceae) (e.g. *Pelargonium rapaceum*)

#### **Schedule 2** (Protected Species):

- All species of the family Mesembryanthemaceae: (e.g. *Antimima pumila*, *Hammeria salteri*, *Cheiridopsis namaquensis*, *Lampranthus spp.*, *Cleretum papulosum subsp. papulosum*, *Drosanthemum spp.*, *Ruschia centrocapsula*)
- All species of the family Amaryllidaceae: (e.g. *Brunsvigia spp* (*B. bosmaniae*), *Haemanthus coccineus*)
- All species of the genus *Colchicum* (Family Colchicaceae): e.g. (*Colchicum coloratum*, *C. Cuspidatum*).
- All species of the family Crassulaceae; e.g. (*Tylecodon wallichii*, *T. ventricosus*, *Crassula deltoidea*, *C. columnaris*, *C. muscosa*, *C. umbella*, *C. glomerata*, *Adromischus filicaulis*)
- All species of the family Iridaceae: (e.g. *Romulea atrandra*, *R. tortuosa*, *komsbergensis*, *Hesperantha acuta*, *Moraea fugax*)
- All species of the genus *Oxalis* (Family: Oxalidaceae): (e.g. *Oxalis obtusa*, *O. melanostica*, *O.palmifrons*)
- All species of the genus *Lachenalia* (Family: Hyacinthaceae): (e.g. *Lachenalia aurioliae*)

It is recommended that a Pre-construction Walk-Through Survey is conducted within the final BESS footprint, to inform search-and-rescue efforts. Species of concern should be recorded and may only be removed, transplanted, destroyed (or any other form of disturbance) after the necessary approval (permits) has been obtained from the relevant authority, i.e. the Department of Agriculture, Environmental Affairs, Rural Development and Land Reform. It is also important to note that species of ecological importance, local endemics and red-listed species should be translocated out of the development footprint, where these have a high probability of survival. These would be identified during the preconstruction walk-through.

### **3.4 FAUNAL COMMUNITIES**

#### **Mammals**

The Gunstfontein BESS site is likely to have moderate to low mammalian species richness. The BESS site falls within or near the edge of the distribution range of at least 44 terrestrial mammals. Within the broader area, the ridges, hills and uplands, with rocky outcrops, rocky bluffs and cliffs provide suitable habitat for species which require or prefer rock cover such as Cape Rock Elephant Shrew, *Elephantulus edwardii*, Hewitt's Red Rock Hare *Pronolagus saundersiae*, Namaqua Rock Mouse *Micaelamys namaquensis* and Rock Hyrax, *Procavia capensis*. Larger species commonly observed in the area include Grey Rhebok, *Pelea capreolus* (Near Threatened) which is likely to use the area on a fairly regular basis and Klipspringer, *Oreotragus oreotragus* which are not likely to be resident in the BESS area as the habitat is not rugged enough to provide cover for this species. The introduced Fallow Deer, *Dama dama* is also common in the area and is likely to occur at the site on occasion. The lower-lying parts of the area are home to species associated with more densely-vegetated lowland habitats on deeper soils and along drainage lines and floodplains, which includes Brants's Whistling Rat *Parotomys brantsii*, the Bush Vlei Rat *Otomys unisulcatus*, Hairy-footed Gerbil *Gerbillurus paeba* and Common Duiker *Sylvicapra grimmia*. Most of these species are likely to be resident within the BESS area or would be using this area on a fairly regular basis.

Listed species which do or may occur at the site include the, Grey Rhebok (Near Threatened) Black-footed Cat *Felis nigripes* (Vulnerable), Leopard *Panthera pardus* (Near Threatened) and Riverine Rabbit *Bunolagus monticularis* (Critically Endangered). All of these species have relatively large ranges across South Africa and the development of the BESS would result in an insignificant extent of habitat loss for these species. Although the Riverine Rabbit *Bunolagus monticularis* is known to occur in the wider area, it is not currently known from the plateau in the affected area and it is considered highly unlikely to be present within the 500m assessment region affected by the BESS. Due to the small footprint of the BESS and its proximity to the approved substation infrastructure, it is not likely that there would be any significant degree of habitat loss for mammals as a result of the construction and operation of the BESS.

Overall there do not appear to be any significant issues regarding mammals and the development of Gunstfontein BESS. In general, the major impact associated with the development of Gunstfontein BESS for mammals would be some localised and minor habitat loss and disturbance during construction and operation.

### **Reptiles**

According to the distribution maps available in the literature, as many as 50 reptiles could occur within the broad area around the BESS. However, according to the records within the Virtual Museum database, only 35 species have been recorded within the 3220 degree square, suggesting that the actual number of reptile species present at the site is likely to be relatively low. In terms of species of conservation concern, the only listed species recorded in the area is the Karoo Padloper *Homopus boulengeri* which is listed as Near Threatened. Although it is possible that this species moves through the BESS area on occasion, it is highly unlikely to be present within the affected area as there not sufficient rock cover for shelter for this species and so it considered highly unlikely to present or be affected by the BESS.

Species commonly observed in the wider area on previous field assessments include the Karoo Tent Tortoise *Psammobates tentorius tentorius*, Angulate Tortoise *Chersina angulata*, Puff Adder *Bitis arietans*, Karoo Girdled Lizard *Cordylus polyzonus*, Southern Rock Agama *Agama atra*, Namaqua Plated Lizard *Gerrhosaurus typicus*, Cape Skink *Mabuya capensis*, Variegated Skink *Trachylepis variegata*, Common Sand Lizard *Pedioplanis lineocellata pulchella* and Cape Cobra *Naja nivea*. While the BESS is likely to result in some localised habitat loss for such typical resident species, there are no parts of the BESS study area that are considered to be especially important or sensitive in terms of reptile abundance or diversity. In terms of impacts of the development on reptiles, the major impact is likely to come from disturbance during the construction phase which would be transient and localised and consequently of low long-term consequence.



**Figure 7.** Common reptiles observed at the site include, from top left, the Variegated Skink, Common Sand Lizard, Ground Agama and Karoo Girdled Lizard.

### **Amphibians**

Only seven amphibians are likely to occur in the area, indicating that the frog diversity of the site is likely to be low. No listed species are likely to occur in the area. All of the species recorded in the area are widespread species of low conservation concern. Species such as the Cape River Frog *Amietia fuscigula* occur along the larger drainage lines in pools and in the farm dams of the area. Species such as Karoo Caco *Cacosternum karooicum*, Karoo Toad *Vandijkophrynus garipeensis* and Cape Sand Frog *Tomopterna delalandii* are less dependent on water and are likely to be more widespread. Given the general aridity and low likely abundance of amphibians within the area, impacts on amphibians are likely to be localised and of a low significance.

### **Avifauna**

Approximately 120 bird species are known to occur within the broader project area (Appendix 5). The bird assemblage of the study area and surrounds is fairly typical of the Succulent Karoo Biome. A number of small passerines that are considered common within the renosterveld and succulent karoo scrub that characterises the area and are considered endemic/near-endemic and biome-restricted (Table 1). Some of these species are nomadic, such as the Black-headed Canary *Serinus alario* and Lark-like Bunting *Emberiza impetuani*, which may be absent in some years. Seemingly cryptic species such as Cinnamon-breasted Warbler is not uncommon along the Great Escarpment and is usually restricted to rocky ridges

and scree with vegetation cover. Other species of some importance include the Karoo Lark *Calendulauda albescens*, Cape Clapper Lark *Mirafra apiata*, and Karoo Long-billed Lark *Certhilauda subcoronata*. While many of these and other species are endemic/near-endemic and biome-restricted, all of these species are widely distributed in the Karoo and Fynbos Biomes.

Species of concern present in the area (Table 1) includes three Endangered species, namely Ludwig's Bustard *Neotis ludwigii*, Martial Eagle *Polemaetus bellicosus* and Black Harrier *Circus maurus*. Species of secondary concern which have also been recorded in the area include Verreaux's Eagle *Aquila verreauxii*, Lanner Falcon *Falco biarmicus* and Black Stork (*Ciconia nigra*). Verreaux's Eagle is the most abundant of the large raptor species in the area, while the latter two species are significantly scarcer. Black Stork often frequent farm dams, not only singly but also in small congregations. The Vulnerable Southern Black Korhaan *Afrotis afra* and the Near-Threatened Karoo Korhaan *Eupodotis vigorsii* are found throughout the region and have also been recorded in the wider area.

**Table 1.** Priority species identified in the project site and surrounds based on their conservation status (Taylor *et al.*, 2015), regional endemism (Birdlife South Africa, 2019), and priority score (Retief *et al.*, 2011).

| Species                    | Cons. Status | Endemic/Near-endemic | Priority Score | SABAP1 reporting rate (%) | Susceptible to                      |
|----------------------------|--------------|----------------------|----------------|---------------------------|-------------------------------------|
| Bustard, Ludwig's          | EN           |                      | 320            | 6                         | Collisions                          |
| Buzzard, Common (Steppe)   |              |                      | 210            | 18                        | Collisions/disturbance              |
| Buzzard, Jackal            |              | NE                   | 250            | 6                         | Collisions/disturbance              |
| Crane, Blue                | NT           |                      | 320            |                           | Collisions                          |
| Eagle, Black-chested Snake |              |                      | 230            |                           | Collisions/disturbance              |
| Eagle, Booted              |              |                      | 230            | 6                         | Collisions/disturbance              |
| Eagle, Martial             | EN           |                      | 350            | 6                         | Collisions/electrocution            |
| Eagle, Verreaux's          | VU           |                      | 360            | 6                         | Collisions/electrocution            |
| Falcon, Lanner             | VU           |                      | 300            |                           | Collisions/disturbance              |
| Flamingo, Greater          | NT           |                      | 290            |                           | Collisions                          |
| Francolin, Grey-winged     |              | SLS                  | 190            | 6                         | Disturbance/habitat loss            |
| Goshawk, Pale Chanting     |              |                      | 200            | 41                        | Disturbance/habitat loss            |
| Harrier, Black             | EN           | NE                   | 345            | 12                        | Collisions/disturbance/habitat loss |
| Harrier-hawk, African      |              |                      | 190            |                           | Disturbance/habitat loss            |
| Kestrel, Rock              |              |                      |                | 59                        | Disturbance/habitat loss            |

|                              |    |     |     |    |                                     |
|------------------------------|----|-----|-----|----|-------------------------------------|
| Kite, Black-winged           |    |     | 174 | 29 | Disturbance/habitat loss            |
| Korhaan, Karoo               | NT |     | 240 |    | Collisions/disturbance/habitat loss |
| Korhaan, Southern Black      | VU | E   | 270 | 18 | Collisions/disturbance/habitat loss |
| Owl, Cape Eagle-             |    |     | 250 |    | Disturbance/habitat loss            |
| Owl, Spotted Eagle-          |    |     | 170 | 6  | Disturbance/habitat loss            |
| Pipit, African Rock          | NT | SLS | 200 |    | Disturbance/habitat loss            |
| Sparrowhawk, Rufous-breasted |    |     | 170 |    | Disturbance/habitat loss            |
| Stork, Black                 | VU |     | 330 | 6  | Collisions/electrocutions           |

The Near-Threatened Blue Crane *Grus paradisea* and Greater Flamingo *Phoenicopterus ruber* are both rare in the region, but may occasionally be present. According to SABAP2 records, Blue Crane has only been recorded in a few pentads within a 50km radius of the project site, while Greater Flamingo have been recorded in a number of pentads, along the plateau of the Great Escarpment where they frequent large farm dams. Both species may however pass through the area *en route* between focal sites, with flamingos possibly commuting in small flocks. African Rock Pipit is not uncommon along the escarpment to the and have also been recorded in the area (EWT, 2014).

The avifauna of the project site and broader area appears fairly typical of the Succulent Karoo Biome. However, due to the presence of a fair number of priority species, the sensitivity of the avifauna can be considered to be of medium significance. In terms of impact, the group of primary concern is the medium to large non-passerines, which include the large terrestrial birds and diurnal raptors. Many of these are also red-listed, such as Ludwig’s Bustard, Martial eagle, and Black Stork. Most of these species are susceptible to collisions with power lines owing to reduced ability to see the power lines and reduced manoeuvrability in flight to avoid collisions (Martin & Shaw, 2010; Jenkins *et al.*, 2010; Jenkin *et al.*, 2011; Shaw, 2013). However, the only species which are highly susceptible include Ludwig’s Bustard and Black Stork (Jenkins *et al.*, 2010). An additional threat faced by the large raptors is electrocution when perched or attempting to perch on power line structures (Lehman *et al.*, 2007), but this depends largely on the type of pylons or towers used. Electrocutions can further be avoided to a large extent by employing suitable mitigation methods. Disturbances during construction of the substation and BESS is also expected to have a negative impact by temporarily displacing birds from foraging habitat.

### **3.5 CRITICAL BIODIVERSITY AREAS & BROAD-SCALE PROCESSES**

An extract of the Northern Cape Critical Biodiversity Areas map for the broader study area is depicted below in Figure 8. The entire BESS study area falls within an area classified as CBA 1. Development within CBAs is undesirable and can potentially lead to loss of biodiversity and negatively affect ecological processes. The impact of the current proposed BESS would be mediated by the proximity of the BESS to the facility substation and the low overall footprint of the BESS.

In terms of the ecological features and processes that may be affected by the development, these are partly described below in Table 1 below, as they relate to the CBAs within the area. But more broadly speaking, the BESS site lies within an area of typical Roggeveld Shale Renosterveld with no features present that would suggest that the study area represents a particularly important corridor or process feature of the wider area. Important features of the area include the escarpment south of the site and the larger drainage features which occur mostly to the north and east of the BESS study area. All of these features are well outside of the BESS site and would not be affected by the BESS.




Figure 8. Extract of the Northern Cape Critical Biodiversity Areas map for the broader study area around the Gunstfontein BESS, showing that the BESS occurs within an area classified as CBA 1.

The loss of an additional ~5ha of habitat near to the approved substation would be very unlikely to compromise the ecological functioning of the affected CBA in any way. A summary of the underlying features associated with the CBA within the site is provided below in Table 1. It is important to note that apart from the process and representivity roles of the CBAs, the features of concern underlying the CBAs are not located within the 500m assessment zone, as these features are well outside of the 500m assessment region. In terms of the representivity of vegetation types, only Roggeveld Shale Renosterveld is within the site and the loss of less than 5ha of this vegetation type would not be significant, especially given that the field assessment indicated that the abundance of SCC within this area was low. In terms of the process features, the development would contribute to cumulative habitat loss and fragmentation in the area to some degree, but the specific contribution of the BESS at less than 5ha would not be sufficient to significantly compromise the functioning of these broad-scale ecological processes. Given the small footprint of the BESS and the avoidance of hydrological features at the site, the impact on the BESS on water runoff quality and quantity



would be minimal and the overall impact on the affected catchment and hydrological processes would be negligible. In terms of the Strategic Water Source Areas (SWSAs) and Freshwater Ecosystem Priority Areas (FEPAs), it is important to note that the 500m assessment region does not occur with either of the SWSA or NFEPA delineated wetland or river systems, and therefore these do not apply to this development. As such, the development of the BESS is considered acceptable in terms of CBAs and ecological processes.

**Table 1.** Reasons underlying the CBA 1 status of the site. The features are obtained from the reasons database associated with the Northern Cape CBA map available on the BGIS database.


| <b>Feature</b>                            | <b>Remarks</b>  |
|---|---|
| Roggeveld Shale Renosterveld              | The development is less than 5ha in extent and would not contribute significantly to the loss of habitat within the Roggeveld Shale Renosterveld vegetation type which is still largely intact.   |
| Threatened Species                        | Although there are some threatened species in the area, the BESS development would not compromise the local populations of any species of concern.  |
| Natural Wetlands                          | There are no significant wetlands within the 500m assessment region. The wetlands underlying the CBA 1 are not within the 500m assessment region.   |
| Rivers                                    | There is a small drainage line within the 500m assessment region, but this can be avoided by placing the infrastructure in low or medium sensitivity regions. The CBA 1 status of the area is based on the presence of the larger drainage lines present in the wider area.   |
| Large high value climate resilience areas | The rugged topography of the area and intact nature of the landscape provides it with climate change resilience. The footprint of the development at less than 5ha would not compromise this function of the landscape. The development would however contribute some degree towards cumulative habitat loss and fragmentation in the area. The specific contribution of the BESS is however insignificant compared to the existing or approved developments. |
| NPAES PA and Focus                        | These have changed since the 2011 NPAES and a new NPAES layer has been developed but has not been released as yet. The loss of less than 5ha to the BESS would not change conservation options in the area as it is already affected by approved and under construction wind farms.   |
| Landscape structural elements             | As with climate change resilience, this feature of the CBA 1 would not be significantly altered by the presence of the BESS. Based on the results of the field assessment it is unlikely that the affected area represents an important movement or migration corridor for any fauna and the presence of the BESS would be very unlikely to compromise the ability of fauna, flora or avifauna to move about the landscape.                                   |
| PA distance buffers 5km & 10km            | There are no mapped protected areas within 10km of the BESS 500m assessment region. There are no known formal protected areas in the vicinity of the 500m assessment region that would be affected by the BESS.   |

### **3.6 CURRENT BASELINE & CUMULATIVE IMPACT**

The BESS is part of the Gunstfontein WEF and there are also several other approved wind energy facilities in the wider area, which together represent a significant potential negative impact on the local environment. However, the majority of wind farms are located below the escarpment and there are few approved projects on the escarpment itself. The Gunstfontein site is however near to the Mainstream Sutherland wind farm which is located ~15km to the east of the current site. As these are all existing developments, they are considered to represent part of the cumulative impact baseline for the area. The primary concern associated with the current development would be the additional contribution of the BESS to cumulative impacts in the area. The footprint of the BESS is estimated at up to 5ha. In context of the generally intact nature of the area, this is seen as a very low contribution. The medium voltage connection from the BESS to the nearby substation would be up to 500m in length and as this would likely be underground, the risk of collisions or electrocution of avifauna would be minimal. Should some or all of the line be overhead cabling, the presence of the BESS itself and the substation would deter many larger species from the immediate area and along with other associated avifaunal mitigation, the additional risk to avifauna from the BESS would be very low. As a result, the contribution of the current proposed BESS infrastructure to overall cumulative impact from wind farm and grid infrastructure development in the wider area is very low and is considered acceptable.

### **3.7 SITE SENSITIVITY ASSESSMENT**

The sensitivity map for the BESS 500m assessment region is illustrated below in Figure 9. The majority of the 500m assessment region is typical, open plains Roggeveld Shale Renosterveld, considered to be low ecological sensitivity. There is however one minor drainage line within the BESS 500m assessment region that is considered to be high ecological sensitivity and unsuitable for development. There are also some areas of rock pavement distributed across the site which are considered medium sensitivity on account of the value of these areas as faunal habitat. Under the layout of the BESS provided for this assessment, the BESS would be restricted to the low sensitivity parts of the site, with the result that the impacts associated with the BESS would be low. Provided that the BESS footprint can be restricted to the low and/or medium sensitivity areas within the 500m assessment zone, the exact placement of the BESS within this area would not result in significant differences in impact. As such, the current placement is considered acceptable but alternative placements within the medium and low sensitivity areas would also be acceptable.



**Figure 9.** Sensitivity map for the Gunstfontein BESS 500m assessment region.

#### 4 IDENTIFICATION & NATURE OF IMPACTS

In this section, the potential impacts and associated risk factors that may be generated by the development are identified and discussed before being assessed in the next section.

#### **4.1 IDENTIFICATION OF IMPACTS**

In this section the potential impacts associated with the establishment of the Gunstfontein BESS are explored in context of the features and characteristics of the development area, the likelihood and extent to which each impact would occur given the characteristics of the development area, and the extent and nature of the development.

##### *Impacts on vegetation and protected plant species*

Several protected species occur in the area and which would potentially be impacted by the development of the Gunstfontein BESS. Vegetation clearing during the construction phase will lead to the loss of currently intact habitat within the footprint and is an inevitable consequence of the establishment of the BESS. As this impact is certain to occur during the construction phase, it is assessed for the construction phase only, as this is when the impact will occur, although the consequences will persist for some time after construction has been completed.

##### *Direct faunal impacts*

Increased levels of noise, pollution, disturbance and human presence during the construction phase will be detrimental to fauna. Sensitive and shy fauna would move away from the development area during the construction phase as a result of the noise and human activities present, while some slow-moving species would not be able to avoid the construction activities and might be killed. Some impact on fauna is highly likely to occur during the construction phase and this impact is therefore assessed for the construction phase only.

##### *Direct Avifaunal impacts*

Vegetation clearing for the BESS and associated infrastructure will impact the local avifauna directly through habitat loss. The presence and operation of construction machinery on site would create a physical impact as well as generate noise, pollution and other forms of disturbance, while increased human presence could lead to poaching, illegal fauna collecting and other forms of disturbance such as fire. Impacts on avifauna during the operational phase would be reduced and the operation of the BESS would generate minor disturbance during maintenance of infrastructure, which may deter some avifauna from the area, especially red-listed avifaunal species which are less tolerant of disturbances. Should the connection between the BESS and the substation require an overhead line, there would be a small risk of collisions with the power line and electrocution from the power line infrastructure (Lehman et al., 2007, Jenkins et al., 2010).

##### *Habitat Degradation due to Erosion and Alien Plant Invasion*

Disturbance within and near the BESS site generated during the construction phase will leave the area vulnerable to erosion and alien plant invasion, which would lead to degradation of the local environment. Although, the disturbance would be created during the construction phase, the major impacts would manifest during the operation phase.

*Impact on CBAs and NPAES Focus Areas*

The development would have an impact on an area classified as CBA 1. However, the BESS is not within an NC-PAES Focus Area, indicating that it has not been identified as being of high significance for future conservation expansion. The impact on the CBAs is assessed as part of the cumulative impacts associated with the development.

## **5 ASSESSMENT OF IMPACTS**

---

The various identified impacts are assessed below for the different phases of the development.

### **5.1 PLANNING & CONSTRUCTION PHASE**

#### **Impact 1. Impacts on vegetation and listed or protected plant species resulting from the BESS construction activities**

|   |   |                        |
|---|---|------------------------|
| <b>Impact Nature:</b> Impacts on vegetation will occur due to disturbance and vegetation clearing associated with the construction of the BESS and associated infrastructure. |   |                        |
|   | <b>Without Mitigation</b>   | <b>With Mitigation</b> |
| <b>Extent</b>   | Local (1)   | Local (1)              |
| <b>Duration</b>   | Long-term (3)   | Long-term (3)          |
| <b>Magnitude</b>  | Low (3)   | Low (2)                |
| <b>Probability</b>  | Highly Likely (4)   | Highly Likely (4)      |
| <b>Significance</b>   | Low (28)  | Low (24)               |
| <b>Status</b>   | Negative  | Negative               |
| <b>Reversibility</b>  | Moderate  | Moderate               |
| <b>Irreplaceable loss of resources</b>  | Low   | Low                    |
| <b>Can impacts be mitigated?</b>  | This impact cannot be well mitigated because some loss of vegetation is unavoidable and is a certain outcome of the development.  |                        |
| <b>Mitigation</b>   | <ul style="list-style-type: none"> <li>Pre-construction walk-through of the final layout in order to locate species of conservation concern that can be translocated as well as comply with the Northern Cape Nature Conservation Act and Northern</li> </ul> |                        |

|                           |  |
|---------------------------|--|
|                           | <p>Cape Department of Agriculture, Environmental Affairs, Rural Development and Land Reform/DAFF permit conditions.</p> <ul style="list-style-type: none"> <li>• Search and rescue for identified species of concern before construction.</li> <li>• Vegetation clearing to commence only after walk-through has been conducted and necessary permits obtained.</li> <li>• Pre-construction environmental induction for all construction staff on site to ensure that basic environmental principles are adhered to. This includes awareness of no littering, appropriate handling of pollution and chemical spills, avoiding fire hazards, minimising wildlife interactions, remaining within demarcated construction areas etc.</li> <li>• Environmental Officer (EO) to provide supervision and oversight of vegetation clearing activities within sensitive areas such as near the drainage lines and wetlands.</li> <li>• Vegetation clearing should be kept to a minimum and restricted to the BESS footprint as closely as possible.</li> <li>• All construction vehicles should adhere to clearly defined and demarcated roads. No off-road driving to be allowed outside of the construction area.</li> <li>• Temporary laydown areas should be located within previously transformed areas or areas that have been identified as being of low sensitivity. These areas should be rehabilitated after use.</li> </ul> |
| <b>Cumulative Impacts</b> | The Gunstfontein BESS will contribute to cumulative impacts on habitat loss and transformation in the area, but the contribution would be very low.  |
| <b>Residual Risks</b>     | The loss of currently intact vegetation is an unavoidable consequence of the development and cannot be entirely mitigated. The residual impact would however be low.   |

***Impact 2. Direct Faunal Impacts Due to Construction Activities***

|   |                           |                        |
|---|---------------------------|------------------------|
| <b>Impact Nature:</b> Disturbance, transformation and loss of habitat will have a negative effect on resident fauna during construction. This will however be transient and restricted to the construction phase. |                           |                        |
|   | <b>Without Mitigation</b> | <b>With Mitigation</b> |
| <b>Extent</b>   | Local (1)                 | Local (1)              |
| <b>Duration</b>   | Short-term (2)            | Short-term (2)         |
| <b>Magnitude</b>  | Low (3)                   | Low (2)                |
| <b>Probability</b>  | Probable (3)              | Probable (3)           |
| <b>Significance</b>   | Low (18)                  | Low (15)               |
| <b>Status</b>   | Negative                  | Negative               |

|  |  |      |
|--|--|------|
| <b>Reversibility</b>                   | High   | High |
| <b>Irreplaceable loss of resources</b> | No   | No   |
| <b>Can impacts be mitigated?</b>       | Partly, although noise and disturbance cannot be well mitigated, impacts on fauna due to human presence such as poaching can be mitigated.   |      |
| <b>Mitigation</b>                      | <ul style="list-style-type: none"> <li>• All personnel should undergo environmental induction with regards to fauna and, in particular, awareness about not harming or collecting species such as snakes, tortoises and owls, which are often persecuted out of superstition.</li> <li>• Any fauna threatened by the construction activities should be removed to safety by an appropriately qualified environmental officer.</li> <li>• All construction vehicles should adhere to a low speed limit on site (40km/h max) to avoid collisions with susceptible species such as snakes and tortoises.</li> <li>• All hazardous materials should be stored in the appropriate manner to prevent contamination of the site. Any accidental chemical, fuel and oil spills that occur at the site should be cleaned up in the appropriate manner as related to the nature of the spill.</li> <li>• If holes or trenches need to be dug for electrical cabling or other facility infrastructure, these should not be left open for extended periods of time as fauna may fall in and become trapped in them. Holes should only be dug when they are required and should be used and filled shortly thereafter.</li> </ul> |      |
| <b>Cumulative Impacts</b>              | During the construction phase the activity would contribute to cumulative fauna disturbance and disruption in the area, but this would be short lived and little long-term impact would be generated.  |      |
| <b>Residual Risks</b>                  | It is probable that some individuals of susceptible species will be lost to construction-related activities despite mitigation. However, this is not likely to impact the viability of the local population of any fauna species.  |      |

***Impact 3. Avifaunal Impact due to Construction Activities***

|   |                           |                        |
|---|---------------------------|------------------------|
| <b>Impact Nature:</b> Direct Avifaunal Impacts During Construction – habitat loss and disturbance |                           |                        |
|   | <b>Without Mitigation</b> | <b>With Mitigation</b> |
| <b>Extent</b>   | Local (1)                 | Local (1)              |
| <b>Duration</b>   | Short-term (1)            | Short-term (1)         |
| <b>Magnitude</b>  | Low to Moderate (4)       | Low (3)                |
| <b>Probability</b>  | Highly likely (4)         | Probable (3)           |
| <b>Significance</b>   | Low (24)                  | Low (15)               |
| <b>Status</b>   | Negative                  | Negative               |
| <b>Reversibility</b>  | High                      | High                   |

|  |   |     |
|--|---|-----|
| <b>Irreplaceable loss of resources</b> | Low   | Low |
| <b>Can impacts be mitigated?</b>       | Although there will be some habitat loss that cannot be well mitigated, impacts on avifauna will be transient and of low magnitude during construction.   |     |
| <b>Mitigation</b>                      | <ul style="list-style-type: none"> <li>• If the connection to the substation is an overhead line then the design of the proposed power line must be of a type or similar structure as endorsed by the Eskom-EWT Strategic Partnership on Birds and Energy, taking into account the mitigation guidelines recommended by Birdlife South Africa (Jenkins et al., 2017).</li> <li>• Where necessary, deterrent devices such as bird guards should be mounted on relevant parts of the pylons to further reduce the possibility of electrocutions.</li> <li>• The power line should be marked with bird diverters in order to make the lines as visible as possible to collision-susceptible species. Recommended bird diverters such as brightly coloured 'aviation' balls, thickened wire spirals, or flapping devices that increase the visibility of the lines should be fitted.</li> <li>• All personnel should undergo environmental induction with regards to avifauna and in particular awareness about not harming, collecting or hunting terrestrial species (e.g. bustards, korhaans, francolin), and owls, which are often persecuted out of superstition.</li> <li>• All construction vehicles should adhere to clearly defined and demarcated roads. No off-road driving to be allowed outside of the construction area.</li> <li>• The use of laydown areas within the footprint of the development should be used where feasible, to avoid habitat loss and disturbance to adjoining areas.</li> <li>• Any avifauna threatened by the construction activities should be removed to safety by the Environmental Officer (EO).</li> <li>• If lights are to be used at night for ensuring that infrastructure on site is lit, this should be done with downward-directed low-UV type lights (such as most HPS bulbs), which do not attract insects and their avian predators., so as to minimise disturbance to birds flying over the site at night.</li> <li>• All vehicles (construction or other) accessing the site should adhere to a low speed limit on site (40km/h max) to avoid collisions with susceptible avifauna, such as nocturnal and crepuscular species (e.g. nightjars and owls) which sometimes forage or rest on roads, especially at night.</li> <li>• If holes or trenches need to be dug for cabling or pylons, these should not be left open and unattended for extended periods (&gt; 1 week) of time as terrestrial avifauna or their flightless young may become entrapped therein. Holes should only be dug when they are required and should be used and filled shortly thereafter, alternately,</li> </ul> |     |



|                           |  |
|---------------------------|--|
|                           | excavated areas should be checked frequently for trapped fauna/ avifauna that require assistance to exit the excavated area.   |
| <b>Cumulative Impacts</b> | The Great karoo BESS will contribute to cumulative impacts on avifaunal habitat loss and fragmentation, as well as collision risk with power line infrastructure in the area. However, given the small footprint of the development and proximity to the approved substation, the contribution would be insignificant. |
| <b>Residual Risks</b>     | There would be some residual habitat loss associated with the development that cannot be avoided.  |

**5.2 OPERATIONAL PHASE IMPACTS**

**Impact 1. Faunal Impacts due to Operation**

|  |  |                        |
|--|--|------------------------|
| <b>Impact Nature:</b> The operation and maintenance of the Gunstfontein BESS may lead to disturbance or persecution of fauna in the vicinity of the development. |  |                        |
|  | <b>Without Mitigation</b>  | <b>With Mitigation</b> |
| <b>Extent</b>  | Local (1)  | Local (1)              |
| <b>Duration</b>  | Long-term (4)  | Long-term (4)          |
| <b>Magnitude</b>   | Minor (2)  | Minor (2)              |
| <b>Probability</b>   | Probable (3)   | Improbable (2)         |
| <b>Significance</b>  | Low (21)   | Low (14)               |
| <b>Status</b>  | Negative   | Negative               |
| <b>Reversibility</b>   | High   | High                   |
| <b>Irreplaceable loss of resources</b>   | No   | No                     |
| <b>Can impacts be mitigated?</b>   | To a large extent, but some low-level residual impact due to noise and human disturbance may occur during maintenance activities.  |                        |
| <b>Mitigation</b>  | <ul style="list-style-type: none"> <li>• Any potentially dangerous fauna such as snakes or fauna threatened by the maintenance and operational activities should be removed to a safe location.</li> <li>• All hazardous materials should be stored in the appropriate manner to prevent contamination of the site. Any accidental chemical, fuel and oil spills that occur at the site should be cleaned up in the appropriate manner as related to the nature of the spill.</li> <li>• All vehicles accessing the site should adhere to a low speed limit on site (40km/h max) to avoid collisions with susceptible species such as snakes and tortoises.</li> </ul> |                        |

|                           |   |
|---------------------------|---|
| <b>Cumulative Impacts</b> | The development would contribute to cumulative disturbance for fauna, but the contribution would be very low and is not considered significant.   |
| <b>Residual Risks</b>     | Disturbance from maintenance activities will occur at a low and infrequent level with the result that no long-term impacts are expected to occur. |

***Impact 2. Avifaunal Impacts due to Operation***

|   |   |                        |
|---|---|------------------------|
| <b>Impact Nature:</b> Direct Avifaunal Impacts During Operation – collisions, electrocution and disturbance |   |                        |
|   | <b>Without Mitigation</b>   | <b>With Mitigation</b> |
| <b>Extent</b>   | Local (1)   | Local (1)              |
| <b>Duration</b>   | Long-term (4)   | Long-term (4)          |
| <b>Magnitude</b>  | Moderate (4)  | Low (3)                |
| <b>Probability</b>  | Likely (3)  | Improbable (2)         |
| <b>Significance</b>   | Low (27)  | Low (16)               |
| <b>Status</b>   | Negative  | Negative               |
| <b>Reversibility</b>  | Medium  | Medium                 |
| <b>Irreplaceable loss of resources</b>  | Low   | Low                    |
| <b>Can impacts be mitigated?</b>  | To a large extent, although bird flappers and other bird diverters are not 100% effective in reducing bird collisions and electrocutions, hence there would still be a low residual impact.   |                        |
| <b>Mitigation</b>   | <ul style="list-style-type: none"> <li>• Any injuries or mortalities of avifauna observed at the BESS should be reported to the EO and recorded for monitoring purposes. Should repeated injuries or fatalities occur, an avifaunal expert should be consulted to identify and remedy the cause of the problem.</li> <li>• movements by vehicles and personnel should remain within the BESS and substation area and should not stray from the approved access and maintenance routes.</li> <li>• Any raptor nests that are discovered on the power line structures should be reported to the Environmental Officer, while utmost care should be taken to not disturb these nests during routine maintenance procedures.</li> </ul> |                        |
| <b>Cumulative Impacts</b>   | The development will contribute to cumulative impacts on avifaunal habitat loss as well as collision and electrocution risk with power line infrastructure in the area, but given the extent of the development, the contribution would be minimal.   |                        |
| <b>Residual Risks</b>   | Deterrent devices such as bird guards to reduce electrocutions, and flight diverters to reduce the risk of collisions with power lines are not 100% effective and some residual impact is likely to occur.  |                        |

**Impact 3. Habitat Degradation due to Erosion and Alien Plant Invasion**

|   |   |                        |
|---|---|------------------------|
| <b>Impact Nature:</b> Disturbance created during the construction phase will leave the site and immediate surroundings vulnerable to erosion and alien plant invasion for several years into the operation phase. |   |                        |
|   | <b>Without Mitigation</b>   | <b>With Mitigation</b> |
| <b>Extent</b>   | Local (1)   | Local (1)              |
| <b>Duration</b>   | Medium-term (2)   | Long-term (3)          |
| <b>Magnitude</b>  | Medium Low (3)  | Minor (2)              |
| <b>Probability</b>  | Probable (3)  | Improbable (2)         |
| <b>Significance</b>   | Low (18)  | Low (12)               |
| <b>Status</b>   | Negative  | Negative               |
| <b>Reversibility</b>  | Medium  | High                   |
| <b>Irreplaceable loss of resources</b>  | Low   | Low                    |
| <b>Can impacts be mitigated?</b>  | Yes, with proper management and avoidance, this impact can be mitigated to a low level.   |                        |
| <b>Mitigation</b>   | <ul style="list-style-type: none"> <li>• Erosion management within the development area should take place according to the Erosion Management Plan and Rehabilitation Plan of the project.</li> <li>• The site access road should have run-off control features which redirect water flow and dissipate any energy in the water which may pose an erosion risk.</li> <li>• Regular monitoring for erosion during operation to ensure that no erosion problems have developed as a result of the disturbance, as per the Erosion Management and Rehabilitation Plans for the project.</li> <li>• All erosion problems observed should be rectified as soon as possible, using the appropriate erosion control structures and revegetation techniques.</li> <li>• There should be follow-up rehabilitation and re-vegetation of any remaining bare areas with indigenous perennial shrubs and succulents from the local area.</li> <li>• Alien management at the site should take place in accordance with the Alien Invasive Management Plan of the project.</li> <li>• Regular monitoring for alien plant proliferation during the operation phase to ensure that no alien invasion problems have developed as result of the disturbance, as per the Alien Invasive Management Plan for the project.</li> <li>• If required, woody alien plant species should be controlled on at least an annual basis using the appropriate alien control techniques as determined by the species present.</li> </ul> |                        |

|                           |   |
|---------------------------|---|
| <b>Cumulative Impacts</b> | Erosion and alien plant invasion would contribute to degradation in the area, but as this can be well-mitigated, the contribution can be minimised. |
| <b>Residual Risks</b>     | Some erosion and alien plant invasion is likely to occur even with the implementation of control measures, but would have a low impact.             |

**5.3 DECOMMISSIONING PHASE**

**Decommissioning Phase Impact 1. Habitat Degradation due to Erosion and Alien Plant Invasion**

|  |   |                        |
|--|---|------------------------|
| <b>Impact Nature:</b> Disturbance created during decommissioning will leave the development area vulnerable to erosion and alien plant invasion for several years. |   |                        |
|  | <b>Without Mitigation</b>   | <b>With Mitigation</b> |
| <b>Extent</b>  | Local (1)   | Local (1)              |
| <b>Duration</b>  | Long-term (4)   | Long-term (3)          |
| <b>Magnitude</b>   | Medium (3)  | Minor (2)              |
| <b>Probability</b>   | Probable (3)  | Improbable (2)         |
| <b>Significance</b>  | Low (24)  | Low (12)               |
| <b>Status</b>  | Negative  | Negative               |
| <b>Reversibility</b>   | Low   | High                   |
| <b>Irreplaceable loss of resources</b>   | Moderate  | Low                    |
| <b>Can impacts be mitigated?</b>   | Yes, with proper management and avoidance, this impact can be mitigated to a low level.   |                        |
| <b>Mitigation</b>  | <ul style="list-style-type: none"> <li>• Erosion management within the development area should take place in accordance with the Erosion Management and Rehabilitation Plan of the project.</li> <li>• All erosion problems observed should be rectified as soon as possible, using the appropriate erosion control structures and revegetation techniques.</li> <li>• There should be follow-up rehabilitation and revegetation of any remaining bare areas with indigenous perennial shrubs, grasses and trees from the local area.</li> <li>• Alien management at the site should take place according to the Alien Invasive Management Plan. This should make provision for alien monitoring and management annually for at least 3 years after decommissioning. Woody aliens should be controlled using the appropriate alien control techniques as determined by the species present. This might include use of herbicides where no practical manual means are feasible.</li> </ul> |                        |
| <b>Cumulative Impacts</b>  | Erosion and alien plant invasion would contribute to degradation in the area, but as this can be well-mitigated, the contribution can be minimised.   |                        |

|                       |  |
|-----------------------|--|
| <b>Residual Risks</b> | Some erosion and alien plant invasion is likely to occur even with the implementation of control measures, but would have a low impact if effectively managed. |
|-----------------------|--|

***Decommissioning Phase Impact 2. Direct Faunal Impacts Due to Decommissioning Activities***

|   |   |                        |
|---|---|------------------------|
| <b>Impact Nature:</b> Due to disturbance, noise and the operation of heavy machinery, faunal disturbance due to decommissioning will extend beyond the footprint and impact adjacent areas to some degree. This will however be transient and restricted to the period while machinery is operational. In the long term, decommissioning should restore the ecological functioning and at least some habitat value to the affected areas. |   |                        |
|   | <b>Without Mitigation</b>   | <b>With Mitigation</b> |
| <b>Extent</b>   | Local (1)   | Local (1)              |
| <b>Duration</b>   | Short-term (1)  | Short-term (1)         |
| <b>Magnitude</b>  | Low (4)   | Minor (3)              |
| <b>Probability</b>  | Probable (3)  | Probable (3)           |
| <b>Significance</b>   | Low (18)  | Low (15)               |
| <b>Status</b>   | Negative  | Negative               |
| <b>Reversibility</b>  | Moderate  | Moderate               |
| <b>Irreplaceable loss of resources</b>  | No  | No                     |
| <b>Can impacts be mitigated?</b>  | Although the noise and disturbance generated at the site during decommissioning is probably largely unavoidable, this will be transient and ultimately the habitat should be restored to something useable by the local fauna.  |                        |
| <b>Mitigation</b>   | <ul style="list-style-type: none"> <li>• All personnel should undergo environmental induction with regards to fauna and, in particular, awareness about not harming or collecting species such as snakes, tortoises and owls, which are often persecuted out of superstition.</li> <li>• Any fauna threatened by the decommissioning activities should be removed to safety by an appropriately qualified environmental officer.</li> <li>• All vehicles should adhere to a low speed limit on site (30km/h for heavy vehicles and 40km/h for light vehicles) to avoid collisions with susceptible species such as snakes and tortoises.</li> <li>• All hazardous materials should be stored in the appropriate manner to prevent contamination of the site and ultimately removed from the site as part of decommissioning. Any accidental chemical, fuel and oil spills that occur at the site should be cleaned up in the appropriate manner as related to the nature of the spill.</li> </ul> |                        |

|                           |  |
|---------------------------|--|
|                           | <ul style="list-style-type: none"> <li>The site should be rehabilitated with locally occurring species to restore ecosystem structure and function.</li> </ul>                               |
| <b>Cumulative Impacts</b> | During the decommissioning, the associated disturbance would contribute to cumulative fauna disturbance and disruption in the area, but this would be transient and not of long-term impact. |
| <b>Residual Risks</b>     | Although some components of disturbance cannot be avoided, the site itself would have low faunal abundance at decommissioning and no significant residual impacts are likely.                |

**5.4 CUMULATIVE IMPACTS**

The following are the cumulative impacts assessed as being a likely consequence of the development of the Gunstfontein BESS. This is assessed in context of the extent of the proposed development area, other developments in the area, as well as general habitat loss and transformation resulting from agriculture and other activities in the area.

**Cumulative Impact 1. Impact on CBAs and broad-scale ecological processes**

|  |   |  |
|--|---|--|
| <b>Nature:</b> The development of Gunstfontein BESS will contribute to cumulative habitat loss within CBAs and other broad-scale cumulative impacts on ecological processes in the wider Roggeveld area. |   |  |
|  | <b>Overall impact of the proposed project considered in isolation</b>   | <b>Cumulative impact of the project and other projects in the area</b> |
| <b>Extent</b>  | Local (1)   | Local (2)  |
| <b>Duration</b>  | Long-term (4)   | Long-term (4)  |
| <b>Magnitude</b>   | Low (2)   | Moderate (5)   |
| <b>Probability</b>   | Improbable (2)  | Probable (3)   |
| <b>Significance</b>  | Low (14)  | Medium (33)  |
| <b>Status</b>  | Negative  | Negative   |
| <b>Reversibility</b>   | High  | Moderate   |
| <b>Irreplaceable loss of resources</b>   | Low   | Low  |
| <b>Can impacts be mitigated</b>  | To some degree, but the majority of the impact results from the presence of the various already approved WEFs which cannot be well mitigated. |  |
| <b>Mitigation:</b>   |   |  |
| <ul style="list-style-type: none"> <li>Ensure that sensitive habitats such as drainage features, are not within the development footprint of the BESS.</li> </ul>  |   |  |

- Ensure that the fencing around the facility is wildlife friendly and does not impede fauna from moving through the area or result in electrocutions.
- Ensure that an alien invasive management plan and erosion management plan compiled for the Wind Energy Facility or the BESS project is effectively implemented at the site.

## **6 CONCLUSION & RECOMMENDATIONS**

---

The vegetation within the Gunstfontein BESS consists of typical Roggeveld Shale Renosterveld which is considered to represent a moderately sensitive vegetation type due to its low total extent and relatively high abundance of plant SCC. The abundance of plant SCC within the study area was low and no species of very high concern were observed. The overall footprint of the BESS would be low and it is highly unlikely to compromise the local populations of any species of concern. In terms of fauna, there are few species of conservation concern that are known to be present in the wider area, and based on the location of the BESS within the low and medium sensitivity areas and the limited extent of the BESS, impact on such species would be minimal. The primary impact of the development on fauna would be minor habitat loss for the more common resident species. As such, no high long-term post-mitigation impacts on fauna are expected to occur. Although several avifaunal species of concern are confirmed present in the area and likely use the BESS location for foraging at least on occasion, the extent of the development is sufficiently low such that it would not generate significant habitat loss for any species of concern. Furthermore, there are no nesting sites or other important habitats within the affected area, with the result that overall impact on avifauna would be low. Consequently, the impacts of the development on fauna and flora are considered acceptable and would be of low significance after mitigation.

Although direct impacts on fauna and flora are considered potentially acceptable, the BESS falls within an area that has been classified as CBA 1. As these are areas that have been identified as being of significance for biodiversity maintenance and ecological processes, development in these areas is undesirable. The footprint of the development would be less than 5ha and would also be located near to the approved WEF substation, with the result that the additional extent of disturbance and habitat loss would be low. As a result, the low overall footprint of the BESS would be very unlikely to compromise the ecological functioning of the affected CBAs in any way beyond that which would already occur as a result of the Gunstfontein WEF.

Cumulative impacts within the broader study area are of potential concern due to the proliferation of WEF energy development in the wider Roggeveld area. The contribution of the BESS would however be very minor and is not considered to represent a significant contributor to cumulative impact in the area. Cumulative impacts associated with the development of the BESS are therefore considered acceptable.

***Impact Statement***

There are no impacts associated with the establishment of Gunstfontein BESS that cannot be mitigated to a low significance. Although cumulative impacts in the area are a concern due to the high density of wind energy developments in the wider area, the contribution of the Gunstfontein BESS would be low and is not considered to be of significance. As such, there are no fatal flaws or high post-mitigation impacts that should prevent the development from proceeding. Based on the location of the BESS as provided for this assessment and the sensitivity determined on site, the Gunstfontein BESS can be supported from a terrestrial and avifaunal ecology point of view at the current location, as well as any other area within the 500m assessment zone, provided no infrastructure is placed within regions of high or very high ecological sensitivity. The Gunstfontein BESS can therefore be supported from terrestrial and avifaunal ecology point of view.



## **7 Activities for Inclusion into the EMPr**

---

An Environmental Management Programme (EMPr) provides a link between the predicted impacts and mitigation measures recommended within the BA and the implementation and operational activities of a project. As the construction and operation of the Gunstfontein BESS may impact the environment, activities that pose a threat should be managed and mitigated so that unnecessary or preventable environmental impacts do not result. The primary objective of the EMPr is to detail actions required to address the impacts identified in the BA during the establishment, operation and rehabilitation of the proposed infrastructure. The EMPr provides an elaboration of how to implement the mitigation measures documented in the BA. As such the purpose of the EMPr can be outlined as follows:

- To outline mitigation measures and environmental specifications which are required to be implemented for the planning, establishment, rehabilitation and operation/maintenance phases of the project in order to minimise and manage the extent of environmental impacts.
- To ensure that the establishment and operation phases of the project do not result in undue or reasonably avoidable adverse environmental impacts, and ensure that any potential environmental benefits are enhanced.
- To identify entities who will be responsible for the implementation of the measures and outline functions and responsibilities.
- To propose mechanisms for monitoring compliance, and preventing long-term or permanent environmental degradation.
- To facilitate appropriate and proactive response to unforeseen events or changes in project implementation that were not considered in the BA process

Below are the ecologically-orientated measures that should be implemented as part of the EMPr for the development to reduce the significance or extent of the above impacts. The measures below do not exactly match with the impacts that have been identified, as certain mitigation measures, such as limiting the loss of vegetation may be effective at combating several different impacts, such as erosion, faunal impact etc.

**Construction Phase Activities**

| Objective: Limit disturbance of vegetation and loss of protected flora during construction  |   |                          |
|---|---|--------------------------|
| Potential Impact  | Loss of plant cover leading to erosion as well as loss of faunal habitat and loss of specimens of protected plants.   |                          |
| Activity/risk source  | Vegetation clearing for the following <ul style="list-style-type: none"> <li>» Clearing for infrastructure establishment.</li> <li>» Access roads.</li> <li>» Laydown areas.</li> <li>» Construction Camps.</li> </ul>  |                          |
| Mitigation:<br>Target/Objective   | <ul style="list-style-type: none"> <li>» Low footprint and low impact on terrestrial environment.</li> <li>» Low impact on protected plant species.</li> </ul>  |                          |
| Mitigation: Action/control  | Responsibility  | Timeframe                |
| <ul style="list-style-type: none"> <li>» Preconstruction walk-through of final BESS footprint must inform final micro-siting and search-and-rescue efforts.</li> <li>» Obtain relevant permits from the Department of Agriculture, Forestry and Fisheries (DEFF) and the Northern Cape Department of Agriculture, Environmental Affairs, Rural Development and Land Reform prior to any construction activities at the site.</li> <li>» Affected individuals of selected protected species which cannot be avoided should be translocated to a safe area on the site prior to construction. This does not include woody species which cannot be translocated and where these are protected by DAFF and a permit for their destruction would be required.</li> <li>» Erosion control measures should be implemented in areas where slopes have been disturbed.</li> <li>» Revegetation of cleared areas or monitoring to ensure that recovery is taking place.</li> <li>» Alien plant clearing where necessary.</li> </ul> | Management/EO   | Construction & Operation |
| Performance Indicator   | <ul style="list-style-type: none"> <li>» Vegetation loss restricted to infrastructure footprint.</li> <li>» Low impact on protected plant species.</li> <li>» Permit obtained to destroy or translocate affected individuals of protected species.</li> </ul> |                          |

|            |  |
|------------|--|
| Monitoring | ECO to monitor construction to ensure that: <ul style="list-style-type: none"> <li>» Vegetation is cleared only within essential areas.</li> <li>» Erosion risk is maintained at an acceptable level through flow regulation structures where appropriate and the maintenance of plant cover wherever possible.</li> </ul> |
|------------|--|

**Objective: Limit direct and indirect terrestrial faunal impacts during construction**

|  |  |              |  |
|--|--|--------------|--|
| Project component/s  | Construction activities especially the following: <ul style="list-style-type: none"> <li>» Vegetation clearing.</li> <li>» Human presence.</li> <li>» Operation of heavy machinery.</li> </ul>   |              |  |
| Potential Impact   | Disturbance of faunal communities due to construction as well as poaching and hunting risk from construction staff.  |              |  |
| Activity/risk source   | <ul style="list-style-type: none"> <li>» Habitat transformation during construction.</li> <li>» Presence of construction crews.</li> <li>» Operation of heavy vehicles.</li> </ul>   |              |  |
| Mitigation:<br>Target/Objective  | Low faunal impact during construction.   |              |  |
| Mitigation: Action/control   | Responsibility   | Timeframe    |  |
| <ul style="list-style-type: none"> <li>» Environmental induction for all construction staff</li> <li>» ECO to monitor and enforce a ban on hunting, collecting etc. of all plants and animals or their products.</li> <li>» Any fauna encountered during construction should be removed to safety by the EO or other suitably qualified person, or allowed to passively vacate the area.</li> <li>» All vehicles to adhere to low speed limits (40km/h max) on the site, to reduce risk of faunal collisions as well as reduce dust.</li> <li>» All night-lighting should use low-UV type lights (such as HPS bulbs), which do not attract insects. The lights should also be directed downward to ensure they do not result in large amounts of light pollution.</li> </ul> | Management/ECO   | Construction |  |
| Performance Indicator  | <ul style="list-style-type: none"> <li>» Low mortality of fauna due to construction machinery and activities.</li> <li>» No poaching etc of fauna by construction personnel during construction.</li> <li>» Removal to safety of fauna encountered during construction.</li> </ul> |              |  |
| Monitoring   | Monitoring for compliance during the construction phase. All incidents to be noted.  |              |  |

**Operational Phase Activities**

| OBJECTIVE: Limit the ecological footprint of the Gunstfontein BESS  |   |           |
|---|---|-----------|
| Project component/s   | Presence and operation of the facility including <ul style="list-style-type: none"> <li>» Movement of vehicles to and from the site.</li> </ul>   |           |
| Potential Impact  | <ul style="list-style-type: none"> <li>» Alien plant invasion</li> <li>» Erosion</li> <li>» Pollution</li> <li>» Faunal Impacts</li> </ul>  |           |
| Activity/risk source  | <ul style="list-style-type: none"> <li>» Alien plant invasion in and around the road.</li> <li>» Unregulated runoff from the access road.</li> <li>» Human presence during road maintenance activities</li> <li>» Pollution from maintenance vehicles due to oil or fuel leaks etc.</li> <li>» Maintenance activities which may lead to negative impacts such as pollution, herbicide drift etc.</li> </ul> |           |
| Mitigation:<br>Target/Objective   | Low ecological footprint of the grid connection infrastructure during operation.  |           |
| Mitigation: Action/control  | Responsibility  | Timeframe |
| Vegetation control should be by manual clearing and herbicides should not be used except to control alien plants in the prescribed manner.  | Management/<br>Contractor   | Operation |
| Annual monitoring for alien plant species - with follow up clearing as needed – or as per the frequency stated in the alien invasive management plan to be developed for the site (or that of the Wind Energy Facility utilised). | Management/<br>Contractor   | Operation |
| Annual site inspection for erosion or water flow regulation problems – with follow up remedial action where problems are identified.  | Management/<br>Contractor   | Operation |
| Performance<br>Indicator  | <ul style="list-style-type: none"> <li>» No erosion problems experience on the site</li> <li>» Low abundance of alien plants.</li> </ul>  |           |
| Monitoring  | <ul style="list-style-type: none"> <li>» Annual monitoring with records of alien species presence and clearing actions.</li> <li>» Annual monitoring with records of erosion problems and mitigation actions taken with photographs.</li> </ul>   |           |

## 8 REFERENCES

---

- Alexander, G. & Marais, J. 2007. *A Guide to the Reptiles of Southern Africa*. Struik Nature, Cape Town.
- Bates, M.F., Branch, W.R., Bauer, A.M., Burger, M., Marais, J., Alexander, G.J. & de Villiers, M. S. 2014. Atlas and Red List of the Reptiles of South Africa, Lesotho and Swaziland. Strelitzia 32. SANBI, Pretoria.
- Branch W.R. 1998. *Field guide to snakes and other reptiles of southern Africa*. Struik, Cape Town.
- Du Preez, L. & Carruthers, V. 2009. *A Complete Guide to the Frogs of Southern Africa*. Struik Nature., Cape Town.
- EWT & SANBI, 2016. Red List of Mammals of South Africa, Lesotho and Swaziland. EWT, Johannesburg.
- Marais, J. 2004. *Complete Guide to the Snakes of Southern Africa*. Struik Nature, Cape Town.
- Nel, J.L., Murray, K.M., Maherry, A.M., Petersen, C.P., Roux, D.J., Driver, A., Hill, L., Van Deventer, H., Funke, N., Swartz, E.R., Smith-Adao, L.B., Mbona, N., Downsborough, L. and Nienaber, S. (2011). Technical Report for the National Freshwater Ecosystem Priority Areas project. WRC Report No. K5/1801.
- Minter LR, Burger M, Harrison JA, Braack HH, Bishop PJ & Kloepfer D (eds). 2004. Atlas and Red Data book of the frogs of South Africa, Lesotho and Swaziland. SI/MAB Series no. 9. Smithsonian Institution, Washington, D.C.
- Mucina L. & Rutherford M.C. (eds) 2006. *The Vegetation of South Africa, Lesotho and Swaziland*. Strelitzia 19. South African National Biodiversity Institute, Pretoria.
- Oosthuysen, E. & Holness, S. 2016. Northern Cape Critical Biodiversity Areas (CBA) Map. Northern Cape Department of Environment and Nature Conservation & Nelson Mandela Metropolitan University. Available at SANBI BGIS <http://bgis.sanbi.org/>.
- Skinner, J.D. & Chimimba, C.T. 2005. The mammals of the Southern African Subregion. Cambridge University Press, Cambridge.

## Appendix 1. Listed Plant Species

List of plant species of conservation concern which are known to occur in the broad vicinity of the Gunstfontein BESS. The list is derived from the SIBIS:SABIF website. Only two of these species can be confirmed present within the BESS 500 assessment region.

| Family         | Species  | Threat status |
|----------------|--|---------------|
| AMARYLLIDACEAE | <i>Brunsvigia josephinae</i> (Redouté) Ker Gawl.                                     | VU            |
|                | <i>Strumaria karooica</i> (W.F.Barker) Snijman                                       | Rare          |
|                | <i>Strumaria pubescens</i> W.F.Barker  | Rare          |
| ANTHERICACEAE  | <i>Chlorophytum lewisiae</i> Oberm.  | Rare          |
| APOCYNACEAE    | <i>Duvalia parviflora</i> N.E.Br.  | VU            |
|                | <i>Hoodia pilifera</i> (L.f.) Plowes subsp. <i>pilifera</i>                          | NT            |
| ASPHODELACEAE  | <i>Astroloba herrei</i> Uitewaal   | VU            |
|                | <i>Bulbine torta</i> N.E.Br.   | Rare          |
|                | <i>Haworthia fasciata</i> (Willd.) Haw.  | NT            |
|                | <i>Gasteria disticha</i>   | CR            |
|                | <i>Haworthia serrata</i>   | CR            |
|                | <i>Haworthia pulchella</i> M.B.Bayer var. <i>pulchella</i>                           | Rare          |
| ASTERACEAE     | <i>Cineraria lobata</i> L'Hér. subsp. <i>lasiocaulis</i> Cron                        | Rare          |
|                | <i>Antithrixia flavicoma</i>   | VU            |
|                | <i>Euryops namaquensis</i>   | VU            |
|                | <b><i>Eriocephalus grandiflorus</i> M.A.N.Müll.</b>                                  | Rare          |
|                | <i>Phymaspermum thymelaeoides</i>  | LC            |
|                | <i>Pteronia hutchinsoniana</i> Compton   | Rare          |
|                | <i>Relhania tricephala</i> (DC.) K.Bremer  | NT            |
| COLCHICACEA    | <i>Wurmbea capensis</i>  | VU            |
| CRASSULACEAE   | <i>Adromischus humilis</i> (Marloth) Poelln.   | Rare          |
|                | <i>Adromischus phillipsiae</i> (Marloth) Poelln.                                     | Rare          |
|                | <i>Adromischus mammillaris</i>   | EN            |
|                | <i>Crassula alpestris</i> Thunb. subsp. <i>massonii</i> (Britten & Baker f.) Toelken | Rare          |
| EUPHORBIACEAE  | <i>Euphorbia nesemannii</i> R.A.Dyer   | NT            |
| FABACEAE       | <i>Amphithalea spinosa</i> (Harv.) A.L.Schutte                                       | VU            |
|                | <i>Amphithalea villosa</i> Schltr.   | VU            |
|                | <i>Lotononis comptonii</i> B.-E.van Wyk  | EN            |
|                | <i>Lotononis gracilifolia</i> B.-E.van Wyk   | EN            |
|                | <i>Lotononis venosa</i> B.-E.van Wyk   | VU            |
| GERANIACEAE    | <i>Pelargonium denticulatum</i> Jacq.  | Rare          |
|                | <i>Pelargonium torulosum</i> E.M.Marais  | Rare          |
| HYACINTHACEAE  | <i>Lachenalia maximiliani</i> Schltr. ex W.F.Barker                                  | Rare          |
| IRIDACEAE      | <i>Geissorhiza inaequalis</i> L.Bolus  | Rare          |
|                | <i>Geissorhiza karooica</i> Goldblatt  | NT            |

Fauna & Flora Specialist Report

---

|                            |  |      |
|----------------------------|--|------|
|                            | <i>Ixia linearifolia</i> Goldblatt & J.C.Manning         | Rare |
|                            | <i>Ixia parva</i> Goldblatt & J.C.Manning                | VU   |
|                            | <i>Moraea aspera</i> Goldblatt                           | VU   |
|                            | <i>Romulea eburnea</i> J.C.Manning & Goldblatt           | VU   |
|                            | <i>Romulea syringodeoflora</i> M.P.de Vos                | VU   |
| <b>MESEMBRYANTHEMACEAE</b> | <b><i>Cleretum lyratifolium</i> Ihlenf. &amp; Struck</b> | Rare |
|                            | <i>Lampranthus amoenus</i> (Salm-Dyck ex DC.) N.E.Br.    | EN   |
| <b>OXALIDACEAE</b>         | <i>Oxalis tenuipes</i> T.M.Salter var. <i>tenuipes</i>   | Rare |
| <b>POACEAE</b>             | <i>Ehrharta eburnea</i> Gibbs Russ.                      | NT   |
| <b>POLYGALACEAE</b>        | <i>Muraltia karroica</i> Levyns                          | VU   |
| <b>RUTACEAE</b>            | <i>Acmadenia argillophila</i> I.Williams                 | NT   |
|                            | <i>Globulariopsis wittebergensis</i> Compton             | Rare |
| <b>SCROPHULARIACEAE</b>    | <i>Oftia glabra</i> Compton                              | Rare |
|                            | <i>Selago albomontana</i> Hilliard                       | Rare |

---

## Appendix 2. List of Mammals

List of Mammals which potentially occur in or near the Gunstfontein BESS site. Taxonomy and habitat notes are derived from Skinner & Chimimba (2005), while conservation status is according to the EWT/SANBI 2016 listing. Confirmed sightings are those for the area and not the site *per se*.

| Scientific Name                          | Common Name              | Status         | Habitat  | Likelihood |
|--|--------------------------|----------------|--|------------|
| <b>Afrosoricida (Golden Moles):</b>      |                          |                |  |            |
| <i>Chlorotalpa sclateri</i>              | Sclater's Golden Mole    | LC             | Montane grasslands, scrub and forested kloofs of the Nama Karoo and grassland biomes   | High       |
| <i>Chrysochloris asiatica</i>            | Cape Golden Mole         | LC             | Coastal parts of the Northern and Western Cape   | High       |
| <b>Macroscledidea (Elephant Shrews):</b> |                          |                |  |            |
| <i>Macroscelides proboscideus</i>        | Round-eared Shrew        | Elephant<br>LC | Species of open country, with preference for shrub bush and sparse grass cover, also occur on hard gravel plains with sparse boulders for shelter, and on loose sandy soil provided there is some bush cover | High       |
| <i>Elephantulus edwardii</i>             | Cape Rock Elephant Shrew | LC             | From rocky slopes, with or without vegetation, from hard sandy ground bearing little vegetation, quite small rocky outcrops  | Confirmed  |
| <b>Tubulentata:</b>                      |                          |                |  |            |
| <i>Orycteropus afer</i>                  | Aardvark                 | LC             | Wide habitat tolerance, being found in open woodland, scrub and grassland, especially associated with sandy soil   | Confirmed  |
| <b>Hyracoidea (Hyraxes)</b>              |                          |                |  |            |
| <i>Procavia capensis</i>                 | Rock Hyrax               | LC             | Outcrops of rocks, especially granite formations and dolomite intrusions in the Karoo. Also erosion gullies  | Confirmed  |
| <b>Lagomorpha (Hares and Rabbits):</b>   |                          |                |  |            |
| <i>Bunolagus monticularis</i>            | Riverine Rabbit          | CR             | Confined to riparian bush on the narrow alluvial fringe of seasonally dry watercourses in the Central Karoo.   | V.Low      |
| <i>Pronolagus saundersiae</i>            | Hewitt's Red Rock Hare   | LC             | Confined to areas of krantzes, rocky hillsides, boulder-strewn koppies and rocky ravines   | Confirmed  |
| <i>Lepus capensis</i>                    | Cape Hare                | LC             | Dry, open regions, with palatable bush and grass   | Confirmed  |
| <i>Lepus saxatilis</i>                   | Scrub Hare               | LC             | Common in agriculturally developed areas, especially in crop-growing areas or in fallow lands where there is some bush development.  | High       |
| <b>Rodentia (Rodents):</b>               |                          |                |  |            |
| <i>Cryptomys hottentotus</i>             | African Mole Rat         | LC             | Wide diversity of substrates, from sandy soils to heavier compact substrates such as decomposed schists and stony soils  | Confirmed  |
| <i>Hystrix africaeaustralis</i>          | Cape Porcupine           | LC             | Catholic in habitat requirements.  | Confirmed  |



*Fauna & Flora Specialist Report*

|                                |                            |    |   |           |
|--------------------------------|----------------------------|----|---|-----------|
| <i>Graphiurus ocularis</i>     | Spectacled Dormouse        | LC | Associated with sandstones of Cape Fold mountains, which have many vertical and horizontal crevices.  | High      |
| <i>Acomys subspinosus</i>      | Cape Spiny Mouse           | LC | Associated with rocky areas on mountain slopes in Fynbos  | Low       |
| <i>Rhabdomys pumilio</i>       | Four-striped Grass Mouse   | LC | Essentially a grassland species, occurs in wide variety of habitats where there is good grass cover.  | High      |
| <i>Mus minutoides</i>          | Pygmy Mouse                | LC | Wide habitat tolerance  | High      |
| <i>Steatomys krebsii</i>       | Kreb's African Fat Mouse   | LC |   |           |
| <i>Micaelamys namaquensis</i>  | Namaqua Rock Mouse         | LC | Catholic in their habitat requirements, but where there are rocky koppies, outcrops or boulder-strewn hillsides they use these preferentially                                     | Confirmed |
| <i>Micaelamys granti</i>       | Grant's Rock Mouse         | LC | Restricted to the karoo where they are associated with rocky terrain.   | High      |
| <i>Parotomys brantsii</i>      | Brants's Whistling Rat     | LC | Associated with a dry sandy substrate in more arid parts of the Nama-karoo and Succulent Karoo. Species selects areas of low percentage of plant cover and areas with deep sands. | High      |
| <i>Parotomys littledalei</i>   | Littledale's Whistling Rat | LC | Riverine associations or associated with Lycium bushes or Psilocaulon absimile  | Low       |
| <i>Otomys unisulcatus</i>      | Bush Vlei Rat              | LC | Shrub and fynbos associations in areas with rocky outcrops Tend to avoid damp situations but exploit the semi-arid Karoo through behavioural adaptation.                          | Confirmed |
| <i>Desmodillus auricularis</i> | Cape Short-tailed Gerbil   | LC | Tend to occur on hard ground, unlike other gerbil species, with some cover of grass or karroid bush   | High      |
| <i>Gerbillurus paebe</i>       | Hairy-footed Gerbil        | LC | Gerbils associated with Nama and Succulent Karoo preferring sandy soil or sandy alluvium with a grass, scrub or light woodland cover  | High      |
| <i>Tatera afra</i>             | Cape Gerbil                | LC | Confined to areas of loose, sandy soils of sandy alluvium. Common on cultivated lands.  | Low       |
| <i>Malacothrix typica</i>      | Gerbil Mouse               | LC | Found predominantly in Nama and Succulent Karoo biomes, in areas with a mean annual rainfall of 150-500 mm.   | High      |
| <i>Dendromus melanotis</i>     | Grey Climbing Mouse        | LC | Often associated with stands of tall grass especially if thickened with bushes and other vegetation   | High      |
| <b>Primates:</b>               |                            |    |   |           |
| <i>Papio hamadryas</i>         | Chacma Baboon              | LC | Can exploit fynbos, montane grasslands, riverine courses in deserts, and simply need water and access to refuges.   | Confirmed |
| <b>Eulipotyphla (Shrews):</b>  |                            |    |   |           |
| <i>Myosorex varius</i>         | Forest Shrew               | LC | Prefers moist, densely vegetated habitat  | High      |

## Fauna & Flora Specialist Report

|                               |                         |    |   |              |
|-------------------------------|-------------------------|----|---|--------------|
| <i>Crocidura cyanea</i>       | Reddish-Grey Musk Shrew | LC | Occurs in relatively dry terrain, with a mean annual rainfall of less than 500 mm. Occur in karroid scrub and in fynbos often in association with rocks.      | High         |
| <b>Carnivora:</b>             |                         |    |   |              |
| <i>Proteles cristatus</i>     | Aardwolf                | LC | Common in the 100-600mm rainfall range of country, Nama-Karoo, Succulent Karoo Grassland and Savanna biomes   | Confirmed    |
| <i>Caracal caracal</i>        | Caracal                 | LC | Caracals tolerate arid regions, occur in semi-desert and karroid conditions   | Confirmed    |
| <i>Felis silvestris</i>       | African Wild Cat        | LC | Wide habitat tolerance.   | Confirmed    |
| <i>Panthera pardus</i>        | Leopard                 | VU | Wide habitat tolerance, associated with areas of rocky koppies and hills, mountain ranges and forest  | Low/Moderate |
| <i>Felis nigripes</i>         | Black-footed cat        | VU | Associated with arid country with MAR 100-500 mm, particularly areas with open habitat that provides some cover in the form of tall stands of grass or scrub. | High         |
| <i>Genetta genetta</i>        | Small-spotted genet     | LC | Occur in open arid associations   | High         |
| <i>Genetta tigrina</i>        | Large-spotted genet     | LC | Fynbos and savanna particularly along riverine areas  | Low          |
| <i>Suricata suricatta</i>     | Meerkat                 | LC | Open arid country where substrate is hard and stony. Occur in Nama and Succulent Karoo but also fynbos  | Confirmed    |
| <i>Cynictis penicillata</i>   | Yellow Mongoose         | LC | Semi-arid country on a sandy substrate  | Confirmed    |
| <i>Galerella pulverulenta</i> | Cape Grey Mongoose      | LC | Wide habitat tolerance  | Confirmed    |
| <i>Vulpes chama</i>           | Cape Fox                | LC | Associated with open country, open grassland, grassland with scattered thickets and coastal or semi-desert scrub  | High         |
| <i>Canis mesomelas</i>        | Black-backed Jackal     | LC | Wide habitat tolerance, more common in drier areas.   | Confirmed    |
| <i>Otocyon megalotis</i>      | Bat-eared Fox           | LC | Open country with mean annual rainfall of 100-600 mm  | Confirmed    |
| <i>Aonyx capensis</i>         | Cape Clawless Otter     | NT | Predominantly aquatic and do not occur far from permanent water   | Medium       |
| <i>Ictonyx striatus</i>       | Striped Polecat         | LC | Widely distributed throughout the sub-region  | Confirmed    |
| <i>Mellivora capensis</i>     | Ratel/Honey Badger      | LC | Catholic habitat requirements   | High         |
| <b>Rumanantia (Antelope):</b> |                         |    |   |              |
| <i>Sylvicapra grimmia</i>     | Common Duiker           | LC | Presence of bushes is essential   | Confirmed    |
| <i>Pelea capreolus</i>        | Grey Rhebok             | NT | Associated with rocky hills, rocky mountainsides, mountain plateaux with good grass cover.  | Confirmed    |
| <i>Antidorcas marsupialis</i> | Springbok               | LC | Arid regions and open grassland.  | Confirmed    |
| <i>Raphicerus campestris</i>  | Steenbok                | LC | Inhabits open country,  | Confirmed    |

*Fauna & Flora Specialist Report*

---

|                              |              |    |  |           |
|------------------------------|--------------|----|--|-----------|
| <i>Raphicerus melanotis</i>  | Cape Grysbok | LC | Thick scrub bush, particularly along the lower levels of hills | Medium    |
| <i>Oreotragus oreotragus</i> | Klipspringer | LC | Closely confined to rocky habitat.                             | Confirmed |

## Appendix 3. List of Reptiles.

List of reptiles which are known from the broad area around the Gunstfontein BESS site, according to the SARCA database. Species in bold are those observed at or near the site.

| Family            | Genus                  | Species               | Subspecies         | Common name                  | Red list category    |
|-------------------|------------------------|-----------------------|--------------------|------------------------------|----------------------|
| <b>Agamidae</b>   | <b>Agama</b>           | <b>atra</b>           |                    | <b>Southern Rock Agama</b>   | <b>Least Concern</b> |
| Agamidae          | <i>Agama</i>           | <i>hispida</i>        |                    | Spiny Ground Agama           | Least Concern        |
| Atractaspididae   | <i>Homoroselaps</i>    | <i>lacteus</i>        |                    | Spotted Harlequin Snake      | Least Concern        |
| Chamaeleonidae    | <i>Bradypodion</i>     | <i>gutturale</i>      |                    | Little Karoo Dwarf Chameleon | Least Concern        |
| Chamaeleonidae    | <i>Chamaeleo</i>       | <i>namaquensis</i>    |                    | Namaqua Chameleon            | Least Concern        |
| Colubridae        | <i>Psammophis</i>      | <i>crucifer</i>       |                    | Cross-marked Grass Snake     | Least Concern        |
| Colubridae        | <i>Pseudaspis</i>      | <i>cana</i>           |                    | Mole Snake                   | Least Concern        |
| Colubridae        | <i>Dasypeltis</i>      | <i>scabra</i>         |                    | Rhombic Egg-eater            | Least Concern        |
| Colubridae        | <i>Dipsina</i>         | <i>multimaculata</i>  |                    | Dwarf Beaked Snake           | Least Concern        |
| Cordylidae        | <i>Cordylus</i>        | <i>minor</i>          |                    | Western Dwarf Girdled Lizard | Least Concern        |
| Cordylidae        | <i>Hemicordylus</i>    | <i>capensis</i>       |                    | Graceful Crag Lizard         | Least Concern        |
| <b>Cordylidae</b> | <b>Karusasaurus</b>    | <b>polyzonus</b>      |                    | <b>Karoo Girdled Lizard</b>  | <b>Least Concern</b> |
| Cordylidae        | <i>Pseudocordylus</i>  | <i>microlepidotus</i> | <i>namaquensis</i> | Nuweveldberg Crag Lizard     | Least Concern        |
| Elapidae          | <i>Hemachatus</i>      | <i>haemachatus</i>    |                    | Rinkhals                     | Least Concern        |
| Elapidae          | <i>Naja</i>            | <i>nigricincta</i>    | <i>woodi</i>       | Black Spitting Cobra         | Least Concern        |
| Elapidae          | <i>Aspidelaps</i>      | <i>lubricus</i>       | <i>lubricus</i>    | Coral Shield Cobra           | Not Listed           |
| Gekkonidae        | <i>Chondrodactylus</i> | <i>angulifer</i>      | <i>angulifer</i>   | Common Giant Ground Gecko    | Least Concern        |
| <b>Gekkonidae</b> | <b>Chondrodactylus</b> | <b>bibronii</b>       |                    | <b>Bibron's Gecko</b>        | <b>Least Concern</b> |
| <b>Gekkonidae</b> | <b>Pachydactylus</b>   | <b>capensis</b>       |                    | <b>Cape Gecko</b>            | <b>Least Concern</b> |
| Gekkonidae        | <i>Pachydactylus</i>   | <i>formosus</i>       |                    | Southern Rough Gecko         | Least Concern        |
| Gekkonidae        | <i>Pachydactylus</i>   | <i>geitje</i>         |                    | Ocellated Gecko              | Least Concern        |
| Gekkonidae        | <i>Pachydactylus</i>   | <i>kladaroderma</i>   |                    | Thin-skinned Gecko           | Least Concern        |
| Gekkonidae        | <i>Pachydactylus</i>   | <i>maculatus</i>      |                    | Spotted Gecko                | Least Concern        |
| Gekkonidae        | <i>Pachydactylus</i>   | <i>mariquensis</i>    |                    | Marico Gecko                 | Least Concern        |
| Gekkonidae        | <i>Pachydactylus</i>   | <i>oculatus</i>       |                    | Golden Spotted Gecko         | Least Concern        |
| Gekkonidae        | <i>Pachydactylus</i>   | <i>purcelli</i>       |                    | Purcell's Gecko              | Least Concern        |
| Gekkonidae        | <i>Pachydactylus</i>   | <i>weberi</i>         |                    | Weber's Gecko                | Least Concern        |

|                     |                           |                             |                         |                                |                      |
|---------------------|---------------------------|-----------------------------|-------------------------|--------------------------------|----------------------|
| Gerrhosauridae      | <i>Cordylosaurus</i>      | <i>subtessellatus</i>       |                         | Dwarf Plated Lizard            | Least Concern        |
| Gerrhosauridae      | <i>Tetradactylus</i>      | <i>tetradactylus</i>        |                         | Cape Long-tailed Seps          | Least Concern        |
| Lacertidae          | <i>Nucras</i>             | <i>tessellata</i>           |                         | Western Sandveld Lizard        | Least Concern        |
| Lacertidae          | <i>Pedioplanis</i>        | <i>burchelli</i>            |                         | Burchell's Sand Lizard         | Least Concern        |
| <b>Lacertidae</b>   | <b><i>Pedioplanis</i></b> | <b><i>laticeps</i></b>      |                         | <b>Karoo Sand Lizard</b>       | <b>Least Concern</b> |
| <b>Lacertidae</b>   | <b><i>Pedioplanis</i></b> | <b><i>lineoocellata</i></b> | <b><i>pulchella</i></b> | <b>Common Sand Lizard</b>      | <b>Least Concern</b> |
| Leptotyphlopidae    | <i>Namibiana</i>          | <i>gracilior</i>            |                         | Slender Thread Snake           | Least Concern        |
| Lamprophiidae       | <i>Boaedon</i>            | <i>capensis</i>             |                         | Brown House Snake              | Least Concern        |
| Lamprophiidae       | <i>Prosymna</i>           | <i>sundevallii</i>          |                         | Sundevall's Shovel-snout       | Least Concern        |
| Lamprophiidae       | <i>Psammophis</i>         | <i>notostictus</i>          |                         | Karoo Sand Snake               | Least Concern        |
| Lamprophiidae       | <i>Psammophylax</i>       | <i>rhombeatus</i>           | <i>rhombeatus</i>       | Spotted Grass Snake            | Least Concern        |
| <b>Scincidae</b>    | <b><i>Trachylepis</i></b> | <b><i>capensis</i></b>      |                         | <b>Cape Skink</b>              | <b>Least Concern</b> |
| <b>Scincidae</b>    | <b><i>Trachylepis</i></b> | <b><i>sulcata</i></b>       | <b><i>sulcata</i></b>   | <b>Western Rock Skink</b>      | <b>Least Concern</b> |
| <b>Scincidae</b>    | <b><i>Trachylepis</i></b> | <b><i>variegata</i></b>     |                         | <b>Variiegated Skink</b>       | <b>Least Concern</b> |
| <b>Testudinidae</b> | <b><i>Chersina</i></b>    | <b><i>angulata</i></b>      |                         | <b>Angulate Tortoise</b>       | <b>Least Concern</b> |
| Testudinidae        | <i>Homopus</i>            | <i>areolatus</i>            |                         | Parrot-beaked Tortoise         | Least Concern        |
| Testudinidae        | <i>Homopus</i>            | <i>boulengeri</i>           |                         | Karoo Padloper                 | Near Threatened      |
| Testudinidae        | <i>Homopus</i>            | <i>femoralis</i>            |                         | Greater Padloper               | Least Concern        |
| <b>Testudinidae</b> | <b><i>Psammobates</i></b> | <b><i>tentorius</i></b>     | <b><i>tentorius</i></b> | <b>Karoo Tent Tortoise</b>     | <b>Not listed</b>    |
| Testudinidae        | <i>Psammobates</i>        | <i>tentorius</i>            | <i>verroxii</i>         | Verrox's Tent Tortoise         | Not listed           |
| Typhlopidae         | <i>Rhinotyphlops</i>      | <i>lalandei</i>             |                         | Delalande's Beaked Blind Snake | Least Concern        |
| <b>Viperidae</b>    | <b><i>Bitis</i></b>       | <b><i>arietans</i></b>      | <b><i>arietans</i></b>  | <b>Puff Adder</b>              | <b>Least Concern</b> |

#### Appendix 4. List of Amphibians

List of amphibians which potentially occur in or near the Gunstfontein BESS site. Taxonomy and habitat notes are from du Preez and Carruthers (2009) and conservation status from the IUCN 2020. (Status: LC = Least Concern, DD = Data Deficient).

| Scientific Name                  | Common Name       | Status         | Habitat   | Distribution  | Likelihood |
|----------------------------------|-------------------|----------------|---|---------------|------------|
| <i>Amietophrynus rangeri</i>     | Raucous Toad      | Not Threatened | Rivers and stream in grassland and fynbos   | Endemic       | High       |
| <i>Vandijkophrynus gariensis</i> | Karoo Toad        | Not Threatened | Karoo Scrub   | Widespread    | High       |
| <i>Xenopus laevis</i>            | Common Platanna   | Not Threatened | Any more or less permanent water  | Widespread    | High       |
| <i>Cacosternum boettgeri</i>     | Common Caco       | Not Threatened | Marshy areas, vleis and shallow pans  | Widespread    | High       |
| <i>Amietia fuscigula</i>         | Cape River Frog   | Not Threatened | Large still bodies of water or permanent streams and rivers.                              | Widespread    | Confirmed  |
| <i>Cacosternum karoicum</i>      | Karoo Caco        | DD             | Dry kloofs and valleys in the Karoo   | Endemic       | High       |
| <i>Cacosternum karoicum</i>      | Karoo Dainty Frog | DD             | Arid areas with unpredictable rainfall. Breeds in small streams as well as man-made dams. | Karoo Endemic | High       |
| <i>Tomopterna delalandii</i>     | Cape Sand Frog    | Not Threatened | Lowlands in fynbos and Succulent Karoo  | Endemic       | High       |
| <i>Tomopterna tandyi</i>         | Tandy's Sand Frog | Not Threatened | Nama karoo grassland and savanna  | Widespread    | High       |

## Appendix 5. List of Avifauna

A consolidated avifaunal list for the Gunstfontein BESS project site and surrounds, including conservation status (Taylor et al., 2015), and SABAP2 reporting rates (%). Codes for conservation status are: EN=Endangered; V = Vulnerable; NT = Near-threatened, and codes for endemism: E=Endemic, NE=Near-endemic.

| Ref | Common group  | Common species | Genus                | Species               | Conservation Status | 3230_2035 | 3230_2030 |
|-----|---------------|----------------|----------------------|-----------------------|---------------------|-----------|-----------|
| 269 | Avocet        | Pied           | <i>Recurvirostra</i> | <i>avosetta</i>       |                     | 4.8       | 0         |
| 432 | Barbet        | Acacia Pied    | <i>Tricholaema</i>   | <i>leucomelas</i>     |                     | 4.8       | 93.8      |
| 674 | Batis         | Pirit          | <i>Batis</i>         | <i>pririt</i>         |                     | 0         | 12.5      |
| 808 | Bishop        | Southern Red   | <i>Euplectes</i>     | <i>orix</i>           |                     | 0         | 6.3       |
| 722 | Bokmakierie   | Bokmakierie    | <i>Telophorus</i>    | <i>zeylonus</i>       |                     | 95.2      | 87.5      |
| 543 | Bulbul        | Cape           | <i>Pycnonotus</i>    | <i>capensis</i>       |                     | 23.8      | 96.9      |
| 871 | Bunting       | Lark-like      | <i>Emberiza</i>      | <i>impetuani</i>      |                     | 19        | 12.5      |
| 873 | Bunting       | Cape           | <i>Emberiza</i>      | <i>capensis</i>       |                     | 66.7      | 93.8      |
| 218 | Bustard       | Ludwig's       | <i>Neotis</i>        | <i>ludwigii</i>       | EN                  | 0         | 0         |
| 152 | Buzzard       | Jackal         | <i>Buteo</i>         | <i>rufofuscus</i>     |                     | 52.4      | 3.1       |
| 154 | Buzzard       | Steppe         | <i>Buteo</i>         | <i>vulpinus</i>       |                     | 0         | 0         |
| 857 | Canary        | Cape           | <i>Serinus</i>       | <i>canicollis</i>     |                     | 9.5       | 28.1      |
| 861 | Canary        | Black-headed   | <i>Serinus</i>       | <i>alario</i>         |                     | 47.6      | 25        |
| 865 | Canary        | White-throated | <i>Crithagra</i>     | <i>albogularis</i>    |                     | 52.4      | 68.8      |
| 866 | Canary        | Yellow         | <i>Crithagra</i>     | <i>flaviventris</i>   |                     | 52.4      | 9.4       |
| 566 | Chat          | Karoo          | <i>Cercomela</i>     | <i>schlegelii</i>     |                     | 14.3      | 37.5      |
| 570 | Chat          | Familiar       | <i>Cercomela</i>     | <i>familiaris</i>     |                     | 0         | 71.9      |
| 571 | Chat          | Tractrac       | <i>Cercomela</i>     | <i>tractrac</i>       |                     | 4.8       | 0         |
| 572 | Chat          | Sickle-winged  | <i>Cercomela</i>     | <i>sinuata</i>        |                     | 66.7      | 12.5      |
| 638 | Cisticola     | Grey-backed    | <i>Cisticola</i>     | <i>subruficapilla</i> |                     | 85.7      | 43.8      |
| 504 | Cliff-swallow | South African  | <i>Hirundo</i>       | <i>spilodera</i>      |                     | 9.5       | 0         |
| 212 | Coot          | Red-knobbed    | <i>Fulica</i>        | <i>cristata</i>       |                     | 42.9      | 0         |
| 47  | Cormorant     | White-breasted | <i>Phalacrocorax</i> | <i>carbo</i>          |                     | 4.8       | 0         |
| 50  | Cormorant     | Reed           | <i>Phalacrocorax</i> | <i>africanus</i>      |                     | 0         | 3.1       |
| 621 | Crombec       | Long-billed    | <i>Sylvietta</i>     | <i>rufescens</i>      |                     | 14.3      | 65.6      |
| 522 | Crow          | Pied           | <i>Corvus</i>        | <i>albus</i>          |                     | 66.7      | 0         |
| 523 | Crow          | Cape           | <i>Corvus</i>        | <i>capensis</i>       |                     | 4.8       | 0         |
| 314 | Dove          | Red-eyed       | <i>Streptopelia</i>  | <i>semitorquata</i>   |                     | 0         | 21.9      |
| 317 | Dove          | Laughing       | <i>Streptopelia</i>  | <i>senegalensis</i>   |                     | 0         | 90.6      |
| 318 | Dove          | Namaqua        | <i>Oena</i>          | <i>capensis</i>       |                     | 0         | 6.3       |
| 95  | Duck          | African Black  | <i>Anas</i>          | <i>sparsa</i>         |                     | 0         | 21.9      |
| 96  | Duck          | Yellow-billed  | <i>Anas</i>          | <i>undulata</i>       |                     | 71.4      | 3.1       |
| 133 | Eagle         | Verreaux's     | <i>Aquila</i>        | <i>verreauxii</i>     | VU                  | 47.6      | 21.9      |
| 139 | Eagle         | Booted         | <i>Aquila</i>        | <i>pennatus</i>       |                     | 0         | 3.1       |

*Fauna & Flora Specialist Report*

---

|      |               |                        |                      |                       |    |      |      |
|------|---------------|------------------------|----------------------|-----------------------|----|------|------|
| 142  | Eagle         | Martial                | <i>Polemaetus</i>    | <i>bellicosus</i>     | EN | 4.8  | 0    |
| 367  | Eagle-owl     | Cape                   | <i>Bubo</i>          | <i>capensis</i>       |    | 4.8  | 0    |
| 600  | Eremomela     | Yellow-bellied         | <i>Eremomela</i>     | <i>icteropygialis</i> |    | 14.3 | 3.1  |
| 626  | Eremomela     | Karoo                  | <i>Eremomela</i>     | <i>gregalis</i>       |    | 9.5  | 0    |
| 707  | Fiscal        | Common (Southern)      | <i>Lanius</i>        | <i>collaris</i>       |    | 47.6 | 65.6 |
| 665  | Flycatcher    | Fiscal                 | <i>Sigelus</i>       | <i>silens</i>         |    | 0    | 9.4  |
| 678  | Flycatcher    | Fairy                  | <i>Stenostira</i>    | <i>scita</i>          |    | 4.8  | 59.4 |
| 176  | Francolin     | Grey-winged            | <i>Scleroptila</i>   | <i>africanus</i>      |    | 38.1 | 34.4 |
| 88   | Goose         | Spur-winged            | <i>Plectropterus</i> | <i>gambensis</i>      |    | 33.3 | 3.1  |
| 89   | Goose         | Egyptian               | <i>Alopochen</i>     | <i>aegyptiacus</i>    |    | 71.4 | 15.6 |
| 165  | Goshawk       | Southern Pale Chanting | <i>Melierax</i>      | <i>canorus</i>        |    | 28.6 | 25   |
| 263  | Greenshank    | Common                 | <i>Tringa</i>        | <i>nebularia</i>      |    | 23.8 | 0    |
| 192  | Guineafowl    | Helmeted               | <i>Numida</i>        | <i>meleagris</i>      |    | 0    | 0    |
| 72   | Hamerkop      | Hamerkop               | <i>Scopus</i>        | <i>umbretta</i>       |    | 4.8  | 31.3 |
| 171  | Harrier-Hawk  | African                | <i>Polyboroides</i>  | <i>typus</i>          |    | 0    | 6.3  |
| 54   | Heron         | Grey                   | <i>Ardea</i>         | <i>cinerea</i>        |    | 14.3 | 15.6 |
| 55   | Heron         | Black-headed           | <i>Ardea</i>         | <i>melanocephala</i>  |    | 4.8  | 0    |
| 418  | Hoopoe        | African                | <i>Upupa</i>         | <i>africana</i>       |    | 0    | 21.9 |
| 81   | Ibis          | African Sacred         | <i>Threskiornis</i>  | <i>aethiopicus</i>    |    | 9.5  | 18.8 |
| 84   | Ibis          | Hadedda                | <i>Bostrychia</i>    | <i>hagedash</i>       |    | 19   | 43.8 |
| 123  | Kestrel       | Rock                   | <i>Falco</i>         | <i>rupicolus</i>      |    | 71.4 | 18.8 |
| 245  | Lapwing       | Blacksmith             | <i>Vanellus</i>      | <i>armatus</i>        |    | 85.7 | 0    |
| 461  | Lark          | Karoo                  | <i>Calendulauda</i>  | <i>albescens</i>      |    | 4.8  | 0    |
| 463  | Lark          | Large-billed           | <i>Galerida</i>      | <i>magnirostris</i>   |    | 81   | 3.1  |
| 474  | Lark          | Spike-heeled           | <i>Chersomanes</i>   | <i>albofasciata</i>   |    | 19   | 0    |
| 488  | Lark          | Red-capped             | <i>Calandrella</i>   | <i>cinerea</i>        |    | 38.1 | 3.1  |
| 4127 | Lark          | Karoo Long-billed      | <i>Certhilauda</i>   | <i>subcoronata</i>    |    | 14.3 | 3.1  |
| 4140 | Lark          | Cape Clapper           | <i>Mirafra</i>       | <i>apiata</i>         |    | 23.8 | 0    |
| 506  | Martin        | Rock                   | <i>Hirundo</i>       | <i>fuligula</i>       |    | 42.9 | 75   |
| 509  | Martin        | Brown-throated         | <i>Riparia</i>       | <i>paludicola</i>     |    | 4.8  | 3.1  |
| 803  | Masked-weaver | Southern               | <i>Ploceus</i>       | <i>velatus</i>        |    | 9.5  | 59.4 |
| 391  | Mousebird     | White-backed           | <i>Colius</i>        | <i>colius</i>         |    | 0    | 78.1 |
| 392  | Mousebird     | Red-faced              | <i>Urocolius</i>     | <i>indicus</i>        |    | 4.8  | 53.1 |
| 531  | Penduline-tit | Cape                   | <i>Anthoscopus</i>   | <i>minutus</i>        |    | 9.5  | 0    |
| 311  | Pigeon        | Speckled               | <i>Columba</i>       | <i>guinea</i>         |    | 9.5  | 78.1 |
| 692  | Pipit         | African                | <i>Anthus</i>        | <i>cinnamomeus</i>    | NT | 23.8 | 0    |
| 237  | Plover        | Kittlitz's             | <i>Charadrius</i>    | <i>pecuarius</i>      |    | 47.6 | 0    |
| 238  | Plover        | Three-banded           | <i>Charadrius</i>    | <i>tricoloris</i>     |    | 76.2 | 53.1 |
| 4139 | Prinia        | Karoo                  | <i>Prinia</i>        | <i>maculosa</i>       |    | 47.6 | 81.3 |
| 524  | Raven         | White-necked           | <i>Corvus</i>        | <i>albicollis</i>     |    | 57.1 | 31.3 |
| 606  | Reed-warbler  | African                | <i>Acrocephalus</i>  | <i>baeticatus</i>     |    | 0    | 15.6 |
| 581  | Robin-chat    | Cape                   | <i>Cossypha</i>      | <i>caffra</i>         |    | 14.3 | 96.9 |
| 307  | Sandgrouse    | Namaqua                | <i>Pterocles</i>     | <i>namaqua</i>        |    | 14.3 | 0    |



## Fauna & Flora Specialist Report

|      |             |                          |                      |                       |    |      |      |
|------|-------------|--------------------------|----------------------|-----------------------|----|------|------|
| 264  | Sandpiper   | Wood                     | <i>Tringa</i>        | <i>glareola</i>       |    | 4.8  | 0    |
| 583  | Scrub-robin | Karoo                    | <i>Cercotrichas</i>  | <i>coryphoeus</i>     |    | 90.5 | 53.1 |
| 90   | Shelduck    | South African            | <i>Tadorna</i>       | <i>cana</i>           |    | 42.9 | 25   |
| 94   | Shoveler    | Cape                     | <i>Anas</i>          | <i>smithii</i>        |    | 23.8 | 0    |
| 146  | Snake-eagle | Black-chested            | <i>Circaetus</i>     | <i>pectoralis</i>     |    | 4.8  | 0    |
| 784  | Sparrow     | House                    | <i>Passer</i>        | <i>domesticus</i>     |    | 0    | 87.5 |
| 786  | Sparrow     | Cape                     | <i>Passer</i>        | <i>melanurus</i>      |    | 52.4 | 84.4 |
| 4142 | Sparrow     | Southern Grey-headed     | <i>Passer</i>        | <i>diffusus</i>       |    | 0    | 18.8 |
| 85   | Spoonbill   | African                  | <i>Platalea</i>      | <i>alba</i>           |    | 4.8  | 0    |
| 181  | Spurfowl    | Cape                     | <i>Pternistis</i>    | <i>capensis</i>       |    | 52.4 | 43.8 |
| 733  | Starling    | Common                   | <i>Sturnus</i>       | <i>vulgaris</i>       |    | 0    | 3.1  |
| 744  | Starling    | Pale-winged              | <i>Onychognathus</i> | <i>nabourop</i>       |    | 47.6 | 78.1 |
| 746  | Starling    | Pied                     | <i>Spreo</i>         | <i>bicolor</i>        |    | 23.8 | 40.6 |
| 270  | Stilt       | Black-winged             | <i>Himantopus</i>    | <i>himantopus</i>     |    | 14.3 | 0    |
| 253  | Stint       | Little                   | <i>Calidris</i>      | <i>minuta</i>         |    | 19   | 0    |
| 79   | Stork       | Black                    | <i>Ciconia</i>       | <i>nigra</i>          | VU | 0    | 3.1  |
| 751  | Sunbird     | Malachite                | <i>Nectarinia</i>    | <i>famosa</i>         |    | 38.1 | 53.1 |
| 760  | Sunbird     | Southern Double-collared | <i>Cinnyris</i>      | <i>chalybeus</i>      |    | 23.8 | 81.3 |
| 764  | Sunbird     | Dusky                    | <i>Cinnyris</i>      | <i>fuscus</i>         |    | 4.8  | 0    |
| 493  | Swallow     | Barn                     | <i>Hirundo</i>       | <i>rustica</i>        |    | 19   | 0    |
| 502  | Swallow     | Greater Striped          | <i>Hirundo</i>       | <i>cucullata</i>      |    | 57.1 | 40.6 |
| 378  | Swift       | Common                   | <i>Apus</i>          | <i>apus</i>           |    | 9.5  | 0    |
| 383  | Swift       | White-rumped             | <i>Apus</i>          | <i>caffer</i>         |    | 23.8 | 31.3 |
| 385  | Swift       | Little                   | <i>Apus</i>          | <i>affinis</i>        |    | 9.5  | 40.6 |
| 386  | Swift       | Alpine                   | <i>Tachymarptis</i>  | <i>melba</i>          |    | 0    | 6.3  |
| 97   | Teal        | Red-billed               | <i>Anas</i>          | <i>erythrorhyncha</i> |    | 19   | 0    |
| 98   | Teal        | Cape                     | <i>Anas</i>          | <i>capensis</i>       |    | 4.8  | 3.1  |
| 305  | Tern        | Whiskered                | <i>Chlidonias</i>    | <i>hybrida</i>        |    | 9.5  | 0    |
| 1104 | Thrush      | Karoo                    | <i>Turdus</i>        | <i>smithi</i>         |    | 0    | 15.6 |
| 525  | Tit         | Grey                     | <i>Parus</i>         | <i>afer</i>           |    | 33.3 | 6.3  |
| 658  | Tit-babbler | Chestnut-vented          | <i>Parisoma</i>      | <i>subcaeruleum</i>   |    | 0    | 3.1  |
| 659  | Tit-babbler | Layard's                 | <i>Parisoma</i>      | <i>layardi</i>        |    | 38.1 | 62.5 |
| 316  | Turtle-dove | Cape                     | <i>Streptopelia</i>  | <i>capicola</i>       |    | 19   | 65.6 |
| 686  | Wagtail     | Cape                     | <i>Motacilla</i>     | <i>capensis</i>       |    | 71.4 | 78.1 |
| 619  | Warbler     | Rufous-eared             | <i>Malcorus</i>      | <i>pectoralis</i>     |    | 19   | 0    |
| 653  | Warbler     | Namaqua                  | <i>Phragmacia</i>    | <i>substriata</i>     |    | 0    | 21.9 |
| 843  | Waxbill     | Common                   | <i>Estrilda</i>      | <i>astrild</i>        |    | 4.8  | 31.3 |
| 799  | Weaver      | Cape                     | <i>Ploceus</i>       | <i>capensis</i>       |    | 9.5  | 93.8 |
| 564  | Wheatear    | Mountain                 | <i>Oenanthe</i>      | <i>monticola</i>      |    | 85.7 | 100  |
| 568  | Wheatear    | Capped                   | <i>Oenanthe</i>      | <i>pileata</i>        |    | 28.6 | 0    |
| 1172 | White-eye   | Cape                     | <i>Zosterops</i>     | <i>virans</i>         |    | 14.3 | 81.3 |
| 445  | Woodpecker  | Ground                   | <i>Geocolaptes</i>   | <i>olivaceus</i>      |    | 52.4 | 12.5 |

