

ECOLOGICAL FAUNA AND FLORA HABITAT SURVEY

**REMAINDER OF THE FARM MATJESSPRUIT
NO. 145, CITY OF MATLOSANA, NORTH WEST
PROVINCE**



New leaves of spring of *Searsia pyroides* (common wild currant/ taaibos) at the site.
Photo: R.F. Terblanche, October 2013

OCTOBER 2013

COMPILED BY:

Reinier F Terblanche

(M.Sc Ecology, *Cum Laude*; Pr.Sci.Nat, Reg. No. 400244/05)

ANTHENE ECOLOGICAL CC

TABLE OF CONTENTS

1. INTRODUCTION	1
2. STUDY AREA	2
3. METHODS	5
4. RESULTS	7
5. DISCUSSION	22
6. IMPACT ASSESSMENT AND MITIGATION MEASURES	28
7. CONCLUSION	32
8. REFERENCES	33
9. ANNEXURE 1 LIST OF PLANT SPECIES	44

1 INTRODUCTION

An ecological habitat survey was required for a proposed photo-voltaic development at Matjesspruit 18 km east-north-east of Leeudoringstad in the North West Province (elsewhere referred to as the site). The survey focused on the possibility that threatened fauna or flora known to occur in North West Province are likely to occur within the proposed development. Species of known high conservation priority that do not qualify for threatened status also received attention in the survey.

1.1 OBJECTIVES OF THE HABITAT STUDY

The objectives of the habitat study are to provide:

- A detailed fauna and flora habitat survey;
- A detailed habitat survey of possible threatened or localised plant species, vertebrates and invertebrates;
- Recording of possible host plants or foodplants of fauna such as butterflies.
- Evaluate the conservation importance and significance of the site with special emphasis on the current status of threatened species;
- Literature investigation of possible species that may occur on site;
- Identification of potential ecological impacts on fauna and flora that could occur as a result of the development; and
- Make recommendations to reduce or minimise impacts, should the development be approved.

1.2 SCOPE OF STUDY

- A survey consisting of two visits to investigate key elements of habitats on the site, relevant to the conservation of fauna and flora.
- Recording of any sightings and/or evidence of existing fauna and flora.
- The selective and careful collecting of voucher specimens of invertebrates where deemed necessary.
- An evaluation of the conservation importance and significance of the site with special emphasis on the current status of threatened species.
- Recording of possible host plants or foodplants of fauna such as butterflies.
- Literature investigation of possible species that might occur on site.
- Integration of the literature investigation and field observations to identify potential ecological impacts that could occur as a result of the development.
- Integration of literature investigation and field observations to make recommendations to reduce or minimise impacts, should the development be approved.

2 STUDY AREA

The study area is 18 km east-north-east Leeudoringstad in the North West Province. The study site is situated at the Grassland Biome which is represented by the Vaal-Vet Sandy Grassland vegetation type (Mucina & Rutherford 2006). A brief overview of the Vaal-Vet Sandy Grassland vegetation type in which the site is located, follows:

Distribution: In South Africa the Vaal-Vet Sandy Grassland is present in the North-West Province and Free State Province. Vaal-Vet Sandy Grassland ranges from south of Lichtenburg and Ventersdorp to Klerksdorp, Leeudoringstad, Bothaville and to the Brandfort areas north of Bloemfontein. Altitude ranges from 1 220 – 1560 m for the entire vegetation type (Mucina & Rutherford 2006).

Vegetation and landscape features: Plains-dominated landscape with some scattered, slightly undulating plains and hills. Mainly low-tussock grasslands with an abundant karroid element are present. Dominance of *Themeda triandra* is an important feature of this vegetation unit. Locally low cover of *Themeda triandra* and the associated increase in *Elionurus muticus*, *Cymbopogon pospischilii* and *Aristida congesta* is attributed to heavy grazing and/or erratic rainfall. **Geology and soils:** Aeolian and colluvial sand overlying sandstone, mudstone, and shale of the Karoo Supergroup (mostly the Ecca group) as well as older Ventersdorp Supergroup and basement gneiss in the north (Mucina & Rutherford 2006).

Climate: Warm-temperate, summer-rainfall climate, with overall mean annual precipitation of 530 mm. High summer temperatures. Severe frost (37 days per year on average) occurs in winter (Mucina & Rutherford 2006).

Important taxa of the Vaal-Vet Sandy Grassland listed by Mucina & Rutherford (2006):
Graminoids: *Antephora pubescens*, *Aristida congesta*, *Chloris virgata*, *Cymbopogon caesius*, *Cynodon dactylon*, *Digitaria argyrograpta*, *Elionurus muticus*, *Eragrostis chloromelas*, *Eragrostis lehmanniana*, *Eragrostis plana*, *Eragrostis trichophora*, *Heteropogon contortus*, *Panicum gilvum*, *Setaria sphacelata*, *Themeda triandra*, *Tragus berteronianus*, *Brachiaria serrata*, *Cymbopogon pospischilii*, *Digitaria eriantha*, *Eragrostis curvula*, *Eragrostis obtusa*, *Eragrostis superba*, *Panicum coloratum*, *Pogonarthria squarrosa*, *Trichoneura grandiglumis*, *Triraphis andropogonoides*. **Herbs:** *Stachys spathulata*, *Barleria macrostegia*, *Berkheya onopordifolia* var. *onopordifolia*, *Chamaesyce inaequilatera*, *Geigeria aspera* var. *aspera*, *Helichrysum caespitium*, *Hermannia depressa*, *Hibiscus pusillus*, *Monsonia burkeana*, *Rhynchosia adenodes*, *Selago densiflora*, *Vernonia*

oligocephala. Geophytic Herbs: *Bulbine narcissifolia*, *Ledebouria marginata*. Succulent Herb: *Tripteris aghillana* var. *integrifolia*. Low shrubs: *Felicia muricata*, *Pentzia globosa*, *Anthospermum rigidum* subsp. *pumilum*, *Helichrysum dregeanum*, *Helichrysum paronychioides*, *Ziziphus zeyheriana*.

According to the National List of Threatened Ecosystems (2011) the Vaal-Sandy is a threatened ecosystem listed as Endangered.

3 METHODS

A desktop study comprised not only an initial phase, but also it was used throughout the study to accommodate and integrate all the data that become available during the field observations.

A survey that consisted of visits by R.F. Terblanche on 30 August 2013 and 8 October 2013 was conducted to note key elements of habitats on the site, relevant to the conservation of fauna and flora. The main purpose of the site visit was ultimately to serve as a habitat survey that concentrated on the possible presence or not of threatened species and other species of high conservation priority.

The following sections highlight the materials and methods applicable to different aspects that were observed.

3.1 HABITAT CHARACTERISTICS AND VEGETATION

The habitat was investigated by noting habitat structure (rockiness, slope, plant structure/physiognomy) as well as floristic composition. Voucher specimens of plant species were only taken where the taxonomy was in doubt and where the plant specimens were of significant relevance for invertebrate conservation. In this case no plant specimens were needed to be collected as voucher specimens or to be send to a herbarium for identification. A wealth of guides and detailed works of plant identifications, ecology and conservation is fortunately available and very useful. Field guides, biogeographic works, species lists, diagnostic outlines, conservation statuses and detail on specific plant groups were sourced from Boon (2010), Court (2010), Germishuizen (2003), Germishuizen, Meyer & Steenkamp (2006), Goldblatt (1986), Goldblatt & Manning (1998), Jacobsen (1983), Manning (2003), Manning (2009), McMurtry, Grobler, Grobler

& Burns (2008), Pooley (1998), Retief & Herman (1997), Smit (2008), Van Ginkel, Glen, Gordon-Gray, Cilliers, Muasya & Van Deventer (2011), Van Jaarsveld (2006), Van Oudtshoorn (1999), Van Wyk (2000), Van Wyk & Smith (2001), Van Wyk & Smith (2003), Van Wyk & Malan (1998) and Van Wyk & Van Wyk (1997). Lists of species, species names and the conservation status of species were mainly sourced from Raimondo, von Staden, Victor, Helme, Turner, Kamundi & Manyama (2009) and updated versions of red lists and species from the Threatened Species Programme of SANBI and the Red List of South African Plants (sanbi.org.za).

3.2 MAMMALS

Mammals were noted as sight records by day. For the identification of species and observation of diagnostic characteristics Smithers (1986), Skinner & Chimimba (2005), Cillié, Oberprieler and Joubert (2004) and Apps (2000) are consulted. Sites have been walked, covering as many habitats as possible. Signs of the presence of mammal species, such as calls of animals, animal tracks (spoor), burrows, runways, nests and faeces were recorded. Walker (1996), Stuart & Stuart (2000) and Liebenberg (1990) were consulted for additional information and for the identification of spoor and signs. Trapping was not done since it proved not necessary in the case of this study. Habitat characteristics were also surveyed to note potential occurrences of mammals. Many mammals can be identified from field sightings but, with a few exceptions bats, rodents and shrews can only be reliably identified in the hand, and even then some species needs examination of skulls, or even chromosomes (Apps, 2000).

3.3 BIRDS

Birds were noted as sight records, mainly with the aid of binoculars (10x30). Nearby bird calls of which the observer was sure of the identity were also recorded. For practical skills of noting diagnostic characteristics, the identification of species and observation techniques Ryan (2001) is followed. For information on identification, biogeography and ecology Barnes (2000), Hockey, Dean & Ryan, P.G. (2005), Cillié, Oberprieler & Joubert (2004), Tarboton & Erasmus (1998) and Chittenden (2007) were consulted. Ringing of birds fell beyond the scope of this survey and was not deemed necessary. Sites have been walked, covering as many habitats as possible. Signs of the presence of bird species such as spoor and nests have additionally been recorded. Habitat characteristics were surveyed to note potential occurrences of birds.

3.4 REPTILES

Reptiles were noted as sight records in the field. Binoculars (10x30) can also be used for identifying reptiles of which some are wary. For practical skills of noting diagnostic characteristics, the identification of species and observation techniques, Branch (1998), Marais (2004), Alexander & Marais (2007) and Cillié, Oberprieler and Joubert (2004) were followed. Sites were walked, covering as many habitats as possible. Smaller reptiles are sometimes collected for identification, but this practice was not necessary in the case of this study. Habitat characteristics are surveyed to note potential occurrences of reptiles.

3.5 AMPHIBIANS

Frogs and toads are noted as sight records in the field or by their calls. For practical skills of noting diagnostic characteristics, the identification of species and observation techniques Carruthers (2001), Du Preez (1996), Conradie, Du Preez, Smith & Weldon (2006) and the recent complete guide by Du Preez & Carruthers (2009) are consulted. CD's with frog calls by Carruthers (2001) and Du Preez & Carruthers (2009) are used to identify species by their calls when applicable. Sites are walked, covering as many habitats as possible. Smaller frogs are often collected by pitfall traps put out for epigeal invertebrates (on the soil), but this practice falls beyond the scope of this survey. Habitat characteristics are also surveyed to note potential occurrences of amphibians.

3.6 BUTTERFLIES

Butterflies were noted as sight records or voucher specimens. Voucher specimens are mostly taken of those species of which the taxa warrant collecting due to taxonomic difficulties or in the cases where species can look similar in the veldt. Many butterflies use only one species or a limited number of plant species as host plants for their larvae. Myrmecophilous (ant-loving) butterflies such as the *Aloeides*, *Chrysothrix*, *Erikssonina*, *Lepidochrysops* and *Orachrysops* species (Lepidoptera: Lycaenidae), which live in association with a specific ant species, require a unique ecosystem for their survival (Deutschländer & Bredenkamp, 1999; Terblanche, Morghental & Cilliers, 2003; Edge, Cilliers & Terblanche, 2008; Gardiner & Terblanche, 2010). Known food plants of butterflies were therefore also recorded. After the visits to the site and the identification of the butterflies found there, a list was also compiled of butterflies that will most probably be

found in the area in all the other seasons because of suitable habitat. The emphasis is on a habitat survey.

3.7 FRUIT CHAFER BEETLES

Different habitat types in the areas were explored for any sensitive or special fruit chafer species. Selection of methods to find fruit chafers depends on the different types of habitat present and the species that may be present. Fruit bait traps would probably not be successful for capturing *Ichnestoma* species in a grassland patch (Holm & Marais 1992). Possible chafer beetles of high conservation priority were noted as sight records accompanied by the collecting of voucher specimens with grass nets or containers where deemed necessary.

3.8 ROCK SCORPIONS

Relatively homogenous habitat / vegetation areas were identified and explored to identify any sensitive or special species. Selected stones that were lifted to search for Arachnids were put back very carefully resulting in the least disturbance possible. All the above actions were accompanied by the least disturbance possible.

3.9 LIMITATIONS

For each site visited, it should be emphasized that surveys can by no means result in an exhaustive list of the plants and animals present on the site, because of the time constraint. The on site invertebrate survey was conducted during August 2013 and October 2013, the latter which is an optimal time of the year to find animals such as invertebrates as well as habitat sensitive plant and vertebrate animal species high conservation priority. Weather conditions during the survey, though dry, were favourable for recording fauna and flora. The focus of the survey remains a habitat survey that concentrates on the possibility that species of particular conservation priority occur on the site or not. It is unlikely that any more visits would reveal information that would change the outcome of this assessment both in terms of ecosystems of special conservation concern or suitable habitats of species of particular conservation concern. Visits that were conducted therefore appear to be sufficient to address the objectives of this study.

4 RESULTS

4.1 HABITAT AND VEGETATION CHARACTERISTICS

Table 4.1 Outline of main landscape and habitat characteristics of the site.

HABITAT FEATURE	DESCRIPTION
Topography	The site proposed for the developments is on very gentle slopes (plain).
Rockiness	No rocky ridges are present at the site.
Presence of wetlands	No wetlands are present at the site. A very small wetland patch is present in the extreme south eastern corner of the site.
Vegetation	Site is characterised by grassland with some dwarf shrubs. Trees occur sparsely or in small bushclumps. Conspicuous key grass species in the area are <i>Themeda triandra</i> , <i>Setaria sphacelata</i> var. <i>torta</i> , <i>Panicum coloratum</i> , <i>Eragrostis chloromelas</i> , <i>Aristida congesta</i> and <i>Elionurus muticus</i> . Herbaceous species include <i>Berkheya onopordifolia</i> , <i>Stachys spathula</i> , <i>Vernonia oligocephala</i> and geophytes such as <i>Bulbine narcissifolia</i> and <i>Ledebouria marginata</i> . Indigenous tree species that are sparsely distributed at the site include <i>Acacia karroo</i> (sweet thorn), <i>Searsia pyroides</i> , <i>Grewia flava</i> , <i>Searsia lancea</i> and <i>Ziziphus mucronata</i> . Clumps of <i>Searsia ciliata</i> are present at the site.
Signs of disturbances	Vegetation at the site is in fair condition.
Connectivity of natural vegetation in the site and between the site and surrounding areas	The site allocated for the proposed development is not part of a corridor of particular conservation importance.



Photo 1 View of site consisting of a grassland plain with sparse cover of trees. Trees in the distance are exotic *Eucalyptus* trees outside the site.
Photo: R.F. Terblanche.



Photo 2 *Acacia karroo*, sweet-thorns, at the site.
Photo: R.F. Terblanche



Photo 3 Exotic *Opuntia ficus-indica*, prickly pear, at the site. In general the cover of indigenous plant species is good at the site, with a few exceptions such as this alien prickly pear species.
Photo: R.F. Terblanche.



Photo 4 *Utetheisa pulchella*, crimson speckled footman, a widespread day-flying moth species (Lepidoptera), at the site.
Photo: R.F. Terblanche.

4.2 ASSESSMENT OF PLANT SPECIES OF PARTICULAR CONSERVATION PRIORITY

4.2.1 Plant species of particular conservation concern according to the red list of plants

Table 4.2 Threatened plant species of the North West Province which are listed in the **Critically Endangered** category. The list here follows the most recent updated red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Brachystelma canum</i>	Critically Endangered	No
<i>Brachystelma gracillimum</i>	Critically Endangered	No

Table 4.3 Threatened plant species of the North West Province which are listed in the **Endangered** category. The list here follows the most recent updated red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Aloe peglerae</i>	Endangered	No
<i>Brachystelma discoideum</i>	Endangered	No

Table 4.4 Threatened plant species of the North West Province which are listed in the **Vulnerable** category. The list here follows the most recent updated red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Brachycorythis conica</i> subsp. <i>transvaalensis</i>	Vulnerable	No
<i>Brachystelma incanum</i>	Vulnerable	No
<i>Ceropegia decidua</i> subsp. <i>pretoriensis</i>	Vulnerable	No
<i>Ceropegia stentiae</i>	Vulnerable	No
<i>Ledebouria atrobrunnea</i>	Vulnerable	No
<i>Marsilea farinosa</i>	Vulnerable	No
<i>Melolobium subspicatum</i>	Vulnerable	No
<i>Prunus africana</i>	Vulnerable	No
<i>Rennera stellata</i>	Vulnerable	No
<i>Searsia maricoan</i>	Vulnerable	No

Table 4.5 Near Threatened plant species of the North West Province. The list here follows the most recent updated red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Andromischus umbraticola</i> subsp. <i>umbraticola</i>	Near Threatened	No
<i>Ceropegia turricula</i>	Near Threatened	No
<i>Cineraria austrotransvaalensis</i>	Near Threatened	No
<i>Cleome conrathii</i>	Near Threatened	No
<i>Delosperma leendertziae</i>	Near Threatened	No
<i>Drimia sanguinea</i>	Near Threatened	No
<i>Elaeodendron transvaalense</i>	Near Threatened	No
<i>Kniphofia typhoides</i>	Near Threatened	No
<i>Lithops leslei</i> subsp. <i>leslei</i>	Near Threatened	No
<i>Nerine gracilis</i>	Near Threatened	No
<i>Sporobolus oxyphyllus</i>	Near Threatened	No

<i>Stenostelma umbelluliferum</i>	Near Threatened	No
-----------------------------------	-----------------	----

Table 4.6 Plant species of the North West Province which are not threatened and not near threatened but which are of particular conservation concern and listed in the **Critically Rare** category (Raimondo *et al.* 2009). The list here follows the most recent red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Conservation status	Resident at the site
<i>Gladiolus filiformis</i>	Critically Rare	No

Table 4.7 Plant species of the North West Province which are not threatened and not near threatened but of which are of particular conservation concern and listed in the **Rare** category (Raimondo *et al.* 2009). The list here follows the most recent red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Brachystelma dimorphum</i> subsp. <i>gratum</i>	Rare	No
<i>Ceropegia insignis</i>	Rare	No
<i>Frithia pulchra</i>	Rare	No
<i>Gnaphalium nelsonii</i>	Rare	No
<i>Habenaria culveri</i>	Rare	No

Table 4.8 Plant species of the North West Province which are not threatened and not near threatened but which are of particular conservation concern and listed in the **Declining** category (Raimondo *et al.* 2009). The list here follows the most recent red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Acacia erioloba</i>	Declining	No
<i>Boophone disticha</i>	Declining	No
<i>Crinum bulbispermum</i>	Declining	No
<i>Crinum macowanii</i>	Declining	No
<i>Drimia altissima</i>	Declining	No
<i>Eucomis autumnalis</i>	Declining	No
<i>Gunnera perpensa</i>	Declining	No
<i>Hypoxis hemerocallidea</i>	Declining	No

<i>Ilex mitis</i>	Declining	No
<i>Pelargonium sidoides</i>	Declining	No

4.2.2 Plant species of particular conservation concern: protected species

Table 4.9 Tree species of the North West Province which are listed as **Protected Species** under the National Forests Act No. 84 of 1998, Section 51(1). No = Plant species is not a resident on the site; Yes = Plant species is a resident at the site.

Species	Conservation status	Resident at the site
<i>Acacia erioloba</i> (Camel Thorn Tree)	Protected	No
<i>Boscia albitrunca</i> (Sheppard's tree)	Protected	No
<i>Sclerocarya birrea</i> (Marula)	Protected	No

4.3 ASSESSMENT OF VERTEBRATE SPECIES OF PARTICULAR HIGH CONSERVATION PRIORITY

4.3.1 Mammals of particular high conservation priority

Table 4.10 Threatened mammal species of the North West Province. Literature sources: Friedman & Daly, (2004), Skinner & Chimimba (2005), Wilson & Reeder (2005). With mammal species which normally needs a large range their residential status does not implicate that they are exclusively dependent on the site or use the site as important shelter or for reproduction. No = Not recorded at site/ Unlikely to be resident at the site. Yes: Recorded at the site/ Likely to be resident at the site.

Species	Threatened Status	Site is part of range	Recorded at site during survey	Likely to be found based on habitat assessment
<i>Chrysospalax villosus</i> Rough-haired golden mole	Vulnerable	No	No	No
<i>Cloeotis percivali</i> Short-eared Trident Bat	Vulnerable/ Near-threatened	No	No	No
<i>Diceros bicornis</i> Black rhinoceros	Critically Endangered	No	No	No

<i>Lycaon pictus</i> African wild dog	Endangered	No	No	No
<i>Loxodonta africana</i> African elephant	Vulnerable	No	No	No
<i>Mystromys albicaudatus</i> White-tailed mouse	Endangered	Yes	No	No
<i>Neamblysomus julianae</i> Juliana's Golden Mole	Critically Endangered	No	No	No
<i>Panthera leo</i> Lion	Vulnerable	No	No	No
<i>Rhinolophus blasii</i> Blasi's Horseshoe Bat	Vulnerable	No	No	No

Table 4.11 Near threatened mammal species known to occur in the North West Province. Literature sources: Skinner & Chimimba (2005). No = Not recorded at site/ unlikely to be resident at the site. Yes: Recorded at the site/ Likely to be resident at the site.

Species	Threatened Status	Site is part of range	Recorded at site during survey	Likely to be found based on habitat assessment
<i>Ceratotherium simum</i> White Rhinoceros	Near threatened	No	No	No
<i>Manis temminckii</i> Ground Pangolin	Near threatened	No	No	No

Table 4.12 Data deficient (or uncertain) mammal species of the North West Province. Literature sources: Skinner & Chimimba (2005). No = Not recorded at site/ unlikely to be resident at the site. Yes: Recorded at the site/ Likely to be resident at the site.

Species	Threatened Status	Recorded at site during survey	Likely be a resident at the site
<i>Myosorex varius</i> Forest shrew	Uncertain	No	No

4.3.2 Birds of particular high conservation priority

Table 4.13 Threatened bird species of the North West Province. Literature sources Barnes (2000), Hockey, Dean & Ryan, P.G. (2005) and Chittenden (2007). No = Not recorded at site/ Unlikely to use site as breeding area or particular habitat on which the species depends. Yes = Recorded at site/ Likely to use site as breeding area or particular habitat on which the species depends.

Species	Common name	Threatened Status	Recorded at site during survey	Likely to use site as breeding area or habitat
<i>Aegypius tracheliotos</i>	Lappet-faced Vulture	Vulnerable	No	No
<i>Anthropoides paradiseus</i>	Blue Crane	Vulnerable	No	No
<i>Aquila rapax</i>	Tawny Eagle	Vulnerable	No	No
<i>Ardeotis kori</i>	Kori Bustard	Vulnerable	No	No
<i>Balearica regulorum</i>	Grey Crowned Crane (Mahem)	Vulnerable	No	No
<i>Botaurus stellaris</i>	Eurasian Bittern	Critically Endangered	No	No
<i>Circus ranivorus</i>	African Marsh-Harrier	Vulnerable	No	No
<i>Crex crex</i>	Corn Crane	Vulnerable	No	No
<i>Eupodotis senegalensis</i>	White-bellied Korhaan	Vulnerable	No	No
<i>Falco naumanni</i>	Lesser Kestrel	Vulnerable	No	No
<i>Geronticus calvus</i>	Southern Bald Ibis	Vulnerable	No	No
<i>Gorsachius leuconotus</i>	White-backed Night-heron	Vulnerable	No	No
<i>Gypaetus barbatus</i>	Bearded Vulture	Endangered	No	No
<i>Gyps africanus</i>	White-backed Vulture	Vulnerable	No	No
<i>Gyps coprotheres</i>	Cape Vulture	Vulnerable	No	No
<i>Pelecanus rufescens</i>	Pink-backed Pelican	Vulnerable	No	No
<i>Polemaetus bellicosus</i>	Martial Eagle	Vulnerable	No	No
<i>Rhynchops flavirostris</i>	African Skimmer	Endangered	No	No
<i>Sarothrura ayresi</i>	White-winged Flufftail	Critically Endangered	No	No
<i>Tyto capensis</i>	African Grass-Owl	Vulnerable	No	No

NB: Though some of these bird species could be visitors to the site, especially some of which will nest at cliffs or artificial cliffs nearby, none of these are likely to be resident at the footprint proposed for the development.

Table 4.14 Near threatened bird species of the North West Province. Literature sources Barnes (2000), Hockey, Dean & Ryan, P.G. (2005) and Chittenden (2007). No = Not recorded at site/ Unlikely to be particularly dependent on the site as breeding area or habitat. Yes = Recorded at site/ Likely to be particularly dependant on the site as breeding area or habitat.

Species	Common name	Threatened Status	Recorded at site during survey	Likely to use site breeding area or habitat
<i>Certhilauda chuana</i>	Short-clawed Lark	Near threatened	No	No
<i>Charadrius pallidus</i>	Chestnut-banded Plover	Near threatened	No	No
<i>Ciconia nigra</i>	Black Stork	Near threatened	No	No
<i>Circus macrourus</i>	Pallid Harrier	Near threatened	No	No
<i>Eupodotis caerulescens</i>	Blue Korhaan	Near threatened	No	No
<i>Falco biarmicus</i>	Lanner Falcon	Near threatened	No	No
<i>Falco peregrinus</i>	Peregrine Falcon	Near threatened	No	No
<i>Glareola nordmanni</i>	Black-winged Pratincole	Near threatened	No	No
<i>Leptoptilos crumeniferus</i>	Marabou Stork	Near threatened	No	No
<i>Mirafra cheniana</i>	Melodious lark	Near threatened	No	No
<i>Mycteria ibis</i>	Yellow-billed Stork	Near threatened	No	No
<i>Phoenicopterus minor</i>	Lesser Flamingo	Near threatened	No	No
<i>Phoenicopterus ruber</i>	Greater Flamingo	Near threatened	No	No
<i>Rostratula benghalensis</i>	Greater Painted-snipe	Near threatened	No	No
<i>Sagittarius serpentarius</i>	Secretarybird	Near threatened*	No	No
<i>Sterna caspia</i>	Caspian Tern	Near threatened	No	No

* Most recent extinction risk assessment for the secretary bird, which may be an occasional visitor at the slopes at the site, is vulnerable.

** NB: Though some of these bird species could be visitors to the site, especially some of which will nest at cliffs or artificial cliffs nearby, none of these are likely to be resident at the footprint proposed for the development.

4.3.3 Reptiles of particular high conservation priority

The following tables list possible presence or absence of threatened reptile or near threatened reptile species in the study area. The Southern African Reptile Conservation Assessment (SARCA) was launched in May 2005 (Branch, Tolley, Cunningham, Bauer, Alexander, Harrison, Turner & Bates, 2006). Its primary aim is to produce a conservation assessment for reptiles of South Africa, Lesotho and Swaziland in the near future (Branch *et al.*, 2006). A full up-dated conservation assessment of reptiles, taking into account the recent IUCN (2001) criteria, can only be used once this work becomes available. Alexander & Marais (2007) and Tolley & Burger (2007) give useful indications of present conservation statuses as well as possible red listings of reptile species and subspecies in the near future.

Table 4.15 Threatened reptile species in North West Province. Sources: Alexander & Marais (2007). No = Reptile species is not a resident on the site; Yes = Reptile species is found to be resident on the site.

Species	Threatened Status	Resident at site	Recorded at site during survey	Likely to be found based on habitat assessment
<i>Cordylus giganteus</i> Sungazer	Vulnerable	No	No	No
<i>Tetradactylus breyeri</i> Breyer's Long-tailed Seps	Vulnerable	No	No	No

Table 4.16 Near threatened reptile species in North West Province. Sources: Alexander & Marais (2007). No = Reptile species is not a resident on the site; Yes = Reptile species is found to be resident on the site.

Species	Threatened Status	Resident at site	Recorded at site during survey	Likely to be found based on habitat assessment
<i>Lamprophis fuscus</i> Yellow-bellied House Snake	Near threatened	No	No	No
<i>Homoroselaps dorsalis</i> Striped Harlequin Snake	Near threatened	No	No	No

4.3.4 Amphibian species of particular high conservation priority

Table 4.17 Near threatened amphibian species in North West Province. No = Amphibian species is not a resident on the site; Yes = Amphibian species is found to be resident on the site.

Species	Threatened Status	Resident at site	Recorded at site during survey	Likely to be found based on habitat assessment
<i>Pyxicephalus adspersus</i> Giant Bullfrog	Near threatened	No	No	No

4.4 ASSESSMENT OF INVERTEBRATE SPECIES OF PARTICULAR HIGH CONSERVATION PRIORITY

4.4.1 Butterflies of particular conservation priority

Table 4.18 Threatened butterfly species in North West Province and Gauteng Province. Sources: Henning, Terblanche & Ball (2009), Mecenero *et al.* (2013). Invertebrates such as threatened butterfly species are often very habitat specific and residential status imply a unique ecosystem that is at stake.

Species	Threatened Status	Recorded at site during survey	Residential status at the site: Yes confirmed, Highly likely, Likely, Medium possibility, Unlikely, Highly unlikely
<i>Aloeides dentatis dentatis</i> Roodepoort Copper	Endangered	No	Highly unlikely
<i>Chrysochrysis aureus</i> Golden Copper	Endangered	No	Highly unlikely
<i>Lepidochrysops praeterita</i> Highveld Blue	Endangered	No	Highly unlikely
<i>Orachrysops mijburghi</i> Mijburgh's Blue	Endangered	No	Highly unlikely

Table 4.19 Butterfly species of the North West Province and Gauteng Province that are not threatened and not near threatened but of which are of particular conservation concern and listed in the **Rare** category (Mecenero *et al.*, 2013). No = Butterfly species is unlikely to be a resident at the study area; Yes = Butterfly species is a resident at the study area.

Species	Threatened Status	Recorded at site during survey	Residential status at the site: Yes confirmed, Highly likely, Likely, Medium possibility, Unlikely, Highly unlikely
<i>Colotis celimene amina</i> Lilac Tip	Rare (Low density)	No	Highly unlikely
<i>Lepidochrysops procera</i> Savanna Blue	Rare (Habitat specialist)	No	Highly unlikely
<i>Metisella meninx</i> Marsh Sylph	Rare (Habitat specialist)	No	Highly unlikely
<i>Platylesches dolomitica</i> Hilltop Hopper	Rare (low density)	No	Highly unlikely

4.4.2 Beetles of particular conservation priority

Table 4.20 Fruit chafer species (Coleoptera: Scarabaeidae: Cetoninae) in the Gauteng Province and North-West Province which are of known high conservation priority.

Species	Threatened Status	Recorded at site during survey	Likely to be resident based on habitat assessment
<i>Ichneustoma stobbiai</i>	Uncertain	No	No
<i>Trichocephala brincki</i>	Uncertain	No	No

4.4.3 Scorpion species of particular conservation priority

Table 4.21 Rock scorpion species (Scorpiones: Ischnuridae) species that are of known high conservation priority in the Gauteng Province and North-West Province.

Species	Threatened Status	Recorded at site during survey	Likely to be resident at site based on habitat assessment
<i>Hadogenes gracilis</i>	Uncertain	No	No
<i>Hadogenes gunningi</i>	Uncertain	No	No

5 DISCUSSION

5.1 HABITAT AND VEGETATION CHARACTERISTICS

An outline of the habitat and vegetation characteristics is given in Table 4.1.

5.2 PLANT SPECIES

Extinct, threatened, near threatened and other plant species of high conservation priority in North West Province are listed in Tables 4.2 – 4.8. Protected tree species are listed in Table 4.9. The presence or not of all the species listed in the tables were investigated during the survey. None of the threatened and near-threatened plant species are likely to occur on the site. None of the declining, data deficient or other plant species of particular conservation priority occurs on the site proposed for development.

5.3 VERTEBRATES

5.3.1 Mammals

Table 4.10, Table 4.11 and Table 4.12 list the possible presence or absence of threatened mammal species, near threatened mammal species and mammal species of which the status is uncertain, respectively, at the site. Literature sources that were used are Friedman & Daly (2004), Skinner & Chimimba (2005) and Wilson & Reeder (2005). Since the site falls outside reserves, threatened species such as the black rhinoceros (*Diceros bicornis*) and the African wild dog (*Lycaon pictus*) are obviously not present. No smaller mammals of particular high conservation significance are likely to be found on the site as well.

5.3.2 Birds

Table 4.13 and Table 4.14 list the possible presence or absence of threatened bird species and near threatened bird species at the site. With bird species which often have a large distributional

range, their presence does not imply that they are particularly dependent on a site as breeding location. Therefore the emphasis in the right hand columns of Table 4.12 and Table 4.13 are on the particular likely dependence or not of bird species on the site. Literature sources that were mainly consulted are Barnes (2000), Hockey, Dean & Ryan, P.G. (2005) and Chittenden (2007). No threat to any threatened bird species or any bird species of particular conservation importance are foreseen.

5.3.3 Reptiles

Table 4.15 and Table 4.16 list the possible presence or absence of threatened and near threatened reptile species on the site. The Southern African Reptile Conservation Assessment (SARCA) was launched in May 2005 (Branch, Tolley, Cunningham, Bauer, Alexander, Harrison, Turner & Bates, 2006). Its primary aim is to produce a conservation assessment for reptiles of South Africa, Lesotho and Swaziland within a four year period, ending 2009 (Branch *et al.*, 2006). Therefore a full up-dated conservation assessment of reptiles, taking into account the recent IUCN (2001) criteria, will only be available in the near future. While the conservation statuses of reptile species are under revision Alexander & Marais (2007) as well as Tolley & Burger 2007 give useful indications of possible red listings in the near future. There appears to be no threat to any reptile species of particular high conservation importance if the site is developed.

5.3.4 Amphibians

No frog species that occur in the North West are threatened as threatened species (vulnerable, endangered or critically endangered) according to Minter, Burger, Harrison, Braack, Bishop and Kloepfer (2004) as well as Du Preez & Carruthers (2009). Table 4.17 lists *Pyxicephalus adspersus* (Giant Bullfrog) as near threatened (Minter *et al.*, 2004; Du Preez & Carruthers, 2009). There is no suitable habitat for *Pyxicephalus adspersus* (Giant Bullfrog) at the site. There appears to be no threat to any amphibian species of particular high conservation importance if the site is developed.

5.4 INVERTEBRATES

5.4.1 Butterflies

Studies about the vegetation and habitat of threatened butterfly species in South Africa showed that ecosystems with a unique combination of features are selected by these often localised threatened butterfly species (Deutschländer and Bredenkamp 1999; Edge 2002, 2005; Terblanche, Morgenthal & Cilliers 2003; Lubke, Hoare, Victor & Ketelaar 2003; Edge, Cilliers & Terblanche, 2008). Threatened butterfly species in South Africa can then be regarded as bio-indicators of rare ecosystems.

Four species of butterfly in Gauteng Province and North West Province combined are listed as threatened in the recent butterfly conservation assessment of South Africa (Mecenero *et al.*, 2013). The expected presence or not of these threatened butterfly species as well as species of high conservation priority that are not threatened, at the site (Table 4.18 and Table 4.19) follows.

5.4.1.1 Assessment of threatened butterfly species

***Aloeides dentatis dentatis* (Roodepoort Copper)**

The proposed global red list status for *Aloeides dentatis dentatis* according to the most recent IUCN criteria and categories is Endangered (Mecenero *et al.*, 2013). *Aloeides dentatis dentatis* colonies are found where one of its host plants *Hermannia depressa* or *Lotononis eriantha* is present. Larval ant association is with *Lepisiota capensis* (S.F. Henning 1983; S.F. Henning & G.A. Henning 1989). The habitat requirements of *Aloeides dentatis dentatis* are complex and not fully understood yet. See Deutschländer and Bredenkamp (1999) for the description of the vegetation and habitat characteristics of one locality of *Aloeides dentatis* subsp. *dentatis* at Ruimsig, Roodepoort, Gauteng Province. There is not an ideal habitat of *Aloeides dentatis* subsp. *dentatis* on the site and it is unlikely that the butterfly is present at the site.

***Chrysoritis aureus* (Golden Opal/ Heidelberg Copper)**

The proposed global red list status for *Chrysoritis aureus* according to the most recent IUCN criteria and categories is Endangered (Mecenero *et al.*, 2013) *Chrysoritis aureus* (Golden Opal/ Heidelberg Copper) is a resident where the larval host plant, *Clutia pulchella* is present. However,

the distribution of the butterfly is much more restricted than that of the larval host plant (S.F. Henning 1983; Terblanche, Morgenthal & Cilliers 2003). One of the reasons for the localised distribution of *Chrysochrysis aureus* is that a specific host ant *Crematogaster liengmei* must also be present at the habitat. Fire appears to be an essential factor for the maintenance of suitable habitat (Terblanche, Morgenthal & Cilliers 2003). Research revealed that *Chrysochrysis aureus* (Golden Opal/ Heidelberg Copper) has very specific habitat requirements, which include rocky ridges with a steep slope and a southern aspect (Terblanche, Morgenthal & Cilliers 2003). Owing to a lack of habitat requirements and ideal habitat the presence of the taxon is highly unlikely.

***Lepidochrysops praeterita* (Highveld Blue)**

The proposed global red list status for *Lepidochrysops praeterita* according to the most recent IUCN criteria and categories is Endangered (G.A. Henning, Terblanche & Ball, 2009; Mecenero *et al.*, 2013). *Lepidochrysops praeterita* is a butterfly that occurs where the larval host plant *Ocimum obovatum* (= *Becium obovatum*) is present (Pringle, G.A. Henning & Ball, 1994), but the distribution of the butterfly is much more restricted than the distribution of the host plant. *Lepidochrysops praeterita* is found on selected rocky ridges and rocky hillsides in parts of Gauteng, the extreme northern Free State and the south-eastern Gauteng Province. No ideal habitat appears to be present for the butterfly on the site. It is unlikely that *Lepidochrysops praeterita* would be present on the site and at the footprint proposed for the development.

***Orachrysops mijburghii* (Mijburgh's Blue)**

The proposed global red status for *Orachrysops mijburghii* according to the most recent IUCN criteria and categories is Endangered (Mecenero *et al.*, 2013). *Orachrysops mijburghii* favours grassland depressions where specific *Indigofera* plant species occur (Terblanche & Edge 2007). The Heilbron population of *Orachrysops mijburghii* in the Free State uses *Indigofera evansiana* as a larval host plant (Edge, 2005) while the Suikerbosrand population in Gauteng uses *Indigofera dimidiata* as a larval host plant (Terblanche & Edge 2007). There is no suitable habitat for *Orachrysops mijburghii* on the site and it is unlikely that *Orachrysops mijburghii* would be present on the site.

Conclusion on threatened butterfly species

There appears to be no threat to any threatened butterfly species if the site is developed.

5.4.1.2 Assessment of butterfly species that are not threatened but also of high conservation priority

***Colotis celimene amina* (Lilac tip)**

Colotis celimene amina is listed as Rare (Low density) by Mecenero *et al.* (2013). In South Africa *Colotis celimene amina* is present from Pietermaritzburg in the south and northwards into parts of Kwa-Zulu Natal, Gauteng, Limpopo, Mpumalanga and the North West Provinces (Mecenero *et al.* In press.). Reasons for its rarity are poorly understood. It is highly unlikely that *Colotis celimene amina* would be present at the site.

***Lepidochrysops procera* (Savanna Blue)**

Lepidochrysops procera is listed as Rare (Habitat specialist) by Mecenero *et al.* (2013). *Lepidochrysops procera* is endemic to South Africa and found in Gauteng, KwaZulu-Natal, Mpumalanga and North West (Mecenero *et al.*, 2013). Owing to a lack of habitat requirements and ideal habitat the presence of the taxon at the site is highly unlikely.

***Metisella meninx* (Marsh Sylph)**

Henning and Henning (1989) in the first South African Red Data Book of Butterflies, listed *Metisella meninx* as threatened under the former IUCN category Indeterminate. Even earlier in the 20th century Swanepoel (1953) raised concern about vanishing wetlands leading to habitat loss and loss of populations of *Metisella meninx*. According to the second South African Red Data Book of butterflies (Henning, Terblanche & Ball, 2009) the proposed global red list status of *Metisella meninx* has been Vulnerable. During a recent large scale atlasing project the *Conservation Assessment of Butterflies of South Africa, Lesotho and Swaziland: Red List and Atlas* (Mecenero *et al.*, 2013) it was found that more *Metisella meninx* populations are present than thought before. Based on this valid new information, the conservation status of *Metisella meninx* is now regarded as Rare (Habitat specialist) (Mecenero *et al.*, 2013). Though *Metisella meninx* is more widespread and less threatened than perceived before, it should be regarded as a localised rare habitat specialist of conservation priority, which is dependent on wetlands with suitable patches of grass at wetlands (Terblanche In prep.). Another important factor to keep in mind for the conservation of *Metisella meninx* is that based on very recent discoveries of new taxa in the group the present *Metisella meninx* is species complex consisting of at least three taxa (Terblanche In prep., Terblanche & Henning In prep.). The ideal habitat of *Metisella meninx* is

treeless marshy areas where *Leersia hexandra* (rice grass) is abundant (Terblanche In prep.). The larval host plant of *Metisella meninx* is wild rice grass, *Leersia hexandra* (G.A. Henning & Roos, 2001). Owing to a lack of habitat requirements and ideal habitat the presence of the taxon at the site is highly unlikely.

***Platylesches dolomitica* (Hilltop Hopper)**

Platylesches dolomitica is listed as Rare (Low density) by Mecenero *et al.* (2013). Historically the conservation status of *Platylesches dolomitica* was proposed to be Vulnerable (Henning, Terblanche & Ball 2009). However this butterfly which is easily overlooked and has a wider distribution than perceived before. *Platylesches dolomitica* has a patchy distribution and is found on rocky ledges where *Parinari capensis* occurs, between 1300 m and 1800m (Mecenero *et al.* 2013, Dobson Pers comm.). Owing to a lack of habitat requirements and ideal habitat the presence of the taxon at the site is highly unlikely.

5.4.2 Fruit chafer beetles

Table 4.20 lists the fruit chafer beetle species (Coleoptera: Scarabaeidae: Cetoninae) that are of known high conservation priority in the North West Province. No *Ichneustoma stobbiai* or *Trichocephala brincki* were found during the surveys. There appears to be no suitable habitat for *Ichneustoma stobbiai* or *Trichocephala brincki* at the site. There appears to be no threat to any of the fruit chafer beetles of particular high conservation priority if the site were developed.

5.4.3 Scorpions

Table 4.21 lists the rock scorpion species (Scorpiones: Ischnuridae) that are of known high conservation priority in the North West Province. None of these rock scorpions have been found at the site and the habitat does not appear to be optimal.

6 IMPACT ASSESSMENT AND MITIGATION MEASURES

The primary cause of loss of biological diversity is habitat degradation and loss (IUCN, 2004; Primack, 2006). Habitats of threatened plants are in danger most often due to urban developments such as is the case for the Gauteng Province (Pfab & Victor, 2002). Habitat conservation is the key to the conservation of invertebrates such as threatened butterflies (Deutschländer and Bredenkamp 1999; Edge 2002, 2005; Terblanche, Morgenthal & Cilliers 2003; Lubke, Hoare, Victor & Ketelaar 2003; Edge, Cilliers & Terblanche, 2008). Though human impacts in few cases have improved the habitat for mammalian species such as greater cane rats, that prosper in sugar cane and maize fields (Apps 2000), for many mammalian habitat specialist species, human impacts has lead to habitat loss. Some mammal species, especially many of the larger species, could adapt to a wide range of habitat types, but then need a large range. Some animals and plants are rare and occupy only one or a few specialised habitats (Primack 2006). Habitat conservation, either as large available land or as specialised habitats is therefore key to the conservation of many threatened plant species and animal species or any other species of high conservation priority (i.e. rare, near threatened species). In addition corridors and linkages may play a significant role in conservation of fauna.

Corridors are important to link ecosystems of high conservation priority. Such corridors or linkages are there to improve the chances of survival of otherwise isolated populations (Samways, 2005). How wide should corridors be? The answer to this question depends on the conservation goal and the focal species (Samways, 2005). Corridors for mammalian species are especially important for migratory species (Mwalyosi, 1991, Pullin 2002). For an African butterfly assemblage this is about 250m when the corridor is for movement as well as being a habitat source (Pryke and Samways 2003). Hill (1995) found a figure of 200m for dung beetles in tropical Australian forest. In the agricultural context, and at least for some common insects, even small corridors can play a valuable role (Samways, 2005). Much more research remains to be done to find refined answers to the width of grassland corridors in South Africa. The width of corridors will also depend on the type of development, for instance the effects of the shade of multiple story buildings will be quite different from that of small houses. Corridors have a number of advantages related to dispersal and gene flow by avoiding isolation of ecological patches. However, corridors could also have potential drawbacks, for example creating gene flow where none has occurred

naturally in the past and also as reservoirs for pathogens or introduced species (Pullin, 2002). Perhault and Lomolino (2000) studied corridors and mammal community structure in an old-growth forest landscape in the United States of America and their data suggest that each corridor should be valued individually. A lot of research remains to be conducted to have a better idea of the value of corridors, but in general corridors would be of considerable value. It appears that a network of wetland corridors and rocky ridges is highly likely to be of considerable benefit in environmental management and planning. Though proper management plans for habitats are not in place, setting aside special ecosystems is in line with the recent Biodiversity Act (2004) of the Republic of South Africa.

To summarise: In practice, as far as any developments are concerned, the key would be to prioritise and plan according to sensitive species and special ecosystems.

In the case of this study the condition of the natural vegetation is natural and not much disturbed. If the development is approved, it is unlikely that there will be loss of any particular ecosystem or corridor of special conservation concern. There appears to be no loss of any particular sensitive species, if the site is developed.

The following potential impacts and mitigation measures with a view to the proposed developments apply:

6.1 Anticipated risks or impacts to the loss of habitat

The following impacts on the loss of habitat apply at the site.

Potential impacts on the available habitat will be of local extent, of permanent duration, of low intensity and high probability. The significance of loss of habitat is expected to be moderate without mitigation and moderate with mitigation.

Impact summary matrix:

Phase	Significance of Impact				
	None	Low	Moderate	High	With mitigation
Operational			X		Moderate

Mitigation measures:

- Exotic and invasive plant species should not be allowed to establish, if the development is approved.

6.2 Anticipated risks or impacts to the loss of sensitive species

Sensitive species are regarded here as those listed in section 5 and constitutes the flora and fauna that are threatened or of other particular high conservation importance. The presence or not of all the species listed in the tables were investigated during the survey. None of the threatened and near-threatened plant species are likely to occur on the site. None of the other plant species of particular conservation priority occur on the site proposed for development.

6.3 Anticipated risks or impacts to habitat connectivity and open space

Potential impacts on connectivity will be of local extent, of permanent duration, of medium intensity and high probability. The significance of the impacts on loss of connectivity is expected to be low without mitigation and low with mitigation.

Impact summary matrix: habitat connectivity

Phase	Significance of Impact				
	None	Low	Moderate	High	With mitigation
Construction		X			Low
Operational		X			Low

Mitigation measures:

- If the development is approved, establishment of exotic and invasive plant species should be avoided and where these have been found at the site continuous eradication should take place.

6.4 Anticipated risks or impacts associated with construction activities

Overall construction activities associated with the development if approved will be of local extent, of medium duration, of medium intensity and high probability. During the construction phase, the significance of the impacts associated with the construction phase is likely to be low without and low with mitigation.

Impact summary matrix:

Phase	Significance of Impact				
	None	Low	Moderate	High	With mitigation
Construction		X			Low
Operational		X			Low

Mitigation measures:

- If the development is approved, contractors must ensure that no animal species are disturbed, trapped, hunted or killed during the construction phase.

7 CONCLUSION

- Vegetation at the site is in fair condition for the vegetation type.
- Vaal-Vet Sandy Grassland of which the site is part is listed as Endangered and the entire vegetation type is an Endangered ecosystem (Mucina & Rutherford 2006). The development if approved should therefore be planned carefully and the footprint restricted to the small area allocated as such.
- Establishment of exotic weeds should be monitored, during construction, if the development is approved, and exotic weeds at the site should be eradicated. By no means should exotic declared invaders such as the mesquite tree (*Prosopis* species), be planted or allowed to establish.
- No wetlands appear to be present at the site, apart from a small wetland patch (0.05 ha) at the south eastern extreme of the site, probably caused by the elevation of the dirt road next to the fence on a flat plain. A buffer zone of 30 m is likely to be sufficient to conserve this small wetland patch. Owing to the small size of the wetland and likely negligible impact of the proposed development, a separate wetland assessment is not recommended.
- A moderate diversity of indigenous plant species and animal species appears to be present at the site proposed for development.
- No loss of particularly sensitive habitat of particular conservation importance is anticipated if the site is developed.
- No loss of corridors or connectivity of ecosystems is anticipated if the sites are developed.
- There appears to be no threat to any protected tree species at the site (National Forests Act No. 84 of 1998).
- It is unlikely that there will be a loss of any plant species of particular high conservation priority, i.e. threatened or near threatened species, if the site is developed.
- It is highly unlikely that there would be a threat to any threatened animal species or any other animal species of particular conservation concern at the site. There is no distinct reason why this relatively small footprint allocated for the development, in the vast countryside of the North West Province is of particular conservation concern for any threatened vertebrate species, including those that roam large areas and which may occasionally or coincidentally visit the site.

8 REFERENCES

- Alexander, G. & Marais, J. 2007. A guide to the reptiles of Southern Africa. Struik, Cape Town.
- Anderson, M.D. & Andersen, T.A. 2001. Too much, too quickly? Doubts about the sustainability of the camelthorn wood harvest. *African Wildlife* 55(3): 21-23.
- Apps, P. 2012. Smithers' mammals of Southern Africa 4th ed: A field guide, revised and updated by Peter Apps. Struik Nature, Cape Town.
- Armstrong, A.J. 1991. On the biology of the marsh owl, and some comparisons with the grass owl. *Honeyguide* 37:148-159.
- Barnes, K.N. ed. 2000. The Eskom Red Data Book of birds of South Africa, Lesotho and Swaziland. BirdLife South Africa, Johannesburg.
- Boon, R. 2010. *Pooley's trees of eastern South Africa: a complete guide 2nd ed.* Flora and Fauna Publications Trust, Durban.
- Branch, B. 1998. Field guide to snakes and other reptiles of southern Africa. 3rd ed. Struik, Cape Town.
- Branch, B. 2008. Tortoises, Terrapins & Turtles of Africa. Struik Nature, Cape Town.
- Branch, W.R. & Patterson, R.W. 1975. Notes on the ecology of the Giant Girdled Lizard, *Cordylus giganteus*. *Journal of Herpetology* 9(4): 364-366.
- Branch, W.R., Tolley, K.A., Cunningham, M., Bauer, A.M., Alexander, G., Harrison, J.A., Turner, A.A. & Bates, M.F. eds. 2006. A plan for phylogenetic studies of southern African reptiles: proceedings of a workshop held at Kirstenbosch, February 2006. Biodiversity Series 5. South African National Biodiversity Institute, Pretoria.
- Bronner, G. 2011. *Mammals*. In: Picker, M. & Griffiths, C. 2011. *Alien & Invasive animals: a South African perspective*. Struik Nature, Cape Town, p 22-35.

- Bromilow, C. 2010. Problem plants and alien weeds of South Africa. Briza Publications, Pretoria.
- Carruthers, V. & Du Preez, 2011. Frogs and frogging in southern Africa 2nd ed. Struik, Cape Town.
- Chittenden, H. 2007. Roberts Bird Guide. John Voelcker Book Fund, Cape Town.
- Cillié, B., Oberprieler, U. & Joubert, C. 2004. Animals of Pilanesberg: an identification guide. Game Parks Publishing, Pretoria.
- Cilliers, S.S., Müller, N. & Drewes, E. 2004. Overview on urban nature conservation: situation in the western-grassland biome of South Africa. *Urban forestry and urban greening* 3: 49-62.
- Coetzee, N. & Monadjem, A. 2008. *Mystromys albicaudatus*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>.
- Conradie, W., Du Preez, L.H., Smith, K. & Weldon, C. 2006. Field guide to the frogs and toads of the Vredefort Dome World Heritage Site. School of Environmental Sciences and Development, Potchefstroom.
- Court, D. 2010. *Succulent Flora of Southern Africa*. Struik Nature, Cape Town.
- Crouch, N.R., Klopper, R.R., Burrows, J.E. & Burrows, S.M. 2011. Ferns of Southern Africa: a comprehensive guide. Struik Nature, Cape Town.
- Del Hoyo, J., Elliot, J. & Sargatal, J. 1992. Handbook of the birds of the world, Vol. 1. Lynx Editions, Barcelona.
- Deutschländer, M.S. & Bredenkamp, C.J. 1999. Importance of vegetation analysis in the conservation management of the endangered butterfly *Aloeides dentatis* subsp. *dentatis* (Swierstra) (Lepidoptera: Lycaenidae). *Koedoe* 42(2): 1-12.

Dippenaar-Schoeman, A.S. 2002. Baboon and trapdoor spiders in southern Africa: an identification manual. Plant Protection Research Institute Handbook No. 13. Agricultural Research Council, Pretoria.

Dippenaar-Schoeman, A.S. & Jocqué, R. 1997. African spiders: an identification manual. Plant Protection Research Institute Handbook No. 9. Agricultural Research Council, Pretoria.

Drinkwater, T.W., Bate, R. & Du Toit, H.A. 1998. A field guide for identification of maize pests in South Africa. Agricultural Research Council: Grain-crops Institute, Potchefstroom.

Du Preez, L.H. 1996. Field guide and key to the frogs and toads of the Free State. Department of Zoology and Entomology, University of the Orange Free State, Bloemfontein.

Du Preez, L.H. & Carruthers, V. 2009. A complete guide to the frogs of southern Africa. Struik Nature, Cape Town. CD with calls included.

Edge, D.A. 2002. Some ecological factors influencing the breeding success of the Brenton Blue butterfly, *Orachrysops niobe* (Trimen) (Lepidoptera: Lycaenidae). *Koedoe*, 45(2): 19-34.

Edge, D.A. 2005. Ecological factors influencing the survival of the Brenton Blue butterfly, *Orachrysops niobe* (Trimen) (Lepidoptera: Lycaenidae). North-West University, Potchefstroom, South Africa (Thesis - D.Phil.).

Edge, D.A., Cilliers, S.S. & Terblanche, R.F. 2008. Vegetation associated with the occurrence of the Brenton blue butterfly. *South African Journal of Science* 104: 505 - 510.

Ferguson-Lees, J. & Christie, D.A. 2001. *Raptors of the world*. Christopher Helm, London.

Filmer, M.R. 1991. Southern African spiders: an identification guide. Struik, Cape Town.

Gardiner, A.J. & Terblanche, R.F. 2010. Taxonomy, biology, biogeography, evolution and conservation of the genus *Erikssonia* Trimmen (Lepidoptera: Lycaenidae). *African Entomology* 18(1): 171 – 191.

Germishuizen, G. 2003. Illustrated guide to the wildflowers of northern South Africa. Briza, Pretoria.

Germishuizen, G., Meyer, N.L. & Steenkamp (eds) 2006. A checklist of South African plants. Southern African Botanical Diversity Network Report No. 41. SABONET, Pretoria.

Goldblatt, P. 1986. The Moraeas of Southern Africa. Annals of Kirstenbosch Botanic Gardens, Volume 14. National Botanic Gardens, Cape Town.

Goldblatt, P. & Manning, J. 1998. Gladiolus in Southern Africa.

Henderson, L. 2001. *Alien weeds and alien invasive plants: a complete guide to the declared weeds and invaders in South Africa*. Plant Protection Research Institute Handbook No. 12. ARC: Plant Protection Research Institute, Pretoria.

Henderson, L. & Cilliers, C.J. 2002. *Invasive aquatic plants: a guide to the identification of the most important and potentially dangerous invasive aquatic and wetland plants in South Africa*. Plant Protection Research Handbook No. 16. Agricultural Research Council, Pretoria.

Henning, G.A. & Roos, P.S. 2001. Threatened butterflies of South African wetlands. *Metamorphosis* 12(1): 26-33.

Henning, G.A., Terblanche, R.F. & Ball, J.B. (eds) 2009. South African Red Data Book: butterflies. *SANBI Biodiversity Series No 13*. South African National Biodiversity Institute, Pretoria.

Henning, S.F. 1983. Biological groups within the Lycaenidae (Lepidoptera). *Journal of the Entomological Society of Southern Africa* 46(1): 65-85.

Henning, S.F. 1987. Outline of Lepidoptera conservation with special reference to ant associated Lycaenidae. *Proceedings of the first Lepidoptera conservation Symposium, Roodepoort: Lepidopterists' Society of southern Africa*: 5-7.

Henning, S.F. & Henning, G.A. 1989. South African Red Data Book: butterflies. *South African National Scientific Programmes Report No. 158*. CSIR, Pretoria.

Herman, P.P.J. 2002. Revision of the *Tarchoanthus camphoratus* complex (Asteraceae-Tarchoanthaeae) in southern Africa. *Bothalia* 32,1: 21-28.

Hill, C.J. 1995. Conservation corridors and rainforest insects. (*In* Watt, A.D., Stork, N.E. & Hunter, M.D. (eds.), *Forests and Insects*. Chapman & Hall, London. p. 381-393.)

Hockey, P. 2011. *Birds*. In: Picker, M. & Griffiths, C. 2011. *Alien & Invasive animals: a South African perspective*. Struik Nature, Cape Town, p 36-44.

Hockey, P.A.R., Dean, W.J.R. & Ryan, P.G. (eds.). 2005. *Roberts Birds of Southern Africa*. John Voelcker Bird Book Fund, Cape Town.

Holm, E. & Marais, E. 1992. *Fruit chafers of southern Africa*. Ekogilde, Hartebeespoort.

IUCN. 2001. *IUCN Red List Categories and Criteria: Version 3.1*. IUCN Species Survival Commission. IUCN, Gland, Switzerland and Cambridge, UK.

IUCN. 2012. *IUCN Red list of Threatened Species*. Version 2012.1)

Jacobsen, W.B.G. 1983. *The ferns and fern allies of Southern Africa*. Butterworths, Durban.

Kemper, N.P. 2001. *RVI: Riparian Vegetation Index, final report*, WRC Report No. 850/3/1. Institute for Water Research, Pretoria.

Kok, J.C. 1998. *Vrystaatse bome, struik en klimplante Kontak-uitgewers*, Pretoria.

Kudrna, O. 1995. Conservation of butterflies in central Europe. (*In* Pullin, A. S. ed. *Ecology and conservation of butterflies*. Chapman & Hall, London. p. 248-257.)

Larsen, T.B. 1995. Butterfly biodiversity and conservation in the Afrotropical region. (*In* Pullin, A.S. ed. *Ecology and conservation of butterflies*. London: Chapman & Hall. p. 290-303.)

Liebenberg, L. 1990. A field guide to the animal tracks of Southern Africa. David Philip Publishers, Cape Town.

Leeming, J. 2003. Scorpions of southern Africa. Struik, Cape Town.

Leroy, A. & Leroy, J. 2003. Spiders of southern Africa. Struik, Cape Town.

Louw, W.J. 1951. *An ecological account of the vegetation of the Potchefstroom Area*. Botanical Survey of South Africa, Memoir No. 24. Government Printer, Pretoria.

Low, A.B. & Rebelo, A.G. (Eds.) 1996. Vegetation of South Africa, Lesotho and Swaziland. Department of Environmental Affairs and Tourism, Pretoria.

Lubke, R.A., Hoare, D., Victor, J. & Ketelaar, R. 2003. The vegetation of the habitat of the Brenton Blue Butterfly, *Orachrysops niobe* (Trimen), in the Western Cape, South Africa. *South African Journal of Science* 99: 201-206.

Manning, J. 2003. Photographic guide to the wild flowers of South Africa. Briza, Pretoria.

Manning, J. 2009. Field guide to the wild flowers of South Africa. Struik, Cape Town.

McMurtry, D., Grobler, L., Grobler, J. & Burns, S. 2008. Field guide to the orchids of northern South Africa and Swaziland. Umdaus Press, Hatfield.

Mecenero, S., Ball, J.B., Edge, D.A., Hamer, M.L., Henning, G.A., Krüger, M., Pringle, E.L., Terblanche, R.F. & Williams, M.C. 2013. *Conservation Assessment of Butterflies of South Africa, Lesotho and Swaziland: Red List and Atlas*. Safronics, Johannesburg & Animal Demography Unit, Cape Town.

Minter, L.R., Burger, M., Harrison, J.A., Braack, H.H., Bishop, P.J. & Kloepfer, D. eds. 2004. Atlas and Red Data Book of the Frogs of South Africa, Lesotho and Swaziland. SI/MAB series 9, Smithsonian Institution, Washington DC.

Mucina, L. & Rutherford, M.C. eds. 2006. The vegetation of South Africa, Lesotho and Swaziland.

Strelitzia 19. Pretoria: South African National Biodiversity Institute.

Mucina, L., Rutherford, M.C., and Powrie, L.W. eds. 2005. Vegetation map of South Africa, Lesotho and Swaziland, 1:1 000 000 scale sheet maps. Pretoria: South African National Biodiversity Institute.

Munguira, M.L. 1995. Conservation of butterfly habitats and diversity in European Mediterranean countries. (*In Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 277- 289.*)

New, T.R. 1993. ed. Conservation biology of *Lycaenidae* (butterflies). *Occasional paper of the IUCN Species Survival Commission No. 8.* .

New, T.R. 1995. Butterfly conservation in Australasia – an emerging awareness and an increasing need. (*In Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 304 – 315.*)

Oates, M.R. 1995. Butterfly conservation within the management of grassland habitats. (*In Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. (p. 98-112.)*)

Opler, P.A. 1995. Conservation and management of butterfly diversity in North America. (*In Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 316-324.*)

Peacock, F. 2006. Pipits of Southern Africa. Published by the author, Pretoria. www.pipits.co.za.

Pfab, M.F. 2002. Priority ranking scheme for Red Data plants in Gauteng, South Africa. *South African Journal of Botany* (68): 299-303.

Pfab, M.F. & Victor, J.E. 2002. Threatened plants of Gauteng, South Africa. *South African Journal of Botany* (68): 370-375.

Picker, M. & Griffiths, C. 2011. Alien & Invasive animals: a South African perspective. Struik Nature, Cape Town.

Picker, M., Griffiths, C. & Weaving, A. 2004. Field guide to insects of South Africa. 2nd ed. Cape Town: Struik.

Pooley, E. 1998. A field guide to wild flowers of KwaZulu-Natal and the eastern region. Natal Flora Publications Trust, Durban.

Pringle, E.L., Henning, G.A. & Ball, J.B. eds. 1994. Pennington's Butterflies of Southern Africa. Struik Winchester, Cape Town.

Pryke, S.R. & Samways, M.J. 2001. Width of grassland linkages for the conservation of butterflies in South African afforested areas. *Biological Conservation* 101: 85-96.

Pullin, A.S. ed. 1995. Ecology and conservation of butterflies. Chapman & Hall, London.

Rautenbach, I.L. 1982. The mammals of the Transvaal. Ecoplan monograph 1: 1-211.

Retief, E. & Herman, P.P.J. 1997. Plants of the northern provinces of South Africa: keys and diagnostic characteristics. Strelitzia 6. National Botanical Institute, Pretoria.

Rutherford, M.C. & Westfall, R.H. 1994. Biomes of southern Africa: An objective categorisation, 2nd ed. Memoirs of the Botanical Survey of South Africa, Vol. 63, pp. 1-94. National Botanical Institute, Pretoria.

Ryan, P. 2001. Practical Birding: A guide to birdwatching in southern Africa. Struik, Cape Town.

Samways, M.J. 2005. Insect diversity conservation. Cambridge University Press, Cambridge.

Skelton, P. 2001. A complete guide to the freshwater fishes of Southern Africa. Struik, Cape Town.

Skelton, P. & Weyl, O. 2011. *Fishes*. In: Picker, M. & Griffiths, C. 2011. *Alien & Invasive animals: a South African perspective*. Struik Nature, Cape Town, p 36-44.

Skinner, J.D. & Chimimba, C.T. 2005. The mammals of the southern African subregion. Cambridge University Press, Cape Town.

Sliwa, A. 2008. *Felis nigripes*. In: IUCN 2012. IUCN Red List of Threatened Species.

Smit, N. 2008. Field guide to the *Acacias* of South Africa. Briza, Pretoria.

Smithers, R.H.N. 1986. South African Red Data Book: Terrestrial mammals. *South African National Scientific Programmes Report* No. 125. CSIR, Pretoria.

South Africa. 2004. National Environmental Management: Biodiversity Act No. 10 of 2004. Government Printer, Pretoria.

Stuart, C. & Stuart, T. 2006. Field guide to the larger mammals of Africa 3rd ed. Struik Nature, Cape Town.

Stuart, C. & Stuart, T. 2013. A field guide to the tracks and signs of Southern, Central and East African wildlife 4th ed. Struik Nature, Cape Town.

Tarboton, W. & Erasmus, R. 1998. Owls and owling in southern Africa. Struik, Cape Town.

Taylor, J.C., Janse Van Vuuren, M.S. & Pieterse, A.J.H. 2007. The application and testing of diatom-based indices in the Vaal and Wilge Rivers, South Africa. *Water SA* 33(1): 51-59.

Terblanche, R.F. & Edge, D.A. 2007. The first record of an *Orachrysops* in Gauteng. *Metamorphosis* 18(4): 131-141.

Terblanche, R.F., Morgenthal, T.L. & Cilliers, S.S. 2003. The vegetation of three localities of the threatened butterfly species *Chrysoritis aureus* (Lepidoptera: Lycaenidae). *Koedoe* 46(1): 73-90.

Terblanche, R.F. & Van Hamburg, H. 2003. The taxonomy, biogeography and conservation of the myrmecophilous *Chrysoritis* butterflies (Lepidoptera: Lycaenidae) in South Africa. *Koedoe* 46(2): 65-81.

Terblanche, R.F. & Van Hamburg, H. 2004. The application of life history information to the conservation management of *Chrysoritis* butterflies (Lepidoptera: Lycaenidae) in South Africa. *Koedoe* 47(1): 55-65.

Thomas, C.D. 1995. Ecology and conservation of butterfly metapopulations in the fragmented British landscape. (*In* Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 46-64.)

Van den Berg, J. & Drinkwater, T.W. 1998. Field guide to identification of sorghum pests in South Africa. Agricultural Research Council: Grain-crops Institute, Potchefstroom.

Van Ginkel, C.E., Glen, R.P., Gordon-Gray, K.D., Cilliers, C.J., Muasya, M. & van Deventer, P.P. 2011. Easy identification of some South African Wetland Plants. WRC Report No TT 479/10. Water Research Commission, Gezina.

Van Jaarsveld, E.J. 2006. The Southern African *Plectranthus* and the art of turning shade to glade.

Van Oudtshoorn, F. 1999. Guide to grasses of southern Africa. Briza, Pretoria.

Van Wyk, B. 2000. A photographic guide to wild flowers of South Africa. Struik, Cape Town.

Van Wyk, B. & Malan, S. 1998. Field Guide to the Wild Flowers of the Highveld. Struik, Cape Town.

Van Wyk, A.E. & Smith, G.F. 2001. Regions of floristic endemism in Southern Africa: a review with emphasis on succulents, Umdaus Press, Pretoria.

Van Wyk, B.E. & Smith, G.F. 2003. Guide to the aloes of South Africa. 2nd ed. Briza, Pretoria.

Van Wyk, B. & Van Wyk, P. 1997. Field guide to trees of southern Africa. Struik, Cape Town.

Walker, C. 1996. Signs of the Wild. 5th ed. Struik, Cape Town.

Warren, M.S. 1995. Managing local microclimates for the high brown fritillary, *Argynnis adipe*. (In Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall.)

Watt, A.D., Stork, N.E. & Hunter, M.D. (eds.), Forests and Insects. London: Chapman & Hall. (p. 381-393.)

Wilson, D.E. & Reeder, D.M. 2005. Mammal species of the world: a taxonomic and geographic reference. 3rd ed. Vol.1 and Vol. 2. John Hopkins University Press, Maryland.

ANNEXURE 1

List of plant species recorded at the study area.

Sources: Germishuizen (2003), Manning (2003), Manning (2009), Van Oudtshoorn (1999), Van Wyk (2000), Van Wyk & Malan (1998), Van Wyk & Van Wyk (1997), Crouch, Klopper, Burrows & Burrows (2011), Goldblatt (1986), Goldblatt & Manning (1998), Jacobsen (1983), McMurtry, Grobler, Grobler & Burns (2008), Smit (2008), Van Ginkel *et al.* (2011), Van Jaarsveld (2006), Van Wyk & Smith (2003).

Plant species are listed alphabetically under main taxonomic groups.

Species marked with an asterisk * are exotic.

TAXON	COMMON NAMES	FAMILY
ANGIOSPERMAE: MONOCOTYLEDONS		
<i>Aristida adscensionis</i>	Annual Three-awn	POACEAE
<i>Aristida congesta</i> subsp. <i>congesta</i>	Tassel Three-awn	POACEAE
<i>Aristida diffusa</i>	Iron Grass	POACEAE
<i>Asparagus laricinus</i>	Common Wild Asparagus	ASPARAGACEAE
<i>Bewsia biflora</i>	False Love Grass	POACEAE
<i>Bulbine narcissifolia</i>		ASPHODELACEAE
<i>Cenchrus ciliaris</i>	Foxtail Buffalo Grass	POACEAE
<i>Cymbopogon pospischillii</i>	Narrow-leaved Turpentine Grass	POACEAE
<i>Cynodon dactylon</i>	Couch Grass	POACEAE
<i>Digitaria eriantha</i>	Common Finger Grass	POACEAE
<i>Enneapogon cenchroides</i>	Nine-awned Grass	POACEAE
<i>Enneapogon scoparius</i>	Bottlebrush Grass	POACEAE
<i>Eragrostis echinochloidea</i>	Tick Grass	POACEAE
<i>Eragrostis lehmanniana</i>	Lehmann's Love Grass	POACEAE
<i>Eragrostis obtusa</i>	Dew Grass	POACEAE
<i>Eragrostis rigidior</i>	Curly Leaf Love Grass	POACEAE
<i>Eragrostis superba</i>	Saw-toothed Love Grass	POACEAE
<i>Fingerhuthia africana</i>	Thimble Grass	POACEAE
<i>Heteropogon contortus</i>	Spear Grass	POACEAE
<i>Hypoxis acuminata</i>		HYPOXIDACEAE
<i>Panicum coloratum</i>	Small Buffalo Grass	POACEAE
<i>Pogonarthria squarrosa</i>	Herringbone Grass	POACEAE
<i>Schmidtia pappophoroides</i>	Sand Quick	POACEAE
<i>Setaria sphacelata</i> var. <i>torta</i>		POACEAE

<i>Sporobolus fimbriatus</i>	Dropseed Grass	POACEAE
<i>Stipagrostis uniplumis</i>	Silky Bushman Grass	POACEAE
<i>Themeda triandra</i>	Red Grass	POACEAE
<i>Tragus racemosus</i>		POACEAE
ANGIOSPERMS: DICOTYLEDONS		
<i>Acacia karroo</i>	Sweet Thorn	(or Fabaceae)
* <i>Argemone ochroleuca</i>	White-flowered Mexican poppy	PAPAVERACEAE
<i>Barleria macrostegia</i>		ACANTHACEAE
<i>Berkheya onopordifolia</i>		ASTERACEAE
* <i>Chenopodium album</i>	White Goosefoot	CHENOPODIACEAE
<i>Chrysocoma ciliata</i>	Bitterbush	ASTERACEAE
<i>Elephantorrhiza elephantina</i>		MIMOSACEAE (or Fabaceae)
<i>Felicia muricata</i>		ASTERACEAE
<i>Gazania krebsiana</i> subsp. <i>krebsiana</i>		ASTERACEAE
<i>Geigeria ornativa</i>		ASTERACEAE
* <i>Gomphrena celosioides</i>	Bachelor's Button	AMARANTHACEAE
<i>Grewia flava</i>	Velvet Raisin	MALVACEAE (or Tiliaceae)
<i>Helichrysum cerastioides</i>		ASTERACEAE
<i>Hermbstaedtia odorata</i>	Wild Cockscomb	AMARANTHACEAE
<i>Hibiscus pusillus</i>		MALVACEAE
<i>Hilliardiella oligocephala</i>		ASTERACEAE
<i>Jamesbrittenia aurantiaca</i>	Cape Saffron	SCROPHULARIACEAE
<i>Lepidium africanum</i>	Pepperweed	BRASSICACEAE
* <i>Lepidium bonariense</i>	Pepperweed	BRASSICACEAE
<i>Lippia scaberrima</i>		VERBENACEAE
<i>Lycium hirsutum</i>		SOLANACEAE
* <i>Malva parviflora</i>	Small Mallow	MALVACEAE
<i>Nidorella anomala</i>		ASTERACEAE
* <i>Oxalis corniculata</i>	Creeping Sorrel	OXALIDACEAE
<i>Pentzia globosa</i>		ASTERACEAE
<i>Persicaria species</i>		POLYGONACEAE
<i>Pollichia campestris</i>	Waxberry	ILLECEBRACEAE
* <i>Polygonum aviculare</i>	Prostrate Knotweed	POLYGONACEAE
<i>Salvia disermas</i>	Large Blue Sage	LAMIACEAE
<i>Salvia stenophylla</i>		LAMIACEAE
* <i>Schkuhria pinnata</i>	Dwarf Marigold	ASTERACEAE
<i>Searsia ciliata</i>	Sour Karree	ANACARDIACEAE
<i>Searsia lancea</i>	Karree	ANACARDIACEAE

<i>Searsia pyroides</i>	Common Wild Currant	ANACARDIACEAE
<i>Selago densiflora</i>		SCROPHULARIACEAE
<i>Senna italica</i>	Wild Senna	CAESALPINIACEAE
<i>Tarchonanthus camphoratus</i>	Wild Camphor Bush	ASTERACEAE
<i>Thesium</i> sp.		SANTALACEAE
<i>Tribulus terrestris</i>	Devil's Thorn	ZYGOPHYLLACEAE
<i>Ziziphus mucronata</i>	Buffalo-thorn	RHAMNACEAE
<i>Ziziphus zeyheriana</i>	Dwarf Buffalo-thorn	RHAMNACEAE