

ECOLOGICAL FAUNA AND FLORA HABITAT SURVEY

**Remaining Extent of the Farm Retreat 671, situated
in the Dr Ruth S Mompoti District Municipality, IN
Registration Division, Province of the North-West,
measuring 297,4480 (two hundred and ninety seven
comma four four eight zero) hectares, Title Deed No.
2676/2009.**



Grewia flava, Velvet Raisin, a widespread woody plant species that is often frequent in the Eastern
Kalahari Bushveld Bioregion.
Photo: November 2015, R.F. Terblanche.

JANUARY 2016

COMPILED BY:

Reinier F Terblanche

(M.Sc Ecology, *Cum Laude*; Pr.Sci.Nat, Reg. No. 400244/05)

ANTHENE ECOLOGICAL CC

TABLE OF CONTENTS

1. INTRODUCTION	1
2. STUDY AREA	2
3. METHODS	4
4. RESULTS	9
5. DISCUSSION	24
6. IMPACT ASSESSMENT AND MITIGATION MEASURES	33
7. CONCLUSION	41
8. REFERENCES	44
9. ANNEXURE 1 LIST OF PLANT SPECIES	56

1 INTRODUCTION

An ecological habitat survey was required for a photo-voltaic development at Remaining Extent of the Farm Retreat 671, situated in the Dr Ruth S Mompati District Municipality, IN Registration Division, Province of the North-West, measuring 297,4480 (two hundred and ninety seven comma four four eight zero) hectares, Title Deed No. 2676/2009, 7 km southwest of Vryburg in the North West Province (elsewhere referred to as the site). The survey focused on the possibility that threatened fauna or flora known to occur in North West Province are likely to occur within the proposed development. Species of known high conservation priority that do not qualify for threatened status also received attention in the survey.

1.1 OBJECTIVES OF THE HABITAT STUDY

The objectives of the habitat study are to provide:

- A detailed fauna and flora habitat survey;
- A detailed habitat survey of possible threatened or localised plant species, vertebrates and invertebrates;
- Recording of possible host plants or foodplants of fauna such as butterflies.
- Evaluate the conservation importance and significance of the site with special emphasis on the current status of threatened species;
- Literature investigation of possible species that may occur on site;
- Identification of potential ecological impacts on fauna and flora that could occur as a result of the development; and
- Make recommendations to reduce or minimise impacts, should the development be approved.

1.2 SCOPE OF STUDY

- Surveys to investigate key elements of habitats on the site, relevant to the conservation of fauna and flora.
- Recording of any sightings and/or evidence of existing fauna and flora.
- The selective and careful collecting of voucher specimens of invertebrates where deemed necessary.
- An evaluation of the conservation importance and significance of the site with special emphasis on the current status of threatened species.
- Recording of possible host plants or foodplants of fauna such as butterflies.
- Literature investigation of possible species that might occur on site.
- Integration of the literature investigation and field observations to identify potential ecological impacts that could occur as a result of the development.
- Integration of literature investigation and field observations to make recommendations to reduce or minimise impacts, should the development be approved.

2 STUDY AREA

The study area is 7 km southwest of Vryburg in the North West Province. The site is situated at the Savanna Biome which is represented by the Ghaap Plateau Vaalbosveld vegetation type (Mucina & Rutherford 2006). A brief overview of SVk 7, the Ghaap Plateau Vaalbosveld vegetation type in which the site is located, follows:

Distribution: In South Africa the Ghaap Plateau Vaalbosveld is found in the Northern Cape and North-West Provinces: Flat plateau from around Campbell in the south, east of Danielskuil through Reivilo to around Vryburg in the north. Altitude at the Ghaap Plateau Vaalbosveld is 1100 – 1500 m.

Vegetation and landscape features: Flat plateau with well-developed shrub layer with *Tarchonanthus camphoratus* and *Acacia karroo*. Open tree layer has *Olea europaea* subsp. *africana*, *Acacia tortilis*, *Ziziphus mucronata* and *Searsia [Rhus] lancea*. *Olea europaea* subsp. *africana* is more important in the southern parts of the unit, while *Acacia tortilis*, *Acacia hebeclada* and *Acacia mellifera* are more important in the north and part of the west of the unit. Much of the south-central part of this unit has remarkably low cover of *Acacia* species for an arid savanna and is dominated by the nonthorny *Tarchonanthus camphoratus*, *Searsia lancea*, and *Olea europaea* subsp. *africana* (Mucina & Rutherford 2006).

Geology and soils: Surface limestone of Tertiary to Recent age, and dolomite and chert of the Campbell Group (Griqualand West Supergroup, Vaalian Erathem) support shallow soils (0.1-0.25 m) of Mispah and Hutton soil forms. Land types mainly Fc with some Ae and Ag (Mucina & Rutherford, 2006).

Climate: Climate is characterized by summer and autumn rainfall and very dry winters. Mean annual precipitation from about 300 mm in the southwest to about 500 mm in the northeast. Frost is frequent to very frequent in winter (Mucina & Rutherford 2006).

Important taxa of the Ghaap Plateau Vaalbosveld listed by Mucina & Rutherford (2006): Tall Tree: *Acacia erioloba*. Small Trees: *Acacia mellifera* subsp. *detinens*, *Searsia lancea*, *Acacia karroo*, *Acacia tortilis* subsp. *heteracantha*, *Boscia albitrunca*. Tall Shrubs: *Olea europaea* subsp. *africana*, *Rhigozum trichotomum*, *Tarchonanthus camphoratus*, *Ziziphus mucronata*, *Diospyros*

pallens, *Ehretia rigida* subsp. *rigida* (this species complex has been revised and the *Ehretia alba* is the species that occurs at the vegetation type, R.F. Terblanche pers. obs.), *Euclea crispa* subsp. *ovata*, *Grewia flava*, *Gymnosporia buxifolia*, *Lessertia frutescens*, *Searsia tridactyla*. Low Shrubs: *Acacia hebeclada* subsp. *hebeclada*, *Aptosimum procumbens*, *Chrysocoma ciliata*, *Helichrysum zeyheri*, *Hermannia comosa*, *Lantana rugosa*, *Leucas capensis*, *Melolobium microphyllum*, *Peliostomum leucorrhizum*, *Pentzia globosa*, *Pentzia viridis*, *Zygophyllum pubescens*. Succulent Shrubs: *Hertia pallens*, *Lycium cinereum*. Semi-parasitic Shrub: *Thesium hystrix*. Woody Climber: *Asparagus africanus*. Graminoids: *Antheophora pubescens*, *Cenchrus ciliaris*, *Digitaria eriantha* subsp. *eriantha*, *Enneapogon scoparius*, *Eragrostis lehmanniana*, *Schmidtia pappophoroides*, *Themeda triandra*, *Aristida adscensionis*, *Aristida congesta*, *Aristida diffusa*, *Cymbopogon pospischilii*, *Enneapogon cenchroides*, *Enneapogon desvauxii*, *Eragrostis echinocloidea*, *Eragrostis obtusa*, *Eragrostis rigidior*, *Eragrostis superba*, *Fingerhutia africana*, *Heteropogon contortus*, *Sporobolus fimbriatus*, *Stipagrostis uniplumis*, *Tragus racemosus*. Herbs: *Barleria macrostegia*, *Geigeria filifolia*, *Geigeria ornativa*, *Gisekia africana*, *Helichrysum cerastoides*, *Heliotropium ciliatum*, *Hermbstaedtia odorata*, *Hibiscus marlothianus*, *Hibiscus pusillus*, *Jamesbrittenia aurantiaca*, *Limeum fenestratum*, *Lippia scaberrima*, *Selago densiflora*, *Vahlia capensis* subsp. *vulgaris*. Succulent herb: *Aloe grandidentata*.

Note: Though many of the above plant species occurs at the site, not all of them necessarily occur at the site.

Ghaap Plateau Vaalbosveld is not listed as threatened according to the National List of Threatened Ecosystems (2011).

3 METHODS

A desktop study comprised not only an initial phase, but also it was used throughout the study to accommodate and integrate all the data that become available during the field observations.

Surveys of a number of study areas in the vicinity of Vryburg, including the site, were conducted from 10 – 16 November 2015 and 23 – 31 January 2016 to note key elements of habitats on the site, relevant to the conservation of fauna and flora. Notes and experience from earlier surveys at the larger study area of the Taung-Vryburg area by R.F. Terblanche that had taken place in July 2011, November 2011, January 2012, February 2012, August 2013, December 2013, January 2014 and November 2014 were also taken into account where applicable. The main purpose of the site visits was ultimately to serve as habitat surveys that concentrated on the possible presence or not of threatened species and other species of high conservation priority.

The following sections highlight the materials and methods applicable to different aspects that were observed.

3.1 HABITAT CHARACTERISTICS AND VEGETATION

The habitat was investigated by noting habitat structure (rockiness, slope, plant structure/physiognomy) as well as floristic composition. Voucher specimens of plant species were only taken where the taxonomy was in doubt and where the plant specimens were of significant relevance for invertebrate conservation. In this case no plant specimens were needed to be collected as voucher specimens or to be send to a herbarium for identification. A wealth of guides and detailed works of plant identifications, ecology and conservation is fortunately available and very useful. Field guides, biogeographic works, species lists, diagnostic outlines, conservation statuses and detail on specific plant groups were sourced from Court (2010), Germishuizen (2003), Germishuizen, Meyer & Steenkamp (2006), Goldblatt (1986), Goldblatt & Manning (1998), Jacobsen (1983), Manning (2003), Manning (2009), McMurtry, Grobler, Grobler & Burns (2008), Pooley (1998), Retief & Herman (1997), Smit (2008), Van Ginkel, Glen, Gordon-Gray, Cilliers, Muasya & Van Deventer (2011), Van Jaarsveld (2006), Van Oudtshoorn (1999), Van Wyk (2000), Van Wyk & Smith (2001), Van Wyk & Smith (2003), Van Wyk & Malan (1998) and Van Wyk & Van Wyk (2013). Lists of species, species names and the conservation status of species were mainly sourced from Raimondo, von Staden, Victor, Helme, Turner, Kamundi & Manyama (2009) and

updated versions of red lists and species from the Threatened Species Programme of SANBI and the Red List of South African Plants (sanbi.org.za).

Indications of abundance and distribution of *Vachellia erioloba* (= *Acacia erioloba*), the Camel Thorn. For areas with where Camel Thorn trees are sparse direct count and positioning of all the individual trees are conducted at the proposed footprint by carefully walking in zigzag fashion through the site. For areas where direct counting become impractical within time and cost limits, counts of individual trees within 50 m x 50 m plots are conducted at zones where higher number of individuals is conspicuous. *Vachellia erioloba* trees sometimes occur in clumps with a number of tree trunks in close proximity which makes the number of individuals difficult to count if the separation of individuals is concerned. As far as possible it was ascertained whether two trunks in close proximity represent different individuals (often the case) or the same individual (when two or more trunks are likely to be part of same tree). Height classes of the trees are taken during the counts.

3.2 MAMMALS

Mammals were noted as sight records by day. For the identification of species and observation of diagnostic characteristics Smithers (1986), Skinner & Chimimba (2005), Cillié, Oberprieler and Joubert (2004) and Apps (2000) are consulted. Sites have been walked, covering as many habitats as possible. Signs of the presence of mammal species, such as calls of animals, animal tracks (spoor), burrows, runways, nests and faeces were recorded. Walker (1996), Stuart & Stuart (2000) and Liebenberg (1990) were consulted for additional information and for the identification of tracks and signs. Because of the type of threatened mammals that are assessed in the local area such as the blackfooted cat and golden moles or rough-haired golden moles which are not to be trapped in normal way, the poor trapping records of species in question such as the White-tailed Mouse as well as the similarity of terrestrial habitats and lack of unique habitats at the sites, trapping was not done since it was not deemed necessary in the case of this study. The focus has been on signs and surveying habitat characteristics to note potential occurrences of mammals of particular conservation concern. Habitat characteristics were also surveyed to note potential occurrences of mammals. Many mammals can be identified from field sightings but, with a few exceptions bats, rodents and shrews can only be reliably identified in the hand, and even then some species needs examination of skulls, or even chromosomes (Apps, 2000).

3.3 BIRDS

A specific avifaunal study has been conducted for the assessment of birds at the site (see avifaunal study).

3.4 REPTILES

Reptiles were noted as sight records in the field. Binoculars (10x30) can also be used for identifying reptiles of which some are wary. For practical skills of noting diagnostic characteristics, the identification of species and observation techniques, Branch (1998), Marais (2004), Alexander & Marais (2007) and Cillié, Oberprieler and Joubert (2004) were followed. The Atlas and Red List of Reptiles of South Africa, Lesotho and South Africa (Bates, Branch, Bauer, Burger, Marais, Alexander & de Villiers, 2014) has been used as the main source to compile the list for assessment. Sites were walked, covering as many habitats as possible. Smaller reptiles are sometimes collected for identification, but this practice was not necessary in the case of this study. Habitat characteristics are surveyed to note potential occurrences of reptiles.

3.5 AMPHIBIANS

Frogs and toads are noted as sight records in the field or by their calls. For practical skills of noting diagnostic characteristics, the identification of species and observation techniques Carruthers (2001), Du Preez (1996), Conradie, Du Preez, Smith & Weldon (2006) and the recent complete guide by Du Preez & Carruthers (2009) are consulted. CD's with frog calls by Carruthers (2001) and Du Preez & Carruthers (2009) are used to identify species by their calls when applicable. Sites are walked, covering as many habitats as possible. Smaller frogs are often collected by pitfall traps put out for epigeal invertebrates (on the soil), but this practice falls beyond the scope of this survey. Habitat characteristics are also surveyed to note potential occurrences of amphibians.

3.6 BUTTERFLIES

Butterflies were noted as sight records or voucher specimens. Voucher specimens are mostly taken of those species of which the taxa warrant collecting due to taxonomic difficulties or in the cases where species can look similar in the veldt. Many butterflies use only one species or a

limited number of plant species as host plants for their larvae. Myrmecophilous (ant-loving) butterflies such as the *Aloeides*, *Chrysoritis*, *Erikssonia*, *Lepidochrysops* and *Orachrysops* species (Lepidoptera: Lycaenidae), which live in association with a specific ant species, require a unique ecosystem for their survival (Deutschländer & Bredenkamp, 1999; Terblanche, Morgenthal & Cilliers, 2003; Edge, Cilliers & Terblanche, 2008; Gardiner & Terblanche, 2010). Known food plants of butterflies were therefore also recorded. After the visits to the site and the identification of the butterflies found there, a list was also compiled of butterflies that will most probably be found in the area in all the other seasons because of suitable habitat. The emphasis of this study remains a habitat survey that focuses on the likelihood of occurrence of threatened, near threatened or rare butterfly species.

3.7 FRUIT CHAFER BEETLES

Different habitat types in the areas were explored for any sensitive or special fruit chafer species. Selection of methods to find fruit chafers depends on the different types of habitat present and the species that may be present. Fruit bait traps would probably not be successful for capturing *Ichneustoma* species in a grassland patch (Holm & Marais 1992). Possible chafer beetles of high conservation priority were noted as sight records accompanied by the collecting of voucher specimens with grass nets or containers where deemed necessary.

3.8 ROCK SCORPIONS

Relatively homogenous habitat / vegetation areas were identified and explored to identify any sensitive or special species. Selected stones that were lifted to search for Arachnids were put back very carefully resulting in the least disturbance possible. All the above actions were accompanied by the least disturbance possible.

3.9 LIMITATIONS

For each site visited, it should be emphasized that surveys can by no means result in an exhaustive list of the plants and animals present on the site, because of the time constraint. Also for the counts of trees an approximation rather than exact abundances are given though some quantitative counts were conducted to groundtruth the estimates. There are many invertebrate

groups with huge taxonomic and biogeographic impediments which further add to limitations of present surveys.

The on site survey was conducted during November 2015 and January 2016 which fall within an optimal time of the season to find sensitive plant and animal species of high conservation priority. Weather conditions during the surveys were favourable for recording fauna and flora. Notes and experience from earlier surveys at the Taung-Vryburg area by R.F. Terblanche that had taken place on July 2011, November 2011, January 2012, February 2012, August 2013, December 2013, January 2014 and November 2014 were also taken into account where applicable, these cover a vast range of conditions including optimal conditions for surveys. The focus of the present survey remains a habitat survey that concentrates on the possibility that species of particular conservation priority occur on the site or not. It is unlikely that any more visits would reveal information that would change the outcome of this assessment both in terms of ecosystems of special conservation concern or suitable habitats of species of particular conservation concern. Visits that were conducted therefore appear to be sufficient to address the objectives of this study.

4 RESULTS

4.1 HABITAT AND VEGETATION CHARACTERISTICS

Table 4.1 Outline of main landscape and habitat characteristics of the site.

HABITAT FEATURE	DESCRIPTION
Topography	The site proposed for the developments is on gentle to very gentle slopes (flat plateau, plain).
Rockiness	No rocky ridges are present at the site.
Presence of wetlands	One small very restricted pan depression is present at the site. No other wetlands are present at the site.
Vegetation	Conspicuous and dense invasions of many parts of the site by the declared alien invasive plant species, <i>Prosopis glandulosa</i> (Mesquite) are encountered at the site. Tree species such as <i>Acacia tortilis</i> subsp. <i>heteracantha</i> (Umbrella Thorn), <i>Searsia lancea</i> (Karee), <i>Acacia mellifera</i> subsp. <i>detinens</i> (Black Thorn) and <i>Acacia erioloba</i> (Camel Thorn) have a dotted distribution in the landscape. Other parts of the site are characterised by typical patches of shrub-height <i>Tarchonanthus camphoratus</i> (Camphor Bush) and <i>Grewia flava</i> (Wild Raisin). Other indigenous trees or shrubs include <i>Searsia lancea</i> (Karee), <i>Diospyros lycioides</i> subsp. <i>lycioides</i> (Karoo Bluebush), <i>Ehretia alba</i> (White Puzzlebush), <i>Ziziphus mucronata</i> (Buffalothorn) and <i>Acacia hebeclada</i> subsp. <i>hebeclada</i> (Candlepod Thorn). Shrublets <i>Ziziphus zeyheriana</i> (Dwarf Buffalothorn) and <i>Gnidia polycephala</i> are present at the site. Dwarf shrubs such as <i>Pentzia calcarea</i> and <i>Pentzia globosa</i> are also found. Indigenous herbaceous plant species at the site include <i>Senna italica</i> subsp. <i>arachoides</i> , <i>Hermibstaedtia odorata</i> , <i>Helichrysum cerastoides</i> , <i>Barleria macrostegia</i> and <i>Salvia disermas</i> . Indigenous grass species include <i>Elionurus muticus</i> , <i>Cynodon dactylon</i> , <i>Aristida congesta</i> , <i>Eragrostis lehmanniana</i> , <i>Aristida adscensionis</i> , <i>Enneapogon cenchroides</i> and <i>Tragus racemosus</i> . Conspicuous exotic plant species at the site include <i>Prosopis glandulosa</i> (Honey Mesquite), <i>Opuntia ficus-indica</i> and <i>Opuntia imbricata</i> .
Signs of disturbances	Visible extensive and dense infestations of declared alien invasive species, <i>Prosopis glandulosa</i> (Mesquite) exist at the site. Some patches with fairly natural vegetation remain but many parts appear ecologically disturbed.
Connectivity of natural vegetation in the site and between the site and surrounding areas	The footprint proposed for the proposed development is not part of a corridor of particular conservation importance. Small and restricted depression at the site could be an important stepping stone in a conservation corridor system if <i>Prosopis glandulosa</i> could be eradicated at its surrounds.



Photo 1 View of part of proposed footprint, with indigenous *Acacia hebeclada* subsp. *hebeclada* (Candlepod Thorn) and the exotic *Opuntia ficus-indica* (Sweet Prickly Pear) in the foreground and grassy patches with bushclumps in the background.
Photo: November 2015, R.F. Terblanche.



Photo 2 Some parts of the site are extensively infested by the declared alien invasive plant species *Prosopis glandulosa* (Honey Mesquite) such as photographed here at site.
Photo: November 2015, R. F. Terblanche.



Photo 3 Alien invasive *Prosopis glandulosa* (Honey Mesquite tree) at the site.
Photo: November 2015, R.F. Terblanche.



Photo 4 *Grewia flava* (Velvet Raisin) a widespread woody plant species that is often frequent in the Eastern Kalahari Bushveld Bioregion.
Photo: November 2015, R.F. Terblanche.



Photo 5 *Ehretia alba* (White Puzzlebush) at the site.
Photo: November 2015, R.F. Terblanche.



Photo 6 *Solanum incanum* (Bitter Apple) a plant species that often establishes at open disturbed areas, at the site.
Photo: November 2015, R.F. Terblanche.

4.2 ASSESSMENT OF PLANT SPECIES OF PARTICULAR CONSERVATION PRIORITY

4.2.1 Plant species of particular conservation concern according to the red list of plants

Table 4.2 Threatened plant species of the North West Province which are listed in the **Critically Endangered** category. The list here follows the most recent updated red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Brachystelma canum</i>	Critically Endangered	No
<i>Brachystelma gracillimum</i>	Critically Endangered	No

Table 4.3 Threatened plant species of the North West Province which are listed in the **Endangered** category. The list here follows the most recent updated red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Aloe peglerae</i>	Endangered	No
<i>Brachystelma discoideum</i>	Endangered	No

Table 4.4 Threatened plant species of the North West Province which are listed in the **Vulnerable** category. The list here follows the most recent updated red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Brachycorythis conica</i> subsp. <i>transvaalensis</i>	Vulnerable	No
<i>Brachystelma incanum</i>	Vulnerable	No
<i>Ceropegia decidua</i> subsp. <i>pretoriensis</i>	Vulnerable	No
<i>Ceropegia stentiae</i>	Vulnerable	No
<i>Ledebouria atrobrunnea</i>	Vulnerable	No
<i>Marsilea farinosa</i>	Vulnerable	No
<i>Melolobium subspicatum</i>	Vulnerable	No
<i>Prunus africana</i>	Vulnerable	No
<i>Rennera stellata</i>	Vulnerable	No
<i>Searsia maricoan</i>	Vulnerable	No

Table 4.5 Near Threatened plant species of the North West Province. The list here follows the most recent updated red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Adromischus umbraticola</i> subsp. <i>umbraticola</i>	Near Threatened	No
<i>Ceropegia turricula</i>	Near Threatened	No
<i>Cineraria austrotransvaalensis</i>	Near Threatened	No
<i>Cleome conrathii</i>	Near Threatened	No
<i>Delosperma leendertziae</i>	Near Threatened	No
<i>Drimia sanguinea</i>	Near Threatened	No
<i>Elaeodendron transvaalense</i>	Near Threatened	No
<i>Kniphofia typhoides</i>	Near Threatened	No
<i>Lithops leslei</i> subsp. <i>leslei</i>	Near Threatened	No
<i>Nerine gracilis</i>	Near Threatened	No
<i>Sporobolus oxyphyllus</i>	Near Threatened	No
<i>Stenostelma umbelluliferum</i>	Near Threatened	No

Table 4.6 Plant species of the North West Province which are not threatened and not near threatened but which are of particular conservation concern and listed in the **Critically Rare** category (Raimondo *et al.* 2009). The list here follows the most recent red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Conservation status	Resident at the site
<i>Gladiolus filiformis</i>	Critically Rare	No

Table 4.7 Plant species of the North West Province which are not threatened and not near threatened but of which are of particular conservation concern and listed in the **Rare** category (Raimondo *et al.* 2009). The list here follows the most recent red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Brachystelma dimorphum</i> subsp. <i>gratum</i>	Rare	No
<i>Ceropegia insignis</i>	Rare	No
<i>Frithia pulchra</i>	Rare	No
<i>Gnaphalium nelsonii</i>	Rare	No
<i>Habenaria culveri</i>	Rare	No

Table 4.8 Plant species of the North West Province which are not threatened and not near threatened but which are of particular conservation concern and listed in the **Declining** category (Raimondo *et al.* 2009). The list here follows the most recent red list of South African plant species (Raimondo *et al.* 2009). No = Plant species is unlikely to be a resident at the site; Yes = Plant species is a resident at the site.

Species	Status: Global status or national status indicated	Resident at the site
<i>Boophone disticha</i>	Declining	Yes
<i>Crinum bulbispermum</i>	Declining	No
<i>Crinum macowanii</i>	Declining	No
<i>Drimia altissima</i>	Declining	No
<i>Eucomis autumnalis</i>	Declining	No
<i>Gunnera perpensa</i>	Declining	No
<i>Hypoxis hemerocallidea</i>	Declining	No
<i>Ilex mitis</i>	Declining	No
<i>Pelargonium sidoides</i>	Declining	No
<i>Vachellia erioloba</i> (= <i>Acacia erioloba</i>)	Declining	Yes

4.2.2 Plant species of particular conservation concern: protected species

Table 4.9 Tree species of the North West Province which are listed as **Protected Species** under the National Forests Act No. 84 of 1998, Section 51(1). No = Plant species is not a resident on the site; Yes = Plant species is a resident at the site.

Species	Conservation status	Resident at the site
<i>Boscia albitrunca</i> (Shepherd's Tree)	Protected	No
<i>Sclerocarya birrea</i> (Marula)	Protected	No
<i>Vachellia erioloba</i> (= <i>Acacia erioloba</i>) (Camel Thorn Tree)	Protected	Yes

4.3 ASSESSMENT OF VERTEBRATE SPECIES OF PARTICULAR HIGH CONSERVATION PRIORITY

4.3.1 Mammals of particular high conservation priority

Table 4.10 Threatened mammal species of the North West Province. Literature sources: Friedman & Daly, (2004), Skinner & Chimimba (2005), Wilson & Reeder (2005). With mammal species which normally needs a large range their residential status does not implicate that they are exclusively dependent on the site or use the site as important shelter or for reproduction. No = Not recorded at site/ Unlikely to be resident at the site. Yes: Recorded at the site/ Likely to be resident at the site.

Species	Threatened Status	Site is part of range	Recorded at site during survey	Likely to be found based on habitat assessment
<i>Chrysoxalax villosus</i> Rough-haired golden mole	Vulnerable	No	No	No
<i>Cloeotis percivali</i> Short-eared Trident Bat	Vulnerable/ Near-threatened	No	No	No
<i>Diceros bicornis</i> Black rhinoceros	Critically Endangered	No	No	No
<i>Lycaon pictus</i> African wild dog	Endangered	No	No	No

<i>Loxodonta africana</i> African elephant	Vulnerable	No	No	No
<i>Mystromys albicaudatus</i> White-tailed mouse	Endangered	Yes	No	No
<i>Neamblysomus julianae</i> Juliana's Golden Mole	Critically Endangered	No	No	No
<i>Panthera leo</i> Lion	Vulnerable	No	No	No
<i>Rhinolophus blasii</i> Blasi's Horseshoe Bat	Vulnerable	No	No	No

Table 4.11 Near threatened mammal species known to occur in the North West Province. Literature sources: Skinner & Chimimba (2005). No = Not recorded at site/ unlikely to be resident at the site. Yes: Recorded at the site/ Likely to be resident at the site.

Species	Threatened Status	Site is part of range	Recorded at site during survey	Likely to be found based on habitat assessment
<i>Ceratotherium simum</i> White Rhinoceros	Near threatened	No	No	No
<i>Manis temminckii</i> Ground Pangolin	Near threatened	No	No	No

Table 4.12 Data deficient (or uncertain) mammal species of the North West Province. Literature sources: Skinner & Chimimba (2005). No = Not recorded at site/ unlikely to be resident at the site. Yes: Recorded at the site/ Likely to be resident at the site.

Species	Threatened Status	Recorded at site during survey	Likely be a resident at the site
<i>Myosorex varius</i> Forest shrew	Uncertain	No	No

4.3.2 Birds of particular high conservation priority

See separate avifaunal report.

4.3.3 Reptiles of particular high conservation priority

The following tables list possible presence or absence of threatened reptile or near threatened reptile species in the study area. The Atlas and Red List of Reptiles of South Africa, Lesotho and South Africa (Bates, Branch, Bauer, Burger, Marais, Alexander & de Villiers, 2014) has been used as the main source to compile the list for assessment.

Table 4.13 Threatened reptile species in North West Province. Main Source: (Bates, Branch, Bauer, Burger, Marais, Alexander & de Villiers, 2014). No = Reptile species is not a resident on the site; Yes = Reptile species is found to be resident on the site.

Species	Threatened Status	Resident at site	Recorded at site during survey	Likely to be found based on habitat assessment
<i>Crocodylus niloticus</i> Nile Crocodile	Vulnerable	No	No	No

Table 4.14 Near threatened reptile species in North West Province. Main Source: Bates, Branch, Bauer, Burger, Marais, Alexander & de Villiers (2014). Though *Homoroselaps dorsalis* has not yet been recorded from the North West Province, its presence in some areas or the Province is anticipated. No = Reptile species is not a resident on the site; Yes = Reptile species is found to be resident on the site.

Species	Threatened Status	Resident at site	Recorded at site during survey	Likely to be found based on habitat assessment
<i>Homoroselaps dorsalis</i> Striped Harlequin Snake	Near threatened	No	No	No

4.3.4. Amphibian species of particular conservation concern

Table 4.15 Near threatened amphibian species in North West Province. No = Amphibian species is not a resident on the site; Yes = Amphibian species is found to be resident on the site.

Species	Threatened Status	Resident at site	Recorded at site during survey	Likely to be found based on habitat assessment
<i>Pyxicephalus adspersus</i> Giant Bullfrog	Near threatened (Currently Least Concern)	No	No	No

4.4 ASSESSMENT OF INVERTEBRATE SPECIES OF PARTICULAR HIGH CONSERVATION PRIORITY

4.4.1 Butterflies of particular conservation priority

Table 4.16 Threatened butterfly species in North West Province, northeastern parts of the Northern Cape and Gauteng Province. Sources: Henning, Terblanche & Ball (2009), Mecenero *et al.* (2013). Invertebrates such as threatened butterfly species are often very habitat specific and residential status imply a unique ecosystem that is at stake.

Species	Threatened Status	Recorded at site during survey	Residential status at the site: Yes confirmed, Highly likely, Likely, Medium possibility, Unlikely, Highly unlikely
<i>Aloeides dentatis dentatis</i> Roodepoort Copper	Endangered	No	Highly unlikely
<i>Anthene lindae</i> Kalahari Hairtail	Vulnerable	No	Unlikely
<i>Chrysochrysis aureus</i> Golden Copper	Endangered	No	Highly unlikely
<i>Lepidochrysops praeterita</i> Highveld Blue	Endangered	No	Highly unlikely
<i>Orachrysops mijburghi</i> Mijburgh's Blue	Endangered	No	Highly unlikely

Table 4.17 Butterfly species of the North West Province and Gauteng Province that are not threatened and not near threatened but of which are of particular conservation concern and listed in the **Rare** category (Mecenero *et al.*, 2013). No = Butterfly species is unlikely to be a resident at the study area; Yes = Butterfly species is a resident at the study area.

Species	Threatened Status	Recorded at site during survey	Residential status at the site: Yes confirmed, Highly likely, Likely, Medium possibility, Unlikely, Highly unlikely
<i>Colotis celimene amina</i> Lilac Tip	Rare (Low density)	No	Highly unlikely
<i>Lepidochrysops procera</i> Savanna Blue	Rare (Habitat specialist)	No	Highly unlikely
<i>Metisella meninx</i> Marsh Sylph	Rare (Habitat specialist)	No	Highly unlikely
<i>Platylesches dolomitica</i> Hilltop Hopper	Rare (low density)	No	Highly unlikely

4.4.2 Beetles of particular conservation priority

Table 4.18 Fruit chafer species (Coleoptera: Scarabaeidae: Cetoninae) in the Gauteng Province and North-West Province which are of known high conservation priority.

Species	Threatened Status	Recorded at site during survey	Likely to be resident based on habitat assessment
<i>Ichneustoma stobbiai</i>	Uncertain	No	No
<i>Trichocephala brincki</i>	Uncertain	No	No

4.4.3 Scorpion species of particular conservation priority

Table 4.19 Rock scorpion species (Scorpiones: Ischnuridae) species that are of known high conservation priority in the Gauteng Province and North-West Province.

Species	Threatened Status	Recorded at site during survey	Likely to be resident at site based on habitat assessment
<i>Hadogenes gracilis</i>	Uncertain	No	No
<i>Hadogenes gunningi</i>	Uncertain	No	No

4.5 ASSESSMENT OF CAMEL THORN TREES AT THE SITE

During the initial surveys it was found that *Vachellia erioloba* (= *Acacia erioloba*), Camel Thorn trees, are present at the site. Additional surveys were conducted to indicate the distribution and abundance of *Vachellia erioloba* at the site.

At the eastern parts of the proposed footprint some *Vachellia erioloba*, including > 5 m – 10 m tall height class appear to have a sparse dotted distribution in the landscape. At the central northern parts of the site possible secondary succession resulted in numerous *Vachellia erioloba* < 5 m where up to 6 individuals taller than 2 m could be found. It is estimated that > 100 but less than 200 individuals of *Vachellia erioloba* are present at the site, possibly owing to secondary succession in the central-northern area at the site.

Table 4.20 Indication of attributes of population of Camel Thorn tree, *Vachellia erioloba* (= *Acacia erioloba*) at the proposed footprint.

Part of proposed footprint	Approximate estimation of number of <i>Vachellia erioloba</i> taller than 2 m, per hectare at the proposed footprint.	Likelihood of presence of individuals taller than 10 m	Notes on individuals less than 2	Presence or absence of particular old and remarkable Camel Thorn trees at proposed footprint	Presence or absence of Camel Thorn Forest of note
Entire footprint	Up to 6 per 50 m x 50 m plot in central northern area at the proposed footprint. Largely absent in western parts of proposed footprint and dotted in landscape at eastern part of proposed footprint. Estimated at more than 100 but less than 200 individuals taller than 2 m.	Highly unlikely	Very few individuals less than 2 m seen at proposed footprint, apart from the central northern-area where some secondary succession possibly is taken place.	No particularly old or remarkable trees found at proposed footprint.	Absent.

4.6 ASSESSMENT OF ALIEN INVASIVE SPECIES WITH PARTICULAR REFERENCE TO *PROSOPIS GLANDULOSA* (HONEY MESQUITE)

Exotic weeds at the site include *Agremone ochroleuca* (White-flowered Mexican Poppy), *Chenopodium album* (Goosefoot), *Opuntia ficus-indica* (Prickly Pear) and *Schkuhria pinnata* (Dwarf Marigold). Though these exotic weeds easily spring up where disturbances such as overgrazing, scraping of an area and diggings are found, at the present study area no severe infestations such as could often be observed in larger urban areas and surrounds in the North West and Gauteng Provinces, are found.

During the initial surveys it was found that *Prosopis glandulosa* (Honey Mesquite tree) thickets occur in some parts of the Naledi Local Municipality and particular consideration has been given to this highly invasive tree species at and near the proposed footprint.

Table 4.21 Summary of assessment of the highly alien invasive *Prosopis glandulosa* (Honey Mesquite tree).

Part of the proposed footprint	Approximate estimation of number of <i>Prosopis glandulosa</i> taller than 2 m	Notes on <i>Prosopis glandulosa</i> individuals of height less than 2 m	Presence or absence of severe invasion by <i>Prosopis glandulosa</i>.
Entire area of proposed footprint	<p>At areas of conspicuous density of <i>Prosopis</i> 12 – 52 individuals taller than 2 m were counted in each of four 50 x 50 m sample plots. At one sample plot 139 <i>Prosopis</i> individuals all together were counted. At least more than 1000 individual <i>Prosopis</i> trees taller than 2 m must be at the proposed footprint and at least as many less than 2 m tall. More than 2000 individuals of <i>Prosopis</i> at the site.</p>	<p>At one sample plot of 50 m x 50 m, 87 individuals less than 2 m tall were counted. Many individuals less than 2 m were seen across the highly infested area.</p>	<p>Extensive invasion by <i>Prosopis glandulosa</i> is present at the proposed footprint area north of the present homestead.</p>

5 DISCUSSION

5.1 HABITAT AND VEGETATION CHARACTERISTICS

An outline of the habitat and vegetation characteristics is given in Table 4.1.

5.2 PLANT SPECIES

Extinct, threatened, near threatened and other plant species of high conservation priority in North West Province are listed in Tables 4.2 – 4.8. Protected tree species are listed in Table 4.9. The presence or not of all the species listed in the tables were investigated during the survey. None of the threatened and near-threatened plant species are likely to occur on the site. Declining occur at the site; these are *Boophane disticha* (Poison Bulb) and *Vachellia erioloba* (= *Acacia erioloba*) (Camel Thorn tree). *Vachellia erioloba* that occurs at the site is also a Protected Tree species. See section 5.5 Protected tree species under the National Forests Act No. 84 of 1998 are listed in Table 4.9. In terms of a part of section 51(1) of Act No. 84 of 1998, no person may cut, disturb, damage or destroy any protected tree or possess, collect, remove, transport, export, purchase, sell, donate or in any other manner acquire or dispose of any protected tree, except under a license granted by the Minister. See section 5.5 for a more detailed discussion of the Camel Thorn tree at the proposed footprint.

5.3 VERTEBRATES

5.3.1 Mammals

Table 4.10, Table 4.11 and Table 4.12 list the possible presence or absence of threatened mammal species, near threatened mammal species and mammal species of which the status is uncertain, respectively, at the site. Literature sources that were used are Friedman & Daly (2004), Skinner & Chimimba (2005) and Wilson & Reeder (2005). Since the site falls outside reserves, threatened species such as the black rhinoceros (*Diceros bicornis*) and the African wild dog (*Lycaon pictus*) are obviously not present. No smaller mammals of particular high conservation significance are likely to be found on the site as well.

5.3.2 Birds

See separate avifaunal report for assessment of birds.

5.3.3 Reptiles

Table 4.13 and Table 4.14 list the possible presence or absence of threatened and near threatened reptile species on the site. The Southern African Reptile Conservation Assessment (SARCA) was launched in May 2005 (Branch, Tolley, Cunningham, Bauer, Alexander, Harrison, Turner & Bates, 2006). Its primary aim is to produce a conservation assessment for reptiles of South Africa, Lesotho and Swaziland within a four year period, ending 2009 (Branch *et al.*, 2006). Therefore a full up-dated conservation assessment of reptiles, taking into account the recent IUCN (2001) criteria, will only be available in the near future. While the conservation statuses of reptile species are under revision Alexander & Marais (2007) as well as Tolley & Burger 2007) give useful indications of possible red listings in the near future. There appears to be no threat to any reptile species of particular high conservation importance if the site is developed.

5.3.4 Amphibians

No frog species that occur in the North West are threatened as threatened species (vulnerable, endangered or critically endangered) according to Minter, Burger, Harrison, Braack, Bishop and Kloepfer (2004) as well as Du Preez & Carruthers (2009). Table 4.15 lists *Pyxicephalus adspersus* (Giant Bullfrog) as near threatened (Minter *et al.*, 2004; Du Preez & Carruthers, 2009). There is no suitable habitat for *Pyxicephalus adspersus* (Giant Bullfrog) at the site. There appears to be no threat to any amphibian species of particular high conservation importance if the site is developed.

5.4 INVERTEBRATES

5.4.1 Butterflies

Studies about the vegetation and habitat of threatened butterfly species in South Africa showed that ecosystems with a unique combination of features are selected by these often localised threatened butterfly species (Deuschländer and Bredenkamp 1999; Edge 2002, 2005; Terblanche, Morgenthal & Cilliers 2003; Lubke, Hoare, Victor & Ketelaar 2003; Edge, Cilliers & Terblanche, 2008). Threatened butterfly species in South Africa can then be regarded as bio-indicators of rare ecosystems.

Five species of butterfly in Gauteng Province, the northeastern parts of the Northern Cape and North West Province combined are listed as threatened in the recent butterfly conservation assessment of South Africa (Mecenero *et al.*, 2013). The expected presence or not of these threatened butterfly species as well as species of high conservation priority that are not threatened, at the site (Table 4.16 and Table 4.17) follows.

5.4.1.1 Assessment of threatened butterfly species

***Aloeides dentatis dentatis* (Roodepoort Copper)**

The proposed global red list status for *Aloeides dentatis dentatis* according to the most recent IUCN criteria and categories is Endangered (Mecenero *et al.*, 2013). *Aloeides dentatis dentatis* colonies are found where one of its host plants *Hermannia depressa* or *Lotononis eriantha* is present. Larval ant association is with *Lepisiota capensis* (S.F. Henning 1983; S.F. Henning & G.A. Henning 1989). The habitat requirements of *Aloeides dentatis dentatis* are complex and not fully understood yet. See Deuschländer and Bredenkamp (1999) for the description of the vegetation and habitat characteristics of one locality of *Aloeides dentatis* subsp. *dentatis* at Ruimsig, Roodepoort, Gauteng Province. There is not an ideal habitat of *Aloeides dentatis* subsp. *dentatis* on the site and it is unlikely that the butterfly is present at the site.

***Anthene lindae* (Kalahari Hairtail)**

Small but distinct butterfly species discovered by R.F. Terblanche in 1990 at the present Witsand Nature Reserve in the Northern Cape. Recent red listing and extinction risk assessments list *Anthene lindae* as Vulnerable (Henning, Terblanche & Ball, 2009; Mecenero *et al.*, 2013).

Because *Acacia erioloba* occurs on the site, the presence or not of *Anthene lindae* (Kalahari hairtail butterfly, Vulnerable) has bearing. The butterfly is intimately associated with *Acacia erioloba* which may prove to be the larval food plant (Terblanche, 1994). Present observations by V. Jessnitz confirmed egg-laying of females of *Anthene lindae* on *Vachellia erioloba* (unpublished). However, all the localities for this butterfly species have been found on what appears to be a unique catchment area and basins with particular high water tables on the western side of the Langberg mountain chain, Northern Cape Province (Terblanche & Taylor, 2000). According to Henning *et al.* (2009) *Anthene lindae* has up to date only been found at an ecotone between Gordonia Plains Shrubland and Olifantshoek Plains Thornveld (Mucina & Rutherford, 2006). *Anthene lindae* is not found everywhere where *Vachellia erioloba* trees are present and based on the present knowledge and the surveys at the site, presence of the butterfly at the site is unlikely.

***Chrysochrysis aureus* (Golden Highveld Opal/ Heidelberg Copper)**

The proposed global red list status for *Chrysochrysis aureus* according to the most recent IUCN criteria and categories is Endangered (Mecenero *et al.*, 2013) *Chrysochrysis aureus* (Golden Opal/ Heidelberg Copper) is a resident where the larval host plant, *Clusia pulchella* is present. However, the distribution of the butterfly is much more restricted than that of the larval host plant (S.F. Henning 1983; Terblanche, Morgenthal & Cilliers 2003). One of the reasons for the localised distribution of *Chrysochrysis aureus* is that a specific host ant *Crematogaster liengmei* must also be present at the habitat. Fire appears to be an essential factor for the maintenance of suitable habitat (Terblanche, Morgenthal & Cilliers 2003). Research revealed that *Chrysochrysis aureus* (Golden Opal/ Heidelberg Copper) has very specific habitat requirements, which include rocky ridges with a steep slope and a southern aspect (Terblanche, Morgenthal & Cilliers 2003). Owing to a lack of habitat requirements and ideal habitat the presence of the taxon is highly unlikely.

***Lepidochrysis praeterita* (Highveld Blue)**

The proposed global red list status for *Lepidochrysis praeterita* according to the most recent IUCN criteria and categories is Endangered (G.A. Henning, Terblanche & Ball, 2009; Mecenero *et al.*, 2013). *Lepidochrysis praeterita* is a butterfly that occurs where the larval host plant *Ocimum obovatum* (= *Becium obovatum*) is present (Pringle, G.A. Henning & Ball, 1994), but the distribution of the butterfly is much more restricted than the distribution of the host plant.

Lepidochrysops praeterita is found on selected rocky ridges and rocky hillsides in parts of Gauteng, the extreme northern Free State and the south-eastern Gauteng Province. No ideal habitat appears to be present for the butterfly on the site. It is unlikely that *Lepidochrysops praeterita* would be present on the site and at the footprint proposed for the development.

***Orachrysops mijburghi* (Mijburgh's Blue)**

The proposed global red status for *Orachrysops mijburghi* according to the most recent IUCN criteria and categories is Endangered (Mecenero *et al.*, 2013). *Orachrysops mijburghi* favours grassland depressions where specific *Indigofera* plant species occur (Terblanche & Edge 2007). The Heilbron population of *Orachrysops mijburghi* in the Free State uses *Indigofera evansiana* as a larval host plant (Edge, 2005) while the Suikerbosrand population in Gauteng uses *Indigofera dimidiata* as a larval host plant (Terblanche & Edge 2007). There is no suitable habitat for *Orachrysops mijburghi* on the site and it is unlikely that *Orachrysops mijburghi* would be present on the site.

Conclusion on threatened butterfly species

There appears to be no threat to any threatened butterfly species if the site is developed.

5.4.1.2 Assessment of butterfly species that are not threatened but also of high conservation priority

***Colotis celimene amina* (Lilac tip)**

Colotis celimene amina is listed as Rare (Low density) by Mecenero *et al.* (2013). In South Africa *Colotis celimene amina* is present from Pietermaritzburg in the south and northwards into parts of Kwa-Zulu Natal, Gauteng, Limpopo, Mpumalanga and the North West Provinces (Mecenero *et al.* In press.). Reasons for its rarity are poorly understood. It is highly unlikely that *Colotis celimene amina* would be present at the site or make use of the site as a particular habitat.

***Lepidochrysops procera* (Savanna Blue)**

Lepidochrysops procera is listed as Rare (Habitat specialist) by Mecenero *et al.* (2013). *Lepidochrysops procera* is endemic to South Africa and found in Gauteng, KwaZulu-Natal, Mpumalanga and North West (Mecenero *et al.*, 2013). Owing to a lack of habitat requirements and ideal habitat the presence of the taxon at the site is highly unlikely.

Metisella meninx (Marsh Sylph)

Henning and Henning (1989) in the first South African Red Data Book of Butterflies, listed *Metisella meninx* as threatened under the former IUCN category Indeterminate. Even earlier in the 20th century Swanepoel (1953) raised concern about vanishing wetlands leading to habitat loss and loss of populations of *Metisella meninx*. According to the second South African Red Data Book of butterflies (Henning, Terblanche & Ball, 2009) the proposed global red list status of *Metisella meninx* has been Vulnerable. During a recent large scale atlasing project the *Conservation Assessment of Butterflies of South Africa, Lesotho and Swaziland: Red List and Atlas* (Mecenero *et al.*, 2013) it was found that more *Metisella meninx* populations are present than thought before. Based on this valid new information, the conservation status of *Metisella meninx* is now regarded as Rare (Habitat specialist) (Mecenero *et al.*, 2013). Though *Metisella meninx* is more widespread and less threatened than perceived before, it should be regarded as a localised rare habitat specialist of conservation priority, which is dependent on wetlands with suitable patches of grass at wetlands (Terblanche *In prep.*). Another important factor to keep in mind for the conservation of *Metisella meninx* is that based on very recent discoveries of new taxa in the group the present *Metisella meninx* is species complex consisting of at least three taxa (Terblanche *In prep.*, Terblanche & Henning *In prep.*). The ideal habitat of *Metisella meninx* is treeless marshy areas where *Leersia hexandra* (rice grass) is abundant (Terblanche *In prep.*). The larval host plant of *Metisella meninx* is wild rice grass, *Leersia hexandra* (G.A. Henning & Roos, 2001). Owing to a lack of habitat requirements and ideal habitat the presence of the taxon at the site is highly unlikely.

Platylesches dolomitica (Hilltop Hopper)

Platylesches dolomitica is listed as Rare (Low density) by Mecenero *et al.* (2013). Historically the conservation status of *Platylesches dolomitica* was proposed to be Vulnerable (Henning, Terblanche & Ball 2009). However this butterfly which is easily overlooked and has a wider distribution than perceived before. *Platylesches dolomitica* has a patchy distribution and is found on rocky ledges where *Parinari capensis* occurs, between 1300 m and 1800m (Mecenero *et al.* 2013, Dobson Pers comm.). Owing to a lack of habitat requirements and ideal habitat the presence of the taxon at the site is highly unlikely.

5.4.2 Fruit chafer beetles

Table 4.18 lists the fruit chafer beetle species (Coleoptera: Scarabaeidae: Cetoninae) that are of known high conservation priority in the North West Province. No *Ichneustoma stobbiai* or *Trichocephala brincki* were found during the surveys. There appears to be no suitable habitat for *Ichneustoma stobbiai* or *Trichocephala brincki* at the site. There appears to be no threat to any of the fruit chafer beetles of particular high conservation priority if the site were developed.

5.4.3 Scorpions

Table 4.19 lists the rock scorpion species (Scorpiones: Ischnuridae) that are of known high conservation priority in the North West Province. None of these rock scorpions have been found at the site and the habitat does not appear to be optimal.

5.5 CAMEL THORN TREES, *VACHELLIA ERIOLOBA* (= *ACACIA ERIOLOBA*)

Camel Thorn trees (*Vachellia erioloba* = *Acacia erioloba*) are found at the proposed footprint. Recent research reinforces the recognition of two genera *Vachellia* and *Senegalia* for hitherto *Acacia* in South Africa (see Kyalangalilwa *et al.*, 2013) hence the name *Vachellia erioloba*. Camel Thorn trees can be described as trees of “great value, beauty and apparent ecological importance” (Anderson & Anderson, 2001).

Table 4.20 gives an indication of attributes of population of Camel Thorn tree, *Vachellia erioloba* (= *Acacia erioloba*) at the proposed footprint. *Vachellia erioloba* individuals at the site are not particularly large and are not part of a camel thorn forest of note. It is estimated that more than 100 but less than 200 individuals of Camel Thorn taller than 2 m are present at the proposed footprint if an arable area is included. It is recommended that a permit at the relevant authorities should be applied for in case of any damage or removal of individual trees and that *Vachellia erioloba* trees (from a nursery or being cultivated) could be planted on site outside the present footprint. These strategies and actions can only be applied in liaison and with the permission of government as outlined above.

5.6 ALIEN INVASIVE SPECIES WITH PARTICULAR REFERENCE TO *PROSOPIS* *GLANDULOSA* (HONEY MESQUITE)

Exotic weeds at the site include *Agremone ochroleuca* (White-flowered Mexican Poppy), *Chenopodium album* (Goosefoot), *Opuntia ficus-indica* (Prickly Pear) and *Schkuhria pinnata* (Dwarf Marigold). Though these exotic weeds easily spring up where disturbances such as overgrazing, scraping of an area and diggings are found, at the present study area no severe infestations such as could often be observed in larger urban areas and surrounds in the North West and Gauteng Provinces, are found.

During the initial surveys it was found that *Prosopis glandulosa* (Honey Mesquite tree) thickets occur in some parts of the Naledi Local Municipality and particular consideration has been given to this highly invasive tree species at and near the proposed footprint. Table 4.21 gives a summary of assessment of the highly alien invasive *Prosopis glandulosa* (Honey Mesquite tree). At areas of conspicuous density of *Prosopis* 12 – 52 individuals taller than 2 m were counted in each of four 50 x 50 m sample plots. At one sample plot 139 *Prosopis* individuals all together were counted.

At least more than 1000 individual *Prosopis* trees taller than 2 m must be at the proposed footprint and at least as many that are less than 2 m tall which constitute that at least more than 2000 individuals of *Prosopis* are present at the proposed footprint.

Many *Prosopis* species have been introduced to South Africa; some taxa and their hybrids have naturalised and become widespread invasive trees. *Prosopis* was introduced to South Africa and has become the second most widespread invasive alien plant taxon in the country (Shackleton et al., 2015c). These invasions have detrimental effects on biodiversity, ecosystem services and human livelihoods (Shackleton et al., 2015a). In South Africa it was found that native woody species density, basal area, richness and diversity all decreased significantly as the basal area of *Prosopis* stands increased (Shackleton et al., 2015a). The cover of native perennial grasses and herbaceous plants declined from 15–20% where the basal area of *Prosopis* was < 2 m²/ha to zero where the basal area of *Prosopis* was > 4.5 m²/ha (Shackleton et al., 2015a). *Prosopis* in South Africa also has higher recruitment (% juvenile plants) than in other areas like Australia. *Prosopis* invasions are having a negative effect on the stability of native tree populations in South

Africa, and are linked to increased mortality of native trees. Improved management of *Prosopis* is needed Shackleton *et al.*, 2015c).

Invasive alien trees impact the environment and human livelihoods. Costs associated with *Prosopis* were perceived to exceed benefits, and most stakeholders wanted to see a reduction in the abundance of *Prosopis* stands (Shackleton *et al.*, 2015b). The mean total cost for the management of *Prosopis* was US\$ 1914 year⁻¹ per farm, where costs ranged from under US\$ 10 to over UD\$ 500 per ha based on invasion densities and objectives for control (Shackleton *et al.*, 2015b).

6 IMPACT ASSESSMENT AND MITIGATION MEASURES

The primary cause of loss of biological diversity is habitat degradation and loss (IUCN, 2004; Primack, 2006). Habitats of threatened plants are in danger most often due to urban developments such as is the case for the Gauteng Province (Pfab & Victor, 2002). Habitat conservation is the key to the conservation of invertebrates such as threatened butterflies (Deutschländer and Bredenkamp 1999; Edge 2002, 2005; Terblanche, Morgenthal & Cilliers 2003; Lubke, Hoare, Victor & Ketelaar 2003; Edge, Cilliers & Terblanche, 2008). Though human impacts in few cases have improved the habitat for mammalian species such as greater cane rats, that prosper in sugar cane and maize fields (Apps 2000), for many mammalian habitat specialist species, human impacts has lead to habitat loss. Some mammal species, especially many of the larger species, could adapt to a wide range of habitat types, but then need a large range. Some animals and plants are rare and occupy only one or a few specialised habitats (Primack 2006). Habitat conservation, either as large available land or as specialised habitats is therefore key to the conservation of many threatened plant species and animal species or any other species of high conservation priority (i.e. rare, near threatened species). In addition corridors and linkages may play a significant role in conservation of fauna.

Corridors are important to link ecosystems of high conservation priority. Such corridors or linkages are there to improve the chances of survival of otherwise isolated populations (Samways, 2005). How wide should corridors be? The answer to this question depends on the conservation goal and the focal species (Samways, 2005). Corridors for mammalian species are especially important for migratory species (Mwalyosi, 1991, Pullin 2002). For an African butterfly assemblage this is about 250m when the corridor is for movement as well as being a habitat source (Pryke and Samways 2003). Hill (1995) found a figure of 200m for dung beetles in tropical Australian forest. In the agricultural context, and at least for some common insects, even small corridors can play a valuable role (Samways, 2005). Much more research remains to be done to find refined answers to the width of grassland corridors in South Africa. The width of corridors will also depend on the type of development, for instance the effects of the shade of multiple story buildings will be quite different from that of small houses. Corridors have a number of advantages related to dispersal and gene flow by avoiding isolation of ecological patches. However, corridors could also have potential drawbacks, for example creating gene flow where none has occurred

naturally in the past and also as reservoirs for pathogens or introduced species (Pullin, 2002). Perhault and Lomolino (2000) studied corridors and mammal community structure in an old-growth forest landscape in the United States of America and their data suggest that each corridor should be valued individually. A lot of research remains to be conducted to have a better idea of the value of corridors, but in general corridors would be of considerable value. It appears that a network of wetland corridors and rocky ridges is highly likely to be of considerable benefit in environmental management and planning. Though proper management plans for habitats are not in place, setting aside special ecosystems is in line with the recent Biodiversity Act (2004) of the Republic of South Africa.

To summarise: In practice, as far as any developments are concerned, the key would be to prioritise and plan according to sensitive species and special ecosystems.

In the case of this study conspicuous and dense invasions of many parts of the site by the declared alien invasive plant species, *Prosopis glandulosa* (Mosquite) are encountered at the site. More natural vegetation patches that remain is a savanna dotted by indigenous trees, with typical patches of shrub-height *Tarchonanthus camphoratus* (Camphor Bush) and *Grewia flava* (Wild Raisin). If the development is approved, it is unlikely that there will be loss of any particular ecosystem or corridor of special conservation concern according to the planned footprint. There appears to be no loss of any particular sensitive species, if the site is developed according to the proposed footprint. Mitigation measures for *Boophone disticha* (Declining) and *Acacia erioloba* (Declining, Protected Tree) apply.

The following potential impacts and mitigation measures with a view to the proposed developments apply:

6.1 Anticipated risks or impacts to the loss of habitat

The following impacts on the loss of habitat apply at the site.

Potential impacts on the available habitat will be of local extent, of permanent duration, of medium intensity and high probability. The significance of loss of habitat is expected to be moderate without mitigation and moderate-low with mitigation.

Impact summary matrix:

Phase	Significance of Impact				
	None	Low	Moderate	High	With mitigation
Operational			X		Moderate-low

Mitigation measures:

- Exotic and invasive plant species should not be allowed to establish, if the development is approved, especially an alien invasive tree species such as *Prosopis*.
- If the development is approved, exclusion of the small restricted pan depression at the site should be upheld and a buffer zone of 32 m applies.

6.2 Anticipated risks or impacts to the loss of sensitive species

Sensitive species are regarded here as those listed in section 5 and constitutes the flora and fauna that are threatened or of other particular high conservation importance. The presence or not of all the species listed in the tables were investigated during the survey. None of the threatened and near-threatened plant species are likely to occur on the site. Apart from one Declining plant species and a Protected Tree species (also listed as Declining), none of the other plant species of particular conservation priority are likely to occur on the footprint proposed for development.

A Protected Tree species, *Acacia erioloba* (also listed as Declining) is found at the site. Protected Tree species under the National Forests Act No. 84 of 1998 are listed in Table 4.9. In terms of a part of section 51(1) of Act No. 84 of 1998, no person may cut, disturb, damage or destroy any protected tree or possess, collect, remove, transport, export, purchase, sell, donate or in any other manner acquire or dispose of any protected tree, except under a license granted by the Minister.

Recommendation for *Vachellia erioloba*:

Vachellia erioloba individuals at the site are not particularly large and are not part of a camel thorn forest of note. It is recommended that a permit at the relevant authorities should be applied for in

case of any damage or removal of individual trees and that new *Vachellia erioloba* trees could be planted on site outside the present footprint.

Mitigation for *Boophone disticha*:

If the development is approved individuals of the Declining plant species *Boophone disticha* need to be relocated to a suitable site nearby before the construction phase. *Boophone disticha* (Poison Bulb) contains highly poisonous substances and the translocation operation should be done with necessary care.

6.3 Anticipated risks or impacts to habitat connectivity and open space

Potential impacts on connectivity will be of local extent, of permanent duration, of medium intensity and high probability. The significance of the impacts on loss of connectivity is expected to be low without mitigation and low with mitigation.

Impact summary matrix: habitat connectivity

Phase	Significance of Impact				
	None	Low	Moderate	High	With mitigation
Construction		X			Low
Operational		X			Low

Mitigation measures:

- If the development is approved, establishment of exotic and invasive plant species should be avoided and where these have been found at the site continuous eradication should take place.
- Alien invasive species could compromise conservation corridors and buffers. It is in particular declared alien invasive species such as *Prosopis glandulosa* (Honey Mesquite) that should not be allowed to establish.
- If the development is approved, exclusion of the small restricted pan depression at the site should be upheld and a buffer zone of 32 m applies, this pan could be an important stepping stone of conservation corridors in the larger area.

6.4 Anticipated risks or impacts associated with construction activities

Overall construction activities associated with the development if approved will be of local extent, of medium duration, of medium intensity and high probability. During the construction phase, the significance of the impacts associated with the construction phase is likely to be moderate without and low with mitigation.

Impact summary matrix:

Phase	Significance of Impact				
	None	Low	Moderate	High	With mitigation
Construction			X		Low
Operational			X		Low

Mitigation measures:

- If the development is approved, contractors must ensure that no animal species are disturbed, trapped, hunted or killed during the construction phase.
- Rubble or waste that could accompany the construction effort, if the development is approved, should be removed during and after construction.
- If the development is approved, measures should be taken to avoid any spills and infiltration of petroleum fuels or any chemical pollutants into the soil during construction phase.
- If the development is approved, the small restricted pan depression at the site should be fenced off during the construction phase, with appropriate tape and any activities at this very small depression avoided.

6.5 Cumulative impacts

Cumulative impacts could arise as other similar projects are constructed in the area.

According to the Energy Blog's database only one other solar PV plant has been granted preferred bidders status within close proximity to the proposed Khubu PV plant:

- *Waterloo Solar Park with a capacity of 75MW near Vryburg, North West Province (Approvals, planning and financing phase) – refer to Figure 10.*

According to the Department's database numerous other solar plants have been proposed in relative close proximity to the proposed activity, namely:

- The proposed Carocraft Solar Park near Vryburg, North West Province (14/12/16/3/3/2/374);*
- Construction of the 75MW Photovoltaic facility and associated infrastructure in Naledi (14/12/16/3/3/2/390).*
- The proposed Tiger Kloof Solar Photovoltaic energy facility near Vryburg, North West Province (14/12/16/3/3/2/535).*
- The proposed Keren Energy Bosh Pan Solar Plant, Northern Cape Province (14/12/16/3/3/1/563);*
- The proposed renewable energy generation project. Carocraft Solar Park in North West Province (14/12/16/3/3/2/699);*
- The proposed Renewable Energy Generation Project rem farm Elda, North West (14/12/16/3/3/2/750);*
- The proposed Renewable Energy Project on Farm Doornbult 29 and Doornbult 33, North West (14/12/16/3/3/2/751);*

Environamics and other environmental consultants are also in the process of applying for Environmental Authorisation for other PV projects in the area, namely:

- The proposed Protea Solar Power Plant near Vryburg, North West Province.*
- The proposed Gamma Solar Power Plant near Vryburg, North West Province.*
- The proposed Alpha Solar Power Plant near Vryburg, North West Province.*
- The proposed Meerkat Solar Power Plant near Vryburg, North West Province.*
- The proposed Sonbesie Solar Power Plant near Vryburg, North West Province.*
- Three PV Solar Energy facilities on the farm Klondike - AMDA Developments*

The potential for cumulative impacts therefore exists. In the Naledi Local Municipality a number of solar power plants are planned and their cumulative impacts to ecosystem and biodiversity need consideration. Several characteristics and development strategies of utility-scale solar energy systems have low environmental impacts relative to other energy systems, including other renewables (Hernandez *et al.*, 2015) so that their cumulative effects may be relatively lower.

Cumulative effects of solar power plants to habitat loss and fragmentation should receive much of the attention. Habitat loss and fragmentation are known to be the main threats to biodiversity (Fahrig, 2003; Wilcove *et al.*, 1998; IUCN, 2004; Primack, 2006). Because some fragmentation will take place if the developments are approved the focus of assessing cumulative effects of solar power plants could be on how the different projects allow for enough corridors and linkages in between the locations of solar power plants to enhance connectivity of biodiversity. An outline of possible cumulative impacts follows.

Possible cumulative effects and mitigation/ avoidance thereof:

1) Cumulative impacts on unique or sensitive habitats:

At the study area cumulative effects on the loss of sensitive habitats are kept to a minimum because such habitats are avoided at large.

2) Cumulative impacts on habitat fragmentation:

Regionally landscape fragmentation could create barriers to the movement of species and their genes (Saunders *et al.*, 1991). The answer to the width and extent of corridors depends on the conservation goal and the focal species (Samways, 2005). Corridors for mammalian species are especially important for migratory species (Mwalyosi, 1991, Pullin 2002). For an African butterfly assemblage this is about 250m when the corridor is for movement as well as being a habitat source (Pryke and Samways 2003). Hill (1995) found a figure of 200m for dung beetles in tropical Australian forest. In the agricultural context, and at least for some common insects, even small corridors can play a valuable role (Samways, 2005).

At the study area of which the site is part:

Corridors and linkages of areas with similar habitat are present in the local district where a number of solar power plants are planned. Watercourses and wetlands are avoided by the proposed footprint so that stepping stone corridors (pans) and a network of linked corridors (active channels with riparian zones) remain. No particular habitats of threatened species that are easily isolated (e.g. beetles with flightless females) are known to be impacted locally in the larger study area where a number of solar power plants are planned to be developed.

3) Emissions and pollutants into the air, soil and water:

Overall because of the restricted nature of power plants and few or no emissions and pollutants into air when operational, soil and water cumulative impacts to the environment are limited (if compared for example to emissions from fossil fuel burning). Ultimately power plants could relieve the pressures to use fossil fuels that are associated with numerous cumulative impacts and habitat losses.

Therefore in the bigger regional context, the vast tracks of relatively similar habitat in the interior are ideal for such power generating facilities. Current developments of solar power plants could therefore not be seen in the same context as many other more developed parts of the world where natural habitats are often severely fragmented. Some fragmentation of habitats will take place and the key issue would be to avoid sensitive habitats and to allow for enough corridors and linkages between habitats such as in the present proposed planned footprints.

The Environmental Impact Assessment (EIA) Report will include a detailed assessment of the potential cumulative impacts associated with the proposed development.

6 CONCLUSION

- Conspicuous and dense invasions of many parts of the site by the declared alien invasive plant species, *Prosopis glandulosa* (Honey Mesquite) are encountered at the site. A conservative estimate is that there are at the very least 2000 individuals of *Prosopis glandulosa* at the proposed footprint. *Prosopis* was introduced to South Africa and has become the second most widespread invasive alien plant taxon in the country (Shackleton et al., 2015c). These invasions have detrimental effects on biodiversity, ecosystem services and human livelihoods Shackleton et al., 2015a). In South Africa it was found that native woody species density, basal area, richness and diversity all decreased significantly as the basal area of *Prosopis* stands increased (Shackleton et al., 2015a). Invasive alien trees impact the environment and human livelihoods. Costs associated with *Prosopis* were perceived to exceed benefits, and most stakeholders wanted to see a reduction in the abundance of *Prosopis* stands (Shackleton et al., 2015b). The mean total cost for the management of *Prosopis* was US\$ 1914 year⁻¹ per farm, where costs ranged from under US\$ 10 to over US\$ 500 per ha based on invasion densities and objectives for control (Shackleton et al., 2015b).
- An opportunity presents itself, if the development is approved to eradicate extensive and dense infestations of the declared alien invasive *Prosopis glandulosa* (Honey Mosquite) at the site.
- Tree species such as *Vachellia tortilis* subsp. *heteracantha* (Umbrella Thorn), *Searsia lancea* (Karee), *Senegalia mellifera* subsp. *detinens* (Black Thorn) and *Vachellia erioloba* (Camel Thorn) have a dotted distribution in the arid savanna landscape, though at some area where secondary succession of woody elements took place more trees are present. Other parts of the site are characterised by typical patches of shrub-height *Tarchonanthus camphoratus* (Camphor Bush) and *Grewia flava* (Wild Raisin).
- The vegetation type to which the site belongs, Ghaap Plateau Vaalbosveld (SVk 7), is not listed as threatened ecosystem according to the National List of Threatened Ecosystems (2011).
- Only wetland that is present and which is excluded in the proposed footprint is a very restricted small pan depression at the northern part of the site. Otherwise no wetlands appear to be present at the proposed footprint.

- No loss of particularly sensitive or localised habitat type of particular conservation importance is anticipated if the site is developed.
- No loss of corridors or connectivity of ecosystems is anticipated if the proposed footprint is developed.
- Ecological sensitivity at the site is medium to low, the latter where high infestations of *Prosopis glandulosa* occur: There are no indications of any particular ecosystems of conservation importance, any particular conservation corridors or a significant impact on any plant, mammal, reptile, amphibian or invertebrate species of particular conservation concern if the site is developed.
- A Protected Tree species, *Vachellia erioloba* (also listed as Declining) is found at the site. Protected Tree species are listed under the National Forests Act No. 84 of 1998. In terms of a part of section 51(1) of Act No. 84 of 1998, no person may cut, disturb, damage or destroy any protected tree or possess, collect, remove, transport, export, purchase, sell, donate or in any other manner acquire or dispose of any protected tree, except under a license granted by the Minister.
- It is estimated that > 100 but less than 200 individuals of *Vachellia erioloba* are present at the site, possibly owing to secondary succession in the central-northern area at the site. Recommendation for Camel Thorn tree at the site if the development is approved. *Vachellia erioloba* individuals at the site are not particularly large (no tree taller than 10 m) and are not part of a camel thorn forest of note (Reference points: large camel thorn forest at Kathu and smaller Camel Thorn forest at Witsand visited by R.F. Terblanche during the time span of the surveys). It is recommended that a permit should be applied for at the relevant authorities in case any removal or damage of Camel Thorn trees. If *Vachellia erioloba* is impacted upon it is also recommended that new (nursery) Camel Thorn trees could be planted on site outside the present footprint.
- A Declining plant species *Boophane disticha* (Poison Bulb) is present at the site but not in any large concentrations. If the development is approved individuals of the Declining plant species *Boophane disticha* need to be relocated to a suitable site nearby before the construction phase, this could be on site outside the proposed footprint. *Boophane disticha* (Poison Bulb) contains highly poisonous substances and the translocation operation should be done with necessary care.
- Cumulative effects of solar power plants are considered with special reference to habitat loss and fragmentation. Habitat loss and fragmentation are known to be the main threats

to biodiversity (Fahrig, 2003; Wilcove *et al.*, 1998; IUCN, 2004; Primack, 2006). Because some fragmentation will take place if the developments are approved the focus of assessing cumulative effects of solar power plants could be on how the different projects allow for enough corridors and linkages in between the locations of solar power plants to enhance connectivity of biodiversity.

- There is no distinct reason why this relatively small footprint allocated for the development, in the vast countryside of the North West Province is of particular conservation concern for any threatened vertebrate species, including those that roam large areas and which may occasionally or coincidentally visit the site.
- It is unlikely that there will be a loss of any known plant, mammal, reptile, amphibian or invertebrate species that are threatened or near threatened, if the site is developed.

8 REFERENCES

- Alexander, G. & Marais, J. 2007. A guide to the reptiles of Southern Africa. Struik, Cape Town.
- Anderson, M.D. & Andersen, T.A. 2001. Too much, too quickly? Doubts about the sustainability of the camelthorn wood harvest. *African Wildlife* 55(3): 21-23.
- Apps, P. 2012. Smithers' mammals of Southern Africa 4th ed: A field guide, revised and updated by Peter Apps. Struik Nature, Cape Town.
- Armstrong, A.J. 1991. On the biology of the marsh owl, and some comparisons with the grass owl. *Honeyguide* 37:148-159.
- Barnes, K.N. ed. 2000. The Eskom Red Data Book of birds of South Africa, Lesotho and Swaziland. BirdLife South Africa, Johannesburg.
- Bates, M.F., Branch, W.R., Bauer, A.M., Burger, M., Marais, J., Alexander, G.J. & De Villiers, M.S. (eds). 2014. Atlas and Red List of the Reptiles of South Africa, Lesotho and Swaziland. *Suricata* 1. South African National Biodiversity Institute, Pretoria.
- Boon, R. 2010. *Pooley's trees of eastern South Africa: a complete guide 2nd ed*. Flora and Fauna Publications Trust, Durban.
- Branch, B. 1998. Field guide to snakes and other reptiles of southern Africa. 3rd ed. Struik, Cape Town.
- Branch, B. 2008. Tortoises, Terrapins & Turtles of Africa. Struik Nature, Cape Town.
- Branch, W.R. & Patterson, R.W. 1975. Notes on the ecology of the Giant Girdled Lizard, *Cordylus giganteus*. *Journal of Herpetology* 9(4): 364-366.

Branch, W.R., Tolley, K.A., Cunningham, M., Bauer, A.M., Alexander, G., Harrison, J.A., Turner, A.A. & Bates, M.F. eds. 2006. A plan for phylogenetic studies of southern African reptiles: proceedings of a workshop held at Kirstenbosch, February 2006. Biodiversity Series 5. South African National Biodiversity Institute, Pretoria.

Bronner, G. 2011. *Mammals*. In: Picker, M. & Griffiths, C. 2011. *Alien & Invasive animals: a South African perspective*. Struik Nature, Cape Town, p 22-35.

Bromilow, C. 2010. Problem plants and alien weeds of South Africa. Briza Publications, Pretoria.

Carruthers, V. & Du Preez, 2011. Frogs and frogging in southern Africa 2nd ed. Struik, Cape Town.

Chittenden, H. 2007. Roberts Bird Guide. John Voelcker Book Fund, Cape Town.

Cillié, B., Oberprieler, U. & Joubert, C. 2004. Animals of Pilanesberg: an identification guide. Game Parks Publishing, Pretoria.

Cilliers, S.S., Müller, N. & Drewes, E. 2004. Overview on urban nature conservation: situation in the western-grassland biome of South Africa. *Urban forestry and urban greening* 3: 49-62.

Coetzee, N. & Monadjem, A. 2008. *Mystromys albicaudatus*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>.

Conradie, W., Du Preez, L.H., Smith, K. & Weldon, C. 2006. Field guide to the frogs and toads of the Vredefort Dome World Heritage Site. School of Environmental Sciences and Development, Potchefstroom.

Court, D. 2010. *Succulent Flora of Southern Africa*. Struik Nature, Cape Town.

Crouch, N.R., Klopper, R.R., Burrows, J.E. & Burrows, S.M. 2011. Ferns of Southern Africa: a comprehensive guide. Struik Nature, Cape Town.

Del Hoyo, J., Elliot, J. & Sargatal, J. 1992. Handbook of the birds of the world, Vol. 1. Lynx Editions, Barcelona.

Deuschländer, M.S. & Bredenkamp, C.J. 1999. Importance of vegetation analysis in the conservation management of the endangered butterfly *Aloeides dentatis* subsp. *dentatis* (Swierstra) (Lepidoptera: Lycaenidae). *Koedoe* 42(2): 1-12.

Dippenaar-Schoeman, A.S. 2002. Baboon and trapdoor spiders in southern Africa: an identification manual. Plant Protection Research Institute Handbook No. 13. Agricultural Research Council, Pretoria.

Dippenaar-Schoeman, A.S. & Jocqué, R. 1997. African spiders: an identification manual. Plant Protection Research Institute Handbook No. 9. Agricultural Research Council, Pretoria.

Drinkwater, T.W., Bate, R. & Du Toit, H.A. 1998. A field guide for identification of maize pests in South Africa. Agricultural Research Council: Grain-crops Institute, Potchefstroom.

Du Preez, L.H. 1996. Field guide and key to the frogs and toads of the Free State. Department of Zoology and Entomology, University of the Orange Free State, Bloemfontein.

Du Preez, L.H. & Carruthers, V. 2009. A complete guide to the frogs of southern Africa. Struik Nature, Cape Town. CD with calls included.

Edge, D.A. 2002. Some ecological factors influencing the breeding success of the Brenton Blue butterfly, *Orachrysops niobe* (Trimen) (Lepidoptera: Lycaenidae). *Koedoe*, 45(2): 19-34.

Edge, D.A. 2005. Ecological factors influencing the survival of the Brenton Blue butterfly, *Orachrysops niobe* (Trimen) (Lepidoptera: Lycaenidae). North-West University, Potchefstroom, South Africa (Thesis - D.Phil.).

Edge, D.A., Cilliers, S.S. & Terblanche, R.F. 2008. Vegetation associated with the occurrence of the Brenton blue butterfly. *South African Journal of Science* 104: 505 - 510.

Fahrig, L. 2003. Effects of habitat fragmentation on biodiversity. *Annual review of Ecology, Evolution and Systematics*, 34: 487 -515.

Ferguson-Lees, J. & Christie, D.A. 2001. *Raptors of the world*. Christopher Helm, London.

Filmer, M.R. 1991. Southern African spiders: an identification guide. Struik, Cape Town.

Gardiner, A.J. & Terblanche, R.F. 2010. Taxonomy, biology, biogeography, evolution and conservation of the genus *Erikssonia* Trimen (Lepidoptera: Lycaenidae). *African Entomology* 18(1): 171 – 191.

Germishuizen, G. 2003. Illustrated guide to the wildflowers of northern South Africa. Briza, Pretoria.

Germishuizen, G., Meyer, N.L. & Steenkamp (eds) 2006. A checklist of South African plants. Southern African Botanical Diversity Network Report No. 41. SABONET, Pretoria.

Goldblatt, P. 1986. The Moraeas of Southern Africa. *Annals of Kirstenbosch Botanic Gardens*, Volume 14. National Botanic Gardens, Cape Town.

Goldblatt, P. & Manning, J. 1998. *Gladiolus in Southern Africa*.

Henderson, L. 2001. *Alien weeds and alien invasive plants: a complete guide to the declared weeds and invaders in South Africa*. Plant Protection Research Institute Handbook No. 12. ARC: Plant Protection Research Institute, Pretoria.

Henderson, L. & Cilliers, C.J. 2002. *Invasive aquatic plants: a guide to the identification of the most important and potentially dangerous invasive aquatic and wetland plants in South Africa*. Plant Protection Research Handbook No. 16. Agricultural Research Council, Pretoria.

Henning, G.A. & Roos, P.S. 2001. Threatened butterflies of South African wetlands. *Metamorphosis* 12(1): 26-33.

Henning, G.A., Terblanche, R.F. & Ball, J.B. (eds) 2009. South African Red Data Book: butterflies. *SANBI Biodiversity Series No 13*. South African National Biodiversity Institute, Pretoria.

Henning, S.F. 1983. Biological groups within the Lycaenidae (Lepidoptera). *Journal of the Entomological Society of Southern Africa* 46(1): 65-85.

Henning, S.F. 1987. Outline of Lepidoptera conservation with special reference to ant associated Lycaenidae. *Proceedings of the first Lepidoptera conservation Symposium, Roodepoort: Lepidopterists' Society of southern Africa: 5-7*.

Henning, S.F. & Henning, G.A. 1989. South African Red Data Book: butterflies. *South African National Scientific Programmes Report No. 158*. CSIR, Pretoria.

Herman, P.P.J. 2002. Revision of the *Tarchonanthus camphoratus* complex (Asteraceae-Tarchonantheae) in southern Africa. *Bothalia* 32,1: 21-28.

Hernandez, R.R., Easter, S.B., Murphy-Mariscal, M.L., Maestre, F.T., Tavassoli, M., Allen, E.B., Barrows, C.W., Belnap, J., Ochoa-Hueso, R., Ravi, S. & Allen, M.F. 2014. Environmental impacts of utility-scale Solar Energy. *Renewable and Sustainable Energy Reviews*, 29: 766-779.

Hill, C.J. 1995. Conservation corridors and rainforest insects. (In Watt, A.D., Stork, N.E. & Hunter, M.D. (eds.), *Forests and Insects*. Chapman & Hall, London. p. 381-393.)

Hockey, P. 2011. *Birds*. In: Picker, M. & Griffiths, C. 2011. *Alien & Invasive animals: a South African perspective*. Struik Nature, Cape Town, p 36-44.

Hockey, P.A.R., Dean, W.J.R. & Ryan, P.G. (eds.). 2005. *Roberts Birds of Southern Africa*. John Voelcker Bird Book Fund, Cape Town.

Holm, E. & Marais, E. 1992. *Fruit chafers of southern Africa*. Ekogilde, Hartebeespoort.

IUCN. 2001. *IUCN Red List Categories and Criteria: Version 3.1*. IUCN Species Survival Commission. IUCN, Gland, Switzerland and Cambridge, UK.

IUCN. 2012. IUCN Red list of Threatened Species. Version 2012.1)

Jacobsen, W.B.G. 1983. The ferns and fern allies of Southern Africa. Butterworths, Durban.

Kemper, N.P. 2001. RVI: Riparian Vegetation Index, final report, WRC Report No. 850/3/1. Institute for Water Research, Pretoria.

Kok, J.C. 1998. Vrystaatse bome, struik en klimplante Kontak-uitgewers, Pretoria.

Kudrna, O. 1995. Conservation of butterflies in central Europe. (*In* Pullin, A. S. ed. Ecology and conservation of butterflies. Chapman & Hall, London. p. 248-257.).

Kyalangalilwa, B., Boatwright, J.S., Daru, B.H., Maurin, O. & van der Bank, M. 2013. Phylogenetic position and revised classification of *Acacia* s.l. (Fabaceae: Mimosoideae) in Africa, including new combinations in *Vachellia* and *Senegalia*. *Botanical Journal of the Linnean Society* 172: 500–523.

Larsen, T.B. 1995. Butterfly biodiversity and conservation in the Afrotropical region. (*In* Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 290-303.)

Liebenberg, L. 1990. A field guide to the animal tracks of Southern Africa. David Philip Publishers, Cape Town.

Leeming, J. 2003. Scorpions of southern Africa. Struik, Cape Town.

Leroy, A. & Leroy, J. 2003. Spiders of southern Africa. Struik, Cape Town.

Louw, W.J. 1951. *An ecological account of the vegetation of the Potchefstroom Area*. Botanical Survey of South Africa, Memoir No. 24. Government Printer, Pretoria.

Low, A.B. & Rebelo, A.G. (Eds.) 1996. Vegetation of South Africa, Lesotho and Swaziland. Department of Environmental Affairs and Tourism, Pretoria.

Lubke, R.A., Hoare, D., Victor, J. & Ketelaar, R. 2003. The vegetation of the habitat of the Brenton

Blue Butterfly, *Orachrysops niobe* (Trimen), in the Western Cape, South Africa. *South African Journal of Science* 99: 201-206.

Manning, J. 2003. Photographic guide to the wild flowers of South Africa. Briza, Pretoria.

Manning, J. 2009. Field guide to the wild flowers of South Africa. Struik, Cape Town.

McMurtry, D., Grobler, L., Grobler, J. & Burns, S. 2008. Field guide to the orchids of northern South Africa and Swaziland. Umdaus Press, Hatfield.

Mecenero, S., Ball, J.B., Edge, D.A., Hamer, M.L., Henning, G.A., Krüger, M, Pringle, E.L., Terblanche, R.F. & Williams, M.C. 2013. *Conservation Assessment of Butterflies of South Africa, Lesotho and Swaziland: Red List and Atlas*. Safronics, Johannesburg & Animal Demography Unit, Cape Town.

Minter, L.R., Burger, M., Harrison, J.A., Braack, H.H., Bishop, P.J. & Kloepfer, D. eds. 2004. Atlas and Red Data Book of the Frogs of South Africa, Lesotho and Swaziland. SI/MAB series 9, Smithsonian Institution, Washington DC.

Mucina, L. & Rutherford, M.C. eds. 2006. The vegetation of South Africa, Lesotho and Swaziland. *Strelitzia* 19. Pretoria: South African National Biodiversity Institute.

Mucina, L., Rutherford, M.C., and Powrie, L.W. eds. 2005. Vegetation map of South Africa, Lesotho and Swaziland, 1:1 000 000 scale sheet maps. Pretoria: South African National Biodiversity Institute.

Munguira, M.L. 1995. Conservation of butterfly habitats and diversity in European Mediterranean countries. (In Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 277- 289.)

New, T.R. 1993. ed. Conservation biology of *Lycaenidae* (butterflies). *Occasional paper of the IUCN Species Survival Commission* No. 8.

New, T.R. 1995. Butterfly conservation in Australasia – an emerging awareness and an increasing need. (*In* Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 304 – 315.)

Oates, M.R. 1995. Butterfly conservation within the management of grassland habitats. (*In* Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. (p. 98-112.)

Opler, P.A. 1995. Conservation and management of butterfly diversity in North America. (*In* Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 316-324.)

Peacock, F. 2006. Pipits of Southern Africa. Published by the author, Pretoria. www.pipits.co.za.

Pfab, M.F. 2002. Priority ranking scheme for Red Data plants in Gauteng, South Africa. *South African Journal of Botany* (68): 299-303.

Pfab, M.F. & Victor, J.E. 2002. Threatened plants of Gauteng, South Africa. *South African Journal of Botany* (68): 370-375.

Picker, M. & Griffiths, C. 2011. Alien & Invasive animals: a South African perspective. Struik Nature, Cape Town.

Picker, M., Griffiths, C. & Weaving, A. 2004. Field guide to insects of South Africa. 2nd ed. Cape Town: Struik.

Pooley, E. 1998. A field guide to wild flowers of KwaZulu-Natal and the eastern region. Natal Flora Publications Trust, Durban.

Pringle, E.L., Henning, G.A. & Ball, J.B. eds. 1994. Pennington's Butterflies of Southern Africa. Struik Winchester, Cape Town.

Pryke, S.R. & Samways, M.J. 2001. Width of grassland linkages for the conservation of butterflies in South African afforested areas. *Biological Conservation* 101: 85-96.

Pullin, A.S. ed. 1995. Ecology and conservation of butterflies. Chapman & Hall, London.

Rautenbach, I.L. 1982. The mammals of the Transvaal. Ecoplan monograph 1: 1-211.

Retief, E. & Herman, P.P.J. 1997. Plants of the northern provinces of South Africa: keys and diagnostic characteristics. *Strelitzia* 6. National Botanical Institute, Pretoria.

Rutherford, M.C. & Westfall, R.H. 1994. Biomes of southern Africa: An objective categorisation, 2nd ed. *Memoirs of the Botanical Survey of South Africa*, Vol. 63, pp. 1-94. National Botanical Institute, Pretoria.

Ryan, P. 2001. *Practical Birding: A guide to birdwatching in southern Africa*. Struik, Cape Town.

Samways, M.J. 2005. *Insect diversity conservation*. Cambridge University Press, Cambridge.

Saunders, D.A., Hobbs, R.J. & Margules, C.R. 1991. Biological consequences of ecosystem fragmentation: A review. *Conservation Biology* 5(1): 18-32.

Shackleton, R.T., le Maitre, D.C., van Wilgen, B.W. & Richardson, D.M. 2015a. The impact of invasive alien *Prosopis* species (mesquite) on native plants in different environments in South Africa. *South African Journal of Botany*, 97: 25-31.

Shackleton, R.T., le Maitre, D.C. & Richardson, D.M. 2015b. Stakeholder perceptions and practices regarding *Prosopis* (mesquite) invasions and management in South Africa. *Ambio: A Journal of the Human Environment*, 44: 569-581.

Shackleton, R.T., le Maitre, D.C. & Richardson, D.M. 2015c. *Prosopis* invasions in South Africa: Population structures and impacts on native tree population stability. *Journal of Arid Environments*, 114: 70 – 78.

Skelton, P. 2001. *A complete guide to the freshwater fishes of Southern Africa*. Struik, Cape Town.

Skelton, P. & Weyl, O. 2011. *Fishes*. In: Picker, M. & Griffiths, C. 2011. *Alien & Invasive animals: a South African perspective*. Struik Nature, Cape Town, p 36-44.

Skinner, J.D. & Chimimba, C.T. 2005. *The mammals of the southern African subregion*. Cambridge University Press, Cape Town.

Sliwa, A. 2008. *Felis nigripes*. In: IUCN 2012. IUCN Red List of Threatened Species.

Smit, N. 2008. *Field guide to the Acacias of South Africa*. Briza, Pretoria.

Smithers, R.H.N. 1986. *South African Red Data Book: Terrestrial mammals*. *South African National Scientific Programmes Report No. 125*. CSIR, Pretoria.

South Africa. 2004. *National Environmental Management: Biodiversity Act No. 10 of 2004*. Government Printer, Pretoria.

Stuart, C. & Stuart, T. 2006. *Field guide to the larger mammals of Africa* 3rd ed. Struik Nature, Cape Town.

Stuart, C. & Stuart, T. 2013. *A field guide to the tracks and signs of Southern, Central and East African wildlife* 4th ed. Struik Nature, Cape Town.

Tarboton, W. & Erasmus, R. 1998. *Owls and owling in southern Africa*. Struik, Cape Town.

Taylor, J.C., Janse Van Vuuren, M.S. & Pieterse, A.J.H. 2007. The application and testing of diatom-based indices in the Vaal and Wilge Rivers, South Africa. *Water SA* 33(1): 51-59.

Terblanche, R.F. & Edge, D.A. 2007. The first record of an *Orachrysops* in Gauteng. *Metamorphosis* 18(4): 131-141.

Terblanche, R.F. 1994. The little hairtail from the Kalahari. *Metamorphosis* 5(4): 173-174.

Terblanche, R.F., Morgenthal, T.L. & Cilliers, S.S. 2003. The vegetation of three localities of the threatened butterfly species *Chrysoritis aureus* (Lepidoptera: Lycaenidae). *Koedoe* 46(1): 73-90.

Terblanche, R.F. & Taylor, J.C. 2000. Notes on the butterflies of Witsand – a unique terrestrial island in the Northern Cape Province, South Africa – with special reference to two RED DATA BOOK butterfly species. *Metamorphosis* 11(3): 122-131.

Terblanche, R.F. & Van Hamburg, H. 2003. The taxonomy, biogeography and conservation of the myrmecophilous *Chrysoritis* butterflies (Lepidoptera: Lycaenidae) in South Africa. *Koedoe* 46(2): 65-81.

Terblanche, R.F. & Van Hamburg, H. 2004. The application of life history information to the conservation management of *Chrysoritis* butterflies (Lepidoptera: Lycaenidae) in South Africa. *Koedoe* 47(1): 55-65.

Thomas, C.D. 1995. Ecology and conservation of butterfly metapopulations in the fragmented British landscape. (*In* Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 46-64.)

Van den Berg, J. & Drinkwater, T.W. 1998. Field guide to identification of sorghum pests in South Africa. Agricultural Research Council: Grain-crops Institute, Potchefstroom.

Van Ginkel, C.E., Glen, R.P., Gordon-Gray, K.D., Cilliers, C.J., Muasya, M. & van Deventer, P.P. 2011. Easy identification of some South African Wetland Plants. WRC Report No TT 479/10. Water Research Commission, Gezina.

Van Jaarsveld, E.J. 2006. The Southern African *Plectranthus* and the art of turning shade to glade.

Van Oudtshoorn, F. 1999. Guide to grasses of southern Africa. Briza, Pretoria.

Van Wyk, B. 2000. A photographic guide to wild flowers of South Africa. Struik, Cape Town.

Van Wyk, B. & Malan, S. 1998. Field Guide to the Wild Flowers of the Highveld. Struik, Cape Town.

Van Wyk, A.E. & Smith, G.F. 2001. Regions of floristic endemism in Southern Africa: a review with emphasis on succulents, Umdaus Press, Pretoria.

Van Wyk, B.E. & Smith, G.F. 2003. Guide to the aloes of South Africa. 2nd ed. Briza, Pretoria.

Van Wyk, B. & Van Wyk, P. 1997. Field guide to trees of southern Africa. Struik, Cape Town.

Walker, C. 1996. Signs of the Wild. 5th ed. Struik, Cape Town.

Warren, M.S. 1995. Managing local microclimates for the high brown fritillary, *Argynnis adipe*. (In Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall.)

Watt, A.D., Stork, N.E. & Hunter, M.D. (eds.), Forests and Insects. London: Chapman & Hall. (p. 381-393.)

Wilcove, D.S., Rothstein, D., Dubow, J. & Losos, E. 1998. Quantifying threats to imperiled species in the United States. *Bioscience*, 48 (8): 607 – 15.

ANNEXURE 1: Plant species list

Plant species recorded or likely to occur at the site.

Plant species are listed alphabetically under life forms that are generally recognizable.

Plant species marked with an asterisk (*) are exotic.

Sources: Germishuizen (2003), Manning (2003), Manning (2009), Van Oudtshoorn (1999), Van Wyk (2000), Van Wyk & Malan (1998), Van Wyk & Van Wyk (2013), Crouch, Klopper, Burrows & Burrows (2011), Goldblatt (1986), Goldblatt & Manning (1998), Jacobsen (1983), McMurtry, Grobler, Grobler & Burns (2008), Smit (2008), Van Ginkel *et al.* (2011), Van Jaarsveld (2006), Van Wyk & Smith (2003).

TAXON	COMMON NAMES	FAMILY
ANGIOSPERMAE: MONOCOTYLEDONS		
<i>Albuca setosa</i>	Fibrous Slime Lily	HYACINTHACEAE
<i>Antheophora pubescens</i>	Wool Grass	POACEAE
<i>Aristida adscensionis</i>	Annual Three-awn	POACEAE
<i>Aristida congesta</i> subsp. <i>congesta</i>	Tassel Three-awn	POACEAE
<i>Aristida meridionalis</i>	Giant Three-awn	POACEAE
<i>Asparagus africanus</i>		ASPARAGACEAE
<i>Asparagus laricinus</i>	Common Wild Asparagus	ASPARAGACEAE
<i>Boophone disticha</i>		AMARYLLIDACEAE
<i>Bulbine frutescens</i>		ASPHODELACEAE
<i>Bulbine narcissifolia</i>		ASPHODELACEAE
<i>Cenchrus ciliaris</i>	Foxtail Buffalo Grass	POACEAE

<i>Chloris virgata</i>	Feather-top Chloris	POACEAE
<i>Chlorophytum fasciculatum</i>		ANTHERICACEAE
<i>Commelina africana</i>		COMMELINACEAE
<i>Cymbopogon pospischilii</i>	Narrow-leaved Turpentine Grass	POACEAE
<i>Cynodon dactylon</i>	Couch Grass	POACEAE
<i>Digitaria eriantha</i>	Common Finger Grass	POACEAE
<i>Eleusine coracana</i>	Goose Grass	POACEAE
<i>Elionurus muticus</i>	Wire Grass	POACEAE
<i>Enneapogon cenchroides</i>	Nine-awned Grass	POACEAE
<i>Enneapogon scoparius</i>	Bottlebrush Grass	POACEAE
<i>Eragrostis curvula</i>	Weeping Love Grass	POACEAE
<i>Eragrostis lehmanniana</i>	Lehmann's Love Grass	POACEAE
<i>Eragrostis superba</i>	Saw-toothed Love Grass	POACEAE
<i>Fingerhuthia africana</i>	Thimble Grass	POACEAE
<i>Heteropogon contortus</i>	Spear Grass	POACEAE
<i>Ledebouria marginata</i>		HYACINTHACEAE
<i>Melinis repens</i>	Natal Red Top	POACEAE
<i>Panicum coloratum</i>	Small Buffalo Grass	POACEAE
<i>Pogonarthria squarrosa</i>	Herringbone Grass	POACEAE
<i>Schmidtia pappophoroides</i>	Sand Quick	POACEAE
<i>Setaria sphacelata</i> var. <i>torta</i>	Creeping Bristle Grass	POACEAE
<i>Setaria verticillata</i>	Bur Bristle Grass	POACEAE
<i>Sporobolus fimbriatus</i>	Dropseed Grass	POACEAE
<i>Stipagrostis uniplumis</i>	Silky Bushman Grass	POACEAE
<i>Themeda triandra</i>	Red Grass	POACEAE
<i>Tragus racemosus</i>		POACEAE
<i>Trichoneura grandiglumis</i>	Small Rolling Grass	POACEAE
ANGIOSPERMS: DICOTYLEDONS		
Acacia indigenous species: see <i>Senegalia</i> and <i>Vachellia</i> below		
<i>Acrotome inflata</i>		LAMIACEAE
* <i>Agave americana</i>	American Agave	AGAVACEAE
* <i>Alternanthera pungens</i>	Duwweltjie	AMARANTHACEAE
<i>Amaranthus thunbergii</i>		AMARANTHACEAE
<i>Aptosimum procumbens</i>	Karoo Violet	SCROPHULARIACEAE
* <i>Argemone ochroleuca</i>	White-flowered Mexican poppy	PAPAVERACEAE
<i>Barleria macrostegia</i>		ACANTHACEAE

<i>Berkheya onopordifolia</i> var. <i>onopordifolia</i>		ASTERACEAE
* <i>Cereus jamacaru</i>	Queen of the Night	CACTACEAE
<i>Ceratotheca triloba</i>	Wild Foxglove	PEDALIACEAE
<i>Chamaecrista</i> species	Cassia	CAESALPINIACEAE
<i>Chamaesyce hirta</i>	Red Milkweed	EUPHORBIACEAE
* <i>Chenopodium album</i>	White Goosefoot	CHENOPODIACEAE
<i>Chrysocoma ciliata</i>	Bitterbush	ASTERACEAE
<i>Clematis brachiata</i>	Traveller's Joy	RANUNCULACEAE
<i>Cleome angustifolia</i>	Yellow Mouse-whiskers	BRASSICACEAE (or Capparaceae)
<i>Convolvulus sagittatus</i>	Wild Bindweed	CONVOLVULACEAE
* <i>Conyza bonariensis</i>	Fleabane	ASTERACEAE
<i>Deverra denudata</i> subsp. <i>aphylla</i>	Wild Celery	APIACEAE
<i>Diospyros austro-africana</i>	Jakkalsbos	EBENACEAE
<i>Diospyros lycioides</i> subsp. <i>lycioides</i>	Karoo Bluebush	EBENACEAE
<i>Ehretia alba</i>	White Puzzle Bush	BORAGINACEAE
<i>Elephantorrhiza elephantina</i>	Eland's Bean	MIMOSACEAE
* <i>Eucalyptus camaldulensis</i>	Red Gum	MYRTACEAE
<i>Felicia muricata</i>		ASTERACEAE
<i>Gazania krebsiana</i> subsp. <i>krebsiana</i>		ASTERACEAE
<i>Geigeria filifolia</i>		ASTERACEAE
<i>Geigeria ornativa</i>		ASTERACEAE
<i>Gisekia africana</i>		GISEKIACEAE
<i>Gnidia polycephala</i>		THYMELAEACEAE
<i>Gomphocarpus tomentosus</i>		APOCYNACEAE
* <i>Gomphrena celosioides</i>	Bachelor's Button	AMARANTHACEAE
<i>Grewia flava</i>	Velvet Raisin	SPARRMANNIACEAE
<i>Helichrysum argyrosphaerum</i>		ASTERACEAE
<i>Helichrysum cerastioides</i>		ASTERACEAE
<i>Helichrysum zeyheri</i>		ASTERACEAE
<i>Heliotropium ciliatum</i>		BORAGINACEAE
<i>Hermbstaedtia odorata</i>	Wild Cockscomb	AMARANTHACEAE
<i>Hertia pallens</i>		
<i>Hilliardiella oligocephala</i> (= <i>Vernonia oligocephala</i>)		ASTERACEAE
<i>Indigofera daleoides</i>		FABACEAE

<i>Ipomoea bolusiana</i>		CONVOLVULACEAE
<i>Jamesbrittenia aurantiaca</i>	Cape Saffron	SCROPHULARIACEAE
<i>Lantana rugosa</i>		VERBENACEAE
<i>Lepidium africanum</i>	Pepperweed	BRASSICACEAE
* <i>Lepidium bonariense</i>	Pepperweed	BRASSICACEAE
<i>Lippia scaberrima</i>		VERBENACEAE
<i>Lycium horridum</i>		SOLANACEAE
* <i>Malva parviflora</i>	Small Mallow	MALVACEAE
<i>Nidorella resedifolia</i>		ASTERACEAE
* <i>Opuntia ficus-indica</i>	Sweet Prickly Pear	CACTACEAE
* <i>Opuntia imbricata</i>	Sweet Prickly Pear	CACTACEAE
<i>Pentzia calcarea</i>		ASTERACEAE
<i>Pentzia globosa</i>		ASTERACEAE
<i>Persicaria</i> species		POLYGONACEAE
<i>Pollichia campestris</i>	Waxberry	ILLECEBRACEAE
* <i>Prosopis glandulosa</i>	Mesquite	FABACEAE
<i>Salvia disermas</i>	Large Blue Sage	LAMIACEAE
<i>Scabiosa columbaria</i>	Wild Scabious, Morning Bride	DIPSACACEAE
* <i>Schkuhria pinnata</i>	Dwarf Marigold	ASTERACEAE
<i>Searsia lancea</i>	Karee	ANACARDIACEAE
<i>Searsia tridactyla</i>		ANACARDIACEAE
<i>Selago densiflora</i>		SCROPHULARIACEAE
<i>Senecio coronatus</i>	Sybossie	ASTERACEAE
<i>Senecio consanguineus</i>	Starvation Senecio	ASTERACEAE
<i>Senegalia mellifera</i> subsp. <i>detinens</i>	Black Thorn	MIMOSACEAE
<i>Senna italica</i> subsp. <i>arachoides</i>	Wild Senna	CAESALPINIACEAE
<i>Sida cordifolia</i>	Heart-leaf Sida/ Flannel Weed	MALVACEAE
<i>Sida rhombifolia</i>	Arrowleaf Sida	MALVACEAE
<i>Solanum incanum</i>	Bitter Apple	SOLANACEAE
<i>Tarchonanthus camphoratus</i>	Wild Camphor Bush	ASTERACEAE
<i>Thesium</i> sp.		SANTALACEAE
<i>Tribulus terrestris</i>	Devil's Thorn	ZYGOPHYLLACEAE
<i>Vachellia erioloba</i> (= <i>Acacia erioloba</i>)	Camel Thorn	MIMOSACEAE
<i>Vachellia hebeclada</i> subsp. <i>hebeclada</i>	Candlepod Thorn	MIMOSACEAE
<i>Vachellia karroo</i>	Sweet Thorn	MIMOSACEAE
<i>Vachellia tortilis</i> subsp. <i>heteracantha</i>	Umbrella Thorn	MIMOSACEAE
<i>Vahlia capensis</i>		VAHLIACEAE

<i>Viscum rotundifolium</i>	Round-leaved Mistletoe	VISCACEAE
<i>Ziziphus mucronata</i>	Buffalo-thorn	RHAMNACEAE
<i>Ziziphus zeyheriana</i>	Dwarf Buffalo-thorn	RHAMNACEAE