


WETLAND ASSESSMENT

**Portion 3 (Portion of Portion 2) of the Farm Vyflings
Pan 598, IN Registration Division, Province of the
North-West, measuring 428.2660 (four hundred and
twenty eight comma two six six zero) hectares, Title
Deed No.: 134/2002**



White-faced Ducks, *Dendrocygna viduata*, a widespread and highly nomadic bird, in flight after being flushed from surface water at wetland depression outside and east of the proposed footprint in the study area.

Photo: February 2016, R.F. Terblanche.

FEBRUARY 2016

COMPILED BY:

Reinier F. Terblanche

(M.Sc, *Cum Laude*; Pr.Sci.Nat, Reg. No. 400244/05)

TABLE OF CONTENTS

| | |
|---|-----------|
| 1. INTRODUCTION | 3 |
| 2. STUDY AREA | 6 |
| 3. METHODS | 9 |
| 4. RESULTS AND DISCUSSION | 17 |
| 5. ANTICIPATED RISKS AND MITIGATIONS TO THE WETLANDS OUTSIDE THE PROPOSED FOOTPRINT BUT WITHIN 500 M FROM THE EDGE OF THE PROPOSED FOOTPRINT | 32 |
| 6. CONCLUSION | 38 |
| 7. REFERENCES | 41 |

1 INTRODUCTION

A wetland assessment is required for a proposed solar power plant development at Portion 3 (Portion of Portion 2) of the Farm Vyflings Pan 598, IN Registration Division, Province of the North-West, measuring 428.2660 (four hundred and twenty eight comma two six six zero) hectares, Title Deed No.: 134/2002, 27 km west-northwest of Vryburg in the North West Province (elsewhere referred to as the site), and if wetlands are present an assessment of these wetlands will take place. Such an assessment would then focus on the hydro-geomorphic setting, an estimate of the properties of the wetlands, an assessment of the functional aspects of wetlands and an impact assessment to wetlands, should the development be approved.

1.1 Wetlands in South Africa

Wetlands are defined by the National Water Act (Act 36 of 1998) as:

“land which is transitional between terrestrial and aquatic ecosystems where the water table is usually at or near the surface, or the land is periodically covered with shallow water, and which land in normal circumstances supports or would support vegetation typically adapted to life in saturated soil”.

According to *A practical field procedure for identification and delineation of wetlands and riparian areas* (DWAF 2005) wetlands must have one or more of the following attributes:

- Wetland (hydromorphic) soils that display characteristics resulting from prolonged saturation
- The presence, at least occasionally, of water loving plants (hydrophytes)
- A high water table that results in saturation at or near the surface, leading to anaerobic conditions developing in the top 50cm of the soil

Wetlands, according to the definition of DWAF (2005) are at the interface of aquatic systems and the terrestrial environment. As such the characteristics of the surface water or near surface water in space and time at this interface between the terrestrial and aquatic environment are fundamental to understand the functioning of a particular wetland. At the higher elevations of South Africa surface water at wetlands are characterised by considerable contrasts between seasons and periodic precipitation events. Generally accepted definitions of wetlands which focus on the wetland attributes of soil and vegetation are therefore useful because of its consistency despite seasonal fluctuations.

The Classification System for Wetlands and other Aquatic Ecosystems in South Africa (Ollis *et al.*, 2013) includes wetland ecosystems defined by the National Water Act (Act 36 of 1998) as well as those “wetland systems” defined in the Ramsar Convention. The broader definition of wetlands, according to the Ramsar Convention is that wetlands are areas of marsh, fen, peatland or water, whether natural or artificial, permanent or temporary, with water that is static or flowing, fresh, brackish or salt, including areas of marine water to the depth of which at low tide does not exceed six metres (cited by Ramsar Convention Secretariat 2011). This Ramsar definition of “wetlands” overlaps broadly with the definition of aquatic systems according to the South African system of classifying wetlands and other aquatic ecosystems. In South Africa an aquatic ecosystem is an ecosystem that is permanently or periodically inundated by flowing or standing water, or which has soils that are permanently or periodically saturated within 0.5 m of the soil surface (Ollis *et al.*, 2013). Therefore an important consideration of the Classification System for Wetlands and other Aquatic Ecosystems in South Africa (Ollis *et al.*, 2013) is that a wetland (narrow definition according to water act and not Ramsar definition) is taken to be a unique type of aquatic system.

1.2 Importance of wetlands

The importance of wetlands for human well-being and the conservation of biodiversity are recognised world-wide. Ecosystem services which directly or indirectly benefit human well-being are of particular importance when wetlands are considered. Wetlands play a major role to enhance supporting services such as nutrient cycling and primary production, which in turn is the basis for other ecosystem services. Wetlands are very important to regulating services such as maintaining water flow and water quality by processing water and regulating water run-off, provisioning services such as providing freshwater, cultural services such as appreciating the landscape and biodiversity. Overall wetlands play a major role in the sustainability of land use from socio-economic and biodiversity conservation perspectives. The setting and function of wetlands at each site should therefore be evaluated to inform land use management.

Wetland vegetation is of significant importance for wetlands to play a role in valuable ecosystem services. Vegetation plays an important role in natural wetland ecosystems. It holds soil together and slows down the flow of water, reducing the risk of erosion and promoting sediment deposition. Plants are the source of organic material in wetland soils, and form the organic soil in peat wetlands.

Vegetation also has an impact on the quality of surface and subsurface water as it (1) provides organic soil matter required by microbes in order to assimilate nutrients and toxicants (2) provides habitat for the microbes in the soil immediately surrounding the roots, and (3) contributes through direct uptake of nutrients and toxicants and incorporation of these into plant tissues (Sieben *et al.* 2009).

1.3 Aims and objectives of the survey

A survey consisting of three visits to investigate key elements of habitats on the site, relevant to the conservation of wetlands are conducted. The importance and significance of the site with special emphasis on the current status of biodiversity and ecological services of the wetland are evaluated. Literature investigations are integrated with field observations to identify potential ecological impacts that could occur as a result of the development and to make recommendations to reduce or minimise impacts, should the development be approved.

The objectives of the wetland habitat assessment are to provide:

- An indication of the existence of wetlands at the site and if so:
- An identification of major aspects of the hydro-geomorphic setting and terrain unit at which the wetland occur;
- An estimate of the size and roughness of the wetland
- An indication of the hydric soils at the site;
- An indication of erodability;
- An indication of the presence or absence of peat at the site;
- An outline of hydrological drivers that support the existence and character of the wetland;
- An assessment of the possible presence or absence of threatened or localised plant species, vertebrates and invertebrates of the region, at the site;
- A description of the functions provided by the wetland at the site;
- An interpretation of the priority of the wetland for local communities in the area;
- An interpretation of the priority of the wetland to biodiversity at the site;

2 STUDY AREA

The study area is at Portion 3 (Portion of Portion 2) of the Farm Vyflings Pan 598, IN Registration Division, Province of the North-West, measuring 428.2660 (four hundred and twenty eight comma two six six zero) hectares, Title Deed No.: 134/2002, 27 km west-northwest of Vryburg in the North West Province (elsewhere referred to as the site).




Figure 1 Location of the site in larger area.

Grid references and altitudes were taken at site with a GPS Garmin E-trex 20 © instrument.
Map information were analysed and depicted on Google images with the aid of Google Earth Pro (US Dept. of State Geographer, MapLink/ Tele Atlas, Google, 2015, licenced software bought by the author).

The site is situated at the Savanna Biome which is represented by the Ghaap Plateau Vaalbosveld vegetation type (Mucina & Rutherford 2006). A brief overview of SVk 7, the Ghaap Plateau Vaalbosveld vegetation type in which the site is located, follows:

Distribution: In South Africa the Ghaap Plateau Vaalbosveld is found in the Northern Cape and North-West Provinces: Flat plateau from around Campbell in the south, east of Danielskuil through Reivilo to around Vryburg in the north. Altitude at the Ghaap Plateau Vaalbosveld is 1100 – 1500 m.

Vegetation and landscape features: Flat plateau with well-developed shrub layer with *Tarchonanthus camphoratus* and *Acacia karroo*. Open tree layer has *Olea europaea* subsp. *africana*, *Acacia tortilis*, *Ziziphus mucronata* and *Searsia [Rhus] lancea*. *Olea europaea* subsp. *africana* is more important in the southern parts of the unit, while *Acacia tortilis*, *Acacia hebeclada* and *Acacia mellifera* are more important in the north and part of the west of the unit. Much of the south-central part of this unit has remarkably low cover of *Acacia* species for an arid savanna and is dominated by the nonthorny *Tarchonanthus camphoratus*, *Searsia lancea*, and *Olea europaea* subsp. *africana* (Mucina & Rutherford 2006).

Geology and soils: Surface limestone of Tertiary to Recent age, and dolomite and chert of the Campbell Group (Griqualand West Supergroup, Vaalian Erathem) support shallow soils (0.1-0.25 m) of Mispah and Hutton soil forms. Land types mainly Fc with some Ae and Ag (Mucina & Rutherford, 2006).

Climate: Climate is characterized by summer and autumn rainfall and very dry winters. Mean annual precipitation from about 300 mm in the southwest to about 500 mm in the northeast. Frost is frequent to very frequent in winter (Mucina & Rutherford 2006).

Important taxa of the Ghaap Plateau Vaalbosveld listed by Mucina & Rutherford (2006): Tall Tree: *Vachellia erioloba* [*Acacia erioloba*]. Small Trees: *Senegalia mellifera* subsp. *detinens* [*Acacia mellifera* subsp. *detinens*], *Searsia lancea*, *Vachellia karroo* [= *Acacia karroo*], *Vachellia tortilis* subsp. *heteracantha* [*Acacia tortilis* subsp. *heteracantha*], *Boscia albitrunca*. Tall Shrubs: *Olea europaea* subsp. *africana*, *Rhigozum trichotomum*, *Tarchonanthus camphoratus*, *Ziziphus mucronata*, *Diospyros pallens*, *Ehretia rigida* subsp. *rigida* (this species complex has been revised and the *Ehretia alba* is the species that occurs at the vegetation type, R.F. Terblanche pers. obs.),

Euclea crispa subsp. *ovata*, *Grewia flava*, *Gymnosporia buxifolia*, *Lessertia frutescens*, *Searsia tridactyla*. Low Shrubs: *Acacia hebeclada* subsp. *hebeclada*, *Aptosimum procumbens*, *Chrysocoma ciliata*, *Helichrysum zeyheri*, *Hermannia comosa*, *Lantana rugosa*, *Leucas capensis*, *Melolobium microphyllum*, *Peliostomum leucorrhizum*, *Pentzia globosa*, *Pentzia viridis*, *Zygophyllum pubescens*. Succulent Shrubs: *Hertia pallens*, *Lycium cinereum*. Semi-parasitic Shrub: *Thesium hystrix*. Woody Climber: *Asparagus africanus*. Graminoids: *Anthehora pubescens*, *Cenchrus ciliaris*, *Digitaria eriantha* subsp. *eriantha*, *Enneapogon scoparius*, *Eragrostis lehmanniana*, *Schmidtia pappophoroides*, *Themeda triandra*, *Aristida adscensionis*, *Aristida congesta*, *Aristida diffusa*, *Cymbopogon pospischilii*, *Enneapogon cenchroides*, *Enneapogon desvauxii*, *Eragrostis echinochloidea*, *Eragrostis obtusa*, *Eragrostis rigidior*, *Eragrostis superba*, *Fingerhutia africana*, *Heteropogon contortus*, *Sporobolus fimbriatus*, *Stipagrostis uniplumis*, *Tragus racemosus*. Herbs: *Barleria macrostegia*, *Geigeria filifolia*, *Geigeria ornativa*, *Gisekia africana*, *Helichrysum cerastoides*, *Heliotropium ciliatum*, *Hermbsstaedtia odorata*, *Hibiscus marlothianus*, *Hibiscus pusillus*, *Jamesbrittenia aurantiaca*, *Limeum fenestratum*, *Lippia scaberrima*, *Selago densiflora*, *Vahlia capensis* subsp. *vulgaris*. Succulent herb: *Aloe grandidentata*.

Note: Though many of the above plant species occurs at the site, not all of them necessarily occur at the site.

Ghaap Plateau Vaalbosveld is not listed as threatened according to the National List of Threatened Ecosystems (2011).

3 METHODS

A desktop study comprised not only an initial phase, but also it was used throughout the study to accommodate and integrate all the data that became available during the field observations.

Surveys of a number of study areas in the vicinity of Vryburg, including the site, were conducted from 10 – 16 November 2015, December 2015 and 23 - 31 January 2016 to note key elements of habitats on the site, relevant to the conservation of fauna and flora. Notes and experience from earlier surveys at the larger study area of the Taung-Vryburg area by R.F. Terblanche that had taken place in July 2011, November 2011, January 2012, February 2012, August 2013, December 2013, January 2014 and November 2014 were also taken into account where applicable.

Classification of any inland wetland systems that could be present at the site is according to the Classification System for Wetlands and other Aquatic Ecosystems in South Africa (Ollis *et al.*, 2013). One of the major advantages of the Classification System for South Africa (Ollis *et al.*, 2013) is that the functional aspects of wetlands are the focal point of the classification. Wetlands are very dynamic systems and their functionality weighs high against the often rapid changes in their appearance (Terblanche *In prep*). In this document the main guideline for the delineation and identification of wetlands where present is the practical field procedure for identification and delineation of wetlands by DWAF (2005).

The following sections highlight the materials and methods applicable to different aspects that were observed.

3.1 Classification of wetlands (SANBI: Ollis *et al.*, 2013)

3.1.1 System, regional setting and landscape unit (Levels 1, 2 and 3)

Three broad types of Inlands Systems are dealt with in the Classification System namely rivers, open waterbodies and wetlands. These Inland Systems are then classified according to a six-tiered structure.

At the systems level (Level 1) of wetland classification, a distinction is made between Marine, Estuarine and Inland ecosystems using the level of connectivity to the open ocean as discriminator of the biophysical character of each (Ollis *et al.*, 2013). Inland wetland systems are aquatic

ecosystems with no existing connection to the ocean (i.e. characterised by the complete absence of marine exchange and/ or tidal influence (Ollis *et al.*, 2013). In this case if any wetland is present it obviously qualifies as an Inland wetland system.

At Level 2 the regional setting is a spatial framework that is preferred by the investigator to allow for gaining an understanding of the broad ecological context within which an aquatic system occurs (Ollis *et al.*, 2013). A regional setting can be identified according to the DWA ecoregion classification of Kleynhans *et al.* (2005).

A distinction is made between four landscape units at Level 3 of the Classification System for Inland Systems on the basis of the landscape setting (i.e. topographical position) (Ollis *et al.*, 2013). Four landscape units are recognized: slope, valley floor, plain and bench.

3.1.2 *Hydrogeomorphic units (Level 4)*

Seven primary hydrogeomorphic (HGM) units are recognised for Inland Systems at Level 4A of the Classification System for Wetlands and other Aquatic Ecosystems in South Africa, on the basis of hydrology and geomorphology (Ollis *et al.*, 2013). These are a River, Channeled valley-bottom wetland, Unchannelled valley-bottom wetland, Floodplain wetland, Depression, Seep and Wetland flat.

3.1.3 *Hydrological regime (Level 5)*

While the hydrogeomorphic unit (HGM) is influenced by the source of water and how it moves into, through and out of an Inland System, the hydrological regime (as catergorised by the Classification System) describes the behaviour fo the water within the system and, for wetlands, in the underlying soil (Ollis *et al.*, 2013). Together with the hydrogeomorphology the hydrological regime are used to describe the wetland as a functional unit (Ollis *et al.*, 2013). In the case of Inland wetlands which are classified as rivers, perennality is an important characteristic to describe the hydrological regime. For Inland Systems other than rivers, five categories relating to the frequency and duration of inundation have been provided: Permanently inundated, Seasonally inundated, Intermittently inundated, Never inundated/ rarely inundated and unknown (Ollis *et al.*, 2013). Period of saturation within the upper 0.5 m of the soil is a very important discriminator that also links to the wetland

delineation system of DWAF (2005). The following categories for saturation of wetland soils are recognised: Permanently saturated, Seasonally saturated, Intermittently saturated and unknown. These categories of period of saturation correspond to the permanent, seasonal and temporary zones of wetlands respectively.

3.1.4 Wetland descriptors (Level 6)

At Level 6 several “descriptors” are included for the structural/ chemical/ biological characterisation of Inland Systems (Ollis *et al.*, 2013). These descriptors are non-hierarchical to one another and can be applied in any order depending on the purpose of a study and the availability of information. Descriptors include natural vs. artificial, salinity, substratum type, pH, geology and vegetation cover (Ollis *et al.*, 2013). Various definitions are given for the descriptors which are likely to increase the consistency and use of the system.

3.2 Delineation of wetland

Together with terrain unit, indirect indicators of prolonged saturation by water: wetland plants (hydrophytes) and wetland (hydromorphic) soils are identified and used to delineate the wetland (DWAF 2005). Three zones, which may not all three be present in all wetlands, namely the permanent zone of wetness, the seasonal zone and the temporary zone are identified. The temporary zone is the outer zone and is saturated for only a short period of the year that is sufficient, under normal circumstances, for the formation of hydromorphic soils and the growth of wetland vegetation (DWAF 2005). Hydromorphic soils must display signs of wetness within 50cm of the soil to qualify as wetland soil that can support hydrophytic vegetation. Grid references and altitudes are taken on site with a GPS Garmin E-trex 20 ® instrument. Map information are analysed and depicted on Google images with the aid of Google Earth Pro (US Dept. of State Geographer, MapLink/ Tele Atlas, Google, 2015).

A few important pointers should be kept in mind when assessing wetlands in South Africa especially for cryptic wetlands and especially for wetlands in their dry conditions. These pointers are then regarded as of particular importance to wetlands in areas with less rainfall in South Africa. No one

indicator provides adequate information about wetland presence, type, hydroperiod, biodiversity, function and principle ecological and hydrological drivers to be useful on its own – particularly with regard to actual or suspected cryptic and/or temporary wetlands (Day *et al.*, 2010). A suite of indicators is required, to build up even a conceptual understanding of wetland ecosystem structure and function (Day *et al.*, 2010). The absence of an indicator does not necessarily equate to the absence of a wetland (Day *et al.*, 2010). DWAF (2005) notes the presence of numerous mottles as indicative of seasonal saturation, while temporary or permanent saturation would both be associated with less abundant mottles. Cryptic wetlands do not usually exhibit mottling, though, often because the soils have naturally low levels of iron, so the absence of mottles does not necessarily indicate the absence of a wetland (Day *et al.*, 2010).

3.3 Vegetation at and near wetland

Though vegetation is a key component of the wetland definition in the Water Act, using vegetation as a primary indicator requires undisturbed conditions and expert knowledge (DWAF 2005). Modern wetland classification systems in South Africa therefore place more emphasis on the soil wetness indicators. It remains however, that plant assemblages undergo distinct changes in species composition from the centre of a wetland to the edge, and into adjacent terrestrial areas (DWAF 2005). This change in species composition of vegetation provides valuable clues for determining the wetland boundary and wetness zones (DWAF 2005).

Apart from botanical aspects which are integrated into the description of a wetland it is imperative to note the existence or not of threatened plant species or other plant species of conservation concern, such as near-threatened, data deficient or declining species in a wetland. Floristic composition is therefore also considered during the wetland assessment. Voucher specimens of plant species are only taken where the taxonomy is in doubt or where the plant specimens are of significant relevance for invertebrate conservation. Field guides such as those by Germishuizen (2003), Manning (2003), Manning (2009), Van Oudtshoorn (2012), Van Wyk (2000), Van Wyk & Malan (1998) and Van Wyk & Van Wyk (2013) were used to confirm the taxonomy of the species. Works on specific plant groups (often genera) such as those by Goldblatt (1986), Goldblatt & Manning (1998), Jacobsen (1983), McMurtry, Grobler, Grobler & Burns (2008), Smit (2008), Van Jaarsveld (2006) and Van Wyk & Smith (2014) were also consulted to confirm the identification of species. An important source of identifications of plant species for the wetland survey is Van Ginkel, Glen, Gordon-Gray, Cilliers, Muasya & Van Deventer (2011). In this case no plant specimens were

needed to be collected as voucher specimens or to be send to a herbarium for identification. For the most recent treatise of scientific plant names and broad distributions, Germishuizen, Meyer & Steenkamp (2006) or Raimondo *et al.* (2009) or updated lists on SANBI websites are followed to compile the lists of species.

3.4 Fauna at and near wetland

Species composition of fauna is not used in wetland characterization and assessments. However, it is important to note species that favour wetlands and especially whether threatened animal species are present at a wetland or not.

Mammals are noted as sight records by day. For the identification of species and observation of diagnostic characteristics Smithers (1986), Skinner & Chimimba (2005), Cillié, Oberprieler and Joubert (2004) and Apps (2000) are consulted. Sites are walked with the aim to cover as many habitats as possible. Signs of the presence of mammal species, such as calls of animals, animal tracks (spoor), burrows, runways, nests and faeces are recorded. Walker (1996), Stuart & Stuart (2000) and Liebenberg (1990) are consulted for additional information and for the identification of spoor and signs. Trapping is only done if necessary. Habitat characteristics are also surveyed to note potential occurrences of mammals. Many mammals can be identified from field sightings but, with a few exceptions bats, rodents and shrews can only be reliably identified in the hand, and even then some species needs examination of skulls, or even chromosomes (Apps, 2000).

Birds are noted as sight records, mainly with the aid of binoculars (10x30). Nearby bird calls of which the observer was sure of the identity were also recorded. For practical skills of noting diagnostic characteristics, the identification of species and observation techniques Ryan (2001) is followed. For information on identification, biogeography and ecology Barnes (2000), Hockey, Dean & Ryan, P.G. (2005), Cillié, Oberprieler & Joubert (2004), Tarboton & Erasmus (1998) and Chittenden (2007) are consulted. Ringing of birds falls beyond the scope of this survey. Sites are walked, covering as many habitats as possible. Signs of the presence of bird species such as spoor and nests are additionally been recorded. Habitat characteristics are surveyed to note potential occurrences of birds.

Reptiles are noted as sight records in the field. Binoculars (10x30) can also be used for identifying reptiles of which some are wary. For practical skills of noting diagnostic characteristics, the

identification of species and observation techniques, Branch (1998), Marais (2004), Alexander & Marais (2007) and Cillié, Oberprieler and Joubert (2004) are followed. Sites are walked, covering as many habitats as possible. Smaller reptiles are sometimes collected for identification, but this practice was not necessary in the case of this study. Habitat characteristics are surveyed to note potential occurrences of reptiles.

Frogs and toads are noted as sight records in the field or by their calls. For practical skills of noting diagnostic characteristics, the identification of species and observation techniques Carruthers (2001), Du Preez (1996), Conradie, Du Preez, Smith & Weldon (2006) and the recent complete guide by Du Preez & Carruthers (2009) are consulted. CD's with frog calls by Carruthers (2001) and Du Preez & Carruthers (2009) are used to identify species by their calls when applicable. Sites are walked, covering as many habitats as possible. Smaller frogs are often collected by pitfall traps put out for epigeal invertebrates (on the soil), but this practice falls beyond the scope of this survey. Habitat characteristics are also surveyed to note potential occurrences of amphibians.

Invertebrates of which enough information is available to be integrated into an assessment, such as butterflies, are recorded as sight records, photographic records or voucher specimens. Voucher specimens are mostly taken of those species of which the taxa warrant collecting due to taxonomic difficulties or in the cases where species can look similar in the veldt. Many butterflies use only one species or a limited number of plant species as host plants for their larvae. Myrmecophilous (ant-loving) butterflies such as the *Aloeides*, *Chrysoritis*, *Erikssonina*, *Lepidochrysops* and *Orachrysops* species (Lepidoptera: Lycaenidae), which live in association with a specific ant species, require a unique ecosystem for their survival (Deutschländer & Bredenkamp, 1999; Terblanche, Morgenthal & Cilliers, 2003; Edge, Cilliers & Terblanche, 2008; Gardiner & Terblanche, 2010). Known food plants of butterflies are therefore also recorded. Other invertebrate groups such as fruit chafer beetles and mygalomorph spiders are also investigated where relevant.

3.5 Present Ecological Status

Ecological status of wetlands are based on models such as the modified Habitat Integrity approach developed by Kleynhans (1996, 1999). Present ecological status PES methodology is then largely based on criteria for assessing the habitat integrity of floodplain wetlands and notes for allocating a

score to attributes and rating the confidence level associated with each score (DWAF 1999). Such criteria are selected on the assumption that anthropogenic modification can generally be regarded as the primary causes of degradation of the ecological integrity of a wetland (see DWAF 1999). This is done by using Table W4-1 given by DWAF (1999):

- Score each attribute according to the guidelines provided in the footnote.
- Calculate a mean score for Table W4-1 using the individual scores for all attributes.
- Provide a confidence rating for each score according to the guidelines provided in the footnote to indicate the areas of uncertainty in the determination.

Table W4-2 provides guidelines for the determination of the Present Ecological Status Class (PESC), based on the mean score determined for Table W4-1. If any of the attributes scores < 2 (i.e., it is considered to be seriously or critically modified) this score and not the mean should be taken into consideration. This approach is based on the assumption that extensive degradation of any of the wetland attributes may determine the Present Ecological Status Category (PESC). In any case, the mean on which the assessment of the PESC is based should be regarded as a guideline and should also be tested against the opinion of local experts (DWAF 1999).

Biological integrity is not directly estimated through this approach though in some systems or parts of systems, information on biological integrity is available. In such cases, the information on biological integrity can be used as a check of the PES Category determination. The mean is used to relate the ecological state of the wetland to a particular PES Category (Table W4-2) (DWAF 1999).

3.6 Ecological Importance and Sensitivity

The assessment of the ecological importance and sensitivity is according to DWAF (1999) which in turn is adapted from Kleynhans (1996) and Kleynhans (1999). "Ecological importance" of a water resource is an expression of its importance to the maintenance of ecological diversity and functioning on local and wider scales. "Ecological sensitivity" refers to the system's ability to resist disturbance and its capability to recover from disturbance once it has occurred. The Ecological Importance and Sensitivity (EIS) provides a guideline for determination of the Ecological Management Class (EMC) DWAF (1999).

In the method outlined here, a series of determinants for EIS according to Table W5-1 of DWAF (1999) are assessed on a scale of 0 to 4, where 0 indicates no importance and 4 indicates very high importance. The method is used as a guideline for the professional judgement of individuals familiar with an area and its wetlands. The assessors must substantiate and document their judgement as far as possible for future reference and revision (DWAF 1999).

3.7 Limitations

Wetlands are very dynamic systems and owing to time constraints a glimpse of conditions at wetlands are taken. However, the hydrogeomorphological setting, soil wetness characteristics and established vegetation constitute some longer term features of a wetland. For each site visited, it should then be emphasized that surveys can by no means result in an exhaustive list of wetland plants and animals present on the site, because of the time constraint. Surveys of a number of study areas in the vicinity of Vryburg, including the site, were conducted from 10 – 16 November 2015, December 2015 and 23 - 31 January 2016 which cover a good range of times of the year to note key elements of habitats on the site, relevant to the conservation of wetland fauna and flora. Notes and experience from earlier surveys at the larger study area of the Taung-Vryburg area by R.F. Terblanche that had taken place in July 2011, November 2011, January 2012, February 2012, August 2013, December 2013, January 2014 and November 2014 were also taken into account where applicable. Weather conditions during the surveys were favourable for recording fauna and flora. The focus of the survey remains a habitat survey that concentrates on the hydrogeomorphological, hydrological and additional descriptors to classify and assess the wetland.

4 RESULTS AND DISCUSSION

4.1 Absence of wetlands at the proposed footprint

Careful search at the proposed footprint led to the conclusion that presence of wetlands at the proposed footprint is highly unlikely.




Figure 2 Indications of the wetland depressions (pans) **MSP 1** and **MSP 2** of which the outer edges are outlined. Indication of outer edge of buffer zone of 50 m is depicted in green around the wetlands. The part of wetland MSP 1, that is south of the dirt road that runs across the wetland, has been ploughed in the past.

Orange area and outline: Proposed footprint

Blue areas and outlines: Wetlands of which the edge are within 500 m from the closest edge of the proposed footprint.

Green lines: Indication of outer edge of 50 m buffer zone.

Grid references and altitudes were taken at site with a GPS Garmin E-trex 20® instrument. Map information was analysed and depicted on Google images with the aid of Google Earth Pro (US Dept. of State Geographer, MapLink/ Tele Atlas, Google).

4.2 Assessment and classification of two wetlands outside the proposed footprint but within 500 m from the edge of the proposed footprint

There are two wetlands within 500 m of the edge of the proposed footprint, one on the western side of the proposed footprint labelled **MSP 1** and one on the eastern side of the proposed footprint labelled **MSP 2** (Figure 2).

4.2.1 Wetland depression MSP 1 on the western side of the proposed footprint

A wetland **MSP 1** of approximately 11 ha is found outside and west of the proposed footprint but within 500 m of the edge of the proposed footprint. Wetland is at a distinct basin that occurs at a plain, the latter with very gentle slopes. No conspicuous inlet or outlet could be found. This depression is probably endorheic, so that water that flows in during rainfall events probably leaves mostly through evaporation and infiltration. Surface water is probably only present following substantial rainfall in a semi-arid zone given that the mean annual rainfall in the area is 500 mm or below.

Shallow brown-greyish soils are present at the wetland. At the periphery soils become lighter and sandy. Vegetation cover is visibly poor at the bottom of the pan owing possibly to salinity but likely also to grazing. In general the pan has been disturbed in the past. A dirt road runs through the wetland and the wetland part south of the dirt road has been ploughed in the past. This wetland is dry for most of the time on its surface. Dwarf shrubs and low-growing herbs and grasses occur at the pan depression. Dwarf shrubs include *Pentzia globosa*. Herbs include *Corchorus asplenifolius*, *Tribulus terrestris* (this plant often at pan fringes even if it is a terrestrial plant elsewhere in the region) and *Falkia oblonga* at the bottom of the depression. At the western fringe of the pan a sandy patch with the herbaceous shrub *Crotalaria cf. orientalis* is present. Grasses include *Eragrostis bicolor*, *Sporobolus rangeri*, *Sporobolus coromandelianus*, *Cynodon dactylon* and *Eragrostis porosa*. In general the grass cover appears to be sparse and poor at centre of the pan.

The present ecological status (PES) of the wetland system CATEGORY C means the system is moderately modified, but with some loss of natural habitats at the site (Table 4.2 and Table 4.3). The ecological importance and sensitivity (EIS) of the wetland system is low to marginal because the wetland has been compromised by a road crossing the wetland, a part of the wetland ploughed and some possible overgrazing or trampling. Furthermore the wetland is not of distinct known ecological importance (Table 4.4 and Table 4.5). Presence or particular association of any threatened wetland plant or animal species at the wetland has not been found and is unlikely.



Photo 1 View towards the south of wetland that is west of the proposed development, MSP 1.
Photo: November 2015, R.F. Terblanche.



Photo 2 Soil in the upper 50 cm at the central parts of pan MSP 1 west of the proposed footprint in the study area.
Photo: January 2016, R.F. Terblanche.

Table 4.1 Classification and outline of characteristics of wetland **MSP 1** at the site according to the Classification System for Wetlands and other Aquatic Ecosystems in South Africa (Ollis *et al.*, 2013).

| CHARACTERISTIC TYPE WETLAND DISCRIMINATORS AND DESCRIPTORS | DESCRIPTION |
|---|---|
| System (level 1) | Inland wetland |
| Regional setting (level 2) | Southern Kalahari Ecoregion (Kleynhans <i>et al.</i> 2005) |
| Landscape unit (level 3) | Plain |
| Hydrogeomorphic unit (level 4) | Depression (Pan) |
| Hydrological regime (Level 5) | Wetland is at a distinct basin that interrupts a plain with very gentle slopes. No conspicuous inlet or outlet could be found. This depression is probably endoreic, so that water that flows in during rainfall events probably leaves mostly through evaporation and infiltration. Surface water is probably only present following substantial rainfall. |
| Additional descriptors (Levels 5,6) | <p>Shallow brown-greyish soils are present at the wetland. At the periphery soils become sandy. Vegetation cover is visibly poor at the bottom of the pan owing possibly to salinity but also likely to grazing. In general the pan has been disturbed in the past. A dirt road runs through the wetland and the wetland part south of the dirt road has been ploughed in the past.</p> <p>Dwarf shrubs and low-growing herbs and grasses occur at the pan depression. Dwarf shrubs include <i>Pentzia globosa</i>. Herbs include <i>Corchorus asplenifolius</i>, <i>Tribulus terrestris</i> (this species often at pan fringes even if it is a terrestrial plant elsewhere) and <i>Falkia oblonga</i> in some patches. At the western fringe of the pan a sandy patch with the herbaceous shrub <i>Crotalaria cf. orientalis</i> is present. Grasses include <i>Eragrostis bicolor</i>, <i>Sporobolus rangeri</i>, <i>Sporobolus coromandelianus</i>, <i>Cynodon dactylon</i> and <i>Eragrostis porosa</i>. In general the grass cover appears to be sparse and poor at centre of the pan.</p> |

Table 4.2 Scoresheet with criteria for assessing habitat integrity of wetland depression **MSP 1** according to DWAF (1999) such as adapted from Kleynhans (1996).

| Criteria and attributes | Relevance | Score | Confidence |
|-------------------------------|---|---------------------------|---------------------------|
| Hydrologic | | | |
| Flow modification | Consequence of abstraction, regulation by impoundments or increased runoff from human settlements or agricultural land. Changes in flow regime (timing, duration, frequency), volumes, velocity which affect inundation of wetland habitats resulting in floristic changes or incorrect cues to biota. Abstraction of groundwater flows to the wetland. | 2 | 4 |
| Permanent inundation | Consequence of impoundment resulting in destruction of natural wetland habitat and cues for wetland biota. | 2 | 4 |
| Water Quality | | | |
| Water quality modification | From point or diffuse sources. Measure directly by laboratory analysis or assessed indirectly from upstream agricultural activities, human settlements and industrial activities. Aggravated by volumetric decrease in flow delivered to the wetland. | 3 | 3 |
| Sediment load modification | Consequence of reduction due to entrapment by impoundments or increase due to land use practices such as overgrazing. Cause of unnatural rates of erosion, accretion or infilling of wetlands and change in habitats. | 2 | 3 |
| Hydraulic/Geomorphic | | | |
| Canalisation | Results in desiccation or changes to inundation patterns of wetland and thus changes in habitats. River diversions or drainage. | 3 | 4 |
| Topographic alteration | Consequence of infilling, ploughing, dykes, trampling, bridges, roads, railway lines and other substrate disruptive activities which reduce or change wetland habitat directly or through changes in inundation patterns. | 1 | 4 |
| Biota | | | |
| Terrestrial encroachment | Consequence of desiccation of wetland and encroachment of terrestrial plant species due to changes in hydrology or geomorphology. Change from wetland to terrestrial habitat and loss of wetland functions. | 3 | 4 |
| Indigenous vegetation removal | Direct destruction of habitat through farming activities, grazing or firewood collection affecting wildlife habitat and flow attenuation functions, organic matter inputs and increases potential for erosion. | 3 | 4 |
| Invasive plant encroachment | Affect habitat characteristics through changes in community structure and water quality changes (oxygen reduction and shading). | 4 | 4 |
| Alien fauna | Presence of alien fauna affecting faunal community structure. | 4 | 4 |
| Overutilisation of biota | Overgrazing, over-fishing etc. | 3 | 4 |
| TOTAL MEAN | | 30 x=2.7 | 42 x=3.8 |

Scoring guidelines per attribute:

natural, unmodified = 5; Largely natural = 4, Moderately modified = 3; largely modified = 2; seriously modified = 1; Critically modified = 0.

Relative confidence of score:

Very high confidence = 4; High confidence = 3; Moderate confidence = 2; Marginal/low confidence = 1.

Table 4.3 Interpretation of scores for determining present ecological status (**PES**) of the wetland depression **MSP 1** according to DWAF (1999) such as adapted from Kleynhans (1999). Present ecological status of the wetland is Category C (indicated in blue font).

| Interpretation of Mean* of Scores for all Attributes: Rating of Present Ecological Status Category (PES Category) |
|--|
| WITHIN GENERALLY ACCEPTABLE RANGE |
| CATEGORY A >4; Unmodified, or approximates natural condition. |
| CATEGORY B >3 and <=4; Largely natural with few modifications, but with some loss of natural habitats. |
| CATEGORY C >2 and <=3; moderately modified, but with some loss of natural habitats. |
| CATEGORY D =2; largely modified. A large loss of natural habitats and basic ecosystem functions has occurred. |
| OUTSIDE GENERAL ACCEPTABLE RANGE |
| CATEGORY E >0 and <2; seriously modified. The losses of natural habitats and basic ecosystem functions are extensive. |
| CATEGORY F 0; critically modified. Modifications have reached a critical level and the system has been modified completely with an almost complete loss of natural habitat. |

* If any of the attributes are rated <2, then the lowest rating for the attribute should be taken as indicative of the PES category and not the mean.

Table 4.4 Score sheet for for determining ecological importance and sensitivity for floodplains for **MSP 1** (DWAF 1999, adapted from Kleynhans 1996, 1999).

| Determinant | Score | Confidence |
|--|--------------|-------------------|
| PRIMARY DETERMINANTS | | |
| 1. Rare & Endangered Species | 0 | 3 |
| 2. Populations of Unique Species | 0 | 3 |
| 3. Species/taxon Richness | 2 | 3 |
| 4. Diversity of Habitat Types or Features | 2 | 3 |
| 5. Migration route/breeding and feeding site for wetland species | 1 | 3 |
| 6. Sensitivity to Changes in the Natural Hydrological Regime | 3 | 3 |
| 7. Sensitivity to Water Quality Changes | 2 | 3 |
| 8. Flood Storage, Energy Dissipation & Particulate/Element Removal | 3 | 3 |
| MODIFYING DETERMINANTS | | |
| 9. Protected Status | 0 | 4 |
| 10. Ecological Integrity | 2 | 4 |
| TOTAL | 15 | 32 |
| MEAN | 1.5 | 3.2 |

Score guideline Very high = 4; High = 3, Moderate = 2; Marginal/Low = 1; None = 0

Confidence rating Very high confidence = 4; High confidence = 3; Moderate confidence = 2; Marginal/low confidence = 1

Table 4.5 Ecological importance and sensitivity categories for **MSP 1**. Interpretation of median scores for biotic and habitat determinants (DWAF 1999, adapted from Kleynhans 1996, 1999).

| Ecological Importance and Sensitivity Category (EIS) | Range of Median | Recommended Ecological Management Class |
|--|-------------------------|---|
| <p><u>Very high</u> Floodplains that are considered ecologically important and sensitive on a national or even international level. The biodiversity of these floodplains is usually very sensitive to flow and habitat modifications. They play a major role in moderating the quantity and quality of water of major rivers.</p> | >3 and <=4 | A |
| <p><u>High</u> Floodplains that are considered to be ecologically important and sensitive. The biodiversity of these floodplains may be sensitive to flow and habitat modifications. They play a role in moderating the quantity and quality of water of major rivers.</p> | >2 and <=3 | B |
| <p><u>Moderate</u> Floodplains that are considered to be ecologically important and sensitive on a provincial or local scale. The biodiversity of these floodplains is not usually sensitive to flow and habitat modifications. They play a small role in moderating the quantity and quality of water of major rivers.</p> | >1 and <=2 | C |
| <p><u>Low/marginal</u> Floodplains which are not ecologically important and sensitive at any scale. The biodiversity of these floodplains is ubiquitous and not sensitive to flow and habitat modifications. They play an insignificant role in moderating the quantity and quality of water of major rivers.</p> | >0 and <=1 | D |

4.2.2 Wetland depression MSP 2 on the eastern side of the proposed footprint

A wetland **MSP 2** of approximately 32 ha is found outside and east of the proposed footprint but within 500 m of the edge of the proposed footprint. Wetland is at a distinct basin that occurs at a plain, the latter with very gentle slopes. No conspicuous inlet or outlet could be found. This depression is probably endorheic, so that water that flows in during rainfall events probably leaves mostly through evaporation and infiltration. Surface water is probably only present following substantial rainfall in a semi-arid zone given that the mean annual rainfall in the area is 500 mm or below.

Though the pan (relatively large in the area) is continuous, habitats vary from typical pan areas where the bottom of depression is flat to small areas where water gathers as pools in darker soils and even a rocky area. Vegetation at the pan responds to differences in habitat and bare areas as well as areas with denser cover of wetland vegetation are present. Hydrophytes such as *Persicaria* species are partially submerged in water following substantial rain in summer though not at all parts of the pan depression. Other plant species at the depression include grass species *Eragrostis bicolor*, *Andropogon appendiculatus*, *Sporobolus rangeri*, *Sporobolus coromandelianus* and *Sporobolus ioclados* as well as sedges *Scirpoides dioecus*, *Schoenoplectus muricinux*, *Cyperus esculentus* and the rush *Juncus rigidus*.

The present ecological status (PES) of the wetland system is CATEGORY B which means the system is largely natural with few modifications, but with some loss of natural habitats (Table 4.6 and Table 4.7). The ecological importance and sensitivity (EIS) of the wetland system is Moderate (Table 4.9 and Table 4.10) because it appears to contain significant indigenous biodiversity but it is not high because the importance to particular threatened species, flocks of migratory species of particular conservation concern or as a conservation area do not distinctly stand out.



Photo 3 Water gathered at pan MSP 2 which is outside the proposed footprint but within 500 m from the edge of the proposed footprint.

Photo: January 2016, R.F. Terblanche.



Photo 4 Hydrophyte, *Persicaria* species, submerged in shallow water at pan MSP 2 at the study area.

Photo: January 2016, R.F. Terblanche.

Table 4.6 Classification and outline of characteristics of wetland **MSP 2** at the site according to the Classification System for Wetlands and other Aquatic Ecosystems in South Africa (Ollis *et al.*, 2013).

| CHARACTERISTIC TYPE WETLAND DISCRIMINATORS AND DESCRIPTORS | DESCRIPTION |
|--|---|
| System (level 1) | Inland wetland |
| Regional setting (level 2) | Southern Kalahari Ecoregion (Kleynhans <i>et al.</i> 2005) |
| Landscape unit (level 3) | Plain |
| Hydrogeomorphic unit (level 4) | Depression (Pan) |
| Hydrological regime (Level 5) | Wetland is at a distinct basin that interrupts a plain with very gentle slopes. No conspicuous inlet or outlet could be found. This depression is probably endorheic, so that water that flows in during rainfall events probably leaves mostly through evaporation and infiltration. Surface water is probably only present following substantial rainfall. |
| Additional descriptors (Levels 5,6) | <p>Though the pan (relatively large in the area) is continuous habitats vary from typical pan areas with bottom of depression flat to small areas where water gathers as pools in darker soils. Vegetation at the pan responds to differences in habitat and bare areas as well as areas with denser cover of wetland vegetation are present.</p> <p>Hydrophytes such as <i>Persicaria</i> species are partially submerged in water following substantial rain in summer. Other plant species at the pan include grass species <i>Eragrostis bicolor</i>, <i>Andropogon appendiculatus</i>, <i>Sporobolus rangeri</i>, <i>Sporobolus coromandelianus</i> and <i>Sporobolus ioclados</i> as well as sedges <i>Scirpoides dioecus</i>, <i>Schoenoplectus muricinux</i>, <i>Cyperus esculentus</i> and the rush <i>Juncus rigidus</i>.</p> |

Table 4.7 Scoresheet with criteria for assessing habitat integrity of wetland depression **MSP 1** according to DWAF (1999) such as adapted from Kleynhans (1996).

| Criteria and attributes | Relevance | Score | Confidence |
|--------------------------------|---|---------------------|---------------------|
| Hydrologic | | | |
| Flow modification | Consequence of abstraction, regulation by impoundments or increased runoff from human settlements or agricultural land. Changes in flow regime (timing, duration, frequency), volumes, velocity which affect inundation of wetland habitats resulting in floristic changes or incorrect cues to biota. Abstraction of groundwater flows to the wetland. | 4 | 4 |
| Permanent inundation | Consequence of impoundment resulting in destruction of natural wetland habitat and cues for wetland biota. | 4 | 4 |
| Water Quality | | | |
| Water quality modification | From point or diffuse sources. Measure directly by laboratory analysis or assessed indirectly from upstream agricultural activities, human settlements and industrial activities. Aggravated by volumetric decrease in flow delivered to the wetland. | 4 | 3 |
| Sediment load modification | Consequence of reduction due to entrapment by impoundments or increase due to land use practices such as overgrazing. Cause of unnatural rates of erosion, accretion or infilling of wetlands and change in habitats. | 3 | 3 |
| Hydraulic/Geomorphic | | | |
| Canalisation | Results in desiccation or changes to inundation patterns of wetland and thus changes in habitats. River diversions or drainage. | 4 | 4 |
| Topographic alteration | Consequence of infilling, ploughing, dykes, trampling, bridges, roads, railway lines and other substrate disruptive activities which reduce or change wetland habitat directly or through changes in inundation patterns. | 4 | 4 |
| Biota | | | |
| Terrestrial encroachment | Consequence of desiccation of wetland and encroachment of terrestrial plant species due to changes in hydrology or geomorphology. Change from wetland to terrestrial habitat and loss of wetland functions. | 4 | 4 |
| Indigenous vegetation removal | Direct destruction of habitat through farming activities, grazing or firewood collection affecting wildlife habitat and flow attenuation functions, organic matter inputs and increases potential for erosion. | 3 | 4 |
| Invasive plant encroachment | Affect habitat characteristics through changes in community structure and water quality changes (oxygen reduction and shading). | 4 | 4 |
| Alien fauna | Presence of alien fauna affecting faunal community structure. | 4 | 4 |
| Overutilisation of biota | Overgrazing, over-fishing etc. | 4 | 4 |
| TOTAL MEAN | | 42 x=3.8 | 42 x=3.8 |

Scoring guidelines per attribute:

natural, unmodified = 5; Largely natural = 4, Moderately modified = 3; largely modified = 2; seriously modified = 1; Critically modified = 0.

Relative confidence of score:

Very high confidence = 4; High confidence = 3; Moderate confidence = 2; Marginal/low confidence = 1.

Table 4.8 Interpretation of scores for determining present ecological status (**PES**) of the wetland depression **MSP 2** according to DWAF (1999) such as adapted from Kleynhans (1999). Present ecological status of the wetland is Category C (indicated in blue font).

| Interpretation of Mean* of Scores for all Attributes: Rating of Present Ecological Status Category (PES Category) |
|--|
| WITHIN GENERALLY ACCEPTABLE RANGE |
| CATEGORY A >4; Unmodified, or approximates natural condition. |
| CATEGORY B >3 and <=4; Largely natural with few modifications, but with some loss of natural habitats. |
| CATEGORY C >2 and <=3; moderately modified, but with some loss of natural habitats. |
| CATEGORY D =2; largely modified. A large loss of natural habitats and basic ecosystem functions has occurred. |
| OUTSIDE GENERAL ACCEPTABLE RANGE |
| CATEGORY E >0 and <2; seriously modified. The losses of natural habitats and basic ecosystem functions are extensive. |
| CATEGORY F 0; critically modified. Modifications have reached a critical level and the system has been modified completely with an almost complete loss of natural habitat. |

* If any of the attributes are rated <2, then the lowest rating for the attribute should be taken as indicative of the PES category and not the mean.

Table 4.9 Score sheet for for determining ecological importance and sensitivity for floodplains for **MSP 2** (DWAF 1999, adapted from Kleynhans 1996, 1999).

| Determinant | Score | Confidence |
|--|--------------|-------------------|
| PRIMARY DETERMINANTS | | |
| 1. Rare & Endangered Species | 0 | 3 |
| 2. Populations of Unique Species | 0 | 3 |
| 3. Species/taxon Richness | 3 | 3 |
| 4. Diversity of Habitat Types or Features | 3 | 3 |
| 5. Migration route/breeding and feeding site for wetland species | 2 | 3 |
| 6. Sensitivity to Changes in the Natural Hydrological Regime | 3 | 3 |
| 7. Sensitivity to Water Quality Changes | 2 | 3 |
| 8. Flood Storage, Energy Dissipation & Particulate/Element Removal | 3 | 3 |
| MODIFYING DETERMINANTS | | |
| 9. Protected Status | 0 | 4 |
| 10. Ecological Integrity | 3 | 4 |
| TOTAL | 19 | 32 |
| MEAN | 1.9 | 3.2 |

Score guideline Very high = 4; High = 3, Moderate = 2; Marginal/Low = 1; None = 0

Confidence rating Very high confidence = 4; High confidence = 3; Moderate confidence = 2; Marginal/low confidence = 1

Table 4.10 Ecological importance and sensitivity categories for **MSP 2**. Interpretation of median scores for biotic and habitat determinants (DWAF 1999, adapted from Kleynhans 1996, 1999).

| Ecological Importance and Sensitivity Category (EIS) | Range of Median | Recommended Ecological Management Class |
|--|--------------------------------|---|
| <p><u>Very high</u> Floodplains that are considered ecologically important and sensitive on a national or even international level. The biodiversity of these floodplains is usually very sensitive to flow and habitat modifications. They play a major role in moderating the quantity and quality of water of major rivers.</p> | <p>>3 and <=4</p> | <p>A</p> |
| <p><u>High</u> Floodplains that are considered to be ecologically important and sensitive. The biodiversity of these floodplains may be sensitive to flow and habitat modifications. They play a role in moderating the quantity and quality of water of major rivers.</p> | <p>>2 and <=3</p> | <p>B</p> |
| <p><u>Moderate</u> Floodplains that are considered to be ecologically important and sensitive on a provincial or local scale. The biodiversity of these floodplains is not usually sensitive to flow and habitat modifications. They play a small role in moderating the quantity and quality of water of major rivers.</p> | <p>>1 and <=2</p> | <p>C</p> |
| <p><u>Low/marginal</u> Floodplains which are not ecologically important and sensitive at any scale. The biodiversity of these floodplains is ubiquitous and not sensitive to flow and habitat modifications. They play an insignificant role in moderating the quantity and quality of water of major rivers.</p> | <p>>0 and <=1</p> | <p>D</p> |

5 ANTICIPATED RISKS AND MITIGATIONS TO THE WETLANDS OUTSIDE THE PROPOSED FOOTPRINT BUT WITHIN 500 M FROM THE EDGE OF THE PROPOSED FOOTPRINT

5.1 Anticipated risks, impacts and mitigation measures of proposed development on the wetland, the latter to be excluded from the proposed footprint

There are two wetlands within 500 m of the edge of the proposed footprint, one on the western side of the proposed footprint labelled **MSP 1** (approximately 11 ha) and one on the eastern side of the proposed footprint labelled **MSP 2** (approximately 32 ha) (Figure 2). Both these wetlands occur as basins that interrupt a plain with very gentle slopes of which the soils are conspicuously sandy. No development or any clearance of vegetation is planned at the wetlands or at the buffer zone of these wetlands and therefore no direct impact is foreseen. Indirect impacts or risks of the development to the two wetlands are considered in the section below.

5.1. Surface flow

Two wetlands outside the proposed footprint are situated from 1981 – 1993 metres and the plain from 1989 to 1996 metres above sea level. Slopes approaching the wetlands are very gentle. Upper soils at the slopes of the plain are also conspicuously sandy which means the infiltration of the soil is high and the water run-off most likely relatively low. Site is also located in a relatively low rainfall area. The amount of runoff and the percentage of runoff as a proportion of rainfall tend to increase with rainfall such that while runoff in the northwest of South Africa is less than 5 mm per annum, there is a fairly systematic increase of runoff towards southwards and eastwards in South Africa (Ellery *et al.*, 2009). Because the site falls in the northwestern parts of the country with a rainfall of 500 mm or less, and the above factors of soil and very gentle slopes the water runoff is relatively very low.

It is anticipated the solar panels could lead to an increase of water runoff during rainfall events and when the solar panels are washed (once a year). This increase in water from the solar panels could however be counteracted by possibly more vegetation at the buffer zone owing to the decrease/absence in grazing activities. It should also be noted that there is no distinct inlets at the depressions at the site therefore the water runoff is a sheet flow into the depression rather than a channeled inflow. Diffusion rather than channelling of water flow towards the excluded wetland depression disperses any water run-off or impact on erosion. Furthermore, the slope at the site is very low, the

buffer zone at the site may have more vegetation owing to the planned decrease in intensity of grazing, revegetation in shaded areas of proposed solar panels is likely to be rapid, so that the buffer zone for a the wetlands on a very gentle slope could be highly effective in minimizing changes in water flow into the wetland. In addition the upper soils at the proposed footprint are visibly sandy with high infiltration that will further limit surface water runoff.

Wetlands are pan depressions which mean that the surface flow at these wetlands when inundated would be diffuse.

Maintaining water quality of the wetlands outside the proposed footprint is essential. Risks of fuel spillages and any sewage or domestic waste that could eventually reach the wetland are low because of mitigation measures to prevent these during the construction phase of the solar power plant. Any such spillages should be avoided and if accidentally taken place be remediated in an immediate time frame.

Mitigation measures:

- A buffer zone of 50 m from the outer edge of the pan depressions MSP 1 and MSP 2, west and east of the proposed footprint respectively, applies.
- Significant decrease in grazing activities to enhance the functioning of the buffer zone around the wetlands.
- Monitoring of any possible sedimentation effects or any significant increases in water runoff owing to the proposed development, if any, should take place. If any such impacts occur necessary steps should be introduced to counteract any significant changes in waterflow and erosion.
- Quick revegetation at soil surface at the solar panel areas should take place.
- Sediment capture and stormwater runoff systems should be planned carefully so that the risk of excessive water runoff towards the wetlands is low.

5.2 Soil erosion

Soil erosion problems occur when the rate of erosion is increased above natural levels (Ashman & Puri 2002). The erosion risk at any site can be determined by considering erosivity and erodability. Erosivity is related to the number of intense storms during a year and erodability depends on the the texture and organic matter content of a soil coupled with site characteristics such as slope and whether the soil is exposed or has vegetative cover (Ashman & Puri 2002). Therefore rainfall

intensity, soil characters related to erodibility, slope angle and length and vegetation cover are all primary factors that could lead to the increase of erosion. Flowing water contains energy which is linked to Mannings equation (also Gauckler-Manning Coefficient (Gauckler 1867, Chanson 2004, Russel 2009) of the velocity of water flow: $v = \frac{R^{2/3} \times s^{1/2}}{n}$ where v is the cross-sectional average velocity, R is the hydraulic radius of the channel, s is the slope of the channel and n is the Manning coefficient (unitless) of roughness. Note Manning's equation is an empirical formula and should be applied as such with care. In practice the relation of Manning's equation means that the flatter the slope, the wider and flatter the area over which the water flows and in addition increased roughness of the area over which the water flows in a wetland, the slower and more diffuse the flow of water will be. Erosion itself is among other factors dependent on the velocity and discharge of water (amount of water), as well as the erodibility of the substrate.

Rainfall is sporadic and low at the site with typical thunderstorm events of the summer rainfall interior of South Africa could lead to a rush of increase of runoff towards the wetland if substantial rain falls, which in fact is one of the reasons why the wetlands exists in a dry area. The slope at the site is low and the roughness of the area could still be high at the soil surface level, especially at the buffer zone where grazing activities would decrease. It should be noted that at the plain and proposed footprint at the site upper soils are conspicuously sandy which means that infiltration is relatively high and the runoff effects are further limited.

Mitigation measures:

- A buffer zone of 50 m from the outer edge of the wetlands west and east of the proposed footprint applies.
- Significant decrease in grazing activities to enhance the functioning of the buffer zones around the wetlands.
- Monitoring of any possible sedimentation effects or any significant increases in water runoff owing to the proposed development, if any, should take place. If any such impacts occur necessary steps should be introduced to counteract any significant changes in waterflow and erosion.
- Quick revegetation at soil surface at the solar panel areas should take place.
- Sediment capture and stormwater runoff systems should be planned carefully so that the risk of excessive water runoff towards the wetland patch remains low.

5.3 Interflow ("Subsurface stormwater flow)

Wetland is very shallow and on a plain in the landscape of an area of which the average annual rainfall is 500 mm or less. There is no distinct indication that interflow plays an important role in the maintenance of the wetlands outside the proposed footprint or that the proposed development will have a significant influence on interflow. Input of water into the small depression appears to be via sheet flow of water during rainfall events, perhaps via some seepage, but overall if surface water is present it appears to be very temporary in a basin which suggests that significant quantities of interflow do not sustain the wetlands.

Mitigation measures:

In the case of this study no risks or mitigation measures could apply to the wetlands are excluded from the proposed footprint.

5.4 Groundwater

Wetland is very shallow and on a plain in the landscape of an area of which the average annual rainfall is 500 mm or less. Input of water into the small depression appears to be via sheet flow of water during rainfall events, perhaps via some seepage, but overall if surface water is present it appears to be very temporary in a basin which suggests that significant quantities of groundwater do not sustain the wetlands. There is no distinct indication that ground water plays an important role in the maintenance of the wetlands at the site.

Mitigation measures:

In the case of this study no risks or mitigation measures could apply for groundwater at the wetlands that are outside the proposed footprint.

5.5 Flow regime

Hydrodynamics at the pan depressions are dominated by vertical water level fluctuations, being “dried out” for lengths of time during the absence of rain. It is unlikely that there would be a significant impact on the flow regime of the wetlands outside the proposed footprint owing to the counterbalances in the surface flow regime. In addition upper soils at the site are conspicuously sandy at the plains and proposed footprints next to the wetland basins, which means that infiltration of water into the soil would be relatively high and water runoff being buffered. The existence of the wetland may be slightly enhanced by possible slight increased surface flow, if at all.

Mitigation measures:

- A buffer zone of 50 m from the outer edge of the wetlands west and east of the proposed footprint applies.
- Significant decrease in grazing activities to enhance the functioning of the buffer zone around the wetlands.
- Monitoring of any possible sedimentation effects or any significant increases in water runoff owing to the proposed development, if any of these would occur, should take place. If any such impacts occur necessary steps should be introduced to counteract any significant changes in waterflow and erosion.
- If the development is approved, quick revegetation at soil surface at the solar panel areas should take place.
- Sediment capture and stormwater runoff systems should be planned carefully so that the risk of excessive water runoff towards the wetlands is limited.

5.6 Geomorphology

Proposed development implies no changes to the geomorphological setting of the wetlands west and east of the proposed footprint. No implications of mitigation measures apply.

5.7 Wetland habitat

It is highly unlikely that any loss of wetland habitat at the site would take place owing to the proposed development.

Mitigation measures:

- A buffer zone of 50 m from the outer edge of the small wetland patch at the southeastern corner of the site applies.
- Exotic and invasive plant species should not be allowed to establish, if the development is approved, especially an alien invasive tree species such as *Prosopis*.
- Monitoring of any possible sedimentation effects or any significant increases in water runoff owing to the proposed development, if any occur, should take place. If any such impacts occur necessary steps should be introduced to counteract any significant changes in waterflow and erosion.
- Fuel spillages and any sewage or domestic waste should be avoided at the proposed footprint if the development is approved so that delayed indirect effects of any such pollution to the wetlands further away are avoided.

5.8 Wetland biota

5.8.1 Wetland biota; general indigenous diversity

Wetlands west and east of the proposed footprint contain some indigenous plant and animal diversity in particular the larger wetland MSP 2 east of the proposed footprint. Standard mitigation measures apply such as prevention of the establishment of exotic vegetation and avoidance of any trapping or killing of mammalian or bird species during the construction phase.

Mitigation measures:

- Exotic and invasive plant species should not be allowed to establish, if the development is approved, especially an alien invasive tree species such as *Prosopis*.

5.8.2 Wetland plant and animal species that are threatened or of any other particular conservation concern

Presence of any threatened animal or plant species, or any other species of particular conservation concern at the site and the wetland are unlikely (see also ecological habitat survey report). No risks or mitigation measures apply for particular sensitive species. No threats to any Threatened or Near Threatened plant or animal species or any distinct impact on migratory species of particular conservation concern at the wetlands are anticipated if the development is approved.

6 CONCLUSION

There appears to be no wetlands at the proposed footprint.

There are two wetlands within 500 m of the edge of the proposed footprint, one on the western side of the proposed footprint (**MSP 1**) of approximately 11 ha and one on the eastern side of the proposed footprint (**MSP 2**) (Figure 2) of approximately 32 ha. Surface water at these pan depressions is clearly temporal and the soil surface appears dry for most of the year.

Both these wetlands contain some diversity of indigenous plants and animals of which MSP 2 appears to have more habitat diversity and subsequently probably a higher diversity. Conspicuous indigenous plant species that occur at wetland MSP 1 include *Pentzia globosa*, *Corchorus asplenifolius*, *Tribulus terrestris* (this species often at pan fringes even if it is a terrestrial plant elsewhere), *Falkia oblonga*, *Eragrostis bicolor*, *Sporobolus rangeri*, *Sporobolus coromandelianus*, *Cynodon dactylon* and *Eragrostis porosa*. Conspicuous indigenous plant species that are present at wetland MSP 2 include *Persicaria* species, *Eragrostis bicolor*, *Andropogon appendiculatus*, *Sporobolus rangeri*, *Sporobolus coromandelianus*, *Sporobolus ioclados*, *Scirpoides dioecus*, *Schoenoplectus muricinux*, *Cyperus esculentus* and *Juncus rigidus*.

The present ecological status (PES) of the wetland MSP 1 is CATEGORY C means the system is moderately modified, but with some loss of natural habitats at the site (Table 4.2 and Table 4.3). The ecological importance and sensitivity (EIS) of the wetland system is low to marginal because the wetland has been compromised by a road crossing the wetland, a part of the wetland ploughed and some possible overgrazing or trampling. Furthermore the wetland is not of distinct known ecological importance (Table 4.4 and Table 4.5).

The present ecological status (PES) of the wetland MSP 2 east of the proposed footprint is CATEGORY B which means the system is largely natural with few modifications, but with some loss of natural habitats (Table 4.6 and Table 4.7). The ecological importance and sensitivity (EIS) of the wetland system is Moderate (Table 4.9 and Table 4.10) because it appears to contain significant indigenous biodiversity but it is not high because there are no distinct indications of importance to particular threatened species, flocks of migratory species of particular conservation concern or as a conservation area.

The site is not part of a Freshwater Ecosystem Priority Area (FEPA) or wetland cluster (Nel, Driver, Strydom, Maherry, Petersen, Hill, Roux, Nienaber, Van Deventer, Swartz, & Smith-Adao, 2011; Nel, Murray, Maherry, Petersen, Roux, Driver, Hill, Van Deventer, Funke, Swartz, Smith-Adao, Mbona,

Downsborough & Nienaber, 2011). Site is part of the Lower Vaal Water Management Area (WMA 10).

The two wetland depressions (pans) west and east of the proposed footprint could be managed in particular to fulfill its role in providing ecosystem services such as:

- Nutrient cycling and primary production, which are basic prerequisites of other ecosystem services of wetlands,
- Attenuation of water and enhancing water quality by processing water and regulating water run-off,
- Counter erosion which remains important even though slopes in the area are not steep,
- Habitat for a number of indigenous plant and animal species that favour of visit wetlands.

It is anticipated that the proposed development would not have a major influence on the hydrological regime of the depressions west and east of the proposed footprint. There appears to be no distinct reason (such as would have been the case for gatherings of large rare water birds; associated unique wetland vegetation; extensive edge effects of impacts; sensitive extensive wetlands) why the buffer zone should be large.

No threatened plant or animal species are suspected to be present at the site.

The type of development proposed, if approved, does not have the same impact as for example a plantation or buildings in terms of shade effects on the flora and fauna, and more importantly, on buffer zones or corridors. Upper soils at the plains where the proposed footprint is situated are conspicuously sandy which because of relatively high infiltration, adds to the limiting of water runoff in combination with very gentle slopes. A buffer zone of 50 m is thought to be adequate to maintain the functioning of the wetland systems west and east of the proposed footprint. Therefore if the development is approved the risks of any significant impacts to the wetlands are decidedly small and most likely to be counteracted or limited owing to the following reasons:

- 1) A presence of a buffer zone of at least 50 m from the perimeter of the wetlands;
- 2) The longitudinal slopes at the plains in which the wetlands are situated is very gentle, so that water runoff is likely to remain relatively low,

- 3) The site is also located in a relatively low rainfall area of which the mean annual precipitation is 500 mm or less. The amount of runoff and the percentage of runoff as a proportion of rainfall tend to increase with rainfall such that while runoff in the northwest of South Africa is less than 5 mm per annum, there is a fairly systematic increase of runoff towards southwards and eastwards in South Africa (Ellery *et al.*, 2009). Because the site falls in the northwestern parts of the country the water runoff is relatively very low,
- 4) Water runoff from the plain toward the wetlands, is a sheet flow into the depressions rather than a channeled inflow. Diffusion rather than channelling of water flow towards the excluded wetland depression disperses any water runoff or impact on erosion,
- 5) Soils at the proposed footprint are conspicuously sandy which means relatively high infiltration of water during rainfall events and an additional limitation to water runoff,
- 6) Mitigation measure of rapid re-vegetation of areas where the proposed solar panels are,
- 7) A significant decrease of grazing activities at the buffer zone of the wetland patch if the development is approved, resulting in higher cover of vegetation at this buffer zone, counteracting rapid water runoff and erosion effects,
- 8) Avoidance of establishment of exotic plant species in particular alien invasive species such as *Prosopis*,
- 9) Maintaining water quality of the wetland patch by practices of avoiding fuel spillages and any sewage or domestic waste onto soils at the plains near the wetlands.

Owing to the lack of a vast body of extensive research of the impacts of solar power plants, a relatively new field the present study could not absolutely guarantee that no risk of changes to the wetland patch are possible. However, given the factors above, it is unlikely that these wetlands would be significantly altered.

7 REFERENCES

- Alexander, G. & Marais, J. 2007. A guide to the reptiles of Southern Africa. Struik, Cape Town.
- Anderson, M.D. & Andersen, T.A. 2001. Too much, too quickly? Doubts about the sustainability of the camelthorn wood harvest. *African Wildlife* 55(3): 21-23.
- Apps, P. 2012. Smithers' mammals of Southern Africa 4th ed: A field guide, revised and updated by Peter Apps. Struik Nature, Cape Town.
- Armstrong, A.J. 1991. On the biology of the marsh owl, and some comparisons with the grass owl. *Honeyguide* 37:148-159.
- Ashman, M.R. & Puri, G. 2002. Essential soil science: a clear and concise introduction to soil science. Blackwell Publishing, Oxford.
- Barnes, K.N. ed. 2000. The Eskom Red Data Book of birds of South Africa, Lesotho and Swaziland. BirdLife South Africa, Johannesburg.
- Bates, M.F., Branch, W.R., Bauer, A.M., Burger, M., Marais, J., Alexander, G.J. & De Villiers, M.S. (eds). 2014. Atlas and Red List of the Reptiles of South Africa, Lesotho and Swaziland. *Suricata* 1. South African National Biodiversity Institute, Pretoria.
- Branch, B. 1998. Field guide to snakes and other reptiles of southern Africa. 3rd ed. Struik, Cape Town.
- Branch, B. 2008. Tortoises, Terrapins & Turtles of Africa. Struik Nature, Cape Town.
- Branch, W.R. & Patterson, R.W. 1975. Notes on the ecology of the Giant Girdled Lizard, *Cordylus giganteus*. *Journal of Herpetology* 9(4): 364-366.
- Branch, W.R., Tolley, K.A., Cunningham, M., Bauer, A.M., Alexander, G., Harrison, J.A., Turner, A.A. & Bates, M.F. eds. 2006. A plan for phylogenetic studies of southern African reptiles:

proceedings of a workshop held at Kirstenbosch, February 2006. Biodiversity Series 5. South African National Biodiversity Institute, Pretoria.

Bronner, G. 2011. *Mammals*. In: Picker, M. & Griffiths, C. 2011. *Alien & Invasive animals: a South African perspective*. Struik Nature, Cape Town, p 22-35.

Bromilow, C. 2010. Problem plants and alien weeds of South Africa. Briza Publications, Pretoria.

Carruthers, V. & Du Preez, 2011. Frogs and frogging in southern Africa 2nd ed. Struik, Cape Town.

Chanson, H. 2004. The hydraulics of open channel flow. Butterworth-Heinemann, Oxford.

Chittenden, H. 2007. Roberts Bird Guide. John Voelcker Book Fund, Cape Town.

Cillié, B., Oberprieler, U. & Joubert, C. 2004. Animals of Pilanesberg: an identification guide. Game Parks Publishing, Pretoria.

Cilliers, S.S., Müller, N. & Drewes, E. 2004. Overview on urban nature conservation: situation in the western-grassland biome of South Africa. *Urban forestry and urban greening* 3: 49-62.

Coetzee, N. & Monadjem, A. 2008. *Mystromys albicaudatus*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>.

Conradie, W., Du Preez, L.H., Smith, K. & Weldon, C. 2006. Field guide to the frogs and toads of the Vredefort Dome World Heritage Site. School of Environmental Sciences and Development, Potchefstroom.

Court, D. 2010. *Succulent Flora of Southern Africa*. Struik Nature, Cape Town.

Crouch, N.R., Klopper, R.R., Burrows, J.E. & Burrows, S.M. 2011. Ferns of Southern Africa: a comprehensive guide. Struik Nature, Cape Town.

- Day, J., Day, E., Ross-Gilespe, V. & Ketley, A. 2010. The assessment of temporary wetlands during dry conditions. Wetland Health and Importance Research Programme. Water Research Commission Report No TT 434/09. Water Research Commission, Gezina.
- Del Hoyo, J., Elliot, J. & Sargatal, J. 1992. Handbook of the birds of the world, Vol. 1. Lynx Editions, Barcelona.
- Deutschländer, M.S. & Bredenkamp, C.J. 1999. Importance of vegetation analysis in the conservation management of the endangered butterfly *Aloeides dentatis* subsp. *dentatis* (Swierstra) (Lepidoptera: Lycaenidae). *Koedoe* 42(2): 1-12.
- Dippenaar-Schoeman, A.S. 2002. Baboon and trapdoor spiders in southern Africa: an identification manual. Plant Protection Research Institute Handbook No. 13. Agricultural Research Council, Pretoria.
- Dippenaar-Schoeman, A.S. & Jocqué, R. 1997. African spiders: an identification manual. Plant Protection Research Institute Handbook No. 9. Agricultural Research Council, Pretoria.
- Drinkwater, T.W., Bate, R. & Du Toit, H.A. 1998. A field guide for identification of maize pests in South Africa. Agricultural Research Council: Grain-crops Institute, Potchefstroom.
- Du Preez, L.H. 1996. Field guide and key to the frogs and toads of the Free State. Department of Zoology and Entomology, University of the Orange Free State, Bloemfontein.
- Du Preez, L.H. & Carruthers, V. 2009. A complete guide to the frogs of southern Africa. Struik Nature, Cape Town. CD with calls included.
- DWAF (Department of Water Affairs and Forestry). 1997. South African Water Quality Guidelines for Aquatic Ecosystems.
- DWAF (Department of Water Affairs and Forestry). 1999. Resource Directed Measures for Protection of Water Resources: Wetland Ecosystems: W4. Department of Water Affairs and Forestry, Pretoria.
- DWAF (Department of Water Affairs and Forestry). 2005. A practical field procedure for identification and delineation of wetland and riparian areas. DWAF, Pretoria.

- Edge, D.A., Cilliers, S.S. & Terblanche, R.F. 2008. Vegetation associated with the occurrence of the Brenton blue butterfly. *South African Journal of Science* 104: 505 - 510.
- Ellery, W., Grenfell, M., Grenfell, S., Kotze, D., McCarthy, T., Tooth, S., Grundling, P-L., Beckedahl, H., Le Maitre, D. & Ramsay, L. 2009. WET-origins: controls on the distribution and dynamics of wetlands in South Africa.
- Ferguson-Lees, J. & Christie, D.A. 2001. *Raptors of the world*. Christopher Helm, London.
- Filmer, M.R. 1991. Southern African spiders: an identification guide. Struik, Cape Town.
- Gardiner, A.J. & Terblanche, R.F. 2010. Taxonomy, biology, biogeography, evolution and conservation of the genus *Erikssonia* Trimen (Lepidoptera: Lycaenidae). *African Entomology* 18(1): 171 – 191.
- Gauckler, P. 1867. Etudes Théoriques et Pratiques sur l'Écoulement et le Mouvement des Eaux. *Comptes Rendues de l'Académie des Sciences*, 64: 818-822.
- Germishuizen, G. 2003. Illustrated guide to the wildflowers of northern South Africa. Briza, Pretoria.
- Germishuizen, G., Meyer, N.L. & Steenkamp (eds) 2006. A checklist of South African plants. Southern African Botanical Diversity Network Report No. 41. SABONET, Pretoria.
- Goldblatt, P. 1986. The *Moraeas* of Southern Africa. *Annals of Kirstenbosch Botanic Gardens*, Volume 14. National Botanic Gardens, Cape Town.
- Goldblatt, P. & Manning, J. 1998. *Gladiolus* in Southern Africa.
- Henderson, L. 2001. *Alien weeds and alien invasive plants: a complete guide to the declared weeds and invaders in South Africa*. Plant Protection Research Institute Handbook No. 12. ARC: Plant Protection Research Institute, Pretoria.

- Henderson, L. & Cilliers, C.J. 2002. *Invasive aquatic plants: a guide to the identification of the most important and potentially dangerous invasive aquatic and wetland plants in South Africa*. Plant Protection Research Handbook No. 16. Agricultural Research Council, Pretoria.
- Henning, G.A. & Roos, P.S. 2001. Threatened butterflies of South African wetlands. *Metamorphosis* 12(1): 26-33.
- Henning, G.A., Terblanche, R.F. & Ball, J.B. (eds) 2009. South African Red Data Book: butterflies. *SANBI Biodiversity Series No 13*. South African National Biodiversity Institute, Pretoria.
- Henning, S.F. 1983. Biological groups within the Lycaenidae (Lepidoptera). *Journal of the Entomological Society of Southern Africa* 46(1): 65-85.
- Henning, S.F. 1987. Outline of Lepidoptera conservation with special reference to ant associated Lycaenidae. *Proceedings of the first Lepidoptera conservation Symposium, Roodepoort: Lepidopterists' Society of southern Africa*: 5-7.
- Henning, S.F. & Henning, G.A. 1989. South African Red Data Book: butterflies. *South African National Scientific Programmes Report No. 158*. CSIR, Pretoria.
- Herman, P.P.J. 2002. Revision of the *Tarchonanthus camphoratus* complex (Asteraceae-Tarchonantheae) in southern Africa. *Bothalia* 32,1: 21-28.
- Hill, C.J. 1995. Conservation corridors and rainforest insects. (In Watt, A.D., Stork, N.E. & Hunter, M.D. (eds.), *Forests and Insects*. Chapman & Hall, London. p. 381-393.)
- Hockey, P. 2011. *Birds*. In: Picker, M. & Griffiths, C. 2011. *Alien & Invasive animals: a South African perspective*. Struik Nature, Cape Town, p 36-44.
- Hockey, P.A.R., Dean, W.J.R. & Ryan, P.G. (eds.). 2005. *Roberts Birds of Southern Africa*. John Voelcker Bird Book Fund, Cape Town.
- Holm, E. & Marais, E. 1992. *Fruit chafers of southern Africa*. Ekogilde, Hartebeespoort.

- IUCN. 2001. *IUCN Red List Categories and Criteria: Version 3.1*. IUCN Species Survival Commission. IUCN, Gland, Switzerland and Cambridge, UK.
- IUCN. 2012. IUCN Red list of Threatened Species. Version 2012.1
- Jacobsen, W.B.G. 1983. *The ferns and fern allies of Southern Africa*. Butterworths, Durban.
- Kemper, N.P. 2001. RVI: Riparian Vegetation Index, final report, WRC Report No. 850/3/1. Institute for Water Research, Pretoria.
- Kok, J.C. 1998. *Vrystaatse bome, struik en klimplante* Kontak-uitgewers, Pretoria.
- Kleynhans, C.J. 1999. A procedure for the determination of the ecological reserve for the purposes of the national water balance model for South African River. Institute of Water Quality Studies, Department of Water Affairs & Forestry, Pretoria.
- Kleynhans, C.J., Thirion, C. & Moolman, J. 2005. A level 1 ecoregion classification system for South Africa, Lesotho and Swaziland. Report No. N/0000/00/REQ0104. Resource Quality Services, Department of Water Affairs and Forestry, Pretoria.
- Kotze, D., Marneweck, G., Batchelor, A., Lindley, D. and Collins, N. 2008. WET-EcoServices: A technique for rapidly assessing ecosystem services supplied by wetlands. Wetland Management Series. Water Research Commission Report TT339/08, Water Research Commission, Pretoria.
- Larsen, T.B. 1995. Butterfly biodiversity and conservation in the Afrotropical region. (*In* Pullin, A.S. ed. *Ecology and conservation of butterflies*. London: Chapman & Hall. p. 290-303.)
- Liebenberg, L. 1990. *A field guide to the animal tracks of Southern Africa*. David Philip Publishers, Cape Town.
- Leeming, J. 2003. *Scorpions of southern Africa*. Struik, Cape Town.
- Leroy, A. & Leroy, J. 2003. *Spiders of southern Africa*. Struik, Cape Town.

- Louw, W.J. 1951. *An ecological account of the vegetation of the Potchefstroom Area*. Botanical Survey of South Africa, Memoir No. 24. Government Printer, Pretoria.
- Low, A.B. & Rebelo, A.G. (Eds.) 1996. *Vegetation of South Africa, Lesotho and Swaziland*. Department of Environmental Affairs and Tourism, Pretoria.
- Lubke, R.A., Hoare, D., Victor, J. & Ketelaar, R. 2003. The vegetation of the habitat of the Brenton Blue Butterfly, *Orachrysops niobe* (Trimen), in the Western Cape, South Africa. *South African Journal of Science* 99: 201-206.
- Manning, J. 2003. *Photographic guide to the wild flowers of South Africa*. Briza, Pretoria.
- Manning, J. 2009. *Field guide to the wild flowers of South Africa*. Struik, Cape Town.
- Marneweck, G.C. & Batchelor, A. 2002. Wetland inventory and classification. In: *Ecological and economic evaluation of wetlands in the upper Olifants River catchment*. Palmer, R.W., Turpie, J., Marneweck, G.C. and Batchelor, A. (eds). Water Research Commission Report No. 1162/02.
- McMurtry, D., Grobler, L., Grobler, J. & Burns, S. 2008. *Field guide to the orchids of northern South Africa and Swaziland*. Umdaus Press, Hatfield.
- Mecenero, S., Ball, J.B., Edge, D.A., Hamer, M.L., Henning, G.A., Krüger, M., Pringle, E.L., Terblanche, R.F. & Williams, M.C. 2013. *Conservation Assessment of Butterflies of South Africa, Lesotho and Swaziland: Red List and Atlas*. Safronics, Johannesburg & Animal Demography Unit, Cape Town.
- Minter, L.R., Burger, M., Harrison, J.A., Braack, H.H., Bishop, P.J. & Kloepfer, D. eds. 2004. *Atlas and Red Data Book of the Frogs of South Africa, Lesotho and Swaziland*. SI/MAB series 9, Smithsonian Institution, Washington DC.
- Mucina, L. & Rutherford, M.C. eds. 2006. *The vegetation of South Africa, Lesotho and Swaziland*. *Strelitzia* 19. Pretoria: South African National Biodiversity Institute.

Mucina, L., Rutherford, M.C., and Powrie, L.W. eds. 2005. Vegetation map of South Africa, Lesotho and Swaziland, 1:1 000 000 scale sheet maps. Pretoria: South African National Biodiversity Institute.

Nel, J.L., Driver, A., Strydom, W.F., Maherry, A.M., Petersen, C.P., Hill, L., Roux, D.J., Nienaber, S., Van Deventer, H., Swartz, E.R. & Smith-Adao, L.B. 2011. Atlas of Freshwater Ecosystem Priority Areas in South Africa: Maps to support sustainable development of water resources. WRC Report No. TT 500/11. Water Research Commission, Pretoria.

Nel, J.L., Murray, K.M., Maherry, A.M., Petersen, C.P., Roux, D.J., Driver, A., Hill, L., Van Deventer, H., Funke, N., Swartz, E.R., Smith-Adao, L.B., Mbona, N., Downsborough, L. & Nienaber, S. 2011. Technical Report for the Freshwater Ecosystem Priority Areas Project. WRC Report No. TT 1801/2/11. Water Research Commission, Pretoria.

New, T.R. 1993. ed. Conservation biology of *Lycaenidae* (butterflies). *Occasional paper of the IUCN Species Survival Commission* No. 8.

New, T.R. 1995. Butterfly conservation in Australasia – an emerging awareness and an increasing need. (*In Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 304 – 315.*)

Ollis, D.J., Snaddon, C.D., Job, N.M. & Mbona, N. 2013. Classification System for Wetlands and other Aquatic Ecosystems in South Africa. User Manual: Inland Systems. *SANBI Biodiversity Series 22*. South African National Biodiversity Institute, Pretoria.

Peacock, F. 2006. Pipits of Southern Africa. Published by the author, Pretoria. www.pipits.co.za.

Pfab, M.F. 2002. Priority ranking scheme for Red Data plants in Gauteng, South Africa. *South African Journal of Botany* (68): 299-303.

Pfab, M.F. & Victor, J.E. 2002. Threatened plants of Gauteng, South Africa. *South African Journal of Botany* (68): 370-375.

- Picker, M. & Griffiths, C. 2011. Alien & Invasive animals: a South African perspective. Struik Nature, Cape Town.
- Picker, M., Griffiths, C. & Weaving, A. 2004. Field guide to insects of South Africa. 2nd ed. Cape Town: Struik.
- Pooley, E. 1998. A field guide to wild flowers of KwaZulu-Natal and the eastern region. Natal Flora Publications Trust, Durban.
- Pringle, E.L., Henning, G.A. & Ball, J.B. eds. 1994. Pennington's Butterflies of Southern Africa. Struik Winchester, Cape Town.
- Pryke, S.R. & Samways, M.J. 2001. Width of grassland linkages for the conservation of butterflies in South African afforested areas. *Biological Conservation* 101: 85-96.
- Pullin, A.S. ed. 1995. Ecology and conservation of butterflies. Chapman & Hall, London.
- Rautenbach, I.L. 1982. The mammals of the Transvaal. Ecoflan monograph 1: 1-211.
- Retief, E. & Herman, P.P.J. 1997. Plants of the northern provinces of South Africa: keys and diagnostic characteristics. Strelitzia 6. National Botanical Institute, Pretoria.
- Russell, W. 2009. *WET-RehabMethods: National guidelines and methods for wetland rehabilitation*. WRC Report TT 341/09, Water Research Commission, Pretoria.
- Rutherford, M.C. & Westfall, R.H. 1994. Biomes of southern Africa: An objective categorisation, 2nd ed. Memoirs of the Botanical Survey of South Africa, Vol. 63, pp. 1-94. National Botanical Institute, Pretoria.
- Ryan, P. 2001. Practical Birding: A guide to birdwatching in southern Africa. Struik, Cape Town.
- Samways, M.J. 2005. Insect diversity conservation. Cambridge University Press, Cambridge.

- Sieben, E.E., Kotze, D.C., Ellery, W.N. & Russell, W.B. 2009. Chapter 6: Using vegetation in wetland rehabilitation. In: Russel, W. 2009. *WET-RehabMethods: National guidelines and methods for wetland rehabilitation*. WRC Report TT 341/09, Water Research Commission, Pretoria, pp. 54-94.
- Skelton, P. 2001. A complete guide to the freshwater fishes of Southern Africa. Struik, Cape Town.
- Skelton, P. & Weyl, O. 2011. *Fishes*. In: Picker, M. & Griffiths, C. 2011. *Alien & Invasive animals: a South African perspective*. Struik Nature, Cape Town, p 36-44.
- Skinner, J.D. & Chimimba, C.T. 2005. The mammals of the southern African subregion. Cambridge University Press, Cape Town.
- Sliwa, A. 2008. *Felis nigripes*. In: IUCN 2012. IUCN Red List of Threatened Species.
- Smit, N. 2008. Field guide to the *Acacias* of South Africa. Briza, Pretoria.
- Smithers, R.H.N. 1986. South African Red Data Book: Terrestrial mammals. *South African National Scientific Programmes Report No. 125*. CSIR, Pretoria.
- South Africa. 2004. National Environmental Management: Biodiversity Act No. 10 of 2004. Government Printer, Pretoria.
- Stuart, C. & Stuart, T. 2006. Field guide to the larger mammals of Africa 3rd ed. Struik Nature, Cape Town.
- Stuart, C. & Stuart, T. 2013. A field guide to the tracks and signs of Southern, Central and East African wildlife 4th ed. Struik Nature, Cape Town.
- Tarboton, W. & Erasmus, R. 1998. Owls and owling in southern Africa. Struik, Cape Town.
- Taylor, J.C., Janse Van Vuuren, M.S. & Pieterse, A.J.H. 2007. The application and testing of diatom-based indices in the Vaal and Wilge Rivers, South Africa. *Water SA* 33(1): 51-59.

- Terblanche, R.F. & Edge, D.A. 2007. The first record of an *Orachrysops* in Gauteng. *Metamorphosis* 18(4): 131-141.
- Terblanche, R.F., Morgenthal, T.L. & Cilliers, S.S. 2003. The vegetation of three localities of the threatened butterfly species *Chrysoritis aureus* (Lepidoptera: Lycaenidae). *Koedoe* 46(1): 73-90.
- Terblanche, R.F. & Van Hamburg, H. 2003. The taxonomy, biogeography and conservation of the myrmecophilous *Chrysoritis* butterflies (Lepidoptera: Lycaenidae) in South Africa. *Koedoe* 46(2): 65-81.
- Terblanche, R.F. & Van Hamburg, H. 2004. The application of life history information to the conservation management of *Chrysoritis* butterflies (Lepidoptera: Lycaenidae) in South Africa. *Koedoe* 47(1): 55-65.
- Thomas, C.D. 1995. Ecology and conservation of butterfly metapopulations in the fragmented British landscape. (*In* Pullin, A.S. ed. Ecology and conservation of butterflies. London: Chapman & Hall. p. 46-64.)
- Van den Berg, J. & Drinkwater, T.W. 1998. Field guide to identification of sorghum pests in South Africa. Agricultural Research Council: Grain-crops Institute, Potchefstroom.
- Van Ginkel, C.E., Glen, R.P., Gordon-Gray, K.D., Cilliers, C.J., Muasya, M. & van Deventer, P.P. 2011. Easy identification of some South African Wetland Plants. WRC Report No TT 479/10. Water Research Commission, Gezina.
- Van Jaarsveld, E.J. 2006. The Southern African *Plectranthus* and the art of turning shade to glade.
- Van Oudtshoorn, F. 2012. Guide to grasses of southern Africa 3rd ed. Briza Publications, Pretoria.
- Van Wyk, B. 2000. A photographic guide to wild flowers of South Africa. Struik, Cape Town.
- Van Wyk, B. & Malan, S. 1998. Field Guide to the Wild Flowers of the Highveld. Struik, Cape Town.

Van Wyk, A.E. & Smith, G.F. 2001. Regions of floristic endemism in Southern Africa: a review with emphasis on succulents, Umdaus Press, Pretoria.

Van Wyk, B.E. & Smith, G.F. 2014. Guide to the aloes of South Africa. 3rd ed. Briza, Pretoria.

Van Wyk, B. & Van Wyk, P. 2013. Field guide to trees of southern Africa 2nd ed. Struik Nature, Cape Town.

Walker, C. 1996. Signs of the Wild. 5th ed. Struik, Cape Town.

Watt, A.D., Stork, N.E. & Hunter, M.D. (eds.), Forests and Insects. London: Chapman & Hall. (p. 381-393.)