

Watercourse Investigation for the Hlabisa Landfill Site (LFS), KwaZulu-Natal Province

Specialist Report

Prepared For: SE Solutions
Email: vici.napier@outlook.com
Tel: +27 (0)12 643 0190
Author: L.E.R. Grobler
Company: Imperata Consulting
Date: November 2015



IMPERATA CONSULTING™

Wetlands • Ecology • Responsibility

CC Reg. No: 2007/043725/23
Sole Member: LER Grobler
Wetland Ecologist (Pr. Sci. Nat)
Reg. No.: 400097/09

P.O. Box 72914
Lynnwood Ridge, 0040
Email: retief@imperata.co.za
Fax: 012 365 3217

Suggested Citation: Grobler, L.E.R.2014. Watercourse Investigation for the Hlabisa Landfill Site (LFS), KwaZulu-Natal Province. Specialist report for SE Solutions. Pretoria.

Approach and Disclaimer

This report provides a brief description of watercourses, as defined by the National Water Act (NWA), Act No. 36 of 1998, that are present within the study area, including wetlands present within a 500m radius of the study area. The latter is undertaken at a secondary level of detail through a mainly desktop approach with limited site surveying. The study area is known as the Hlabisa landfill site (LFS), an operational LFS, located in the north-western section of Hlabisa in the Hlabisa Local Municipality, in northern KwaZulu-Natal. The investigation furthermore provides a description of selected aspects of the study area and identifies potential project related impacts, recommended mitigation measures and an impact assessment table.

This study does not provide detailed descriptions of the local geology, agricultural potential, climatic conditions, hydrology of the aquatic environments (including volumes and flow patterns), surface and ground water quality, aquatic and terrestrial flora and fauna, or a detailed review of the legal constraints associated with potential project related impacts on the environment. It has been assumed for the purposes of this report that these aspects have been the subject of separate specialist studies should they be required as part of the environmental authorisation process. The following refers to general limitations that affect the accuracy of information represented within this report:

- A soil specific wetland delineation approach had to be adopted as hydrophyte indicators were not readily available, due to the survey period (October 2015), overlapping with the non-growing season when plant identification to the species level is constrained by the general absence of reproductive structures. In addition, plant identification was further hindered by excavation and stockpile disturbances associated with the operation of a landfill site within the study area, while a high grazing pressure was present in the surrounding 500m study area buffer.
- Hydrophyte wetland indicators were therefore of limited use during the field surveys and a soil specific wetland delineation approach had to be adopted. Soil indicators were, however, also partially obscured within the study area by soil stockpiles located in between excavated areas.

Table of Contents

List of Figures	4
List of Tables	4
1. Introduction	5
1.1. Background and Terms of Reference	5
1.2. Experience of the author	6
1.4. General assumptions	6
1.5. Overview of wetlands and riparian habitat.....	6
1.5.1. What are wetlands?.....	6
2. Methods	8
2.1. General	8
2.2. Limitations.....	10
3. Study Area Description	11
3.1. Location and existing land use	11
3.3. Study Area catchment and surface hydrology.....	11
3.4. Local climate, regional vegetation and soils	13
4. Watercourse Delineation and Assessment.....	14
4.1. Delineated and classified watercourses.....	14
4.2. Present Ecological State & Ecological Importance and Sensitivity Assessments ..	19
5. Discussion and Impact Assessment.....	19
5.1. General discussion	19
5.2. Project-related impact identification	21
5.2.1. Loss of watercourse habitat and indigenous species.....	21
5.2.2. Water quality degradation in watercourses and drainage lines	23
5.2.3. Erosion and sedimentation	23
5.3. Water Use License requirements.....	24
6. References and Further Reading.....	25

List of Figures

- Figure 1: Locality map of the study area illustrating the study area boundary, 500 m study area buffer, wetlands indicated on the NFEPA dataset, and drainage lines from the 1:50 000 topographical map (2831BB). 12
- Figure 2: Illustrates the northern section of the site located on a crest landscape portion with a low gradient (top) and the southern section of the site, which is located on a steep slope (bottom). 13
- Figure 3: Illustrates landscape positions in the study area and its surroundings that are more likely to contain wetlands and other watercourses, as determined by a Topographical Wetness Index model created with the SAGA GIS program. Areas with increased wetness have only been modeled and may therefore differ from actual site conditions. 15
- Figure 4: Illustrates delineated watercourses (wetland and riparian habitat) along with demarcated drainage lines within the study area and its 500 m buffer. Map label numbers from 1 to 7 are used to indicate different watercourse and drainage line features, which are referred to in the main text. 17
- Figure 5: Illustrates a soil sample and profile within the study area that lack wetland indicators (top row); and a soil sample and profile that do contain signs of hydromorphism in the form of mottles in a low chroma (grey) matrix, sampled in the channelled valley bottom wetland (bottom row). 18
- Figure 6: Illustrates erosion along a livestock path located southwest of the study area. Continuous erosion has resulted in a 'meandering' path, which leads to further erosion over a wider area. 18
- Figure 7: Illustrates a 32 m buffer around all delineated wetlands, riparian habitat and drainage lines. Map label numbers from 1 to 7 are used to indicate different watercourse and drainage line features, which are referred to in the main text. 20

List of Tables

- Table 1: Provides a summary of surface area sizes of delineated watercourses and the length of drainage lines. 19
- Table 2: Pre-mitigation impact assessment table for the Hlabisa LFS. 22
- Table 3: Post-mitigation impact assessment table for the Hlabisa LFS. 22

1. Introduction

1.1. Background and Terms of Reference

SE Solutions appointed Imperata Consulting to conduct a watercourse specialist investigation for the existing and operational Hlabisa landfill Site (LFS), located in the Hlabisa Local Municipality in the KwaZulu-Natal Province. The study was undertaken by Mr. L.E.R Grobler from Imperata Consulting who compiled the report on the findings of the commissioned watercourse assessment, which included a desktop component as well as a fieldwork survey component. Ms. L. Van Rooyen (nee Pretorius), a PhD student at the University of the Free State, was also present during the site survey. Mrs. Van Rooyen is currently undertaking a PhD study that evaluates soil organic matter and other features as possible indicators of wetland conditions in northern KwaZulu-Natal. The terms of reference for the specialist study include the following:

- The delineation and assessment of wetlands and other watercourses present within the study area, including the delineation of wetlands within a 500m radius around the property (henceforth referred to as the 500m study area buffer or 500m buffer). The delineation of wetlands within the 500m buffer will be undertaken at a secondary level of detail through a mainly desktop approach with limited site surveying.
- Watercourses identification will be based on definitions specified in the National Water Act (Act No. 36 of 1998) (NWA). Watercourse definitions used as part of the investigation include the following:
 - A river or spring.
 - A natural channel in which water flows regularly or intermittently.
 - A wetland, lake or dam into which, or from which, water flows.
- The description and classification of delineated wetlands into corresponding hydro-geomorphic (HGM) units according to Ollis *et al.* (2013).
- Present Ecological State (PES) assessment of identified wetlands within the LFS study area. PES assessments for wetlands and other watercourses located outside of the study area, but inside the 500m buffer, are excluded from this study.
- Ecological Importance and Sensitivity (EIS) assessment of identified wetlands present within the study area. EIS assessments for wetlands and other watercourses located outside of the study area, but inside the 500m buffer, are excluded from this study.
- The identification of potential project-related impacts along with an impact assessment and the recommendation of appropriate mitigation measures.

1.2. Experience of the author

Mr. Grobler has undergraduate majors in Botany (UP) and Soil Science (UP), an honours degree in Botany from the University of Pretoria (cum laude), and a MSc (cum laude) from the Department of Plant Sciences (UP) with a focus on peatland wetland systems. He is a registered Pr. Sci. Nat professional natural scientist in the fields of Botanical Science and Ecological Science (Reg. no. 400097/09). He has been working as a consultant based in Pretoria, with work experience in Gauteng, Mpumalanga, North-West, Eastern Cape, Northern Cape, Free State, Limpopo and KwaZulu-Natal Provinces over the last eight years. Areas of specialisation include wetland, riparian and headwater drainage line assessments, with a special interest in peat wetlands.

1.4. General assumptions

- This study assumes that the project proponents will always strive to *avoid, mitigate or offset* potentially negative project related impacts on the environment, with impact avoidance being considered the most successful approach, followed by mitigation and offset. It further assumes that the project proponents will seek to enhance potential positive impacts on the environment.
- Spatial GIS shapefiles received from the client were used to demarcate the landfill site boundaries are deemed accurate.
- The project proponents will commission an additional study to assess the impact(s) if there is a change in the size and/or extent of the study area or proposed infrastructure that is likely to have a potentially significant and/ or unavoidable impact on watercourses (e.g. wetlands).

1.5. Overview of wetlands and riparian habitat

1.5.1. What are wetlands?

In terms of the Ramsar Convention on Wetlands (Iran 1971), to which South Africa is a contracting party, "... wetlands include a wide variety of habitats such as marshes, peatlands, floodplains, rivers and lakes, and coastal areas such as salt marshes, mangroves, and sea grass beds, but also coral reefs and other marine areas no deeper than six meters at low tide, as well as human-made wetlands such as waste-water treatment ponds and reservoirs" (Ramsar Convention Secretariat 2007).

In South Africa, wetlands are defined as "...land which is transitional between terrestrial and aquatic systems where the water table is usually at or near the surface, or the land is periodically covered with shallow water, and which land in normal circumstances supports or

would support vegetation typically adapted to life in saturated soil" (National Water Act, 1998 (Act No. 36 of 1998)). Wetlands are also included in the definition of a watercourse within the NWA, which implies that whatever legislation refers to the aforementioned will also be applicable to wetlands.

In addition, the NWA stipulates that "...reference to a watercourse includes, where relevant, its bed and banks...". This has important implications for the management of watercourses and encroachment on their boundaries, as discussed further on in this document.

The NWA defines riparian areas as "...the physical structure and associated vegetation of the areas associated with a watercourse which are commonly characterized by alluvial soils, and which are inundated or flooded to an extent and with a frequency sufficient to support vegetation of species with a composition and physical structure distinct from those of adjacent land areas..." Note that this does not imply that the plant species within a riparian zone must be aquatic, only that the species composition of plant assemblages must be different within the riparian area and adjacent uplands.

In terms of the wetland delineation document available from the Department of Water Affairs and Forestry (DWAf), now known as the Department of Water and Sanitation (DWS), "wetlands must have one or more of the following attributes" (DWAf 2005):

- Wetland (hydromorphic) soils that display characteristics resulting from prolonged saturation.
- The presence, at least occasionally, of water loving plants (hydrophytes).
- A high water table that results in saturation at or near the surface, leading to anaerobic conditions developing in the top 50 cm of the soil.

It follows that the level of confidence associated with a specific area being considered as a wetland is proportionate to the number of confirmed indicators that positively correlate with wetland habitat. Not all indicators are always present within a specific biophysical and land use setting, while not all indicators are always reliable and/or useful under all conditions. The use of additional wetness indicators from different disciplines that are internationally applied therefore adds value and confidence in the identification and delineation of wetland habitats, especially in challenging environments. These types of environments include urban settings where disturbances to the natural soil and vegetation are common.

2. Methods

2.1. General

The following methods and approaches were applied as part of the wetland investigation:

- Existing spatial datasets that indicate potential watercourses and ecologically important areas were used as part of an initial desktop approach:
 - The 1:50 000 river and drainage line data of the study area and its surroundings was used, as illustrated on the relevant topographic map (2831BBHlabisa).
 - The recently completed National Freshwater Ecosystem Priority Areas (NFEPA) spatial database was used to identify potential wetland areas within the study area and its immediate surroundings. This wetland layer has been formed by combining information from the National Land Cover 2000 data set (NLC 2000), 1:50 000 topographic maps and sub national data (Van Deventer *et al.* 2010).
 - The National Spatial Biodiversity Assessment (NSBA) spatial dataset, which is based on the DWA 1:500 000 rivers GIS layer (Driver *et al.* 2004). The GIS layer was obtained via the BGIS website hosted by the South African National Biodiversity Institute (SANBI).
 - The KZN Freshwater Systematic Conservation Plan 2007 and the KZN Terrestrial Systematic Conservation Plan 2010.
- A Topographic Wetness Index model was performed with SAGA GIS software to help indicate the potential occurrence of wetlands and other watercourses within the study area and its surrounding 500m buffer (also referred to as the 500 m study area buffer). Sample points were targeted in areas with expected increased wetness, as indicated on the modeled map, which were regarded as more likely to contain wetlands.
- A wetland site survey was undertaken on 27 October 2015 by Mr. LER Grobler & Ms. L Van Rooyen.
- Watercourses were identified and delineated through the procedure described by the Department of Water and Sanitation (DWS; previously also known as DWAF and DWA) in their document entitled: "A Practical field procedure for the identification and delineation of wetlands and riparian areas" (DWAF 2005).
- Available wetland indicators that were investigated included hydromorphic (wetland soil) features, the presence of wetland plant species (e.g. hydrophytes), riparian species and vegetation features, alluvial soil features, and terrain unit indicators.

- A strong emphasis was placed on the identification of hydromorphic features to identify and delineated wetland areas. Investigated hydromorphic features typically included the presence of mottling, gleying, localised iron depletion, low chroma matrix colours, and organic enrichment in the A horizon (DWAF, 2005 & Nobel *et al.* 2005).
- Sample points were generally arranged along transects perpendicular and parallel to areas with convergent contour lines where drainage lines or flow paths were expected, in order to record gradients of change between terrestrial and watercourse habitats.
- The field surveys primarily focussed on the delineation of watercourses within the study area, while selected areas were investigated for the presence of wetland habitat within a 500 m radius of the site. Any wetland habitat located within the 500m study area buffer was primarily delineated and classified through a desktop approach with limited site surveying.
- Identified wetland areas and other watercourses were delineated into GIS polygon shapefiles, which were used for map creation.
- All natural wetlands identified within the study area and 500 m buffer were classified according to the recently completed 'Classification System for Wetlands and other Aquatic Ecosystems in South Africa' up to the hydrogeomorphic (HGM) unit level (Ollis *et al.* 2013).
- The HGM classification system is based on three key parameters pertaining to the wetland: the geomorphic setting of the wetland, the source of water inputs into the wetland, and its hydrodynamics (how does water move through the wetland), (Brinson 1993; Kotze *et al.* 2008).
- The Present Ecological State (PES) of any wetland that may occur within the study area was to be assessed according to the method developed by Kleynhans (DWA 1999) or the Wetland IHI method developed by DWA (2007).
- The PES method compares the current condition of a wetland, or other watercourse type, to its perceived reference condition, in order to determine the extent to which the watercourse had been modified from its pristine (reference) condition.
- Results from the PES assessments are rated into one of six categories ranging from unmodified/ pristine wetlands (Class A) to critically/ totally modified HGM wetland units (Class F).
- An Ecological Importance and Sensitivity (EIS) assessment of any identified wetlands that may occur within the study area were undertaken to provide an indication of the conservation value and sensitivity of these watercourses. The applied EIS wetland assessment was based on the following criteria derived from the method proposed by Rountree & Malan (2010):

- Habitat uniqueness
- Species of conservation concern
- Habitat fragmentation with regards to ecological corridors
- Prominent ecosystem services

2.2.Limitations

The following refers to general limitations that affect the applicability of information represented within this report (also refer to the Approach and Disclaimer section):

- Wetland assessments are based on a selection of available techniques that have been developed through the Department of Water and Sanitation (DWS), as well as the Water Research Council (WRC) based on site conditions and applicability. These techniques are, however, largely qualitative in nature with associated limitations due to the range of interdisciplinary aspects that have to be taken into consideration.
- Wetland areas within transformed landscapes, such as urban, agricultural settings, or other areas with existing disturbances, such as landfill sites, are often affected by disturbances that restrict the use of available wetland indicators, such as hydrophytic vegetation or soil indicators (e.g. as a result of the dominance of alien vegetation, cultivation, hard surfaces, and dumping and infilling). Hence, a wide range of available indicators are considered, to help determine wetland boundaries more accurately.
- A soil specific wetland delineation approach had to be adopted as hydrophyte indicators were not readily available due to the survey period overlapping with the non-growing season (dry season) when plant identification is constrained by the general absence of reproductive structures. In addition, plant identification was further hindered by excavation and stockpiling activities within the study area.

3. Study Area Description

3.1. Location and existing land use

- The Hlabisa landfill site (LFS), henceforth referred as the study area or site, is located within the Hlabisa Local Municipality in KwaZulu-Natal Province (coordinates 28°8 37'.034"S, 31°51, 52'. 366"E).
- Hluhluwe-iMfolozi Game Reserve, managed by Ezemvelo KZN Wildlife (EKZNW), is located approximately 5.25 km southeast of the site, while Hluhluwe town is located approximately 43 km to the east-northeast. Other nearby towns include St Lucia and Mtubatuba that are located respectively ±60 km and ±45 km to the southeast from the site.
- The study area is situated in the north-western portion of Hlabisa, centered between dirt roads that converged into urban tar roads to the east, such as the R618 provincial road, located approximately 1 km southeast of the site (Figure 1).
- The study area and its surroundings are located on a low mountain topography and characterised by grassland, bushland, forested woodland and fragmented patches. The site itself is located on a steep slope, especially its southern section (Figure 2).
- The existing land use of the study area and its immediate surrounding area include cultivated land (subsistence farming), plantations (agricultural facility) and homesteads. Homesteads are primarily located on crests and other high-lying areas rather than on steep slopes and narrow valley floors. Livestock grazing is also common in the 500 m buffer area.
- The study area is surrounded by the Good Shepherd Mission and Bazaneni, Hlambanyathi, Amatshamnyama and Sitezi residential areas.

3.3. Study Area catchment and surface hydrology

- The study area is located within the Usuthu toMhlathuze Water Management Area (WMA) and falls within Mfolozi and Pongolo Quaternary Catchment W32E.
- The Mfolozi and Pongolo Quaternary catchment W32E has a Very High conservation status and has a Natural, Unmodified Condition (Class A), Present Ecological State (PES) as provided by (Middleton and Bailey, 2008).
- A non-perennial tributary of the Bazaneni River is located approximately 100 m southwest of the study area and flows in a south-eastern direction within the 500m study area buffer (Figure 1).




Figure 1: Locality map of the study area illustrating the study area boundary, 500 m study area buffer, wetlands indicated on the NFEPA dataset, and drainage lines from the 1:50 000 topographical map (2831BB).



Figure 2: Illustrates the northern section of the site located on a crest landscape portion with a low gradient (top) and the southern section of the site, which is located on a steep slope (bottom).

3.4. Local climate, regional vegetation and soils

- Annual rainfall is approximately 769 mm and falls mostly during the summer months (Middleton & Bailey 2008). The mean annual temperature (MAT) is 18.8°C and days with frost are infrequent to occasional during winter (Mucina & Rutherford, 2006).
- The vegetation map for South Africa, Lesotho and Swaziland (VEGMAP) indicate that the study area forms part of the Savanna Biome and Lowveld Bioregion (subgroup). The entire study area and its immediate surrounds falls within the Northern Zululand Sourveld vegetation unit (Mucina & Rutherford, 2006).
- This vegetation type is classified as Vulnerable due to vegetation transformation and low levels of protection. Almost 22% of the vegetation has been transformed due to cultivation and plantations, however, 4% is statutorily conserved within the Ithala Game Reserve and Hluhluwe-iMfolozi Park (Mucina & Rutherford, 2006).
- A recent revision of VEGMAP for KwaZulu-Natal by Scott-Shaw & Escott (2011) delineated a new vegetation unit within the 500 m study area buffer, namely Scarp Forest, which is located south and southeast of the study area. This vegetation unit has a Least Threatened conservation status in KwaZulu-Natal.

- The study area falls within Land Type Ab81 and features Red-yellow apedal, freely drained soils, while the geology of the area contains Dolerite and Sandstone of the Vryheid Formation (Ecca Group), (Mucina & Rutherford, 2006).
- The study area does not overlap with any listed Threatened Ecosystem areas according to the 2011 Schedule (Government Gazette of December 2011) of the National Environmental Management: Biodiversity Act, 2004 (Act No. 10 of 2004) (NEMBA).
- The Ezemvelo KwaZulu-Natal Wildlife (EKZNW, 2010) Terrestrial Systematic Conservation Plan: Minimum Selection Surface (MINSET), was consulted to identify important aspects regarding the study area and its surroundings:
 - The study area and its entire 500m buffer do not overlap with any of the three Critical Biodiversity Area categories, but are associated with a Biodiversity Area (map legend OCO).
 - Land that falls within the Biodiversity Area category is not open for wholesale development, as important species are still located within them and these should be accounted for in the EIA process.
 - The site and its buffer are associated with a Terrestrial Surface
- According to (EKZNW, 2007) Freshwater Systematic Conservation Plan the following important aspects are defined for the study area and its surroundings:
 - The entire study area and majority of the 500 m buffer are located within an Earmarked Planning Unit, while an Available Planning Unit overlaps within the northern-most section of the 500 m study area buffer.

4. Watercourse Delineation and Assessment

4.1. Delineated and classified watercourses

- The study area nor its 500m buffer overlaps with wetland habitat indicated on the NFEPA wetland spatial dataset (Figure 1).
- No drainage lines, as indicated on the 1:50000 topographical map 2831BB, overlap with the study area, but they are indicated in the 500 m buffer (Figure 1).
- A Topographical Wetness Index model was created to illustrate potential areas with increase soil moisture conditions within the site and its surroundings (Figure 3). This map was used to help target surveys during the site visit. Areas with expected increased wetness and therefore possible wetland conditions do not overlap with the study area, but are associated with the drainage lines from the topographical map in the 500 m study area buffer (Figure 3).




Figure 3: Illustrates landscape positions in the study area and its surroundings that are more likely to contain wetlands and other watercourses, as determined through a Topographical Wetness Index model created with the SAGA GIS program. Areas with increased wetness have only been modeled and may therefore differ from actual site conditions.

- The site survey confirmed the absence of wetland habitat within the study area as no hydromorphic features were identified, nor were any obligate hydrophytes recorded in the study area (Figure 4 & 5). Some facultative hydrophytes, such as *Centella asiatica*, was identified in the southern section (lowest area) of the landfill site, but did not correspond with wetland soil indicators.
- Channelled valley bottom wetland habitat, as described by Ollis *et al.* (2013), was recorded and delineated in a valley floor landscape setting, located southwest of the study area (Figure 4). The watercourse is expected to have existed as an unchannelled valley bottom that have become eroded over time.
- The wetland contains distinct mottling with low chroma matrix colours (grey) in the top 0.5 m of the soil profile and also contained obligate hydrophytes in the form of *Mariscus* sp. (Figure 5).
- Other watercourses identified in the 500m study area buffer include riparian habitat located immediately downstream of the channelled valley bottom. The delineated riparian habitat is characterised by an incised channel with riparian habitat, but also contains woody vegetation that is not associated with riparian habitat along its outer margins. The woody non-riparian vegetation is expected to be associated with the Scarp Forest vegetation unit described by Mucina & Rutherford (2006) and revised by Scott-Shaw & Escott in 2011 in KwaZulu-Natal.
- Areas mapped as riparian habitat therefore represent a conservative estimation of riparian habitat within the 500 m buffer, as it includes both riparian and non-riparian habitat. Distinction was not made as the area was primarily assessed as part of a desktop evaluation.
- The closest delineated riparian area (map label 1) is located approximately 75 m south of the study area (Figure 5).
- Drainage lines that represent areas where waterflow is concentrated, as a result of convergent contour lines and valley floor landscape settings, have also been delineated (Figure 4). These areas are not regarded as watercourses, as defined in the NWA, as no wetland features, channel features or riparian habitat characteristics were identified during the site visit. They do, however, form part of the drainage network and are regarded as areas where surface flow occurs during runoff events following sufficient rainfall. These areas have therefore been mapped as drainage lines that should be considered as areas where potential watercourses may still occur.
- Erosion features are common in the 500 m buffer and livestock paths are especially prone to erosion (Figure 6).




Figure 4: Illustrates delineated watercourses (wetland and riparian habitat) along with demarcated drainage lines within the study area and its 500 m buffer. Map label numbers from 1 to 7 are used to indicate different watercourse and drainage line features, which are referred to in the main text.



Figure 5: Illustrates a soil sample and profile within the study area that lack wetland indicators (top row); and a soil sample and profile that do contain signs of hydromorphism in the form of mottles in a low chroma(grey) matrix, sampled in the channelled valley bottom wetland (bottom row).



Figure 6: Illustrates erosion along a livestock path located southwest of the study area. Erosion has led to the development of a 'meandering' livestock path, which leads to further erosion over a wider area.

4.2. Present Ecological State & Ecological Importance and Sensitivity Assessments

No Present Ecological State (PES) or Ecological Importance and Sensitivity (EIS) assessments were undertaken as no wetland or other type of watercourse were identified within the study area.

5. Discussion and Impact Assessment

5.1. General discussion

- No wetlands or other watercourses are present within the study area, while only a single channelled valley bottom wetland was identified within the 500 m study area buffer, along with three riparian areas that are connected to one another, and headwater drainage lines (Figure 4).
- A total wetland area of 1.92 ha and a combined riparian area of 5.40 ha have been delineated within the 500 m buffer (Table 5; Figure 7).
- No watercourse is located within 32 m of the study area (Figure 7).
- The impact of the landfill site on the channelled valley bottom wetland is regarded as negligible, as the wetland is located in a separate catchment, located upstream of the LFS site (Figure 7).

Table 1: Provides a summary of surface area sizes of delineated watercourses and the length of drainage lines.

Feature	Map interpretation notes (see Figures 4 & 7).	Surface area size in hectares or length in meters
Study area	-	0.71 ha
500 m study area buffer	-	95.74 ha
Riparian habitat	Map label 1	4.45 ha
Channelled valley bottom wetland	Map label 2	1.92 ha
Drainage line	Map label 3	441.92 m
Drainage line	Map label 4	125.18 m
Drainage line	Map label 5	89.71 m
Drainage line	Remaining drainage line	131.26 m
Riparian habitat	Map label 6	0.72 ha
Riparian habitat	Map label 7	0.23 ha




Figure 7: Illustrates a 32 m buffer around all delineated wetlands, riparian habitat and drainage lines. Map label numbers from 1 to 7 are used to indicate different watercourse and drainage line features, which are referred to in the main text.

5.2. Project-related impact identification

No new developments in the form of infrastructure construction are expected to be associated with the landfill site (LFS), as it is likely to continue in its current operational state for a further 5 years thereafter it will be closed. The current operation of the LFS does, however, require environmental authorisation. The environmental authorisation process is currently ongoing and this watercourse assessment study forms part thereof.

No proposed development impacts are therefore threatening watercourses adjacent to the study area, but existing watercourse-related impacts have to be identified, assessed and mitigated through appropriate recommendations. An impact assessment rating for each identified impact is provided in Table 2, while impact descriptions and mitigation measures are discussed below. The impact assessment method is based on the template received from SE Solutions and can be made available on request.

5.2.1. Loss of watercourse habitat and indigenous species

Natural habitat on the landfill site is largely transformed by active operations leading to a loss of indigenous species. No wetland or other watercourses are present within the study area, but do occur in the surrounding 500 m buffer. The closest of these is riparian habitat located approximately 75 m south of the study area (Figure 7). It is therefore highly unlikely that wetland and riparian habitat will be affected through clearing or other activities related to the landfill site as long as activities are confined to the existing study area boundaries. The impact is considered to be Neutral and of Low significance (Tables 2& 3).

Recommended mitigation:

- No landfill activities should occur outside of the existing study area boundaries.
- Any future expansion plans should consider delineated watercourses, drainage lines and their combined 32 m buffer as sensitive features (Figure 7), which should be further investigated and verified during environmental planning and authorisation phases.
- The fence around the study area should be repaired and maintained.
- Vehicle movement must be restricted to the study area and existing vehicle tracks outside of the study area.

Table 2: Pre-mitigation impact assessment table for the Hlabisa LFS

Nature	Extent	Duration	Intensity	Probability	Status	Significance	Confidence	Reversibility	Replaceability
Loss of wetland habitat and indigenous species	Footprint	Short term	Low	Possible	Neutral	No impact	High	Partially reversible	Moderate - High
Water quality degradation in watercourses and drainage lines	Local	Long term	Medium-High	Highly likely	Negative	Medium - High	Medium	Partially reversible	Moderate - High
Erosion and Sedimentation	Local	Long term	Medium	Likely	Negative	Medium	Medium	Partially reversible	Low - Moderate

Table 3: Post-mitigation impact assessment table for the Hlabisa LFS

Nature	Extent	Duration	Intensity	Probability	Status	Significance	Confidence	Reversibility	Replaceability
Loss of wetland habitat and indigenous species	Footprint	Short term	Low	Improbable	Neutral	No impact	High	Partially reversible	Moderate - High
Water quality degradation in watercourses and drainage lines	Local	Long term	Medium	Possible	Negative	Low	Medium	Partially reversible	Moderate - High
Erosion and Sedimentation	Local	Long term	Low-Medium	Possible	Negative	Low	Medium	Partially reversible	Low - Moderate

5.2.2. Water quality degradation in watercourses and drainage lines

The spread of litter can lead to surface pollution in delineated watercourses. Leachate from the landfill site is another source of potential pollution that may be transported as runoff or interflow into downstream wetlands. No drainage pipes or pipe outlets were observed downslope of the study area, between the site and the riparian habitat, and it remains unknown whether a drainage system is in place. No signs of a lining (sealing layer) was recorded within the LFS, but a vegetated buffer of terrestrial land of approximately 75 m is present between the LFS and its nearest watercourse (Figure 7). The impact is considered to be Negative and of Medium-High significance (Table 2), but can be reduced to Low with mitigation (Table 3).

Additional information related to the risks, impacts and effective mitigation of water quality impacts, especially groundwater quality impacts, should be obtained from the groundwater study, which was not available at the time of report compilation.

Recommended mitigation:

- The fence around the landfill site should be maintained to contain solid waste.
- Dumping activities must be confined to the active operational area only.
- Vehicle movement must be restricted to the fenced area and the access roads leading to the landfill.
- The operational landfill site should not be expanded without relevant environmental authorisation/s.
- A lining or sealing layer should be applied to the excavated cavity in order to reduce the risk of leachate contaminating the groundwater table.
- Trenches and other stormwater management interventions as part of a stormwater management plan is recommended in order to restrict water accumulation in the waste body, as water ingress into the waste body can result in more leachate. Trenches and related stormwater control infrastructure should be located outside of buffered watercourses and drainage lines (Figure 7).
- Further details related to water quality mitigation should be obtained from the groundwater study.

5.2.3. Erosion and sedimentation

Earthmoving activities associated with landfill operations, such as digging, trenching and stockpile movement can lead to increased erosion and the alteration of sedimentation regime. Erosion disturbances were recorded within the channelled valley bottom wetland in the form of headcut development and channel incision, as well as the formation of lateral headcuts. Erosion is also prominent along livestock and pedestrian paths. Sediment mobilisation within the study area along its steep slope in the southern section of the site can, however, result in sediment

influx into lower-lying watercourses. The impact is considered to be Negative and of Medium significance (Table 2), but can be reduced to Low with mitigation (Table 3).

Recommended mitigation:

- Earthmoving activities outside the footprint area within the study area and vehicle movement outside of existing access roads should be prohibited.
- Monitor all stockpiles for signs of erosion.
- Any areas where active erosion and sedimentation is observed must be rehabilitated and berms must be utilised to slow movement of water, where necessary.
- No new access road crossings to the study area across wetland or other watercourse features should be constructed without environmental authorisation/s, as they have the potential to concentrate surface flow and result in erosion.

5.3. Water Use License requirements

- Wetlands and other watercourses are protected water resources in terms of the NWA. Development or transformation of the watercourses is regarded as a *water use*, which can only be allowed through an approved Water Use License, irrespective of the condition of the affected watercourse.
- The implication is that authorization will have to be obtained from DWS before water use activities can be initiated in demarcated wetlands and other watercourses.
- Section 21 of the NWA defines different types of water use in a watercourse. Examples of water use activities that may be applicable to the artificial wetland in the study area include the following
 - (c) impeding or diverting the flow of water in a watercourse.
 - (i) altering the bed, banks, course or characteristics of a watercourse.
- A DWS stipulation published in Government Gazette No 32805 (December 2009) also require that a Water Use License should be applied for when any wetlands are present within a 500 m radius (buffer) of Section 21 (c) and/or Section 21 (i) water use activities.

6. References and Further Reading

Anon, 2000. Rehabilitation recommendation after alien plant control. Plant Protection Research Institute, Agriculture Research Council, Pretoria.

Anon. 2004. A guide to the use of herbicides (18th Edition). National Department of Agriculture, Directorate of Food Safety and Quality Assurance: Pretoria.

Berner, J.T., Thiesing M.A., Simpson R. & Jantz C. 2008. Alternative Futures for Headwater Stream and Wetland Landscapes in the Upper Delaware Basin, New York, USA. Unpublished document.

Brinson M. 1993. A hydro-geomorphic classification for wetlands. Wetland Research Programme Technical Report WRP-DE-4. US Army Corps of Engineers, Waterways Experiment Station, Vicksburg, MS.

Broderson, J.M.. 1973. Sizing buffer strips to maintain water quality. M.S. Thesis, University of Washington, Seattle.

Bullock, A. and Acreman, M. 2003. The role of wetlands in the hydrological cycle. *Hydrology and Earth System Sciences*, 7, 3, 358-389.

Calhoun, A.J.K., Miller, N.A. and Klemens, M.W. 2005. Conserving pool-breeding amphibians in human-dominated landscapes through local implementation of Best Management Practices. *Wetlands Ecology and Management*, 13, 291-304.

Castelle, A.J., Conolly, C., Emers, M., Metz, E.D., Meyer, S., Witter, M., Mauermann, S., Erickson, T. and Cooke, S.S. 1992. Wetland Buffers: use and effectiveness. In Adolfsen Associates, Inc., Shorelands and Coastal Zone Management Program, Washington Department of Ecology, Olympia, Publication No. 92-10.

Clark, J.R. 1977. Coastal Ecosystem Management: A Technical Manual for the Conservation of Coastal Zone Resources. John Wiley and Sons, New York, New York.

CSIRO 2006. Urban Stormwater – Best Practice Environmental Management Guidelines. Commonwealth Scientific and Industrial Research Organisation. Collingwood, Australia.

Department of Environmental Affairs (DEA). 2014. Government Notice 599 National Environmental Management: Biodiversity Act (10/2004): Alien and Invasive Species List. Gazette number 37886.

Department of Water Affairs and Forestry. 1996. Aquatic ecosystems. Volume 7. South African Water quality guidelines. Department of Water Affairs and Forestry, Pretoria.

Department of Water Affairs and Forestry (DWAf). 1999. Resource Directed Measures for Protection of Water Resources. Wetland Ecosystems. Version 1.0, September 1999.

Department of Water Affairs and Forestry. 2005. A practical field procedure for identification and delineation of wetlands and riparian areas. Edition 1. Department of Water Affairs and Forestry, Pretoria.

Department of Water Affairs and Forestry. 2007. Manual for the assessment of a Wetland Index of Habitat Integrity for South African floodplain and channelled valley bottom wetland types by M. Rountree (ed); C.P. Todd, C. J. Kleynhans, A. L. Batchelor, M. D. Louw, D. Kotze, D. Walters, S. Schroeder, P. Illgner, M. Uys. and G.C. Marneweck. Report no. N/0000/00/WEI/0407. Resource Quality Services, Department of Water Affairs and Forestry, Pretoria, South Africa.

Department of Water Affairs and Forestry. 2009. Government Gazette No. 32805. Impeding or Diverting the Flow of Water in a Watercourse [Section 21(c)] and Altering the Bed, Banks, Course or Characteristics of a Watercourse [Section 21(i)]. Pp66-71, Pretoria.

Dodds, W.K. & Oaks, R.M. 2008. Headwater influences on downstream water quality. *Environmental Management* 41:367–377.

Driver, A., Maze, K., Lombard, A.T., Nel, J., Rouget, M., Turpie, J.K., Cowling, R.M., Desmet, P., Goodman, P., Harris, J., Jonas, Z., Reyers, B., Sink, K. & Strauss, T. 2004. South African National Spatial Biodiversity Assessment 2004: Summary Report. Pretoria: South African National Biodiversity Institute.

EKZNW (2007) Freshwater Systematic Conservation Plan: Best Selected Surface (Marxan). Unpublished GIS Coverage [Freshwater_cons_plan_2007], Biodiversity Conservation Planning Division, Ezemvelo KZN Wildlife, P. O. Box 13053, Cascades, Pietermaritzburg, 3202.

EKZNW (2010) Terrestrial Systematic Conservation Plan: Minimum Selection Surface (MINSET). Unpublished GIS Coverage [tscp_minset_dist_2010_wll.zip], Biodiversity Conservation Planning Division, Ezemvelo KZN Wildlife, P. O. Box 13053, Cascades, Pietermaritzburg, 3202.

Erman, D.C., Newbold, J.D. and Roby, K.B. 1977. Evaluation of streamside bufferstrips for protecting aquatic organisms. Technical Completion Report, Contribution No.165. California Water Resources Center, University of California, Davis.

Environmental Law Institute 2008. Planner's Guide to Wetland Buffers for Local Governments. Washington D.C., USA. <http://www.eli.org>

EPA 1996. Protecting Natural Wetlands – A Guide to Stormwater Best Management Practices. Environmental Protection Agency (EPA). Washington, United States of America.

Gomi, T., Sidl, R.C., Richardson, J.S. 2002. Understanding processes and downstream linkages of headwater systems. *BioScience*, 52, 10, 905-916.

Kleynhans, C.J. 1999. Comprehensive habitat integrity assessment. In: Water resources protection policy implementation. Resource Directed Measures for Protection of Water Resources. River Ecosystems Version 1.0. Department of Water Affairs and Forestry.

- Kleynhans C.J., MacKenzie J. & Louw M.D. 2007. Module F: Riparian Vegetation Response Assessment Index in River EcoClassification: Manual for EcoStatus Determination (version 2). Joint Water Research Commission and Department of Water Affairs and Forestry report. WRC Report No. TT 333/08.
- Kotze, D.C., Marneweck, G.C., Batchelor, A.L., Lindley, D.S. & Collins, N.B. 2008. Wetland-Ecoservices: A rapid assessment procedure for describing wetland benefits. WRC Report TT 339/08. Water Research Commission, Pretoria.
- Macfarlane D.M, Kotze D, Walters D, Ellery W, Koopman V, Goodman P, and Goge C. 2008. WET-Health: A Technique for Rapidly Assessing Wetland Health. WRC Report TT 340/08. Water Research Commission, Pretoria.
- Meyer, J.L. & Wallace, J.B. 2001. Lost linkages and lotic ecology: Rediscovering small streams. Pages 295–317 in Press MC, Huntly, N.J. & Levin S, (eds). *Ecology: Achievement and Challenge*. Oxford (United Kingdom): Blackwell Scientific.
- Middleton, B.J. & Bailey, A.K. 2008. Water Resources of South Africa, 2005 Study (WR2005). Water Research Commission (WRC) Report TT380/08, Pretoria.
- Mucina, L. & Rutherford, M.C. (eds). 2006. *The vegetation of South Africa, Lesotho and Swaziland*. Strelitzia 19, South African National Biodiversity Institute, Pretoria.
- Nanson, G.C. and Croke, J.C. 1992. A genetic classification of floodplains. *Geomorphology*, 4, 459-486.
- NC Division of Water Quality. 2005. Identification Methods for the Origins of Intermittent and Perennial streams, Version 3.1. North Carolina Department of Environment and Natural Resources, Division of Water Quality. Raleigh, NC.
- Newbold, J.D., Erman, D.C., Roby, K.B.. 1980. Effects of logging on macroinvertebrates in streams with and without buffer strips. *Can. J. Fish Aquat. Sci.*, 37,1076-1085.
- Noble, C. V., Martel, D. J., & Wakeley, J. S. 2005. A national survey of potential wetland hydrology regional indicators, WRAP Technical Notes Collection (ERDC TNWRAP- 05-1). U.S. Army Engineer Research and Development Center, Vicksburg, MS.
- Ollis, D.J., Snaddon, C.D., Job, N.M. & Mboma, N. 2013. Classification System for Wetlands and other Aquatic Ecosystems in South Africa. User Manual: Inland Systems. SANBI Biodiversity Series 22. South African National Biodiversity Institute, Pretoria.
- Rountree, M.W. and H. Malan (editors). 2010. Rapid Ecological Reserve Determination Methods for Wetlands (Version 2.0). Joint Department of Water Affairs and Water Research Commission, Pretoria.
- VAN WYK, A.E. & SMITH, G.F. 2001. Regions of floristic endemism in southern Africa. A review with emphasis on succulents. Umdaus Press, Pretoria.

Young, R.A., Huntrods, T. and Anderson, W.. 1980. Effectiveness of vegetated buffer strips in controlling pollution from feedlot runoff. *J Environ. Qual.*, 9, 483-497.