



SiVEST SA (PTY) LTD

**PROPOSED CONSTRUCTION OF THE KRAALTJIES
WIND ENERGY FACILITY AND ASSOCIATED
INFRASTRUCTURE, NEAR BEAUFORT WEST,
WESTERN CAPE PROVINCE, SOUTH AFRICA**

Terrestrial Biodiversity Assessment

DEFF Reference:	TBA
Report Prepared by:	EnviroSci (Pty) Ltd
Issue Date:	20 November 2022
Version No.:	1

SiVEST SA (PTY) LTD

PROPOSED CONSTRUCTION OF THE KRAALTJIES WIND ENERGY FACILITY AND ASSOCIATED INFRASTRUCTURE, NEAR BEAUFORT WEST, WESTERN CAPE PROVINCE, SOUTH AFRICA

TERRESTRIAL BIODIVERSITY ASSESSMENT

EXECUTIVE SUMMARY

South Africa Mainstream Renewable Power Developments (Pty) Ltd (hereafter referred to as “Mainstream”), has appointed SiVEST SA (Pty) Ltd (hereafter referred to as “SiVEST”) to undertake the required EIA processes for the proposed construction of the 240MW Kraaltjies Wind Energy Facility (WEF), Battery Energy Storage System (BESS), 11-33kv portion / yard of the shared 11-33kv/132kv onsite substation and associated infrastructure near Beaufort West in the Western Cape Province.

The overall objective of the development is to generate electricity by means of renewable energy technology capturing wind energy to feed into the National Grid.

It is anticipated that the proposed Kraaltjies WEF will comprise up to sixty (60) wind turbines with a maximum total energy generation capacity of up to approximately 240MW. The electricity generated by the proposed WEF development will be fed into the national grid via a 132kV overhead power line and will form part of a separate environment assessment and authorisation process

In terms of the Environmental Impact Assessment (EIA) Regulations, which were published on 04 December 2014 [GNR 982, 983, 984 and 985) and amended on 07 April 2017 [promulgated in Government Gazette 40772 and Government Notice (GN) R326, R327, R325 and R324 on 7 April 2017], various aspects of the proposed development are considered listed activities under GNR 327 and GNR 324 which may have an impact on the environment and therefore require authorisation from the National Competent Authority (CA), namely the Department of Environment, Forestry and Fisheries (DFFE), prior to the commencement of such activities. Specialist studies have been commissioned to assess and verify the project under the new Gazetted specialist protocols.

The regulatory requirements are also discussed with regards to the NEMA and the National Water Act in Section 4 of this report. The PROTOCOL FOR SPECIALIST ASSESSMENT AND MINIMUM REPORT CONTENT REQUIREMENTS FOR THE ENVIRONMENTAL IMPACTS ON BIODIVERSITY and Appendix 6 of the NEMA EIA Regulations, have been adhered to.

This report fulfils the Biodiversity Specialist Assessment Report criteria for assessment listed under the various Theme Sensitivity Protocols, where the following sensitivity ratings were contained in the Screening Tool Report

- Animal Species Combined – Medium related to a potential occurrence of the Critically Endangered Riverine Rabbit (*Bunolagus monticularis*) and the Endangered reptile *Cherobius boulengeri* (“Dwarf” Karoo padloper).

- Plant Species – Medium sensitivity due to the potential presence of *Tritonia florentiae*, *Peersia frithii*, Species 383, Species 1039 (Protocol does not allow for the listing of the names of species under threat within public documents and is only made known to the specialist conducting the assessment).
- Terrestrial Biodiversity – Very High sensitivity related to the presence of Critical Biodiversity Areas (CBAs), Ecological Support Areas (ESAs) and National Freshwater Ecosystem Priority Areas (NFEPAs).

The verification of any of the Very High Sensitivity rated habitats / species localities is thus critical as the proposed development should then avoid these areas. During the screening assessment, a four-day site visit of the area was conducted in November 2021 and again in February 2022, in which the habitats / species listed above were considered, together with a description of the general environment and species assemblages found present. This spatial data is then supplied to the Applicant to develop the layout outside of these areas (inclusive of suitable buffers) as a mechanism of impact avoidance using fine scale mapping data.

The study area had received some much-needed winter rainfall, which aided in critically assessing the ecological character of the site, with particular reference to any linkages between the aquatic and terrestrial environment as indicated in the Screening Tool Results (CBA, ESA & FEPAs). The information collected, was also compared to previous assessments within the region by members of EnviroSci, used in the assessment of the wind farms that have been completed.

In summary two key terrestrial habitats and the aquatic habitats were observed and then rated based on their sensitivity to the proposed development. These habitats included:

1. *Gamka Karoo vegetation unit*
 - a. *Shale / Mudrock Plains*
 - b. Small ridges / inselbergs
2. Alluvial rivers with and without riparian vegetation (discussed in Aquatic Assessment)

The Gamka-Karoo vegetation spans the entire site / study area with the presence / absence and abundance of plant species dependent on the slope and stability of the soils found present.

Thus, the flat plains areas contained most of the plant and animal species known to occur within region, while the steep rock cliffs were more devoid of species. This is possibly linked to the fact that the soils in these areas are composed mostly of mudrock (shales), that weather and decompose easily, thus unstable, only allowing more hardy succulent species to colonise these areas.

The sensitivity assessment mentioned considered the habitats observed and these were categorised or rated based on the presence/absence of the following:

- Unique or sensitive habitats
- Presence of importance or listed taxa (faunal & floral)
- Intact and functional habitat associated with sensitive areas indicated in the DFFE Screening Tool results

Several High Sensitivity Habitats were observed and mapped, and these were then considered No-Go for any new infrastructure, while Moderate and Low sensitivity areas could be considered for development. The only exception being road crossings and transmission lines which would be considered acceptable within No-Go areas, if these areas are spanned and/ or located within existing disturbance footprints (e.g. roads within existing farm tracks) and/or suitably mitigated.

The following direct impacts were identified, which are aligned with those contained in the Biodiversity Assessment Protocol and will be assessed in greater detail in the EIA phase of the assessment: Construction and to a degree the Operational and Decommissioning Phases where relevant, as per below:

Construction & Decommissioning Phases

- Impact 1: Loss of species of special concern
- Impact 2: Loss of terrestrial habitats – flora and vegetation
- Impact 3: Loss of terrestrial species - fauna

Operational phase

- Impact 4: Loss of terrestrial species - fauna

The project overall has a small footprint spread out over a large area, allowing for retention of much of the natural environment so that the systems should remain largely unaffected. Therefore, the wind farm is such that it carries a low intensity impact on terrestrial resources but requiring the clearing of areas with terrestrial vegetation.

A variety of environmental features were observed within the study areas and these were mapped and buffered as necessary for their protection. The current layout has, to a large degree, avoided the most sensitive features and the associated buffer areas, greatly reducing the potential overall impact and environmental risk. Noting that these are mostly linked to the CBA/ESAs which are directly linked to the aquatic environment (Alluvial rivers and watercourses) that dominate the majority of the site.

However several sensitive species were observed within the site, which included both plants and reptiles, the former being found throughout the site, while the latter are highly mobile, thus core sensitive areas could not be mapped as these species are thus encountered throughout the site. Thus once the layout for roads, hardstands and buildings has been refined a micro-siting walkdown must be conducted to ensure that any populations of these species are avoided.

The overall and cumulative impacts, as assessed, are linked to instances where complete avoidance was not possible, or the nature of the activities involve a potential risk. Overall, it is expected that the impact on the environment would be Low (-). Noteworthy areas, that should be avoided, include the Very High Sensitivity areas as shown in this report.

Based on the findings of this study, the specialist finds no reason to withhold to an authorisation for any of the proposed activities, assuming that key mitigation measures are implemented. Due consideration must be given to all the substations (Option 1 and 2) due to being located within High Sensitivity watercourses in particular option 1 (See aquatic report)

However, this must all still be assessed once the roads layout has been provided, coupled to a microsite walkdown once all information is available.

NATIONAL ENVIRONMENTAL MANAGEMENT ACT, 1998 (ACT NO. 107 OF 1998) AND ENVIRONMENTAL IMPACT REGULATIONS, 2014 (AS AMENDED) - REQUIREMENTS FOR SPECIALIST REPORTS (APPENDIX 6)

Regulation GNR 326 of 4 December 2014, as amended 7 April 2017, Appendix 6	Section of Report
1. (1) A specialist report prepared in terms of these Regulations must contain- a) details of- i. the specialist who prepared the report; and ii. the expertise of that specialist to compile a specialist report including a curriculum vitae;	Appendix 1 CV
b) a declaration that the specialist is independent in a form as may be specified by the competent authority;	Attached to Report
c) an indication of the scope of, and the purpose for which, the report was prepared;	Section 1.1 and 1.3 of this report
(cA) an indication of the quality and age of base data used for the specialist report;	Section 1.3
(cB) a description of existing impacts on the site, cumulative impacts of the proposed development and levels of acceptable change;	Section 5
d) the date and season of the site investigation and the relevance of the season to the outcome of the assessment;	Section 1.3 and 5
e) a description of the methodology adopted in preparing the report or carrying out the specialised process inclusive of equipment and modelling used;	Section 1.3
f) details of an assessment of the specific identified sensitivity of the site related to the proposed activity or activities and its associated structures and infrastructure, inclusive of a site plan identifying site alternatives;	Section 5.1
g) an identification of any areas to be avoided, including buffers;	Section 5 & 6
h) a map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	Section 5
i) a description of any assumptions made and any uncertainties or gaps in knowledge;	Section 2
j) a description of the findings and potential implications of such findings on the impact of the proposed activity, (including identified alternatives on the environment) or activities;	Section 6 & 8

k) any mitigation measures for inclusion in the EMPr;	Section 6
l) any conditions for inclusion in the environmental authorisation;	Section 5. 6 and 8
m) any monitoring requirements for inclusion in the EMPr or environmental authorisation;	Section 6
n) a reasoned opinion- <ul style="list-style-type: none"> i. (as to) whether the proposed activity, activities or portions thereof should be authorised; <ul style="list-style-type: none"> (iA) regarding the acceptability of the proposed activity or activities; and ii. if the opinion is that the proposed activity, activities or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan; 	Section 8
o) a description of any consultation process that was undertaken during the course of preparing the specialist report;	N/A
p) a summary and copies of any comments received during any consultation process and where applicable all responses thereto; and	N/A
q) any other information requested by the competent authority.	N/A
2) Where a government notice <i>gazetted</i> by the Minister provides for any protocol or minimum information requirement to be applied to a specialist report, the requirements as indicated in such notice will apply.	Yes - Appendix 2

SIVEST SA (PTY) LTD

PROPOSED CONSTRUCTION OF THE KRAALTJIES WIND ENERGY FACILITY AND ASSOCIATED INFRASTRUCTURE, NEAR BEAUFORT WEST, WESTERN CAPE PROVINCE, SOUTH AFRICA

TERRESTRIAL BIODIVERSITY ASSESSMENT

Contents

1.	INTRODUCTION	1
1.1	Terms of Reference	1
1.2	Specialist Credentials	1
1.3	Assessment Methodology	2
2.	ASSUMPTIONS AND LIMITATIONS	3
	TECHNICAL DESCRIPTION	4
3.	LEGAL REQUIREMENT AND GUIDELINES	5
4.	DESCRIPTION OF THE RECEIVING ENVIRONMENT	5
5.	SPECIALIST FINDINGS / IDENTIFICATION AND ASSESSMENT OF IMPACTS	9
5.1	Construction & Decommissioning Phase	12
5.2	Operation	15
5.3	No go Impact	16
5.4	Cumulative Impacts	18
5.5	Overall Impact Rating	1
6.	COMPARATIVE ASSESSMENT OF ALTERNATIVES	1
6.1	Wind Energy Facility	1
6.2	Grid components	Error! Bookmark not defined.
6.3	No-Go Alternative	2
7.	CONCLUSION AND SUMMARY	2
7.1	Summary of Findings	2
7.2	Conclusion and Impact Statement	3

8.	REFERENCES.....	4
8.1	Waterbody classification systems	14
8.2	Wetland definition	15
8.3	National Wetland Classification System method	16
8.4	Waterbody condition	1
8.5	Aquatic ecosystem importance and function	1

List of TablesA

Table 1: Results of the sensitivity rating / constraints assessment.....	10
Table 2: Rating of impacts for the construction and decommissioning phase.....	12
Table 3: Rating of impacts for the operational phase	15
Table 4: Rating of impacts (No-go)	16
Table 5: <i>Rating of cumulative impacts</i>	19
Table 6: Overall Impact Significance for the WEF (Pre- and Post-Mitigation).....	1
Table 7: Comparative assessment of WEF components.....	1

List of Figures

Figure 1: Regional Context Map	Error! Bookmark not defined.
Figure 2: Karee WEF Site Locality	Error! Bookmark not defined.
Figure 3: Proposed 132kV Power Line Route Alignment	Error! Bookmark not defined.
Figure 4: Preliminary Turbine layout and development area	Error! Bookmark not defined.
Figure 5: Proposed Substation and Power line options.....	Error! Bookmark not defined.

List of Appendices

Appendix 1 Specialist CV
Appendix 2 DFFE Site verification statement
Appendix 3 Species Checklists
Appendix 4 Detailed study methodology

Glossary of Terms

- **Alien vegetation:** Plants that do not occur naturally within the area but have been introduced either intentionally or unintentionally. Vegetation species that originate from outside of the borders of the biome -usually international in origin.
- **Biome:** A broad ecological unit representing major life zones of large natural areas – defined mainly by vegetation structure and climate.
- **Indigenous vegetation:** Vegetation occurring naturally within a defined area.
- **RDL (Red Data listed):** Species Organisms that fall into the Extinct in the Wild (EW), critically endangered (CR), Endangered (EN), Vulnerable (VU) categories of ecological status.
- **SCC (Species of Conservation Concern):** The term SCC in the context of this report refers to all RDL (Red Data) and IUCN (International Union for the Conservation of Nature) listed species as well as protected species of relevance to the project.

List of Abbreviations

AER	Along Existing Roads – cables that are included in existing road servitudes
CARA	Conservation of Agricultural Resources Act
CBA	Critical Biodiversity Area
CSIR	Council for Scientific and Industrial Research
DDD	Data Deficient
ECO	Environmental Control Officer
EIA	Environmental Impact Assessment
EIS	Ecological Importance and Sensitivity
EMPr	Environmental Management Programme Report
EN	Endangered
EO	Environmental Office
ESA	Ecological Support Area
GA	General Authorisation (WUA type)
GBIF	Global Biodiversity Information Facility
GIS	Geographic Information System
LC	Least Concern
NFEPA	National Freshwater Ecosystem Priority Atlas (Nel, <i>et al.</i> 2011).
NT	Near Threatened
OHL	Overhead Line – transmission line cable that is not buried
ORC	Off road cable – underground or overhead transmission cable not within a road reserve
PES	Present Ecological State
SANBI	South African National Biodiversity Institute
SQ	Subquaternary catchment = Quinary catchment
VU	Vulnerable
WEF	Wind Energy Facility

SiVEST SA (PTY) LTD

PROPOSED CONSTRUCTION OF THE KRAALTJIES WIND ENERGY FACILITY AND ASSOCIATED INFRASTRUCTURE, NEAR BEAUFORT WEST, WESTERN CAPE PROVINCE, SOUTH AFRICA

TERRESTRIAL BIODIVERSITY ASSESSMENT

1. INTRODUCTION

South Africa Mainstream Renewable Power Developments (Pty) Ltd (hereafter referred to as “Mainstream”), has appointed SiVEST SA (Pty) Ltd (hereafter referred to as “SiVEST”) to undertake the required BA Processes for the proposed construction of the 240MW Kraaltjies Wind Energy Facility (WEF) and associated infrastructure near Beaufort West in the Western Cape Province.

The overall objective of the development is to generate electricity by means of renewable energy technology capturing wind energy to feed into the National Grid.

It is anticipated that the proposed Kraaltjies WEF will comprise up to sixty (60) wind turbines with a maximum total energy generation capacity of up to approximately 240MW. The electricity generated by the proposed WEF development will be fed into the national grid via a 132kV overhead power line and will form part of a separate environment assessment and authorisation process.

In terms of the Environmental Impact Assessment (EIA) Regulations, which were published on 04 December 2014 [GNR 982, 983, 984 and 985] and amended on 07 April 2017 [promulgated in Government Gazette 40772 and Government Notice (GN) R326, R327, R325 and R324 on 7 April 2017], various aspects of the proposed development are considered listed activities under GNR 327 and GNR 324 which may have an impact on the environment and therefore require authorisation from the National Competent Authority (CA), namely the Department of Forestry, Fisheries and the Environment (DFFE), prior to the commencement of such activities. Specialist studies have been commissioned to assess and verify the project under the new Gazetted specialist protocols.

1.1 Terms of Reference

Please refer to Specialist ToR provided in the Scoping Report.

1.2 Specialist Credentials

Please see Appendix 1 - Specialist CVs

1.3 Assessment Methodology

These assessments were conducted using the following assessment process based on 4 days field work conducted in November 2021, early summer, and again in February 2022, but after several good winter months with rainfall, therefore many of the plants were showing improved growth and most had or were flowering after a prolonged period of drought in the region:

Methodology summary

(Excluding bats and avifauna)

A desktop and literature review of the area under investigation was conducted to collate as much information as possible prior to any detailed fieldwork. The purpose of the desktop assessment is to rank relevant areas according to their ecological sensitivity and to identify areas of ecological risk prior to the site visit.

Other relevant literature for e.g. Global Biodiversity Information Facility, Virtual Atlas Projects, iNaturalist, relevant Red Data books, ordinances and all systematic bioregional / conservation plans.

Fieldwork was limited to visual sightings by means of transect walks and plot-based sampling, while particular attention was also paid to the occurrence Red Data species or Protected species as follows:

Vegetation units was sampled by means of the following techniques as per each site:

- Data collection was plot-based and in the form of vegetation samples within selected reference areas to categorise the various vegetation units.
- Results from the data analysis provided a description of the dominant and typical species occurring on the site(s), and included:
 - Threatened, endemic or rare species, with an indication of the relative functionality and conservation importance of the specific community in the area under investigation
 - Invasive or exotic species present and localities in the area
 - The functional and conservation importance of all vegetation communities in the investigation area

Mammals were sampled by means of the following techniques:

- Fieldwork included visual sightings by means of transect walks to evaluate the presence of mammal taxa. During the site visit, specific attention was given to signs (droppings, burrows, vocalisations, etc.) of taxa and the presence of suitable habitat
- Camera traps were deployed for the maximum possible time with important or strategic habitat, thus any images collected will form part of the EIA phase of the assessment
- A full list of species observed and expected to occur will then be included
- Specific reference will be made to the occurrence of Red Data species

Herpetofauna (reptiles & amphibians) were sampled by means of the following techniques:

- Visual observations (including nocturnal surveys)
- Installation of pitfall traps and two drift fence arrays. Data collected from these will also be included in the EIA phase
- Active searching techniques; and
- Vocalisations (for amphibians)

Invertebrates will be sampled by means of the following techniques:

- Random linear transects using standard hand nets while focussing on specific indicator groups;
- All taxa caught, were identified to species level if appropriate literature is available (as in the case of butterflies), otherwise the concept known as RTU's (Recognisable Taxonomic Units) or morphospecies was applied;
- The presence of conservation important taxa was also be verified by intensive searching of likely habitat types or burrows.
- Additional information of faunal community residing in the area of investigation were sourced from distributional data/records (both recent and historical), relevant literature, the private sector and other atlas projects.

Habitat areas (based on the species compositions of the vegetation analysis, topography and soils) was then ranked into Very High / No-Go, High, Medium or Low classes in terms of their significance based on the Ecological Sensitivity and Conservation Importance. A sensitivity and habitat map (including buffer zones if applicable) was produced based on the above information. This combined with the aquatic sensitivity map will then be used by the proponent to finalise the development layout in the remaining phases of the project.

2. ASSUMPTIONS AND LIMITATIONS

To obtain a comprehensive understanding of the dynamics of both the flora and fauna of communities within a study site, as well as the status of endemic, rare or threatened species in any area, assessments should always consider investigations at different time scales (across seasons/years) and through replication. However, due to time constraints these long-term studies are not feasible and are thus mostly based on instantaneous sampling. This limitation is common to many impact assessment type studies, but the findings are deemed adequate for the purposes of decision-making regarding project acceptability, unless otherwise stated.

Therefore, due to the scope of the work presented in this report, a long-term investigation of the proposed site was not possible and as such not perceived as part of the Terms of Reference. However, a concerted effort was made to sample and assess as much of the potential site, as well as make use of any supporting literature, species distribution data and aerial photography.

It should be emphasised that information, as presented in this document, only has reference to the study area as indicated on the accompanying maps. Therefore, this information cannot be applied to any other area without detailed investigation.

TECHNICAL DESCRIPTION

At this stage it is anticipated that the proposed Kraaltjies WEF will comprise of a maximum of sixty (60) wind turbines with a maximum total energy generation capacity of up to 240MW. The electricity generated by the proposed WEF development will be fed into the national grid via a 132kV overhead power line. The 132kV overhead power line will however require a separate EA and is subject to a BA process, which is currently being undertaken in parallel to this EIA process but as a separate process and is not assessed herein.

- Up to sixty (60) wind turbines, each between, with a maximum export capacity of approximately 240MW. This will be subject to allowable limits in terms of the Renewable Energy Independent Power Producer Procurement Programme (REIPPPP). The final number of turbines and layout of the WEF will, however, be dependent on the outcome of the Specialist Studies conducted during the EIA process.
- Each wind turbine will have a hub height of between 120m and 200m and rotor diameter of up to approximately 200m.
- Permanent compacted hardstand areas / platforms (also known as crane pads) of approximately 90m x 50m (total footprint of approx. 4 500m²) per turbine during construction and for on-going maintenance purposes for the lifetime of the proposed development.
- Each wind turbine will consist of a foundation of up to approximately 15m x 15m in diameter. In addition, the foundations will be up to approximately 3m in depth.
- Electrical transformers (690V/33kV) adjacent to each wind turbine (typical footprint of up to approximately 2m x 2m) to step up the voltage to 11-33kV.
- Associated infrastructure of approximately 25ha which includes:
 - One (1) new 11-33kV/132kV IPP on-site substation including associated equipment and infrastructure the proposed substation will be a step-up substation and will include an Eskom portion and an IPP portion, hence the substation has been included in the WEF EIA and in the grid infrastructure (substation and 132kV overhead power line) BA to allow for handover to Eskom. Following construction, the substation will be owned and managed by Eskom.
 - A Battery Energy Storage System (BESS) will be located next to the onsite 11-33kV/132kV substation. The storage capacity and type of technology would be determined at a later stage during the development phase, but most likely comprise an array of containers, outdoor cabinets and/or storage tanks.
 - One (1) construction laydown / staging area. It should be noted that no construction camps will be required in order to house workers overnight as all workers will be accommodated in the nearby town.
 - Operation and Maintenance (O&M) buildings, including offices, a guard house, operational control centre, O&M area / warehouse / workshop and ablution facilities to be located on the site identified for the substation.
- The wind turbines will be connected to the proposed substation via medium voltage (11-33kV) underground cabling and overhead power lines.
- Internal roads with a servitude up to approximately 8m wide will provide access to each wind turbine. Existing site roads will be used wherever possible, although new site roads will be constructed where necessary. Turns will have a radius of up to 50m for abnormal loads (especially turbine blades) to access the various wind turbine positions. It should be noted that the proposed application site will be accessed via the N12 National Route.
- A wind measuring lattice (approximately 140m in height) mast has already been strategically placed within the wind farm application site in order to collect data on wind conditions.

- No new fencing is envisaged at this stage. Current fencing is standard farm fence approximately 1-1.5m in height. Fencing might be upgraded (if required) to be up to approximately 2m in height; and
- Water will either be sourced from existing boreholes located within the application site or will be trucked in, should the boreholes located within the application site be limited.

3. LEGAL REQUIREMENT AND GUIDELINES

The following is pertinent to this study:

- Section 24 of The Constitution of the Republic of South Africa;
- Agenda 21 – Action plan for sustainable development of the Department of Environmental Affairs and Tourism (DEAT) 1998;
- National Environmental Management Act (NEMA), 1998 (Act No. 107 of 1998) inclusive of all amendments, as well as the NEM: Biodiversity Act;
- National Water Act, 1998 (Act No. 36 of 1998);
- Conservation of Agricultural Resources Act, 1983 (Act No. 43 of 1983);
- Minerals and Petroleum Resources Development Act, 2002 (Act No. 28 of 2002);
- Nature and Environmental Conservation Ordinance (No. 19 of 1974);
- National Forest Act (No. 84 of 1998); and
- National Heritage Resources Act (No. 25 of 1999) – could apply if cultural use or heritage is linked to any natural resources.

4. DESCRIPTION OF THE RECEIVING ENVIRONMENT

The site is dominated by one terrestrial vegetation type, according to Mucina and Rutherford (2007 – amended 2018) namely Gamka-Karoo (NKI1) (Figure 6)

Based on observations made during the site visits, two key terrestrial habitats and the aquatic habitats (Figure 7) were observed and then rated based on their sensitivity to the proposed development. These habitats included:

1. *Gamka Karoo vegetation unit*
 - a. *Shale / Mudrock Plains*
 - b. *Small ridges / inselbergs*
2. Alluvial rivers with and without riparian vegetation (discussed in Aquatic Assessment)

The Gamka-Karoo vegetation spans the entire site / study area with the presence / absence and abundance of plant species dependent on the slope and stability of the soils present.

Thus, the flat plains areas contained most of the plant and animal species known to occur within the region, while the steep rock cliffs were more devoid of species. This is possibly linked to the fact that the soils in these areas are composed mostly of mudrock (shales), that weather and decompose easily, thus unstable, only allowing more hardy succulent species to colonise these areas.

None of these are listed as a Threatened Ecosystem as per the National Environmental Management Biodiversity Act, this is due to the vast area these vegetation units occupy, with little in terms of human / agricultural use.

Appendix 3 lists the typical species assemblages based on previous observations made within the region which include species records for both flora and fauna, housed in Global Biodiversity Information Facility (GBIF accessed December 2021). A potential 900 species has been previously recorded in the Quarter Degree Square grid that cover the site (3222DA), of which ca. 80% are plant species. The remainder, which excludes birds and bats as these are assessed separately, include the following taxa:

- Mammals 36 Species
- Reptiles 16 Species
- Amphibians 5 Species
- Fish 0 Species
- Insects 74 Species
- Spiders / Scorpions 2 Species
- Fungi 2 Species

This was then compared to observations made within a 4 day site-specific assessment conducted in November 2021, and again in February 2022, conducted after a period of significant winter rains (more than previous years, of which some response by flora was observed. However, the prolonged drought in the region has affected the growth of the plants, especially those in the low-lying plains areas, that have shallow soils.

Vegetation and flora

The vegetation observed within the study area corresponds with the descriptions associated with the National Vegetation Map (NSBA, 2018) and Mucina and Rutherford (2007) (Figure 6). In other words, an area that is covered by sparse dwarf shrubland on undulating plains (Plate 1), dominated by *Chrysocoma oblongifolia*, *Eriocephalus microphyllus* *E. ericoides* and *Searsia undulata*.

Other species observed included:

Aridaria noctiflora (L.) Schwantes
ssp. straminea (Haw.) Gerbaulet
Aristida congesta Roem. & Schult.
Aristida diffusa Trin.
Asparagus burchellii Baker
Blepharis mitrata C.B. Clarke
Cadaba aphylla (Thunb.) Wild
Cenchrus ciliaris L.
Chrysocoma oblongifolia DC.
Crassula lanceolata (Eckl. & Zeyh.)
Endl. ex Walp. ssp. lanceolata
Crassula muscosa L.
Dicoma capensis Less.
Digitaria argyrograpta (Nees) Stapf

Drimia intricata (Baker) J.C. Manning
 & Goldblatt
Drosanthemum lique (N.E.Br.)
 Schwantes
Enneapogon desvauxii P.Beauv.
Enneapogon scaber Lehm.
Eragrostis homomalla Nees
Eragrostis lehmanniana Nees
Eragrostis obtusa Munro ex Ficalho
 & Hiern
Eriocephalus karooicus M.A.N.Müll.
Eriocephalus microphyllus DC. var.
pubescens (DC.) M.A.N.Müll.
Euphorbia inaequilatera Sond. var.
inaequilatera

Felicia filifolia (Vent.) Burt Davy ssp.
filifolia
Felicia lasiocarpa DC.
Felicia muricata (Thunb.) Nees
Felicia muricata (Thunb.) Nees ssp.
cinerascens Grau
Fingerhuthia africana Lehm.
Galenia fruticosa (L.f.) Sond.
Galenia glandulifera Bittrich
Galenia secunda (L.f.) Sond.
Garuleum schinzii O.Hoffm. ssp.
crinitum (Dinter) Merxm.
Gazania lichtensteinii Less.
Gomphocarpus filiformis (E.Mey.)
 D.Dietr.

Helichrysum lucilioides Less.
Hermannia desertorum Eckl. & Zeyh.
Hoodia dregei N.E.Br.
Jamesbrittenia tenuifolia (Bernh.) Hilliard
Kleinia longiflora DC.
Lepidium africanum (Burm.f.) DC. ssp. *africanum*
Lepidium desertorum Eckl. & Zeyh.
Leysera tenella DC.
Limeum aethiopicum Burm.f.
Lycium cinereum Thunb.
Microcoma armatum (Thunb.) Schltr.
Monechma spartioides (T.Anderson) C.B.Clarke

Moraea polystachya (Thunb.) Ker Gawl.
Nolletia tenuifolia Mattf.
Osteospermum microphyllum DC.
Pentzia incana (Thunb.) Kuntze
Pentzia pinnatisecta Hutch.
Plinthus karooicus I.Verd.
Pteronia adenocarpa Harv.
Pteronia viscosa Thunb.
Rhigozum obovatum Burch.
Rhinephyllum luteum (L.Bolus) L.Bolus
Searsia burchellii Sond. ex Engl.
Rosenia humilis (Less.) K.Bremer

Ruschia beaufortensis L.Bolus
Ruschia spinosa (L.) Dehn
Salsola gemmifera Botsch.
Salsola tuberculata (Moq.) Fenzl
Sesamum capense Burm.f.
Stipagrostis obtusa (Delile) Nees
Tetragonia microptera Fenzl
Thesium lineatum L.f.
Tragus berteronianus Schult.
Tritonia tugwelliae L.Bolus
Ursinia nana DC.
Vachellia karroo Hayne
Zygophyllum microcarpum Licht. ex Cham. & Schltl.
Zyrphelis lasiocarpa (DC.) Kuntze




Figure 6: National Vegetation Map as per Mucina and Rutherford (2007) amended NBSA 2018




Figure 7: Spatial representation of the observed habitats, with the open remaining areas dominated by the plains Nama-Karoo vegetation type



Plate 1: A view from the northern of the site, dominated by the shale /mudstone (purple) soils and isolated ridges/cliffs in the background. The site is further dominated by the minor watercourses as shown in the foreground



Plate 2: A view of one the few cliffs within the site, that will be avoided by the placement of any turbines, and its assumed that any of the proposed roads will also avoid any areas with such slopes

No rare or listed plant species were observed during the survey period within the proposed turbine positions; however, several species are protected in terms of the Western Cape legislation (Provincial Nature Conservation Ordinance). The disturbance, destruction and/or relocation, whichever is more relevant, of these species would require the relevant permits from the provincial authority. It is highly recommended that a detailed walkdown of the final layout is conducted, during a suitable time of the year (Spring / early summer). This will result in a complete species list for the actual footprints and / or assist with any micro-siting that may be required to avoid any important habitat, as the relocation of certain species during a search and rescue operation is not always successful, thus avoidance is found to be a better solution.

Several of these species are shown in Plate 3 and where found throughout the site.

The DFFE Screening Tool lists Plants *Tritonia florentiae*, *Peersia frithii*, Species 383, Species 1039, which were actively searched for, but suitable habitat and / or the presence / absence of these species was not confirmed. Most of these species habitat is associated with rocky outcrop environments and can be avoided by the proposed development areas once the layout has been micro-sited.



Plate 3: Several interesting plants occur within the region and include (in a clockwise direction), *Eriospermum paradoxum*, remnants of a *Gonialoe variegata* (syn. *Aloe variegata*) specimen, also known as tiger aloes and partridge-breasted aloes, *Mesembryanthemum resurgens* and *Hoodia dregei*, an endemic species to the greater study area.

Fauna

As previously mentioned approximately 135 animal species have been previously observed within the quarter degrees square area associated with the study area. These are predominantly Mammal (26%), Reptile (12%) and Insect species (55%), which for the most part are highly mobile and / or habitat specific. These, as listed in the Species Checklist created for the assessment (Appendix 3), were then searched for during the site visit. The only exception being the fish and amphibian species as no permanent or suitable habitat was observed within the study area, although aquatic habitats do occur downstream of the project area.

The DFFE screening tool results only include one important species (High Sensitivity), namely the Critically Endangered Riverine Rabbit (*Bunolagus monticularis*). Riverine rabbits are habitat-specific associated with dense patches of riverine bush along seasonal rivers similar to those found downstream of the site. The Riverine rabbit is the only indigenous burrowing species in Africa, and thus requires deep, soft alluvial soils. It is therefore important that the Alluvial Wash Floodplains with riparian areas, which also contain both *Lycium* and *Salsola* plant species, a favoured food source for this rabbit, are avoided as far as possible by the proposed development.

Two of the Endangered reptile *Cherobius boulengeri* ("Dwarf" Karoo padloper) were observed outside of the proposed wind farm boundary, but within the adjacent farm portions, thus it must be assumed that this species will occur within the site. Therefore, considerable caution is advised during the construction period for the potential disturbance of this small animal species, especially during the vegetation clearing process, thus an Environmental Control Officer / Environmental Officer (whichever is most applicable) must during construction to ensure that none are present during this phase within the proposed works area for the day.

In terms of fauna, the following are species which potentially occur at the site and are listed as protected species, with those species highlighted in BOLD being observed in this and past assessments:

Schedule 1: Specially Protected Fauna as per the Western Cape Nature Conservation Ordinance (No. 3 of 2000) that may occur within the region or have suitable habitat present:

- *Felis nigripes* - Black-footed cat/Miersshooptier
- *Felis silvestris* - African wild cat/Afrika wildekat
- ***Ictonyx striatus* - Striped polecat/Stinkmuishond**
- *Mellivora capensis* - Honey badger/Ratel
- ***Otocyon megalotis* - Bat-eared fox/Bakoovos**
- *Proteles cristatus* – Aardwolf/Maanhaarjakkals
- *Vulpes chama* - Cape fox / Silver jackal Silwervos
- *Orycteropus afer* - Aardvark / Ant-bear Erdvark / Aardvark
- *Atelerix frontalis* – South African hedgehog
- Family: Chamaeleonidae - Chamaeleons, all species
- Family: Cordylidae Girdled lizards, all species

Virtually all indigenous fauna which do not fall under Schedule 1 are classified under Schedule 2, except those species classified as pests. In terms of mammals most rodents, shrews, elephant shrews, bats, hares and rabbits, carnivores such as mongoose, genets, and meerkat, antelope such as klipspringer, steenbok, Mountain reedbuck and duiker are included. In terms of other vertebrates, all tortoises, lizards, most harmless snakes and all frogs are listed under Schedule 2. The full list is contained within the Schedule and are not repeated here.

In terms of fauna, the following, *inter alia*, are protected and may not be hunted, captured or harmed without a permit:

- All tortoises [2 species observed which include Angulate tortoise (*Chersina angulate* - Plate 3), Karoo Padloper (*Cherobius boulengeri*);
- All lizards;
- All frogs;
- Most snakes [2 species have been observed in the past on site, namely Cape cobra (*Naja nivea*) and Mole snake (*Pseudoaspis cana*);
- All indigenous antelope;
- Aardvark;
- Most small carnivores such as Honey Badger, Cape Fox, Bat-eared Fox;
- Large Grey Mongoose etc.; and

With the exception of the tortoises, lizards and snakes, the species listed above typically leave the area once construction commences, thus permits for the relocation of lizards, snakes and tortoises must be obtained.



Plate 3: A common site with the study area, namely Angulate tortoises (*Chersina angulata*) and Corncricket (*Hetrodes pupus pupus*)

The Western Cape Provincial Biodiversity Spatial Plan Critical Biodiversity Area (CBA) spatial layers (Figure 8). Noting that with the exception of the substations, infrastructure are located outside of any of the areas shown in Figure 8 and the reader is referred to the aquatic assessment in this regards as the CBA/ESA areas are related to river / aquatic zones.

The study area is also not located within an Important Bird Area (IBA) or a Strategic Water Resource Area and did not contain any wetland clusters or listed Threatened Ecosystems.




Figure 8: The Critical Biodiversity Areas as per the Western Cape Biodiversity Spatial Plan (2017)

5. SPECIALIST FINDINGS / IDENTIFICATION AND ASSESSMENT OF IMPACTS

Using the baseline description and field data while considering the current disturbances and site characteristics, the following features were identified, then categorized into one of a number pre-determined sensitivity categories to provide protect and/or guide the layout planning and design processes. Sensitivity areas (with their buffers) were categorized as follows:

Very High = No Go	Legislated “no go” areas or setbacks and areas or features that are considered of such significance that impacting them may be regarded as fatal flaw or strongly influence the project impact significance profile
High	Areas or features that are considered to have a high sensitivity or where project infrastructure would be highly constrained and should be avoided as far as possible. Infrastructure located in these areas are likely to drive up impact significance ratings and mitigations
Medium	Buffer areas and or areas that are deemed to be of medium sensitivity
Low	Areas of low sensitivity or constraints
Neutral	Unconstrained areas (left blank in mapping)

Table 1 below provides an overview of the sensitivity of various features (with buffers distances where relevant) as it relates to the main project component types for the project. The features are shown spatially in Figures 9. The sensitivity ratings of Very High (No-go), High, Medium and Low were determined through an assessment of the habitat sensitivity and related constraints. However, these No-Go areas relate in general terms to the project and there are areas where encroachment on these areas would occur (i.e., existing road crossings within Very High sensitivity areas) but this is only considered acceptable if these areas have already been impacted.

Table 1: Results of the sensitivity rating / constraints assessment

Development Component	Habitat type	Sensitivity rating of the respective waterbody type against the development type and the required buffer	Sensitivity rating override if an impact such as a road already occurs within the proposed footprint
WTG areas	Steep slopes / cliffs and small inselbergs	No-go	N/a
	Remaining areas	Low – thus acceptable	N/a
Hardstands , Buildings / Substations & BESS	Steep slopes / cliffs and small inselbergs	No-go	N/a
	Remaining areas	Low – thus acceptable	N/a
Roads	Steep slopes / cliffs and small inselbergs	No-go	LOW if an existing crossing / road or impact is already present, that must then be included in the potential road network inclusive of any upgrades
	Remaining areas	Low – thus acceptable	N/a
Overhead Lines	Steep slopes / cliffs and small inselbergs	These should be spanned with no towers	
	Remaining areas	Low – thus acceptable	




Figure 9: Habitat sensitivity map inclusive of terrestrial and aquatic habitats assessed

The following impacts were then assessed, which are aligned with those contained in the Biodiversity Assessment Protocols and included in the table below and assessed against the proposed and potential activities:

Biodiversity Assessment Protocol Impacts found applicable to this project	Impacts assessed in this report below
Faunal and vegetation communities inhabiting the site	Impact 1, 2, 3 and 4
Fragmentation (physical loss of ecological connectivity and or CBA corridors)	Impact 1, 2, 3 and 4
Changes in numbers and density of species	Impact 1, 2, 3 and 4
No-Go Impact	Impact 5
Cumulative Impacts	Impact 6

As highlighted above, the following impacts on the environment have been identified and will be assessed in greater detail as follows, as well as separately the No-Go and Cumulative impacts:

Construction & Decommissioning Phases

- Impact 1: Loss of species of special concern
- Impact 2: Loss of terrestrial habitats – flora and vegetation
- Impact 3: Loss of terrestrial species - fauna

Operational phase

- Impact 4: Loss of terrestrial species - fauna

5.1 Construction & Decommissioning Phase

Table 2: Rating of impacts for the construction and decommissioning phase

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION								
		E	P	R	L	D	I / M	TOTAL	STATUS	S		E	P	R	L	D	I / M	TOTAL	STATUS	S
Construction/ Decommissioning Phase																				
Impact 1: Loss of species of special concern	The construction activities will result in the disturbance of terrestrial habitats that may contain listed and or protected plant or animal species. However, none of the plant species were observed during this assessment within the low sensitivity areas	1	1	1	1	1	1	5	-	LOW (-ve)	Develop and implement an Rehabilitation and Monitoring plan post Environmental Authorisation. This must be developed following the finalisation of the turbine / road layout and a walk down has been completed. This plan should include relocation of suitable plant species, but more importantly protect any topsoil stores and promote the collection of vegetative material and propagules / seed to assist with the revegetation of the site Where possible, temporary construction lay-down or assembly areas should be sited on transformed areas; and Rapid regeneration of plant cover must be encouraged by setting aside topsoil during earthmoving and replacing onto areas where the re-	1	1	1	1	1	1	5	-	LOW (-ve)

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION								
		E	P	R	L	D	I / M	TOTAL	STATUS	S		E	P	R	L	D	I / M	TOTAL	STATUS	S
Impact 3: Loss of terrestrial species - fauna	Although most of the species observed are mobile, the increase in vehicle movement could result in an increase in road mortalities.	2	3	2	2	3	2	24	-	MEDIUM (-ve)	<p>Clear demarcation during the construction phase of all undisturbed sensitive areas that are not within the direct footprint of the WEF to ensure that there is no uncontrolled access by construction vehicles and labourers;</p> <p>ECO/EO (whichever is applicable) must be present on a daily basis to remove any reptiles such as the Karoo Padloper if present.</p> <p>Educate contractors as to the importance of the undisturbed conservation areas and importance of avoiding them;</p> <p>All vehicles must stick to designated and prepared roads and adhere to the speed limit on site of 40km/hr;</p> <p>Mitigating the risk of poaching by fencing in the accommodation compounds of the construction crews, to prevent individuals from wandering in the veld after hours; banning the possession of dogs on site</p>	1	3	2	1	2	2	18	-	LOW (-ve)

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION								
		E	P	R	L	D	I / M	TOTAL	STATUS	S		E	P	R	L	D	I / M	TOTAL	STATUS	S
											by construction and maintenance staff.									

5.2 Operation

Table 3: Rating of impacts for the operational phase

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION								
		E	P	R	L	D	I / M	TOTAL	STATUS	S		E	P	R	L	D	I / M	TOTAL	STATUS	S
Operation Phase																				
Impact 4: Loss of terrestrial species - fauna	Although most of the species observed are mobile, the increase in vehicle movement could result in an increase in road mortalities.	2	3	2	2	3	2	24	-	MEDIUM (-ve)	Clear demarcation during the construction phase of all undisturbed sensitive areas that are not within the direct footprint of the WEF to ensure that there is no uncontrolled access by construction vehicles and labourers; Educate contractors as to the importance of the undisturbed conservation areas and importance of avoiding them; All vehicles must stick to designated and prepared roads and	1	3	2	1	2	2	18	-	LOW (-ve)

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION								
		E	P	R	L	D	I / M	TOTAL	STATUS	S		E	P	R	L	D	I / M	TOTAL	STATUS	S
											adhere to the speed limit on site of 40km/hr; Mitigating the risk of poaching by fencing in the accommodation compounds of the construction crews, to prevent individuals from wandering in the veld after hours; banning the possession of dogs on site by construction and maintenance staff.									

5.3 No go Impact

Table 4: Rating of impacts (No-go)

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION								
		E	P	R	L	D	I / M	TOTAL	STATUS	S		E	P	R	L	D	I / M	TOTAL	STATUS	S
No-Go																				
Impact on terrestrial resources should the project not go ahead (i.e. the No Go Alternative)	Should the project not proceed, then current status quo with regard the environment would remain unchanged. Overall, the area is largely in a natural	1	3	2	1	2	2	18	-	LOW (-ve)	Improve current grazing management, although this is occurring within the surrounding conservation areas and / or areas that	1	3	2	1	2	2	18	-	LOW (-ve)

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION								
		E	P	R	L	D	I / M	TOTAL	STATUS	S		E	P	R	L	D	I / M	TOTAL	STATUS	S
	state. But present day impacts do occur in localised areas and include the following: <ul style="list-style-type: none"> • Increase in unpalatable flora species due to past grazing activities crossings; • Erosion as a result of road crossings; • Several farm dams; and • Undersized culverts within the existing road crossings. 										are used for any hunting / game farming. Improve the current stormwater and energy dissipation features not currently found along the tracks and roads within the region. Install properly sized culverts with erosion protection measures at the existing road / track crossings									

5.4 Cumulative Impacts

A cumulative impact assessment was conducted by assessing this project in relation to any other proposed projects within a 35km radius, as shown in Figure 13.

The report author has been involved in the assessment of all the listed projects within the exception of the Kwagga & Jessa projects. However, all of the reports were review and these are based on the premise that all layouts were developed with impact avoidance in mind, with particular reference to the avoidance of Very High & High Sensitivity areas.

Consequently, all the impacts that remain could be mitigated mostly through revegetation and / or proper stormwater management. Thus, all the impacts would be Medium to Low depending on the scale of the sites, but found acceptable.




Figure 13: Renewable energy projects within a 35km radius

Table 5: Rating of cumulative impacts

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION										RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION									
		E	P	R	L	D	I/M	TOTAL	STATUS	S	E		P	R	L	D	I/M	TOTAL	STATUS	S		
Cumulative Phase																						
Cumulative Impact of various proposed wind farms and associated grid lines on the natural environment	The cumulative assessment considers the various proposed renewable projects that occur within a 35km radius of this site, where the author has either been involved in the assessment of these projects and / or review of the past assessments as part of any required Water Use Licenses.	1	1	1	1	1	1	5	-	LOW (-ve)	The premise of all the reviewed or assessed projects has been the avoidance of impacts on the Very High Sensitivity environments, which have been achieved by the various proposed layouts. The only remaining impacts will be the crossing of internal roads and MV overhead powerlines over minor watercourse / drainage lines or areas rated as LOW sensitivity.	1	3	2	1	2	2	18	-	LOW (-ve)		

5.5 Overall Impact Rating

Table 6: Overall Impact Significance for the WEF (Pre- and Post-Mitigation)

Nature of impact and Phase	Overall Impact Significance (Pre - Mitigation)	Proposed mitigation	Overall Impact Significance (Post - Mitigation)
Construction Phase			
Impact 1: Loss of species of special concern	Low	<p>Develop and implement a Rehabilitation and Monitoring plan post Environmental Authorisation. This must be developed following the finalisation of the turbine / road layout and a walk down has been completed. This plan should include relocation of suitable plant species, but more importantly protect any topsoil stores and promote the collection of vegetative material and propagules / seed to assist with the revegetation of the site.</p> <p>Where possible, temporary construction lay-down or assembly areas should be sited on transformed areas; and</p> <p>Rapid regeneration of plant cover must be encouraged by setting aside topsoil during earthmoving and replacing onto areas where the re- establishment of plant cover is desirable to prevent erosion.</p>	Low
Impact 2: Loss of terrestrial habitats – flora and vegetation	Medium	<p>A pre-construction walkthrough by the ecologist is recommended, who can assist with the development of the Rehabilitation and Monitoring plan, coupled to micro-siting of the final layout. All alien plant re-growth, which is currently low within the greater region must be monitored and should it occur, these plants must be eradicated within the project footprints.</p> <p>Where possible, temporary construction lay-down or assembly areas should be sited on transformed areas; and</p> <p>Rapid regeneration of plant cover must be encouraged by setting aside topsoil during earthmoving and replacing onto areas where the re- establishment of plant cover is desirable to prevent erosion.</p>	Low

Impact 3: Loss of terrestrial species - fauna	Medium	<p>Clear demarcation during the construction phase of all undisturbed sensitive areas that are not within the direct footprint of the WEF to ensure that there is no uncontrolled access by construction vehicles and labourers;</p> <p>Educate contractors as to the importance of the undisturbed conservations areas and importance of avoiding them;</p> <p>All vehicles must stick to designated and prepared roads and adhere to the speed limit on site of 40km/hr;</p> <p>Mitigating the risk of poaching by fencing in the accommodation compounds of the construction crews, to prevent individuals from wandering in the veld after hours; banning the possession of dogs on site by construction and maintenance staff.</p>	Low
Operation Phase			
Impact 4: Loss of terrestrial species - fauna	Medium	<p>Clear demarcation during the construction phase of all undisturbed sensitive areas that are not within the direct footprint of the WEF to ensure that there is no uncontrolled access by construction vehicles and labourers;</p> <p>Educate contractors as to the importance of the undisturbed conservations areas and importance of avoiding them;</p> <p>All vehicles must stick to designated and prepared roads and adhere to the speed limit on site of 40km/hr;</p> <p>Mitigating the risk of poaching by fencing in the accommodation compounds of the construction crews, to prevent individuals from wandering in the veld after hours; banning the possession of dogs on site by construction and maintenance staff.</p>	Low
No-Go	Low	Improve current grazing management, although this is occurring within the surrounding conservation areas and or areas that are used for any hunting / game farming.	Low

		<p>Improve the current stormwater and energy dissipation features not currently found along the tracks and roads within the region.</p> <p>Install properly sized culverts with erosion protection measures at the existing road / track crossings</p>	
Cumulative Impacts	Low	<p>The premise of all the reviewed or assessed projects has been the avoidance of impacts on the Very High Sensitivity environments, which have been achieved by the various proposed layouts. The only remaining impacts will be the crossing of internal roads and overhead powerlines over minor watercourse / drainage lines or areas rated as LOW sensitivity.</p>	Low

6. COMPARATIVE ASSESSMENT OF ALTERNATIVES

Key

PREFERRED	The alternative will result in a low impact / reduce the impact / result in a positive impact
FAVOURABLE	The impact will be relatively insignificant
LEAST PREFERRED	The alternative will result in a high impact / increase the impact
NO PREFERENCE	The alternative will result in equal impacts

6.1 Wind Energy Facility

Table 7: Comparative assessment of WEF components

Alternative	Preference	Reasons (incl. potential issues)
SUBSTATION SITE ALTERNATIVES		
Substation Option 1	Impacts on an aquatic system	With minor layout adjustment, the water course can be avoided
Substation Option 2	Impacts on an aquatic system	Need significant adjustment to avoid the system (See aquatic report)

6.2 No-Go Alternative

Should the project not proceed, then current status quo with regard the aquatic environment would remain unchanged.

Land owners should undertake the following:

- Improve grazing management practices
- Improve the current stormwater and energy dissipation features not currently found along some of the tracks and roads within the region
- Install properly sized culverts or drifts with erosion protection measures at the present road / track crossings

7. CONCLUSION AND SUMMARY

7.1 Summary of Findings

The project overall has a small footprint spread out over a large area, allowing for retention of much of the natural environment so that the systems should remain largely unaffected. Therefore, the wind farm is such that it carries a low intensity impact on terrestrial resources but requiring the clearing of areas with terrestrial vegetation within the footprint of the wind farm.

A variety of environmental features were observed within the study areas and these were mapped and buffered as necessary for their protection. The current layout has, to a large degree, avoided the most sensitive features and buffer areas, greatly reducing the potential overall impact and environmental risk. Noting that these are mostly linked to the CBA/ESAs which are directly linked to the aquatic environment (alluvial rivers and watercourses) that dominate the majority of the site.

However several sensitive species were observed within the site, which included both plants and reptiles, the former being found throughout the site, while the later are highly mobile, thus core sensitive areas could not be mapped as these species are thus encountered throughout the site. Thus once the layout for roads, hardstands and buildings has been refined, a micro-siting walkdown must be conducted to ensure that any populations of these sensitive species are avoided. Due care must be carried out during construction period with respect the presence of listed reptiles and an ECO / EO must be present on a daily basis to remove any reptiles such as the Karoo Padloper.

The overall and cumulative impacts, as assessed, are linked to instances where complete avoidance was not possible, or the nature of the activities involve a potential risk. Overall, it is expected that the impact on the environment would be Low (-). Noteworthy areas, that should be avoided, include the Very High Sensitivity areas as shown in this report.

7.2 Conclusion and Impact Statement

Based on the findings of this study, the specialist finds no reason to withhold to an authorisation for any of the proposed activities, assuming that key mitigations measures are implemented. Due consideration must be given to all the substations (Option 1 and 2) due to being located within High Sensitivity watercourses in particular option 1(see Aquatic report).

However, this must all still be assessed once the roads layout has been provided, coupled to a micro-siting walkdown once all information is available

8. REFERENCES

- GBIF.org (07 December 2021) GBIF Occurrence Download <https://doi.org/10.15468/dl.hgt8qf>
- Agenda 21 – Action plan for sustainable development of the Department of Environmental Affairs and Tourism (DEAT) 1998.
- Agricultural Resources Act, 1983 (Act No. 43 of 1983).
- Davies, B. and Day J., (1998). *Vanishing Waters*. University of Cape Town Press.
- Department of Water Affairs and Forestry - DWAF (2005). *A practical field procedure for identification and delineation of wetland and riparian areas Edition 1*. Department of Water Affairs and Forestry , Pretoria.
- Department of Water Affairs and Forestry - DWAF (2008). *Manual for the assessment of a Wetland Index of Habitat Integrity for South African floodplain and channelled valley bottom wetland types* by M. Rountree (ed); C.P. Todd, C. J. Kleynhans, A. L. Batchelor, M. D. Louw, D. Kotze, D. Walters, S. Schroeder, P. Illgner, M. Uys. and G.C. Marneweck. Report no. N/0000/00/WEI/0407. Resource Quality Services, Department of Water Affairs and Forestry, Pretoria, South Africa.
- Driver A., Sink, K.J., Nel, J.N., Holness, S., Van Niekerk, L., Daniels, F., Jonas, Z., Majiedt, P.A., Harris, L. & Maze, K. 2012. *National Biodiversity Assessment 2011: An assessment of South Africa's biodiversity and ecosystems*. Synthesis Report. South African National Biodiversity Institute and Department of Environmental Affairs, Pretoria.
- Du Preez, L. And Carruthers, V. 2009. *A Complete Guide To Frogs Of Southern Africa*. Struik Nature, Cape Town
- Ewart-Smith J.L., Ollis D.J., Day J.A. and Malan H.L. (2006). *National Wetland Inventory: Development of a Wetland Classification System for South Africa*. WRC Report No. KV 174/06. Water Research Commission, Pretoria.
- IUCN (2019). *Red List of Threatened Species*. IUCN Species Survival Commission, Cambridge Available: <http://www.iucnredlist.org/>
- Kleynhans C.J., Thirion C. and Moolman J. (2005). *A Level 1 Ecoregion Classification System for South Africa, Lesotho and Swaziland*. Report No. N/0000/00/REQ0104. Resource Quality Services, Department of Water Affairs and Forestry, Pretoria.
- Kotze D.C., Marneweck G.C., Batchelor A.L., Lindley D.S. and Collins N. (2008). *WET-EcoServices A technique for rapidly assessing ecosystem services supplied by wetlands*. WRC Report No: TT 339/08.
- Macfarlane, D.M. & Bredin, I.P. 2017. *Buffer Zone Guidelines for Rivers, Wetlands and Estuaries Buffer Zone Guidelines for Rivers, Wetlands and Estuaries*. WRC Report No TT 715/1/17 Water Research Commission, Pretoria.
- Minerals and Petroleum Resources Development Act, 2002 (Act No. 28 of 2002), as amended.
- Mitsch, J.G. and Gosselink, G. (2000). *Wetlands 3rd End*, Wiley, NewYork, 2000, 920 pg.
- Mucina, L., & Rutherford, M.C., 2006. *The Vegetation of South Africa, Lesotho and Swaziland*, Strelitzia 19, South Africa.

National Environmental Management Act, 1998 (Act No. 107 of 1998), as amended.

National Water Act, 1998 (Act No. 36 of 1998), as amended

Nel, J., Maree, G., Roux, D., Moolman, J., Kleynhans, N., Silberbauer, M. and Driver, A. 2004. South African National Spatial Biodiversity Assessment 2004: Technical Report. Volume 2: River Component. CSIR Report Number ENV-S-I-2004-063. Council for Scientific and Industrial Research, Stellenbosch.

Nel, J.L., Murray, K.M., Maherry, A.M., Petersen, C.P., Roux, D.J., Driver, A., Hill, L., Van Deventer, H., Funke, N., Swartz, E.R., Smith-Adao, L.B., Mbona, N., Downsborough, L. and Nienaber, S. (2011). Technical Report for the National Freshwater Ecosystem Priority Areas project. WRC Report No. K5/1801.

Nel, J., Colvin, C., Le Maitre, D., Smith, J. & Haines, I. (2013). South Africa's Strategic Water Source Areas. CSIR Report No: CSIR/NRE/ECOS/ER/2013/0031/A. Report for WWF South Africa

Ollis, D.J., Snaddon, C.D., Job, N.M. & Mbona, N. 2013. Classification System for Wetlands and other Aquatic Ecosystems in South Africa. User Manual: Inland Systems. SANBI Biodiversity Series 22. South African National Biodiversity Institute, Pretoria.

Parsons R. (2004). Surface Water – Groundwater Interaction in a Southern African Context. WRC Report TT 218/03, Pretoria.

Ramsar Convention, (1971) including the Wetland Conservation Programme (DEAT) and the National Wetland Rehabilitation Initiative (DEAT, 2000).

Rowntree, K., Wadesone, R. and O'Keeffe, J. 2000. The development of a geomorphological classification system for the longitudinal zonation of South African rivers. South African Geographical Journal 82(3): 163-172.

South African Bird Atlasing Project 2 (SABAP2). 2017. Animal Demographic Unit. Available online: <http://sabap2.adu.org.za/>

Stuart, C and Stuart, T. 2007. A field guide to the mammals of Southern Africa. Struik Nature, Cape Town.

van Deventer H., Smith-Adao, L. Petersen C., Mbona N., Skowno A., Nel, J.L. (2020) Review of available data for a South African Inventory of Inland Aquatic Ecosystems (SAIIAE). Water SA 44 (2) 184-199

Appendix 1 Specialist CV

CURRICULUM VITAE
Dr Brian Michael Colloty
7212215031083

1 Rossini Rd
Pari Park
Gqeberha, 6070
083 498 3299

Profession: Ecologist & Environmental Assessment Practitioner (Pr. Sci. Nat. 400268/07)
Member of the South African Wetland Society
Specialisation: Ecology and conservation importance rating of inland habitats, wetlands, rivers & estuaries
Years experience: 25 years

SKILLS BASE AND CORE COMPETENCIES

- 25 years experience in environmental sensitivity and conservation assessment of aquatic and terrestrial systems inclusive of Index of Habitat Integrity (IHI), WET Tools, Riparian Vegetation Response Assessment Index (VEGRAI) for Reserve Determinations, estuarine and wetland delineation throughout Africa. Experience also includes biodiversity and ecological assessments with regard sensitive fauna and flora, within the marine, coastal and inland environments. Countries include Mozambique, Kenya, Namibia, Central African Republic, Zambia, Eritrea, Mauritius, Madagascar, Angola, Ghana, Guinea-Bissau and Sierra Leone. Current projects also span all nine provinces in South Africa.
- 15 years experience in the coordination and management of multi-disciplinary teams, such as specialist teams for small to large scale EIAs and environmental monitoring programmes, throughout Africa and inclusive of marine, coastal and inland systems. This includes project and budget management, specialist team management, client and stakeholder engagement and project reporting.
- GIS mapping and sensitivity analysis

TERTIARY EDUCATION

- 1994: B Sc Degree (Botany & Zoology) - NMU
- 1995: B Sc Hon (Zoology) - NMU
- 1996: M Sc (Botany - Rivers) - NMU
- 2000: Ph D (Botany – Estuaries & Mangroves) – NMU

EMPLOYMENT HISTORY

- 1996 – 2000 Researcher at Nelson Mandela University – SAB institute for Coastal Research & Management. Funded by the WRC to develop estuarine importance rating methods for South African Estuaries
- 2001 – January 2003 Training development officer AVK SA (reason for leaving – sought work back in the environmental field rather than engineering sector)
- February 2003- June 2005 Project manager & Ecologist for Strategic Environmental Focus (Pretoria) – (reason for leaving – sought work related more to experience in the coastal environment)
- July 2005 – June 2009 Principal Environmental Consultant Coastal & Environmental Services (reason for leaving – company restructuring)
- June 2009 – August 2018 Owner / Ecologist of Scherman Colloty & Associates cc
- August 2018 Owner / Ecologist - EnviroSci (Pty) Ltd

SELECTED RELEVANT PROJECT EXPERIENCE

World Bank IFC Standards

- Botswana South Africa 400kv transmission line (400km) biodiversity assessment on behalf of Aurecon - current
- Farim phosphate mine and port development, Guinea Bissau – biodiversity and estuarine assessment on behalf of Knight Piesold Canada – 2016.
- Tema LNG offshore pipeline EIA – marine and estuarine assessment for Quantum Power (2015).
- Colluli Potash South Boulder, Eritrea, SEIA marine baseline and hydrodynamic surveys co-ordinator and coastal vegetation specialist (coastal lagoon and marine) (on-going).
- Wetland, estuarine and riverine assessment for Addax Biofuels Sierra Leone, Makeni for Coastal & Environmental Services: 2009
- ESHIA Project manager and long-term marine monitoring phase coordinator with regards the dredge works required in Luanda bay, Angola. Monitoring included water quality and biological changes in the bay and at the offshore disposal outfall site, 2005-2011

South African

- Plant and animal search and rescue for the Karusa and Soetwater Wind Farms on behalf of Enel Green Power, Current
- Plant and animal search and rescue for the Nxuba, Oyster Bay and Garob Wind Farms on behalf of Enel Green Power, 2018 - 2019
- Plant and Animal Search and Rescue for the Port of Ngqura, Transnet Landside infrastructure Project, with development and management of on site nursery, Current
- Plant and Animal Search and Rescue for the Port of Ngqura, OTGC Tank Farm Project (2019)
- Plant search and rescue, for NMBM (Driftsands sewer, Glen Hurd Drive), Department of Social Development (Military veterans housing, Despatch) and Nxuba Wind Farm, - current
- Wetland specialist appointed to update the Eastern Cape Biodiversity Conservation Plan, for the Province on behalf of EOH CES appointment by SANBI – current. This includes updating the National Wetland Inventory for the province, submitting the new data to CSIR/SANBI.
- CDC IDZ Alien eradication plans for three renewable projects Coega Wind Farm, Sonop Wind Farm and Coega PV, on behalf of JG Afrika (2016 – 2017).
- Nelson Mandela Bay Municipality Baakens River Integrated Wetland Assessment (Inclusive of Rehabilitation and Monitoring Plans) for CEN IEM Unit - Current
- Rangers Biomass Gasification Project (Uitenhage), biodiversity and wetland assessment and wetland rehabilitation / monitoring plans for CEM IEM Unit – 2017
- Gibson Bay Wind Farm implementation of the wetland management plan during the construction and operation of the wind farm (includes surface / groundwater as well wetland rehabilitation & monitoring plan) on behalf of Enel Green Power - 2018
- Gibson Bay Wind Farm 133kV Transmission Line wetland management plan during the construction of the transmission line (includes wetland rehabilitation & monitoring plan) on behalf of Eskom – 2016.
- Tsitsikamma Community Wind Farm implementation of the wetland management plan during the construction of the wind farm (includes surface / biomonitoring, as well wetland rehabilitation & monitoring plan) on behalf of Cennergi – completed May 2016.
- Alicedale bulk sewer pipeline for Cacadu District, wetland and water quality assessment, 2016
- Mogalakwena 33kv transmission line in the Limpopo Province, on behalf of Aurecon, 2016
- Cape St Francis WWTW expansion wetland and passive treatment system for the Kouga Municipality, 2015
- Macindane bulk water and sewer pipelines wetland and wetland rehabilitation plan 2015
- Eskom Prieska to Copperton 132kV transmission line aquatic assessment, Northern Cape on behalf of Savannah Environmental 2015.
- Joe Slovo sewer pipeline upgrade wetland assessment for Nelson Mandela Bay Municipality 2014
- Cape Recife Waste Water Treatment Works expansion and pipeline aquatic assessment for Nelson Mandela Bay Municipality 2013
- Pola park bulk sewer line upgrade aquatic assessment for Nelson Mandela Bay Municipality 2013
- Transnet Freight Rail – Swazi Rail Link (Current) wetland and ecological assessment on behalf of Aurecon for the proposed rail upgrade from Ermelo to Richards Bay
- Eskom Transmission wetland and ecological assessment for the proposed transmission line between Pietermaritzburg and Richards Bay on behalf of Aurecon (2012).
- Port Durnford Exarro Sands biodiversity assessment for the proposed mineral sands mine on behalf of Exxaro (2009)
- Fairbreeze Mine Exxaro (Mtunzini) wetland assessment on behalf of Strategic Environmental Services (2007).
- Wetland assessment for Richards Bay Minerals (2013) – Zulti North haul road on behalf of RBM.
- Biodiversity and aquatic assessments for 118 renewable projects in the past 9 years in the Western, Eastern, Northern Cape, KwaZulu-Natal and Free State provinces. Clients included RES-SA, Red Cap, ACED Renewables, Mainstream Renewable, GDF Suez, Globeleq, ENEL, Abengoa amongst others. Particular aquatic sensitivity assessment and Water Use License Applications on behalf of Mainstream Renewable Energy (8 wind farms and 3 PV facilities.), Cennergi / Exxaro (2 Wind farms), WKN Wind current (2 wind farms & 2 PV facilities), ACED (6 wind farms) and Windlab (3 Wind farms) were also conducted. Several of these projects also required the assessment of the proposed transmission lines and switching stations, which were conducted on behalf of Eskom.
- Vegetation assessments on the Great Brak rivers for Department of Water and Sanitation, 2006 and the Gouritz Water Management Area (2014)
- Proposed FibreCo fibre optic cable vegetation assessment along the PE to George, George to Graaf Reinet, PE to Colesburg, and East London to Bloemfontein on behalf of SRK (2013-2015).

Appendix 2 – Site Verification Report

SITE SENSITIVITY VERIFICATION (IN TERMS OF PART A OF THE ASSESSMENT PROTOCOLS PUBLISHED IN GN 320 ON 20 MARCH 2020

INTRODUCTION

In accordance with Appendix 6 of the National Environmental Management Act (Act 107 of 1998, as amended) (NEMA) Environmental Impact Assessment (EIA) Regulations of 2014, a site sensitivity verification has been undertaken in order to confirm the current land use and environmental sensitivity of the proposed project area as identified by the National Web-Based Environmental Screening Tool (Screening Tool).

SITE SENSITIVITY VERIFICATION

Using the result of the specialist ecological impact assessment, that made use of past and current spatial databases, aerial images and field work conducted within and adjacent to the site over a number of years / seasons, various habitats were delineated and the rated in terms of their sensitivity.

OUTCOME OF SITE SENSITIVITY VERIFICATION

Similar to the results of the Screening Tool, the study area contained three types of sensitivity, namely Very High Medium and Low (Figure 1-3). However, the extent of the Very High Sensitivity areas was found be greater in extent as shown in Figure 4.

NATIONAL ENVIRONMENTAL SCREENING TOOL

Based on the DFFE Screening Tool, the site contains areas of very high sensitivity due to the presence of CBAs, Ecological Support Areas and NFEPAS (Terrestrial Theme). The remaining area within the development footprint is deemed to be of Medium (Animals & Plants) or Low sensitivity (Figure 1-3).




Figure 1. DFFE Screening Tool outcome for the animal biodiversity theme




Figure 2. DFFE Screening Tool outcome for the Plant biodiversity theme




Figure 3. DFFE Screening Tool outcome for the Terrestrial biodiversity theme

Figure 4 below shows the sensitivity map produced following the ecological assessment as well as a ground-truthing exercises, with mapping of the observed features at a finer scale.




Figure 4. Environmental sensitivity map produced by the aquatic specialist

CONCLUSION

In conclusion, the DFFE Screening Tool identified three sensitivity ratings within the development study area, very high, medium and low. Although there is some overlap with the findings on site and the Screening Tool's outcome, the extent of the Very High sensitivity areas was found to be greater than the extent in the Screening Tool.

However and appropriate layout can be developed to minimise the impact on the Very High areas, but must be verified once the final layout inclusive of roads has been developed.

Appendix 3 – Species checklists

Species					Red list
#	code	Family	Scientific name	Common name	category
1	310	Bufonidae	<i>Vandijkophrynus gariepensis gariepensis</i>	Karoo Toad (subsp. gariepensis)	
2	890	Pyxicephalidae	<i>Amietia fuscigula</i>	Cape River Frog	Least Concern (2017)
3	400	Pyxicephalidae	<i>Cacosternum boettgeri</i>	Common Caco	Least Concern (2013)
4	850	Pyxicephalidae	<i>Pyxicephalus adspersus</i>	Giant Bull Frog	Near Threatened
5	1000	Pyxicephalidae	<i>Tomopterna delalandii</i>	Cape Sand Frog	Least Concern

Species					Red list
#	code	Family	Scientific name	Common name	category
1	212190	Bovidae	<i>Antidorcas marsupialis</i>	Springbok	Least Concern (2016)
2	213320	Bovidae	<i>Raphicerus campestris</i>	Steenbok	Least Concern (2016)
3	127750	Gliridae	<i>Graphiurus (Graphiurus) ocularis</i>	Spectacled African Dormouse	Least Concern
4	197770	Hyaenidae	<i>Proteles cristata</i>	Aardwolf	Least Concern (2016)
5	158240	Leporidae	<i>Lepus saxatilis</i>	Scrub Hare	Least Concern
6	144330	Muridae	<i>Desmodillus auricularis</i>	Cape Short-tailed Gerbil	Least Concern (2016)
7	151210	Muridae	<i>Parotomys brantsii</i>	Brants's Whistling Rat	Least Concern (2016)
8	195120	Viverridae	<i>Genetta genetta</i>	Common Genet	Least Concern (2016)

Species					Red list
#	code	Family	Scientific name	Common name	category
1	1450	Agamidae	<i>Agama aculeata aculeata</i>	Common Ground Agama	Least Concern (SARCA 2014)
2	3050	Cordylidae	<i>Karusasaurus polyzonus</i>	Karoo Girdled Lizard	Least Concern (SARCA 2014)
3	5340	Elapidae	<i>Naja nivea</i>	Cape Cobra	Least Concern (SARCA 2014)
4	202	Gekkonidae	<i>Chondrodactylus angulifer</i>	Giant Ground Gecko	Least Concern (IUCN 2009)
5	480	Gekkonidae	<i>Chondrodactylus bibronii</i>	Bibron's Gecko	Least Concern (SARCA 2014)
6	620	Gekkonidae	<i>Pachydactylus latirostris</i>	Quartz Gecko	Least Concern (SARCA 2014)
7	600	Gekkonidae	<i>Pachydactylus maculatus</i>	Spotted Gecko	Least Concern (SARCA 2014)
8	610	Gekkonidae	<i>Pachydactylus mariquensis</i>	Marico Gecko	Least Concern (SARCA 2014)
9	1890	Lacertidae	<i>Pedioplanis lineoocellata pulchella</i>	Common Sand Lizard	Least Concern (SARCA 2014)
10	1900	Lacertidae	<i>Pedioplanis namaquensis</i>	Namaqua Sand Lizard	Least Concern (SARCA 2014)

11	5781	Pelomedusidae	<i>Pelomedusa galeata</i>	South African Marsh Terrapin	Not evaluated
12	2470	Scincidae	<i>Trachylepis sulcata sulcata</i>	Western Rock Skink	Least Concern (SARCA 2014)
13	5530	Testudinidae	<i>Chersina angulata</i>	Angulate Tortoise	Least Concern (SARCA 2014)
14	5691	Testudinidae	<i>Psammobates tentorius subsp. ?</i>	Tent Tortoise (subsp. ?)	Least Concern (SARCA 2014)
15	5670	Testudinidae	<i>Psammobates tentorius tentorius</i>	Karoo Tent Tortoise	
16	5540	Testudinidae	<i>Stigmochelys pardalis</i>	Leopard Tortoise	Least Concern (SARCA 2014)

Species

#	code	Family	Scientific name	Common name	Red list category
---	------	--------	-----------------	-------------	-------------------

1	668420	Libellulidae	<i>Sympetrum fonscolombii</i>	Red-veined Darter or Nomad	LC
---	--------	--------------	-------------------------------	----------------------------	----

Species

#	code	Family	Scientific name	Common name	Red list category
---	------	--------	-----------------	-------------	-------------------

1	620400	CRAMBIDAE	<i>Loxostege frustalis</i>		
2	457090	LYCAENIDAE	<i>Chrysoritis chrysaor</i>	Burnished opal	Least Concern (SABCA 2013)
3	438050	NYMPHALIDAE	<i>Vanessa cardui</i>	Painted lady	Least Concern (SABCA 2013)
4	407450	PIERIDAE	<i>Belenois aurota</i>	Pioneer caper white	Least Concern (SABCA 2013)
5	405610	PIERIDAE	<i>Pontia helice helice</i>	Southern meadow white	Least Concern (SABCA 2013)

Appendix 4: Detailed aquatic assessment methodology

This study followed the approaches of several national guidelines with regards to wetland assessment. These have been modified by the author, to provide a relevant mechanism of assessing the present state of the study area aquatic systems, applicable to the specific environment and, in a clear and objective manner, identify and assess the potential impacts associated with the proposed development site based on information collected within the relevant farm portions.

Current water resource classification systems make use of the Hydrogeomorphic (HGM) approach, and for this reason, the National Wetland Classification System (NWCS) approach will be used in this study. It is also important to understand the legal definition of a wetland, the means of assessing wetland conservation and importance and the relevant legislation aimed at protecting wetlands. These aspects will be discussed in greater depth in this section of the report, as they form the basis of the study approach to assessing wetland impacts.

For reference the following definitions are as follows:

- **Drainage line:** A drainage line is a lower category or order of watercourse that does not have a clearly defined bed or bank. It carries water only during or immediately after periods of heavy rainfall i.e. non-perennial, and riparian vegetation may not be present.
- **Perennial and non-perennial:** Perennial systems contain flow or standing water for all or a large proportion of any given year, while non-perennial systems are episodic or ephemeral and thus contains flows for short periods, such as a few hours or days in the case of drainage lines.
- **Riparian:** The area of land adjacent to a stream or river that is influenced by stream-induced or related processes. Riparian areas which are saturated or flooded for prolonged periods would be considered wetlands and could be described as riparian wetlands. However, some riparian areas are not wetlands (e.g. an area where alluvium is periodically deposited by a stream during floods but which is well drained).
- **Wetland:** Land which is transitional between terrestrial and aquatic systems where the water table is usually at or near the surface, or the land is periodically covered with shallow water, and which under normal circumstances supports or would support vegetation typically adapted to life in saturated soil (Water Act 36 of 1998); land where an excess of water is the dominant factor determining the nature of the soil development and the types of plants and animals living at the soil surface (Cowardin *et al.*, 1979).
- **Water course:** As per the National Water Act means -

(a) a river or spring;

(b) a natural channel in which water flows regularly or intermittently;

(c) a wetland, lake or dam into which, or from which, water flows; and

(d) any collection of water which the Minister may, by notice in the Gazette, declare to be a watercourse, and a reference to a watercourse includes, where relevant, its bed and banks

8.1 Waterbody classification systems

Since the late 1960's, wetland classification systems have undergone a series of international and national revisions. These revisions allowed for the inclusion of additional wetland types, ecological and conservation rating metrics, together with a need for a system that would allude to the functional requirements of any given wetland (Ewart-Smith *et al.*, 2006). Wetland function is a consequence of biotic and abiotic factors, and wetland classification should strive to capture these aspects.

Coupled to this was the inclusion of other criteria within the classification systems to differentiate between river, riparian and wetland systems, as well as natural versus artificial waterbodies.

The South African National Biodiversity Institute (SANBI) in collaboration with several specialists and stakeholders developed the newly revised and now accepted National Wetland Classification Systems (NWCS) (Ollis *et al.*, 2013). This system comprises a hierarchical classification process of defining a wetland based on the principles of the hydrogeomorphic (HGM) approach at higher levels, with including structural features at the finer or lower levels of classification (Ollis *et al.*, 2013).

Wetlands develop in a response to elevated water tables, linked either to rivers, groundwater flows or seepage from aquifers (Parsons, 2004). These water levels or flows then interact with localised geology and soil forms, which then determines the form and function of the respective wetlands. Water is thus the common driving force, in the formation of wetlands (DWA, 2005). It is significant that the HGM approach has now been included in the wetland classifications as the HGM approach has been adopted throughout the water resources management realm with regards to the determination of the Present Ecological State (PES) and Ecological Importance and Sensitivity (EIS) and WET-Health assessments for aquatic environments. All these systems are then easily integrated using the HGM approach in line with the Eco-classification process of river and wetland reserve determinations used by the Department of Water and Sanitation (DWS). The Ecological Reserve of a wetland or river is used by DWS to assess the water resource allocations when assessing WULAs

The NWCS process is provided in more detail in the methods section of the report, but some of the terms and definitions used in this document are present below:

Definition Box

SiVEST Environmental Aquatic Impact Assessment Version No. 1
--

Prepared by: EnviroSci (Pty) Ltd

Date: 20 November 2022

Present Ecological State is a term for the current ecological condition of the resource. This is assessed relative to the deviation from the Reference State. Reference State/Condition is the natural or pre-impacted condition of the system. The reference state is not a static condition, but refers to the natural dynamics (range and rates of change or flux) prior to development. The PES is determined per component - for rivers and wetlands this would be for the drivers: flow, water quality and geomorphology; and the biotic response indicators: fish, macroinvertebrates, riparian vegetation and diatoms. PES categories for every component would be integrated into an overall PES for the river reach or wetland being investigated. This integrated PES is called the EcoStatus of the reach or wetland.

EcoStatus is the overall PES or current state of the resource. It represents the totality of the features and characteristics of a river and its riparian areas or wetland that bear upon its ability to support an appropriate natural flora and fauna and its capacity to provide a variety of goods and services. The EcoStatus value is an integrated ecological state made up of a combination of various PES findings from component EcoStatus assessments (such as for invertebrates, fish, riparian vegetation, geomorphology, hydrology, and water quality).

Reserve: The quantity and quality of water needed to sustain basic *human needs* and *ecosystems* (e.g. estuaries, rivers, lakes, groundwater and wetlands) to ensure ecologically sustainable development and utilisation of a water resource. The *Ecological Reserve* pertains specifically to aquatic ecosystems.

Reserve requirements: The quality, quantity and reliability of water needed to satisfy the requirements of basic human needs and the Ecological Reserve (inclusive of instream requirements).

Ecological Reserve determination study: The study undertaken to determine Ecological Reserve requirements.

Licensing applications: Water users are required (by legislation) to apply for licenses prior to extracting water resources from a water catchment or any other activity that qualifies as a water use.

Ecological Water Requirements: This is the quality and quantity of water flowing through a natural stream course that is needed to sustain instream functions and ecosystem integrity at an acceptable level as determined during an EWR study. These then form part of the conditions for managing achievable water quantity and quality conditions as stipulated in the **Reserve Template**

Water allocation process (compulsory licensing): This is a process where all existing and new water users are requested to reapply for their licenses, particularly in stressed catchments where there is an over-allocation of water or an inequitable distribution of entitlements.

Ecoregions are geographic regions that have been delineated in a top-down manner on the basis of physical/abiotic factors. • NOTE: For purposes of the classification system, the 'Level I Ecoregions' for South Africa, Lesotho and Swaziland (Kleynhans *et al.* 2005), which have been specifically developed by the Department of Water Affairs & Forestry (DWAF) for rivers but are used for the management of inland aquatic ecosystems more generally, are applied at Level 2A of the classification system. These Ecoregions are based on physiography, climate, geology, soils and potential natural vegetation.

8.2 Wetland definition

Although the National Wetland Classification System (NWCS) (Ollis *et al.*, 2013) is used to classify wetland types it is still necessary to understand the definition of a wetland. Terminology currently strives to characterise a wetland not only on its structure (visible form), but also to relate this to the function and value of any given wetland.

The Ramsar Convention definition of a wetland is widely accepted as "areas of marsh, fen, peatland or water, whether natural or artificial, permanent or temporary, with water that is static or flowing, fresh, brackish or salt, including areas of marine water the depth of which at low tide does not exceed six metres" (Davis 1994). South Africa is a signatory to the Ramsar Convention and therefore its extremely broad definition of wetlands has been adopted for the proposed NWCS, with a few modifications.

Whereas the Ramsar Convention included marine water to a depth of six metres, the definition used for the NWCS extends to a depth of ten metres at low tide, as this is recognised as the seaward boundary of the shallow photic zone (Lombard *et al.*, 2005). An additional minor adaptation of the definition is the removal of the term 'fen' as fens are considered a type of peatland. The adapted definition for the NWCS is, therefore, as follows (Ollis *et al.*, 2013):

WETLAND: *an area of marsh, peatland or water, whether natural or artificial, permanent or temporary, with water that is static or flowing, fresh, brackish or salt, including areas of marine water the depth of which at low tide does not exceed ten metres.*

This definition encompasses all ecosystems characterised by the permanent or periodic presence of water other than marine waters deeper than ten metres. The only legislated definition of wetlands in South Africa, however, is contained within the National Water Act (Act No. 36 of 1998) (NWA), where wetlands are defined as "land which is transitional between terrestrial and aquatic systems, where the water table is usually at, or near the surface, or the land is periodically covered with shallow water and which land in normal circumstances supports, or would support, vegetation adapted to life in saturated soil." This definition is consistent with more precise working definitions of wetlands and therefore includes only a subset of ecosystems encapsulated in the Ramsar definition. It should be noted that the NWA definition is not concerned with marine systems and clearly distinguishes wetlands from estuaries, classifying the latter as a watercourse (Ollis *et al.*, 2013). Table 1 below provides a comparison of the various wetlands included within the main sources of wetland definitions used in South Africa. Although a subset of Ramsar-defined wetlands was used as a starting point for the compilation of the first version of the National Wetland Inventory (i.e. "wetlands", as defined by the NWA, together with open waterbodies), it is understood that

subsequent versions of the Inventory include the full suite of Ramsar-defined wetlands in order to ensure that South Africa meets its wetland inventory obligations as a signatory to the Convention (Ollis *et al.*, 2013).

Wetlands must therefore have one or more of the following attributes to meet the above definition (DWAF, 2005):

- A high-water table that results in the saturation at or near the surface, leading to anaerobic conditions developing in the top 50 cm of the soil.
- Wetland or hydromorphic soils that display characteristics resulting from prolonged saturation, i.e. mottling or grey soils
- The presence of, at least occasionally, hydrophilic plants, i.e. hydrophytes (water loving plants).

It should be noted that riparian systems that are not permanently or periodically inundated are not considered true wetlands, i.e. those associated with the drainage lines and rivers.

Table 2: Comparison of ecosystems considered to be ‘wetlands’ as defined by the proposed NWCS, the NWA and ecosystems included in DWAF’s (2005) delineation manual.

Ecosystem	NWCS “wetland”	National Water Act wetland	DWAF (2005) delineation manual
Marine	YES	NO	NO
Estuarine	YES	NO	NO
Waterbodies deeper than 2 m (i.e. limnetic habitats often described as lakes or dams)	YES	NO	NO
Rivers, channels and canals	YES	NO ¹	NO
Inland aquatic ecosystems that are not river channels and are less than 2 m deep	YES	YES	YES
Riparian ² areas that are permanently / periodically inundated or saturated with water within 50 cm of the surface	YES	YES	YES ³
Riparian ³ areas that are not permanently / periodically inundated or saturated with water within 50 cm of the surface	NO	NO	YES ³

¹ Although river channels and canals would generally not be regarded as wetlands in terms of the National Water Act, they are included as a ‘watercourse’ in terms of the Act

² According to the National Water Act and Ramsar, riparian areas are those areas that are saturated or flooded for prolonged periods and would be considered riparian wetlands, as opposed to non –wetland riparian areas that are only periodically inundated and the riparian vegetation persists due to having deep root systems drawing on water many meters below the surface.

³ The delineation of ‘riparian areas’ (including both wetland and non-wetland components) is treated separately to the delineation of wetlands in DWAF’s (2005) delineation manual.

8.3 National Wetland Classification System method

Due to the nature of the wetlands and watercourses observed, it was determined that the newly accepted NWCS should be adopted. This classification approach has integrated aspects of the HGM approach used in the WET-Health system as well as the widely accepted eco-classification approach used for rivers.

The NWCS (Ollis *et al.*, 2013) as stated previously, uses hydrological and geomorphological traits to distinguish the primary wetland units, i.e. direct factors that influence wetland function. Other wetland assessment techniques, such as the DWAF (2005) delineation method, only infer wetland function based on abiotic and biotic descriptors (size, soils & vegetation) stemming from the Cowardin approach (Ollis *et al.*, 2013).

The classification system used in this study is thus based on Ollis *et al.* (2013) and is summarised below:

The NWCS has a six-tiered hierarchical structure, with four spatially nested primary levels of classification (Figure 2). The hierarchical system firstly distinguishes between Marine, Estuarine and Inland ecosystems (**Level 1**), based on the degree of connectivity the particular system has with the open ocean (greater than 10 m in depth). Level 2 then categorises the regional wetland setting using a combination of biophysical attributes at the landscape level, which operate at a broad bioregional scale.

This is opposed to specific attributes such as soils and vegetation. **Level 2** has adopted the following systems:

- Inshore bioregions (marine)
- Biogeographic zones (estuaries)
- Ecoregions (Inland)

Level 3 of the NWCS assess the topographical position of inland wetlands as this factor broadly defines certain hydrological characteristics of the inland systems. Four landscape units based on topographical position are used in distinguishing between Inland systems at this level. No subsystems are recognised for Marine systems, but estuaries are grouped according to their periodicity of connection with the marine environment, as this would affect the biotic characteristics of the estuary.

Level 4 classifies the hydrogeomorphic (HGM) units discussed earlier. The HGM units are defined as follows:

- Landform – shape and localised setting of wetland
- Hydrological characteristics – nature of water movement into, through and out of the wetland
- Hydrodynamics – the direction and strength of flow through the wetland

These factors characterise the geomorphological processes within the wetland, such as erosion and deposition, as well as the biogeochemical processes.

Level 5 of the assessment pertains to the classification of the tidal regime within the marine and estuarine environments, while the hydrological and inundation depth classes are determined for inland wetlands. Classes are based on frequency and depth of inundation, which are used to determine the functional unit of the wetlands and are considered secondary discriminators within the NWCS.

Level 6 uses six descriptors to characterise the wetland types based on biophysical features. As with Level 5, these are non-hierarchical in relation to each other and are applied in any order, dependent on the availability of information. The descriptors include:

- Geology;
- Natural vs. Artificial;
- Vegetation cover type;
- Substratum;
- Salinity; and
- Acidity or Alkalinity

It should be noted that where sub-categories exist within the above descriptors, hierarchical systems are employed, and these are thus nested in relation to each other.

The HGM unit (Level 4) is the focal point of the NWCS, with the upper levels (Figure 3 Figure – Inland systems only) providing means to classify the broad bio-geographical context for grouping functional wetland units at the HGM level, while the lower levels provide more descriptive detail on the particular wetland type characteristics of a particular HGM unit. Therefore Level 1 – 5 deals with functional aspects, while Level 6 classifies wetlands on structural aspects.




Figure 2: Basic structure of the NWCS, showing how ‘primary discriminators’ are applied up to Level 4 to classify Hydrogeomorphic (HGM) Units, with ‘secondary discriminators’ applied at Level 5 to classify the tidal/hydrological regime, and ‘descriptors’ applied




Figure 3: Illustration of the conceptual relationship of HGM Units (at Level 4) with higher and lower levels (relative sizes of the boxes show the increasing spatial resolution and level of detail from the higher to the lower levels) for Inland Systems (from Ollis *et al.*, 2013)

8.4 Waterbody condition

To assess the PES or condition of the observed wetlands, a modified Wetland Index of Habitat Integrity (DWAf, 2007) was used. The Wetland Index of Habitat Integrity (WETLAND-IHI) is a tool developed for use in the National Aquatic Ecosystem Health Monitoring Programme (NAEHMP), formerly known as the River Health Programme (RHP). The output scores from the WETLAND-IHI model are presented in the standard DWAf A-F ecological categories (Table) and provide a score of the PES of the habitat integrity of the wetland system being examined. The author has included additional criteria into the model-based system to include additional wetland types. This system is preferred when compared to systems such as WET-Health – wetland management series (WRC 2009), as WET-Health (Level 1) was developed with wetland rehabilitation in mind and is not always suitable for impact assessments. This coupled with the degraded state of the wetlands in the study area, indicated that a complex study approach was not warranted, i.e. conduct a Wet-Health Level 2 and WET-Ecosystems Services study required for an impact assessment.

Table 3: Description of A – F ecological categories based on Kleynhans *et al.*, (2005)

ECOLOGICAL CATEGORY	ECOLOGICAL DESCRIPTION	MANAGEMENT PERSPECTIVE
A	Unmodified, natural.	Protected systems; relatively untouched by human hands; no discharges or impoundments allowed
B	Largely natural with few modifications. A small change in natural habitats and biota may have taken place but the ecosystem functions are essentially unchanged.	Some human-related disturbance, but mostly of low impact potential
C	Moderately modified. Loss and change of natural habitat and biota have occurred, but the basic ecosystem functions are still predominantly unchanged.	Multiple disturbances associated with need for socio-economic development, e.g. impoundment, habitat modification and water quality degradation
D	Largely modified. A large loss of natural habitat, biota and basic ecosystem functions has occurred.	
E	Seriously modified. The loss of natural habitat, biota and basic ecosystem functions is extensive.	Often characterized by high human densities or extensive resource exploitation. Management intervention is needed to improve health, e.g. to restore flow patterns, river habitats or water quality
F	Critically / Extremely modified. Modifications have reached a critical level and the system has been modified completely with an almost complete loss of natural habitat and biota. In the worst instances the basic ecosystem functions have been destroyed and the changes are irreversible.	

The WETLAND-IHI model is composed of four modules. The “Hydrology”, “Geomorphology” and “Water Quality” modules all assess the contemporary driving processes behind wetland formation and maintenance. The last module, “Vegetation Alteration”, provides an indication of the intensity of human land use activities on the wetland surface itself and how these may have modified the condition of the wetland. The integration of the scores from these 4 modules provides an overall PES score for the wetland system being examined. The WETLAND-IHI model is an MS Excel-based model, and the data required for the assessment are generated during a site visit.

Additional data may be obtained from remotely sensed imagery (aerial photos; maps and/or satellite imagery) to assist with the assessment. The interface of the WETLAND-IHI has been developed in a format which is similar to DWA’s River EcoStatus models which are currently used for the assessment of PES in riverine environments.

8.5 Aquatic ecosystem importance and function

South Africa is a Contracting Party to the Ramsar Convention on Wetlands, signed in Ramsar, Iran, in 1971, and has thus committed itself to this intergovernmental treaty, which provides the framework for the national protection of wetlands and

the resources they could provide. Wetland conservation is now driven by the South African National Biodiversity Institute, a requirement under the National Environmental Management: Biodiversity Act (No 10 of 2004).

Wetlands are among the most valuable and productive ecosystems on earth, providing important opportunities for sustainable development (Davies and Day, 1998). However, wetlands in South Africa are still rapidly being lost or degraded through direct human induced pressures (Nel *et al.*, 2004).

The most common attributes or goods and services provided by wetlands include:

- Improve water quality;
- Impede flow and reduce the occurrence of floods;
- Reeds and sedges used in construction and traditional crafts;
- Bulbs and tubers, a source of food and natural medicine;
- Store water and maintain base flow of rivers;
- Trap sediments; and
- Reduce the number of water-borne diseases.

In terms of this study, the wetlands provide ecological (environmental) value to the area acting as refugia for various wetland associated plants, butterflies and birds.

In the past wetland conservation has focused on biodiversity as a means of substantiating the protection of wetland habitat. However not all wetlands provide such motivation for their protection, thus wetland managers and conservationists began assessing the importance of wetland function within an ecosystem.

Table below summarises the importance of wetland function when related to ecosystem services or ecoservices (Kotze *et al.*, 2008). One such example is emergent reed bed wetlands that function as transformers converting inorganic nutrients into organic compounds (Mitsch and Gosselink, 2000).

Table 4: Summary of direct and indirect ecoservices provided by wetlands from Kotze *et al.*, 2008

Ecosystem services supplied by wetlands	<i>Indirect benefits</i>	Hydro-geochemical benefits	Flood attenuation	
			Stream flow regulation	
			Water quality enhancement benefits	Sediment trapping
				Phosphate assimilation
				Nitrate assimilation
				Toxicant assimilation
				Erosion control
	Carbon storage			
	Biodiversity maintenance			
	<i>Direct benefits</i>	<i>Provision of water for human use</i>		
		<i>Provision of harvestable resources²</i>		
		<i>Provision of cultivated foods</i>		
		<i>Cultural significance</i>		
<i>Tourism and recreation</i>				
<i>Education and research</i>				

Conservation importance of the individual wetlands was based on the following criteria:

- Habitat uniqueness;
- Species of conservation concern;
- Habitat fragmentation or rather, continuity or intactness with regards to ecological corridors; and
- Ecosystem service (social and ecological).

The presence of any or a combination of the above criteria would result in a HIGH conservation rating if the wetland was found in a near natural state (high PES). Should any of the habitats be found modified the conservation importance would rate as MEDIUM, unless a Species of Conservation Concern (SCC) was observed, in which case it would receive a HIGH rating. Any system that was highly modified (low PES) or had none of the above criteria, received a LOW conservation

importance rating. Wetlands with HIGH and MEDIUM ratings should thus be excluded from development with incorporation into a suitable open space system, with the maximum possible buffer being applied. Natural wetlands or Wetlands that resemble some form of the past landscape but receive a LOW conservation importance rating could be included into stormwater management features and should not be developed to retain the function of any ecological corridors.