

# AGRICULTURAL COMPLIANCE STATEMENT FOR THE PROPOSED GRASPAN PV DEVELOPMENT

# Witput, Northern Cape Province

November 2021

Client



# Prepared by:

The Biodiversity Company

Cell: +27 81 319 1225
Fax: +27 86 527 1965
info@thebiodiversitycompany.com
www.thebiodiversitycompany.com



Report Name	Agricultural Compliance Statement for the proposed Graspan PV Facility							
Submitted to	Savannah Environmental							
	Andrew Husted	HAX						
Report Reviewer	with a MSc in Zoology. He, is Pr Sci Nat register Science, Environmental Science and Aquatic S Specialist with 12 years' experience in the enviro	ializing in freshwater systems and wetlands, who graduated red (400213/11) in the following fields of practice: Ecological cience. Mr Husted is an Aquatic, Wetland and Biodiversity nmental consulting field. In addition to his ecological working I and soil assessments, this includes the consideration of land						
	Ivan Baker							
Report Writer	wetland and ecosystem service specialist, a hydrogenialist studies ranging from basic assessmen following FC standards. Ivan completed training	in environmental science and geological science. Ivan is a dropedologist and pedologist that has completed numerous ts to EIAs. Ivan has carried out various international studies g in Tools for Wetland Assessments with a certificate of ental science and hydropedology at the North-West University						
Declaration	South African Council for Natural Scientific Profe financial interests in the proponent, other than for Regulations, 2017. We have no conflicting interes secondary developments resulting from the auth	perate as independent consultants under the auspice of the ssions. We declare that we have no affiliation with or vested work performed under the Environmental Impact Assessment sts in the undertaking of this activity and have no interests in norisation of this project. We have no vested interest in the rvice within the constraints of the project (timing, time and						



# **Table of Contents**

1		Intro	duction	1
	1.1	l	Background	1
2		Proje	ect Area	1
3	;	Scop	pe of Work	3
4		Limit	tations	3
5		Ехре	ertise of the Specialists	4
	5.1		Andrew Husted	4
	5.2	2	Ivan Baker	4
6		Liter	ature Review	4
	6.1		Land Capability	4
7		Meth	nodology	5
	7.1	l	Desktop Assessment	5
	7.2	2	Agricultural Potential	
	As	sses	sment	5
	7.3	3	Climate Capability	7
	7.4	1	Current Land Use	8
8		Desl	ktop Findings	8
	8.1		Climate	8
	8.2	2	Soils and Geology	9
	8.3	3	Terrain10	0
9		Resi	ults and Discussion1	2
	9.1	l	Baseline Findings	2
	9.2	2	Sensitivity Verification1	3
1(	)	Impa	act Statement1	4
	10	.1	Loss of Land Capability14	4
	10	.2	Construction Phase	4
	10	.3	Operational Phase14	4
	10	.4	Cumulative Impacts19	5
	10	.5	Specialist Opinion	5
11		Rec	ommendations and Mitigation Measures1	6
	11	.1	General Mitigation	6
	11	.2	Restoration of Vegetation Cover1	6
		11.2	.1 Ripping Compacted Areas1	6
		11.2	.2 Revegetate Degraded Areas10	6
	11	.3	Specialist Recommendation1	7
12	2	Con	clusion1	7
13	3	Refe	erences1	8



# **Figures**

Figure 2-1	Map illustrating the location of the proposed project area	2
Figure 8-1	Climate for the region	9
Figure 8-2	Illustration of the Ag 146 land type terrain units (Land Type Survey Staff, 1972 - 200	)6)9
Figure 8-3	Slope percentage map for the project area	11
Figure 8-4	Elevation of the project area (metres above sea level)	11
Figure 9-1	Example of a shallow Glenrosa soil form with a depth of less than 20 cm	12
Figure 9-2	Land Capability Sensitivity (DAFF, 2017)	13
	Tables	
Table 6-1	Land Capability (DAFF, 2017)	4
Table 7-1	Land capability class and intensity of use (Smith, 2006)	5
Table 7-2	The combination table for land potential classification	6
Table 7-3	The Land Potential Classes.	7
Table 7-4	Climatic capability (step 1) (Scotney et al., 1987)	7
Table 8-1	Soils expected at the respective terrain units within the Ag 146 land type (Land T Survey Staff, 1972 - 2006)	
Table 10-1	Impact assessment related to the loss of land capability during the construction ph of the proposed PV area	
Table 10-2	Impact assessment related to the loss of land capability during the operational phas the proposed PV area	
Table 10-3	Impact assessment related cumulative impacts	15



# **Declaration**

#### I, Ivan Baker declare that:

- I act as the independent specialist in this application;
- I will perform the work relating to the application in an objective manner, even if this results in views and findings that are not favourable to the applicant;
- I declare that there are no circumstances that may compromise my objectivity in performing such work;
- I have expertise in conducting the specialist report relevant to this application, including knowledge of the Act, regulations and any guidelines that have relevance to the proposed activity;
- I will comply with the Act, regulations and all other applicable legislation;
- I have no, and will not engage in, conflicting interests in the undertaking of the activity;
- I undertake to disclose to the applicant and the competent authority all material information in
  my possession that reasonably has or may have the potential of influencing any decision to be
  taken with respect to the application by the competent authority; and the objectivity of any
  report, plan or document to be prepared by myself for submission to the competent authority;
- all the particulars furnished by me in this form are true and correct; and
- I realise that a false declaration is an offence in terms of Regulation 71 and is punishable in terms of Section 24F of the Act.

D

Ivan Baker

**Soil Specialist** 

The Biodiversity Company

November 2021



## 1 Introduction

The Biodiversity Company was appointed to conduct a pedology (agricultural potential, land capability and land use) baseline and impact assessment for the establishment of a solar photovoltaic (PV) project, namely Graspan Solar PV. A site assessment was conducted during November 2021.

ENGIE Graspan Solar Project (Pty) Ltd obtained an Environmental Authorisation for the proposed Graspan PV Facility and associated infrastructure, located on remaining extent of Farm Graspan (No. 172), situated in the Siyancuma Local Municipality in the Northern Cape province in April 2013 (DFFE Reference No.: 14/12/16/3/3/276). The project has been selected as a Preferred Bidder project under Round 5 of the Renewable Energy Independent Power Producers Procurement Programme (REIPPPP).

The proposed facility will have a contracted capacity of 90MW and will include the following infrastructure:

- PV solar panels/modules (arranged in arrays);
- PV module mountings;
- DC-AC current inverters and transformers;
- An on-site 132kV Independent Power Producer (IPP) substation to facilitate the grid connection.
- Underground cabling/ overhead power lines;
- On-site buildings (including an operational control centre, office, ablutions and a guard house);
- Access roads and internal road network; and
- Ancillary infrastructure.

A developmental footprint of 150 ha in extent is authorised for the facility and associated infrastructure. In order to implement the project, an additional 19.9ha is required. This additional area is immediately adjacent to the authorised area.

The approach adopted for the assessment has taken cognisance of the recently published Government Notice 320 in terms of NEMA dated 20 March 2020: "Procedures for the Assessment and Minimum Criteria for Reporting on Identified Environmental Themes in terms of Sections 24(5)(a) and (h) and 44 of the National Environmental Management Act, 1998, when applying for Environmental Authorisation".

This report aims to present and discuss the findings from the soil resources identified on-site, the agricultural and land potential of these resources, the land uses within the project area as well as the risks associated with the proposed PV facility.

#### 1.1 Background

The following specialist report was reviewed and considered to supplement the project findings:

 Soil and agricultural assessment for the proposed Graspan PV Solar Power Facility (SiVest, 2012).

# 2 Project Area

The project is found 23 km north east from Witput and 38 km south of Modderrivier in the Northern Cape Presently, the project area is adjacent to the N12 road and surrounded by some open natural areas and in close proximity to the Driekopspan (Figure 2-1).






Figure 2-1 Map illustrating the location of the proposed project area



# 3 Scope of Work

The following tasks were completed in fulfilment of the terms of reference for this assessment:

- To conduct a soil assessment which includes a description of the physical properties which
  characterise the soil within the proposed area of development of the relevant portions of the
  affected properties;
- Using the findings from the soil assessment to determine the existing land capability/potential and current land use of the entire surface area of the relevant portions of the project area;
- To delineate soil resources;
- To determine the sensitivity of the baseline findings;
- The soil classification was done according to the Taxonomic Soil Classification System for South Africa, 1991. The following attributes must be included at each observation:
  - Soil form and family (Taxonomic Soil Classification System for South Africa, 1991);
  - Soil depth;
  - Estimated soil texture;
  - Soil structure, coarse fragments, calcareousness;
  - Buffer capacities;
  - Underlying material;
  - Current land use; and
  - Land capability.
- To complete an impact statement;
- Discussing the feasibility of the proposed activities;
- Confirmation that no agricultural segregation will take place and that all options have been considered to avoid segregation; and
- Recommend relevant mitigation measures to limit all associated impacts.

# 4 Limitations

The following limitations are relevant to this agricultural compliance statement;

- It has been assumed that the extent of the properties to be assessed together with the locations of the proposed components are correct and final; and
- The handheld GPS used potentially could have inaccuracies up to 5 m. Any and all delineations therefore could be inaccurate within 5 m.



# 5 Expertise of the Specialists

#### 5.1 Andrew Husted

Mr. Andrew Husted is an aquatic ecologist, specializing in freshwater systems and wetlands, who graduated with a MSc in Zoology. He, is Pr Sci Nat registered (400213/11) in the following fields of practice: Ecological Science, Environmental Science and Aquatic Science. Mr Husted is an Aquatic, Wetland and Biodiversity Specialist with 12 years' experience in the environmental consulting field. In addition to his ecological working experience, Andrew has experience in agricultural and soil assessments, this includes the consideration of land uses and land cover.

#### 5.2 Ivan Baker

Ivan Baker is Cand. Sci Nat registered (119315) in environmental science and geological science. Ivan is a wetland and ecosystem service specialist, a hydropedologist and pedologist that has completed numerous specialist studies ranging from basic assessments to EIAs. Ivan has carried out various international studies following FC standards. Ivan completed training in Tools for Wetland Assessments with a certificate of competence and completed his MSc in environmental science and hydropedology at the North-West University of Potchefstroom.

#### 6 Literature Review

# 6.1 Land Capability

According to Smith (2006), the capability of land concerns the wise use of land to ensure economical production on a sustained basis, under specific uses and treatments. The object of land classification is the grouping of different land capabilities, to indicate the safest option for use, to indicate permanent hazards and management requirements. These land capability classes decrease in capability from I to VIII and increase in risk from I to VIII. DAFF (2017) further defines land capability as "the most intensive long-term use of land for purposes of rainfed farming, determined by the interaction of **climate**, **soil** and **terrain**.

DAFF (2017) has further modelled the land capability on a rough scale for the entire of South Africa and has divided these results into 15 classes (see Table 6-1). Terrain, climate and soil capability was used as the building blocks for this exercise to ensure a national land capability data set.

Table 6-1 Land Capability (DAFF, 2017)

Land Capability Class (DAFF, 2017)	Description of Capability			
1	Vondow			
2	Very Low			
3	Very Low to Low			
4	very Low to Low			
5	Low			
6	Low to Moderate			
7	Low to inductate			
8	Moderate			
9	Moderate to High			





10	
11	High
12	High to Vary High
13	High to Very High
14	VoncHigh
15	Very High

It is worth noting that this nation-wide data set has some constraints of its own. According to DAFF (2017), inaccuracies and the level of detail of these datasets are of concern. Additionally, the scale used to model these datasets are large (1:50 000 to 1:100 000) and is not suitable for farm level planning. Furthermore, it is mentioned by DAFF (2017) that these datasets should not replace any site-based assessments given the accuracies perceived.

# 7 Methodology

The pedology assessment was conducted using the Provincial and National Departments of Agriculture recommendations. The assessment was broken into two phases. Phase 1 was a desktop assessment to determine the following:

- Historic climatic conditions;
- The base soils information from the land type database (Land Type Survey Staff, 1972 2006); and
- The geology for the proposed project site.

Phase 2 of the assessment was to conduct a soil survey to determine the actual agricultural potential. During this phase the current land use was also surveyed.

#### 7.1 Desktop Assessment

As part of the desktop assessment, baseline soil information was obtained using published South African Land Type Data. Land type data for the site was obtained from the Institute for Soil Climate and Water (ISCW) of the Agricultural Research Council (ARC) (Land Type Survey Staff, 1972 - 2006). The land type data is presented at a scale of 1:250 000 and comprises of the division of land into land types.

#### 7.2 Agricultural Potential Assessment

Land capability and agricultural potential will be determined by a combination of soil, terrain and climate features. Land capability is defined by the most intensive long-term sustainable use of land under rain-fed conditions. At the same time an indication is given about the permanent limitations associated with the different land use classes.

Land capability is divided into eight classes and these may be divided into three capability groups. Table 7-1 shows how the land classes and groups are arranged in order of decreasing capability and ranges of use. The risk of use and sensitivity increases from class I to class VIII (Smith, 2006).

Table 7-1 Land capability class and intensity of use (Smith, 2006)

Land		Land
Capability	Increased Intensity of Use	Capability
Class	·	Groups





1	W	F	LG	MG	IG	LC	MC	IC	VIC	
T.	W	F	LG	MG	IG	LC	MC	IC		
III	W	F	LG	MG	IG	LC	MC			Arable Land
IV	W	F	LG	MG	IG	LC				
V	W	F	LG	MG						
VI	W	F	LG	MG						Grazing Land
VII	W	F	LG							
VIII	W									Wildlife
W - Wildlife		MG - N	loderate Gra	zing	MC - Moderate Cultivation		ation			
F- Forestry		IG - Int	ensive Graz	ing	IC - Intensive Cultivation					
LG - Light Graz	zing	LC - Li	ght Cultivati	on	VIC - Very	Intensive (	Cultivation			

Land capability has been classified into 15 different categories by DAFF (2017) which indicates the national land capability category and associated sensitivity related to soil resources. Given the fact that ground truthing and DSM exercises have indicated anomalies in the form of high sensitivity soil resources (which was not indicated by the DAFF (2017) raster file), the ground-truthed baseline delineations and sensitivities were used for this assessment rather than that of DAFF (2017).

The land potential classes are determined by combining the land capability results and the climate capability of a region as shown in Table 7-2. The final land potential results are then described in

Table 7-3. These land potential classes are regarded as the final delineations subject to sensitivity, given the comprehensive addition of climatic conditions as those relevant to the DAFF (2017) land capabilities. The main contributors to the climatic conditions as per Smith (2006) is that of Mean Annual Precipitation (MAP), Mean Annual Potential Evaporation (MAPE), mean September temperatures, mean June temperatures and mean annual temperatures. These parameters will be derived from Mucina and Rutherford (2006) for each vegetation type located within the relevant project area. This will give the specialist the opportunity to consider micro-climate, aspect, topography etc.

Table 7-2 The combination table for land potential classification

Land capability class	Climate capability class							
Lanu capability class	C1	C2	C3	C4	C5	C6	<b>C</b> 7	C8
I	L1	L1	L2	L2	L3	L3	L4	L4
II	L1	L2	L2	L3	L3	L4	L4	L5
III	L2	L2	L3	L3	L4	L4	L5	L6
IV	L2	L3	L3	L4	L4	L5	L5	L6
V	Vlei	Vlei	Vlei	Vlei	Vlei	Vlei	Vlei	Vlei
VI	L4	L4	L5	L5	L5	L6	L6	L7
VII	L5	L5	L6	L6	L7	L7	L7	L8
VIII	L6	L6	L7	L7	L8	L8	L8	L8





Table 7-3 The Land Potential Classes.

Land potential	Description of land potential class
L1	Very high potential: No limitations. Appropriate contour protection must be implemented and inspected.
L2	High potential: Very infrequent and/or minor limitations due to soil, slope, temperatures or rainfall. Appropriate contour protection must be implemented and inspected.
L3	Good potential: Infrequent and/or moderate limitations due to soil, slope, temperatures or rainfall. Appropriate contour protection must be implemented and inspected.
L4	Moderate potential: Moderately regular and/or severe to moderate limitations due to soil, slope, temperatures or rainfall. Appropriate permission is required before ploughing virgin land.
L5	Restricted potential: Regular and/or severe to moderate limitations due to soil, slope, temperatures or rainfall.
L6	Very restricted potential: Regular and/or severe limitations due to soil, slope, temperatures or rainfall. Non-arable
L7	Low potential: Severe limitations due to soil, slope, temperatures or rainfall. Non-arable
L8	Very low potential: Very severe limitations due to soil, slope, temperatures or rainfall. Non-arable

# 7.3 Climate Capability

According to Smith (2006), climatic capability is determined by taking into consideration various steps pertaining to the temperature, rainfall and Class A-pan of a region. The first step in this methodology is to determine the Mean Annual Precipitation (MAP) to Class A-pan ratio.

Table 7-4 Climatic capability (step 1) (Scotney et al., 1987)

Climatic Capability Class	Limitation Rating	Description	MAP: Class A- pan Class
C1	None to Slight	Local climate is favourable for good yields for a wide range of adapted crops throughout the year.	0.75-1.00
C2	Slight	Local climate is favourable for a wide range of adapted crops and a year-round growing season. Moisture stress and lower temperature increase risk and decrease yields relative to C1.	0.50-0.75
C3	Slight to Moderate	Slightly restricted growing season due to the occurrence of low temperatures and frost. Good yield potential for a moderate range of adapted crops.	0.47-0.50
C4	Moderate	Moderately restricted growing season due to the occurrence of low temperatures and severe frost. Good yield potential for a moderate range of adapted crops but planting date options more limited than C3.	0.44-0.47
C5	Moderate to Severe	Moderately restricted growing season due to low temperatures, frost and/or moisture stress. Suitable crops at risk of some yield loss.	0.41-0.44
C6	Severe	Moderately restricted growing season due to low temperatures, frost and/or moisture stress. Limited suitable crops that frequently experience yield loss.	0.38-0.41
<b>C</b> 7	Severe to Very Severe	Severely restricted choice of crops due to heat and moisture stress.	0.34-0.38
C8	Very Severe	Very severely restricted choice of crops due to heat and moisture stress. Suitable crops at high risk of yield loss.	0.30-0.34

In the event that the MAP: Class A-pan ratio is calculated to fall within the C7 or C8 class, no further steps are required, and the climatic capability can therefore be determined to be C7 or C8. In cases where the above-mentioned ratio falls within C1-C6, steps 2 to 3 will be required to further refine the climatic capability.



# Step 2

Mean September temperatures;

- $<10^{\circ}C = C6$ ;
- 10 11 °C = C5;
- 11 12 °C = C4;
- 12 13  $^{\circ}$ C = C3; and
- >13°C = C1.

# Step 3

Mean June temperatures;

- <9°C = C5;
- 9 10 °C = C4;
- $10 11^{\circ}C = C3$ ; and
- $11 12 ^{\circ} C = C2$ .

#### 7.4 Current Land Use

A generalised land-use will be derived for the larger project area considering agricultural productivity.

- Mining;
- Bare areas;
- Agriculture crops;
- Natural veld;
- Grazing lands;
- Forest;

- Plantation;
- Urban;
- Built-up;
- · Waterbodies; and
- · Wetlands.

# 8 Desktop Findings

#### 8.1 Climate

This region's rainfall peaks during autumn months, especially March. The Mean Annual Precipitation (MAP) ranges from 190 to 400 mm with the mean minimum and maximum monthly temperatures for Britstown being -3.6 °C and 37.9 °C for July and January respectively (also see Figure 8-1 for more information).






Figure 8-1 Climate for the region

# 8.2 Soils and Geology

According to the land type database (Land Type Survey Staff, 1972 - 2006) the development falls within the Ag 146 land type. The Ag land type is characterised by freely drained Red or Yellow-Brown Apedal soils with red soils being dominant. These soils are characterised by a high base status and is likely to be less than 300 mm deep. The Ag 146 land type terrain units and expected soils are illustrated in Figure 8-2 and Table 8-1 respectively.




Figure 8-2 Illustration of the Ag 146 land type terrain units (Land Type Survey Staff, 1972 - 2006)

Table 8-1 Soils expected at the respective terrain units within the Ag 146 land type (Land Type Survey Staff, 1972 - 2006)

Terrain units							
1 (21%)		3 (49%)		4 (25%)		5 (5%)	
Bare Rock	65%	Bare Rock	50%	Hutton	65%	Oakleaf	30%
Hutton	20%	Hutton	25%	Glenrosa	10%	Valsrivier	25%
Glenrosa	10%	Glenrosa	20%	Oakleaf	10%	Streambeds	25%
Swartland	5%	Swartland	5%	Valsrivier	10%	Hutton	10%
				Swartland	5%	Glenrosa	5%
						Swartland	5%

The geology of this area is characterised by the Volksrust Formation shales as well as the Prince Albert Formation and the Dwyka Group diamictites (Mucina and Rutherford, 2006). The Jurassic Karoo Dolerite





sills and sheets support the vegetation in this area soils varying from shallow to deep. Red and yellow-brown apedal soils are common in this region with the Ae, Fc and Ag land types prominently featuring.

#### 8.3 Terrain

The slope percentage of the project area has been calculated and is illustrated in Figure 8-3. The majority of the project area is characterised by a slope percentage between 0 and 2%, with some smaller patches within the project area characterised by a slope percentage up to 4.3%. This illustration indicates a gentle and uniform topography. The elevation of the project area (Figure 8-4) indicates an elevation of 1 228 to 1 241 Metres Above Sea Level (MASL).






Figure 8-3 Slope percentage map for the project area




Figure 8-4 Elevation of the project area (metres above sea level)



#### 9 Results and Discussion

# 9.1 Baseline Findings

One main soil form was identified throughout the 50 m regulated area, namely the Glenrosa soil form (see Figure 9-1). The Glenrosa soil forms consists of an orthic topsoil on top of a lithic horizon and doesn't exceed depths of 40 cm.

The land capability of the abovementioned soil has been determined to have a land capability class of "VI" and a climate capability level 8 given the low Mean Annual Precipitation (MAP) and the high Mean Annual Potential Evapotranspiration (MAPE) rates. The combination between the determined land capabilities and climate capabilities results in a land potential "L7". The "L7" land potential level is characterised by a low potential. Severe limitations exist due to soil, slope, temperatures, or rainfall. This region is therefore regarded as being non-arable.



Figure 9-1 Example of a shallow Glenrosa soil form with a depth of less than 20 cm



# 9.2 Sensitivity Verification

The following land potential level has been determined;

Land potential level 7 (this land potential level is characterised by a low potential. Severe limitations
exist due to soil, slope, temperatures or rainfall. This region is therefore regarded as being nonarable).

Fifteen land capabilities have been digitised by (DAFF, 2017) across South Africa, of which eight potential land capability classes are located within the proposed footprint area's assessment corridor, including;

- Land Capability 1 to 5 (Very Low to Low Sensitivity); and
- Land Capability 6 to 8 (Low/Moderate to Moderate Sensitivity).

The baseline findings and the sensitivities as per the Department of Agriculture, Forestry and Fisheries (DAFF, 2017) national raster file concur with one another. It therefore is the specialist's opinion that the land capability and land potential of the resources in the regulated area is characterised by a maximum of "Moderate" sensitivities (see Figure 9-2), which conforms to the requirements of an agricultural compliance statement only.




Figure 9-2 Land Capability Sensitivity (DAFF, 2017)





# 10 Impact Statement

# 10.1 Loss of Land Capability

The impact assessment will consider the calculated sensitivities associated with the soil resources expected to be impacted upon by the relevant components. This impact assessment will purely focus on the impacts expected towards natural resources (in specific, the soil and associated land capability).

#### 10.2 Construction Phase

During the construction phase heavy vehicles (trucks) will be used to transport PV structures throughout the footprint area with reliance on manual labour for finer refinement. Potential erosion is possible during the construction phase.

It is evident from the impact calculations in Table 10-1 that in a pre-mitigation state, moderate impacts are expected. The main mitigation objective would be to limit the area to be impacted upon by means of not using concrete pylons but rather installing pylons directly into the soil surface. In the event that this recommendation be adhered to, lower impacts are foreseen which ultimately results in a post-mitigation significance rating of "Low".

Table 10-1 Impact assessment related to the loss of land capability during the construction phase of the proposed PV area

Nature: Loss of land capability							
	Without mitigation	With mitigation					
Extent	Low (2)	Low (2)					
Duration	Short Term (2)	Short Term (2)					
Magnitude	Moderate (6)	Low (4)					
Probability	Probable (3)	Probable (3)					
Significance	Medium	Low					
Status (positive or negative)	Negative	Negative					
Reversibility	High	High					
Irreplaceable loss of resources?	No	No					
Can impacts be mitigated?	Yes						
Mitigation: See Section 11							
Residual Impacts:							
Limited residual impacts will be associ adhered to.	Limited residual impacts will be associated with these activities, assuming that all prescribed mitigation measures be strictly adhered to.						

# 10.3 Operational Phase

During the operational phase, very little impacts are foreseen. Maintenance of vegetation as well as the occasional maintenance of PV structures will have to be carried out throughout the life of the project. It is expected that these maintenance practices can be undertaken by means of manual labour. Overland flow dynamics are expected t be affected, although only slightly, due to access and maintenance routes.

Considering the low magnitude of impacts as well as the fact that pylons will not be cemented to the surface, very little impacts are expected post-mitigation for the proposed operational phase.





**Table 10-2** Impact assessment related to the loss of land capability during the operational phase of the proposed PV area

Nature: Loss of land capability		
	Without mitigation	With mitigation
Extent	Low (2)	Low (2)
Duration	Long Term (4)	Long Term (4)
Magnitude	Moderate (6)	Minor (2)
Probability	Probable (3)	Probable (3)
Significance	Medium	Low
Status (positive or negative)	Negative	Negative
Reversibility	High	High
Irreplaceable loss of resources?	No	No
Can impacts be mitigated?	Yes	
Mitigation: See Section 11		
Residual Impacts:		

Limited residual impacts will be associated with these activities, assuming that all prescribed mitigation measures be strictly adhered to.

# 10.4 Cumulative Impacts

Cumulative impacts within the proposed PV area and its surroundings have been determined to be low. Soil resources in the area has been impacted upon predominantly by means of erosion although to a lesser extent.

**Table 10-3** Impact assessment related cumulative impacts

Nature: Loss of land capability		
	Without mitigation	With mitigation
Extent	Low (2)	Low (2)
Duration	Permanent (5)	Permanent (5)
Magnitude	Minor (2)	Minor (2)
Probability	Improbable (2)	Improbable (2)
Significance	Low	Low
Status (positive or negative)	Negative	Negative
Reversibility	High	High
Irreplaceable loss of resources?	No	No
Can impacts be mitigated?	Yes	
Mitigation: See Section 11		
Residual Impacts:		
Limited residual impacts will be associ	iated with these activities, assuming the	hat all prescribed mitigation measures be strictly

# 10.5 Specialist Opinion

It is the specialist's opinion that the baseline findings do not concur with the land capabilities identified by means of the DAFF (2017) desktop findings in regard to land capability sensitivities. Even though



adhered to.



the land capability, in theory, is similar to that portrayed by (DEA, 2021), the climatic conditions have been deemed to be extremely poor. These poor climatic conditions have resulted in a land potential level characterised by "Low" sensitivity throughout the project area. No "High" land capability sensitivities were identified within proximity to any of the proposed activities. Considering the lack of sensitivity and the measures put in place in regard to stormwater management and erosion control, it is the specialist's opinion that all activities will have an acceptable impact on agricultural productivity. Furthermore, no measures in regard to moving components in their micro-setting were required to avoid or minimise fragmentation and disturbances of agricultural activities.

# 11 Recommendations and Mitigation Measures

# 11.1 General Mitigation

General mitigations will ensure the conservation of all soil resources, regardless of the sensitivity of resources and the intensity of impacts.

- Prevent any spills from occurring. Machines must be parked within hard park areas and must be checked daily for fluid leaks;
- Proper invasive plant control must be undertaken quarterly; and
- All excess soil (soil that are stripped and stockpiled to make way for foundations) must be stored, continuously rehabilitated to be used for rehabilitation of eroded areas.

# 11.2 Restoration of Vegetation Cover

Restoring vegetation cover is the first step to successful rehabilitation. Vegetation cover decreases flow velocities and minimises erosion.

# 11.2.1 Ripping Compacted Areas

All areas outside of the footprint areas that will be degraded (by means of vehicles, laydown yards etc.) must be ripped where compaction has taken place. According to the Department of Primary Industries and Regional Development (Agriculture and Food) (2017), ripping tines must penetrate to just below the compacted horizons (approximately 300 – 400 mm) with soil moisture being imminent to the success of ripping. Ripping must take place within 1-3 days after seeding, and also following a rain event to ensure a higher moisture content. To summarise:

- Rip all compacted areas outside of the developed areas that have been compacted;
- This must be done by means of a commercial ripper that has at least two rows of tines; and
- Ripping must take place between 1 and 3 days after seeding and following a rainfall event (seeding must therefore be carried out directly after a rainfall event).

#### 11.2.2 Revegetate Degraded Areas

Vegetation within the footprint areas will be cleared to accommodate the excavation activities coupled with the proposed footprint areas' foundations. This impact will degrade soil resources, ultimately decreasing the land capability of resources and increasing erosion. According to Russell (2009), areas characterised by a loss of soil resources should be revegetated by means of vegetation with vigorous growth, stolons or rhizomes that more or less resembles the natural vegetation in the area.

It is recommended that all areas surrounding the development footprint areas that have been degraded by traffic, laydown yards etc. must be ripped and revegetated by means of indigenous grass species. Mixed stands or monocultures will work sufficiently for revegetation purposes. Mixed stands tend to blend in with indigenous vegetation species and are more natural. Monocultures however could achieve



## Graspan PV



high productivity. In general, indigenous vegetation should always be preferred due to various reasons including the aesthetical presence thereof as well as the ability of the species to adapt to its surroundings.

Plant phase plants which are characterised by fast growing and rapid spreading conditions. Seed germination, seed density and seed size are key aspects to consider before implementing revegetation activities. The amount of seed should be limited to ensure that competition between plants are kept to a minimum. During the establishment of seed density, the percentage of seed germination should be taken into consideration. *E curvula* is one of the species recommended due to the ease of which it germinates. This species is also easily sown by means of hand propagation and hydro seeding.

The following species are recommended for rehabilitation purposes;

- Eragrostis teff;
- Cynodon species (Indigenous and altered types);
- · Chloris gayana;
- Panicum maximum;
- Digitaria eriantha;
- · Anthephora pubescens; and
- Cenchrus ciliaris.

# 11.3 Specialist Recommendation

The proposed activities may proceed as have been planned without the concern of loss of high sensitivity land capabilities or agricultural productivity.

# 12 Conclusion

One soil form was identified within the project area, namely the Glenrosa soil form. This soil form has been determined to be associated with one land capabilities, namely LCVI. This land capability class was then further refined to a land potential level 7 by comparing land capability of climatic capabilities of the project area.

This land potential level was used to determine the sensitivities of soil resources. Only "Low" sensitivities were determined throughout the project area by means of baseline findings. Considering the low sensitivities associated with land potential resources, it is the specialist's opinion that the proposed activities will have an acceptable impact on soil resources and that the proposed activities should proceed as have been planned.





# 13 References

Department of Primary Industries and Regional Development. 2017. Deep ripping for soil compaction. https://www.agric.wa.gov.au/soil-compaction/deep-ripping-soil-compaction.

Land Type Survey Staff. 1972 - 2006. Land Types of South Africa: Digital Map (1:250 000 Scale) and Soil Inventory Databases. Pretoria: ARC-Institute for Soil, Climate, and Water.

Mucina, L., & Rutherford, M. C. 2006. The Vegetation of South Africa, Lesotho, and Swaziland. Strelitzia 19. Pretoria: National Biodiversity Institute.

Russell, W. 2009. WET-RehabMethods. National guidelines and methods for wetland rehabilitation.

Smith, B. 2006. The Farming Handbook. Netherlands & South Africa: University of KwaZulu-Natal Press & CTA.

Soil Classification Working Group. 1991. Soil Classification A Taxonomic system for South Africa. Pretoria: The Department of Agricultural Development.

Soil Classification Working Group. 2018. Soil Classification A Taxonomic system for South Africa. Pretoria: The Department of Agricultural Development.