



TERRESTRIAL ECOLOGY BASLINE & IMPACT ASSESSMENT FOR THE PROPOSED SBPM SOLAR FACILITY

Northam, Limpopo & North West Provinces

May 2022

CLIENT

savannah
environmental

Prepared by:

The Biodiversity Company

Cell: +27 81 319 1225

Fax: +27 86 527 1965

info@thebiodiversitycompany.com

www.thebiodiversitycompany.com




Table of Contents

1	Introduction.....	1
1.1	Background	1
1.1.1	Project Description	1
1.1.2	SCSC PV RE project.....	1
1.2	Specialist Details	5
1.3	Terms of Reference	6
1.4	Assumptions and Limitations	6
1.5	Key Legislative Requirements.....	6
2	Methods.....	7
2.1	Desktop Baseline	7
2.1.1	Ecologically Important Landscape Features	7
2.1.2	Desktop Flora Baseline	9
2.1.3	Desktop Faunal Baseline	9
2.2	Biodiversity Field Assessment	10
2.2.1	Flora Survey	10
2.2.2	Fauna Survey	10
2.3	Terrestrial Site Ecological Importance	11
3	Results & Discussion	14
3.1	Desktop Baseline	14
3.1.1	Ecologically Important Landscape Features	14
3.1.2	Flora Assessment.....	22
3.1.3	Faunal Assessment.....	24
3.1.4	Review of previous reports.....	Error! Bookmark not defined.
3.2	Field Assessment.....	28
3.2.1	Flora Assessment.....	28
3.2.2	Faunal Assessment.....	32
4	Habitat Assessment and Site Ecological Importance	38
4.1	Habitat Assessment	38
4.2	Site Ecological Importance (SEI)	43
5	Impact Risk Assessment.....	47
5.1	Biodiversity: Risk Assessment Method	47
5.1.1	Present Impacts to Biodiversity	47

5.1.2	Identification of Additional Potential Impacts	49
5.1.3	Assessment of Impact Significance	52
5.1.4	Biodiversity Management Outcomes	56
6	Conclusion and Impact Statement	62
6.1	Conclusion.....	62
6.1.1	Terrestrial Biodiversity.....	62
6.2	Impact Statement	62
7	References	64
8	Appendix Items.....	66
8.1	Appendix A – Specialist Declaration of Independence	66

List of Tables

Table 1-1	A list of key legislative requirements relevant to biodiversity and conservation in the Limpopo and North West Provinces.....	6
Table 2-1	Summary of Conservation Importance (CI) criteria.....	11
Table 2-2	Summary of Functional Integrity (FI) criteria	11
Table 2-3	Matrix used to derive Biodiversity Importance (BI) from Functional Integrity (FI) and Conservation Importance (CI)	12
Table 2-4	Summary of Receptor Resilience (RR) criteria	12
Table 2-5	Matrix used to derive Site Ecological Importance from Receptor Resilience (RR) and Biodiversity Importance (BI)	13
Table 2-6	Guidelines for interpreting Site Ecological Importance in the context of the proposed development activities	13
Table 3-1	Summary of relevance of the proposed project to ecologically important landscape features	14
Table 3-2	Threatened flora species that may occur within the project area.....	24
Table 3-3	Threatened amphibian species that are expected to occur within the project area.....	24
Table 3-4	Threatened reptile species that are expected to occur within the project area.....	24
Table 3-5	Threatened mammal species that are expected to occur within the project area.	25
Table 3-6	Trees, shrub and herbaceous plant species recorded in the project area.....	28
Table 3-7	Summary of herpetofauna species recorded within the project area.....	33
Table 3-8	Summary of mammal species recorded within the project area. Mammal species are considered 'captive' species as these were only present within the game farm areas, marked in green text.....	35
Table 4-1	Summary of habitat types delineated within the project area	44
Table 5-1	Potential impacts to biodiversity associated with the proposed activity	49
Table 5-2	Impacts to biodiversity associated with the proposed construction phase.	52
Table 5-3	Impacts to biodiversity associated with the proposed construction phase.	52
Table 5-5	Impacts to biodiversity associated with the proposed construction phase.	53
Table 5-6	Impacts to biodiversity associated with the proposed operational phase	54
Table 5-7	Impacts to biodiversity associated with the proposed operational phase.	54
Table 5-8	Impacts to biodiversity associated with the proposed operational phase	55
Table 5-9	Mitigation measures including requirements for timeframes, roles and responsibilities for this report	57

List of Figures

Figure 1-1	Proposed location of the project area in relation to the nearby towns	3
Figure 1-2	The various components of the project	4
Figure 2-1	Map illustrating extent of area used to obtain the expected flora species list from the Plants of South Africa (POSA) database. Yellow dot indicates approximate location of the project area. The red squares are cluster markers of botanical records as per POSA data	9
Figure 3-1	Map illustrating the ecosystem threat status associated with the project area.	15
Figure 3-2	Map illustrating the ecosystem protection level associated with the project area	16
Figure 3-3	Map illustrating the locations of CBAs in the project area	17
Figure 3-4	The project area in relation to the protected areas	18
Figure 3-5	The project area in relation to the National Protected Area Expansion Strategy	19
Figure 3-6	Map illustrating ecosystem threat status of rivers and wetland ecosystems in the project area	20
Figure 3-7	The project area in relation to the National Freshwater Ecosystem Priority Areas.	21
Figure 3-8	The project area in relation to the renewable energy database projects in the area.	22
Figure 3-9	Map illustrating the vegetation type associated with the project area	23
Figure 3-10	Map illustrating ecological sensitivity according to the literature review	27
Figure 3-11	Photographs illustrating some of the flora recorded within the assessment area. A) <i>Hermbstaedtia odorata</i> , B) <i>Vachellia nilotica</i> , C) <i>Nymphaea lotus</i> , D) <i>Heliotropium amplexicaule</i> and E) <i>Hibiscus pusillus</i>	31
Figure 3-13	Photographs illustrating the reptile species recorded within the assessment area associated with the project area during the survey period. A) <i>Tomopterna cryptotis</i> , B) <i>Lygodactylus capensis</i> , and C) <i>Varanus niloticus</i> , D) <i>Stigmochelys pardalis</i> and E) <i>Acanthocercus atricollis</i>	34
Figure 3-14	Photographs illustrating the mammal species recorded within the project area during the survey period. A) <i>Equus quagga</i> and <i>Alcelaphus buselaphus</i> , B) <i>Mungos mungo</i> , C) <i>Damaliscus pygargus phillipsi</i> , D) <i>Chlorocebus pygerythrus</i> and E) <i>Giraffa camelopardalis</i>	36
Figure 3-15	Photographs illustrating a mammal SCC recorded within the project area during the survey period: <i>Leptailurus serval</i>	37
Figure 4-1	Habitats identified in the project area	39
Figure 4-2	A typical example of degraded Bushveld habitat from the project area	40
Figure 4-3	A typical example of fragmented Bushveld habitat from the project area	40
Figure 4-4	A typical example of disturbed Bushveld habitat from the project area.	41
Figure 4-5	Illustration of transformed habitat from the project area.	41
Figure 4-6	Illustration of transformed habitat from the project area.	42
Figure 4-7	Illustration of wetland habitat from the project area (Sefathlane River)	42

Figure 4-8	Illustration of wetland habitat from the project area	43
Figure 4-9	Terrestrial Biodiversity Theme Sensitivity, TBC Screening Report.....	44
Figure 4-10	Sensitivity of the project area	46
Figure 5-1	Photographs illustrating impacts to biodiversity A) Roads and fencing), B) Powerlines and associated servitude C) Wood harvesting D) Livestock, roads and railway and E) Livestock and litter.....	48
Figure 5-2	Proposed infrastructure layout overlaid with the SEI.	51

1 Introduction

1.1 Background

The Biodiversity Company was appointed to undertake a biodiversity impact assessment for the proposed SBPM Solar Facility for Siyanda Bakgatla Platinum Mine in Northam, Limpopo Province. The project infrastructure is located in both the Limpopo and also North West provinces. The project is located 6.5 km west from Northam. The Northam focus area has been identified by the potential development area for the construction and operation of solar and battery facilities consisting of 251 Ha (Figure 1-2).

The approach was informed by the Environmental Impact Assessment Regulations, 2014 (GNR 326, 7 April 2017) of the National Environmental Management Act, 1998 (Act No. 107 of 1998) (NEMA). The approach has taken cognisance of the recently published Government Notices 320 (20 March 2020) in terms of NEMA, dated 20 March: “*Procedures for the Assessment and Minimum Criteria for Reporting on Identified Environmental Themes in terms of Sections 24(5)(a) and (h) and 44 of the National Environmental Management Act, 1998, when applying for Environmental Authorisation*” (Reporting Criteria). The National Web based Environmental Screening Tool has characterised the terrestrial sensitivity of the project area as “Very High”.

This report, after taking into consideration the findings and recommendations provided by the specialist herein, should inform and guide the Environmental Assessment Practitioner (EAP) and regulatory authorities, enabling informed decision making, as to the ecological viability of the proposed project.

1.2 Project Description

1.2.1 SBPM PV RE project, Limpopo Province

Main Street 1886 Proprietary Limited proposes the development of the Solar Photovoltaic (PV) facility and associated infrastructure on a site bordering the eastern end of the Siyanda Bakgatla Platinum Mine area near Northam. The solar PV facility will comprise several arrays of PV panels, a Battery Energy Storage System (BESS), and associated infrastructure with a contracted capacity of up to 100MW.

The purpose of the proposed project is to generate electricity for exclusive use by the Siyanda Mine, following which any excess power produced will be distributed to the national grid, if applicable. The construction of the PV facility aims to reduce the Siyanda Mine’s dependency on direct supply from Eskom’s national grid for operation activities, while simultaneously decreasing the mine’s carbon footprint.

A preferred project site with an extent of ~1138 ha and a development area of 574 ha has been identified by Main Street 1886 Proprietary Limited as a technically suitable area for the development of the Solar PV Facility. The study area is located on Portion 4 of Farm Grootkuil 409. The project site falls within the Thabazimbi Local Municipality within the Waterberg District Municipality in the Limpopo Province. The site is located ~6.5 km west of the town of Northam and is accessible via the Swartklip Road which branches off the R510 provincial route.

Infrastructure associated with the solar PV facility will include:

- 100MW Solar PV array comprising PV modules and mounting structures.
- Inverters and transformers.
- Cabling between the project components.
- Battery Energy Storage System.
- On-site facility substation and power lines between the solar PV facility and the Mine and Eskom substation.
- Site offices, Security office, operations and control, and maintenance and storage laydown areas.
- Access roads, internal distribution roads.

Grid connection solution.

To evacuate the generated power to the Siyanda Mine, the grid connection solution consisting of the following is proposed:

The power generated by the solar PV facility will be transferred to the three step up transformers at the on-site/plant substation. Power will then be delivered from each step-up transformer as follows:

- two 6.6 km, 33 kV transmission lines to the Mortimer substation with four step down transformers (33/6.6 kV; 10 MVA).
- two 4.7 km, 33 kV transmission lines to the Fridge substation with two step down transformers (33/6.6 kV; 10 MVA).
- two 2.9 km, 33 kV transmission lines to the Ivan substation with three step down transformers (33/11 kV; 10 MVA).

The grid connection is proposed on the following properties:

- Portion 3 of Farm Grootkuil 409.
- Portion 4 of Farm Grootkuil 409.
- Portion 5 of Farm Grootkuil 409.

The development area of 574ha is larger than the area needed for the construction of a 100MW PV facility and will provide the opportunity for the optimal placement of the infrastructure, ensuring avoidance of major identified environmental sensitivities by the development footprint of ~240 ha¹.

¹ The development footprint is the defined area (located within the development area) where the PV panel array and other associated infrastructure for Solar PV will be planned to be constructed. This will be the actual footprint of the facility, and the area which would be disturbed. The extent of the development footprint will be determined in the EIA Phase.




Figure 1-1 Proposed location of the project area in relation to the nearby towns








Figure 1-2 The various components of the project

1.3 Specialist Details

Report Name	TERRESTRIAL ECOLOGY BASLINE & IMPACT ASSESSMENT FOR THE PROPOSED SBPM SOLAR FACILITY
Reference	Siyanda PV
Submitted to	
Report Writer	Lindi Steyn 
Desktop	Dr Lindi Steyn has completed her PhD in Biodiversity and Conservation from the University of Johannesburg. Lindi is a terrestrial ecologist with a special interest in ornithology. She has completed numerous studies ranging from basic Assessments to Environmental Impact Assessments following IFC standards.
Report Writer (Screening) and Fieldwork	Lusanda Matee 
	Lusanda Matee is a registered scientist (119257/2018) in the fields of Biological Science (Pr.Sci Nat.) and Ecological Science (Cand Nat.). He is a specialist terrestrial ecologist and botanist who conducts floral surveys faunal surveys which include mammals, birds, amphibians, and reptiles. He has 4 years of experience in environmental consulting. He received a Bachelor of Science, Honours, and MSc in Biological Sciences from the University of KwaZulu-Natal.
Report Writer and Fieldwork	Marnus Erasmus 
	Martinus Erasmus obtained his B-Tech degree in Nature Conservation in 2016 at the Tshwane University of Technology. Martinus has been conducting EIAs, IFC standard surveys, basic assessments and assisting specialists in field during his studies since 2015. Martinus is Cand. Sci. Nat. registered (118630) is a specialist terrestrial ecologist and botanist which conducts floral surveys faunal surveys which include mammals, birds, amphibians and reptiles.
Reviewer	Andrew Husted 
	Andrew Husted is Pr Sci Nat registered (400213/11) in the following fields of practice: Ecological Science, Environmental Science and Aquatic Science. Andrew is an Aquatic, Wetland and Biodiversity Specialist with more than 12 years' experience in the environmental consulting field. Andrew has completed numerous wetland training courses, and is an accredited wetland practitioner, recognised by the DWS, and also the Mondi Wetlands programme as a competent wetland consultant.
Declaration	The Biodiversity Company and its associates operate as independent consultants under the auspice of the South African Council for Natural Scientific Professions. We declare that we have no affiliation with or vested financial interests in the proponent, other than for work performed under the Environmental Impact Assessment Regulations, 2017. We have no conflicting interests in the undertaking of this activity and have no interests in secondary developments resulting from the authorisation of this project. We have no vested interest in the project, other than to provide a professional service within the constraints of the project (timing, time and budget) based on the principals of science.

1.4 Terms of Reference

The principle aim of the assessment was to provide information to guide the risk of the proposed activity to the flora and fauna communities of the associated ecosystems within the project area/corridor. This was achieved through the following:

- Desktop assessment to identify the relevant ecologically important geographical features within the project area;
- Desktop assessment to compile an expected species list and possible threatened flora and fauna species that occur within the project area;
- Field survey to ascertain the species composition of the present flora and fauna community within the project area;
- Delineate and map the habitats and their respective sensitivities that occur within the project area;
- Identify the manner that the proposed project impacts the flora and fauna community and evaluate the level of risk of these potential impacts; and
- The prescription of mitigation measures and recommendations for identified risks.

1.5 Assumptions and Limitations

The following assumptions and limitations are applicable for this assessment:

- The assessment area was based on the area provided by the client and any alterations to the footprint and/or missing GIS information pertaining to the assessment area would have affected the area surveyed;
- The species likelihood of occurrence is based on desktop information and might be changed after the assessment;
- The impact assessment included is preliminary and is solely based on the screening survey and desktop information; and
- No decommissioning phase impacts have been considered for this project. The life of operation is unknown and expected for perpetuity.

1.6 Key Legislative Requirements

The legislation listed below in Table 1-1 are applicable to the current project. The list below, although extensive, may not be complete and other legislation, policies and guidelines may apply in addition to those listed below.

Table 1-1 *A list of key legislative requirements relevant to biodiversity and conservation in the Limpopo and North West Provinces*

Region	Legislation / Guideline
National	Constitution of the Republic of South Africa (Act No. 108 of 1996)
	The National Environmental Management Act (NEMA) (Act No. 107 of 1998)
	The National Environmental Management: Protected Areas Act (Act No. 57 of 2003)
	The National Environmental Management: Biodiversity Act (Act No. 10 of 2004), Threatened or Protected Species Regulations
	Procedures for the Assessment and Minimum Criteria for Reporting on Identified Environmental Themes in terms of Sections 24(5)(a) and (h) and 44 of the National Environmental Management Act, 1998, GNR 320 of Government Gazette 43310 (March 2020)
	Procedures for the Assessment and Minimum Criteria for Reporting on Identified Environmental Themes in terms of Sections 24(5)(a) and (h) and 44 of the National Environmental Management Act, 1998, GNR 1150 of Government Gazette 43855 (October 2020)
	The National Environmental Management: Waste Act, 2008 (Act 59 of 2008);
	National Forest Act (Act No. 84 of 1998)
	National Veld and Forest Fire Act (101 of 1998)

Solar Facility

	National Water Act (NWA) (Act No. 36 of 1998)
	Alien and Invasive Species Regulations and, Alien and Invasive Species List 2014/2020, published under NEMBA
	Conservation of Agricultural Resources Act, 1983 (Act 43 of 1983) (CARA)
Provincial	Limpopo Conservation Plan (2018)
	Limpopo Environmental Management Act (2003)
	North-West Biodiversity Sector Plan of 2015 (READ, 2015).
	The North West Biodiversity Management Amendment Bill, 2017

2 Methods

2.1 Desktop Baseline

The desktop assessment was principally undertaken using a Geographic Information System (GIS) to access the latest available spatial datasets to develop digital cartographs and species lists. These datasets and their date of publishing are provided below.

2.1.1 Ecologically Important Landscape Features

Existing ecologically relevant data layers were incorporated into a GIS to establish how the proposed project might interact with any ecologically important entities. Emphasis was placed around the following spatial datasets:

- National Biodiversity Assessment 2018 (Skowno et al, 2019) (NBA) - The purpose of the NBA is to assess the state of South Africa's biodiversity based on best available science, with a view to understanding trends over time and informing policy and decision-making across a range of sectors. The NBA deals with all three components of biodiversity: genes, species, and ecosystems; and assesses biodiversity and ecosystems across terrestrial, freshwater, estuarine and marine environments. The two headline indicators assessed in the NBA are:
 - *Ecosystem Threat Status* – indicator of an ecosystem's wellbeing, based on the level of change in structure, function or composition. Ecosystem types are categorised as Critically Endangered (CR), Endangered (EN), Vulnerable (VU), Near Threatened (NT) or Least Concern (LC), based on the proportion of the original extent of each ecosystem type that remains in good ecological condition.
 - *Ecosystem Protection Level* – indicator of the extent to which ecosystems are adequately protected or under-protected. Ecosystem types are categorised as Well Protected (WP), Moderately Protected (MP), Poorly Protected (PP), or Not Protected (NP), based on the proportion of the biodiversity target for each ecosystem type that is included within one or more protected areas. NP, PP or MP ecosystem types are collectively referred to as under-protected ecosystems.
- Protected areas - South Africa Protected Areas Database (SAPAD) (DEA, 2021) – The SAPAD Database contains spatial data pertinent to the conservation of South African biodiversity. It includes spatial and attribute information for both formally protected areas and areas that have less formal protection. SAPAD is updated on a continuous basis and forms the basis for the Register of Protected Areas, which is a legislative requirement under the National Environmental Management: Protected Areas Act, Act 57 of 2003.
- National Protected Areas Expansion Strategy (NPAES) (SANBI, 2016) – The NPAES provides spatial information on areas that are suitable for terrestrial ecosystem protection. These focus areas are large, intact and unfragmented and therefore, of high importance for biodiversity, climate resilience and freshwater protection.
- Conservation/Biodiversity Sector Plans:

The **Limpopo Conservation Plan** was completed in 2018 for the Limpopo Department of Economic Development, Environment & Tourism (LEDET) (Desmet *et al.*, 2013). The purpose of the LCPv2 was to develop the spatial component of a bioregional plan (i.e., map of Critical Biodiversity Areas and associated land-use guidelines). The previous Limpopo Conservation Plan (LCPv1) was

completely revised and updated (Desmet et al., 2013). A Limpopo Conservation Plan map was produced as part of this plan and sites were assigned to the following CBA categories based on their biodiversity characteristics, spatial configuration, and requirement for meeting targets for both biodiversity pattern and ecological processes:

- Critical Biodiversity Area 1 (CBA1);
- Critical Biodiversity Area 2 (CBA2);
- Ecological Support Area 1 (ESA1);
- Ecological Support Area 2 (ESA2);
- Other Natural Area (ONA);
- Protected Area (PA); and
- No Natural Remaining (NNR).

Critical Biodiversity Areas (CBAs) are terrestrial and aquatic areas of the landscape that need to be maintained in a natural or near-natural state to ensure the continued existence and functioning of species and ecosystems and the delivery of ecosystem services. Thus, if these areas are not maintained in a natural or near natural state then biodiversity targets cannot be met. Maintaining an area in a natural state can include a variety of biodiversity compatible land uses and resource uses (Desmet et al., 2013).

Ecological Support Areas (ESA's) are not essential for meeting biodiversity targets but play an important role in supporting the ecological functioning of Critical Biodiversity Areas and/or in delivering ecosystem services (SANBI, 2017). Critical Biodiversity Areas and Ecological Support Areas may be terrestrial or aquatic.

Other Natural Areas (ONAs) consist of all those areas in good or fair ecological condition that fall outside the protected area network and have not been identified as CBAs or ESAs. A biodiversity sector plan or bioregional plan must not specify the desired state/management objectives for ONAs or provide land-use guidelines for ONAs (Driver *et al.*, 2017).

Areas with No Natural Habitat Remaining (NNR) are areas in poor ecological condition that have not been identified as CBAs or ESAs. They include all irreversibly modified areas (such as urban or industrial areas and mines), and most severely modified areas (such as cultivated fields and forestry plantations). A biodiversity sector plan or bioregional plan must not specify the desired state/management objective or provide land-use guidelines for NNR areas (Driver *et al.*, 2017).

The **North-West Department of Rural, Environment, and Agricultural Development (READ)**, as custodian of the environment in the North West, is the primary implementing agent of the Biodiversity Sector Plan. The spatial component of the Biodiversity Sector Plan is based on systematic biodiversity planning undertaken by READ. The purpose of a Biodiversity Sector Plan is to inform land use planning, environmental assessments, land and water use authorisations, as well as natural resource management, undertaken by a range of sectors whose policies and decisions impact on biodiversity. This is done by providing a map of biodiversity priority areas, referred to as Critical Biodiversity Areas (CBAs) and Ecological Support Areas (ESAs), with accompanying land use planning and decision-making guidelines (READ, 2015).

- Important Bird and Biodiversity Areas (IBAs) (BirdLife South Africa, 2015) – IBAs constitute a global network of over 13 500 sites, of which 112 sites are found in South Africa. IBAs are sites of global significance for bird conservation, identified through multi-stakeholder processes using globally standardised, quantitative and scientifically agreed criteria; and
- South African Inventory of Inland Aquatic Ecosystems (SAIIAE) (Van Deventer *et al.*, 2018) – A SAIIAE was established during the NBA of 2018. It is a collection of data layers that represent the extent of river and inland wetland ecosystem types and pressures on these systems.

2.1.2 Desktop Flora Baseline

The Vegetation of South Africa, Lesotho and Swaziland (Mucina & Rutherford, 2006) and SANBI (2019) was used to identify the vegetation type that would have occurred under natural or pre-anthropogenically altered conditions. Furthermore, the Plants of Southern Africa (POSA) database was accessed to compile a list of expected flora species within the project area (Figure 2-1). The Red List of South African Plants (Raimondo *et al.*, 2009; SANBI, 2020) was utilized to provide the most current national conservation status of flora species.




Figure 2-1 Map illustrating extent of area used to obtain the expected flora species list from the Plants of South Africa (POSA) database. Yellow dot indicates approximate location of the project area. The red squares are cluster markers of botanical records as per POSA data

2.1.3 Desktop Faunal Baseline

The faunal desktop assessment comprised of the following, compiling an expected:

- Amphibian list, generated from the IUCN spatial dataset (2017) and AmphibianMap database (Fitzpatrick Institute of African Ornithology, 2021a), using the 2427 quarter degree square;
- Reptile list, generated from the IUCN spatial dataset (2017) and ReptileMap database (Fitzpatrick Institute of African Ornithology, 2021b), using the 2427 quarter degree square;
- Avifauna list, generated from the SABAP2 dataset by looking at pentads 2450_2700; 2450_2705; 2455_2700; 2455_2700; 2455_2705; 2455_2710; 2500_2700_2500_2705); and
- Mammal list from the IUCN spatial dataset (2017).

2.2 Biodiversity Field Assessment

The following methodologies will be implemented for the baseline assessment (phase) of the project.

2.2.1 Flora Survey

The fieldwork and sample sites will be placed within targeted areas (i.e., target sites) perceived as ecologically sensitive based on the preliminary interpretation of satellite imagery (Google Corporation) and GIS analysis (which will include the latest applicable biodiversity datasets) available prior to the fieldwork. The focus of the fieldwork will therefore be to maximise coverage and navigate to each target site in the field, to perform a rapid vegetation and ecological assessment at each sample site. Emphasis will be placed on sensitive habitats, especially those overlapping with the proposed project area.

Homogenous vegetation units will be subjectively identified using satellite imagery and existing land cover maps. The floristic diversity and search for flora SCC will be conducted through timed meanders within representative habitat units delineated during the fieldwork. Emphasis will be placed mostly on sensitive habitats overlapping with the proposed project areas.

The timed random meander method is highly efficient for conducting floristic analysis, specifically in detecting flora SCC and maximising floristic coverage. In addition, the method is time and cost effective and highly suited for compiling flora species lists and therefore gives a rapid indication of flora diversity. The timed meander search will be performed based on the original technique described by Goff *et al.* (1982). Suitable habitat for SCC will be identified according to Raimondo *et al.* (2009) and targeted as part of the timed meanders.

At each sample site notes will be made regarding current impacts (e.g., livestock grazing, erosion etc.), subjective recording of dominant vegetation species, and any sensitive features (e.g., wetlands, outcrops etc.). In addition, opportunistic observations will be made while navigating through the project area.

2.2.2 Fauna Survey

The faunal assessment within this report pertains to herpetofauna (amphibians and reptiles), avifauna and mammals. The faunal field survey will be comprised of the following techniques:

- Visual and auditory searches - This typically comprises of meandering and using binoculars to view species from a distance without them being disturbed; and listening to species calls;
- Active hand-searches - Used for species that shelter in or under particular micro-habitats (typically rocks, exfoliating rock outcrops, fallen trees, leaf litter, bark etc.)
- Camera trapping;
- Point counts for the avifauna; and
- Utilization of local knowledge.

Relevant field guides and texts that will be consulted for identification purposes included the following:

- Field Guide to Snakes and other Reptiles of Southern Africa (Branch, 1998);
- A Complete Guide to the Snakes of Southern Africa (Marais, 2004);
- Atlas and Red List of the Reptiles of South Africa, Lesotho and Swaziland (Bates *et al.*, 2014);
- A Complete Guide to the Frogs of Southern Africa (du Preez and Carruthers, 2009);
- Smithers' Mammals of Southern Africa (Apps, 2000);
- A Field Guide to the Tracks and Signs of Southern and East African Wildlife (Stuart and Stuart, 2000);
- Book of birds of South Africa, Lesotho and Swaziland (Taylor *et al.*, 2015); and
- Roberts – Birds of Southern Africa (Hockey *et al.*, 2005).

2.3 Terrestrial Site Ecological Importance

The different habitat types within the project area will be delineated and identified based on observations during the field assessment, and available satellite imagery. These habitat types will be assigned Ecological Importance (EI) categories based on their ecological integrity, conservation value, the presence of species of conservation concern and their ecosystem processes.

Site Ecological Importance (SEI) is a function of the Biodiversity Importance (BI) of the receptor (e.g., SCC, the vegetation/fauna community or habitat type present on the site) and Receptor Resilience (RR) (its resilience to impacts) as follows.

BI is a function of Conservation Importance (CI) and the Functional Integrity (FI) of the receptor as follows. The criteria for the CI and FI ratings are provided in Table 2-1 and Table 2-2, respectively.

Table 2-1 Summary of Conservation Importance (CI) criteria

Conservation Importance	Fulfilling Criteria
Very High	Confirmed or highly likely occurrence of Critically Endangered (CR), Endangered (EN), Vulnerable (VU) or Extremely Rare or CR species that have a global extent of occurrence (EEO) of < 10 km ² . Any area of natural habitat of a CR ecosystem type or large area (> 0.1% of the total ecosystem type extent) of natural habitat of an EN ecosystem type. Globally significant populations of congregatory species (> 10% of global population).
High	Confirmed or highly likely occurrence of CR, EN, VU species that have a global EEO of > 10 km ² . IUCN threatened species (CR, EN, VU) must be listed under any criterion other than A. If listed as threatened only under Criterion A, include if there are less than 10 locations or < 10 000 mature individuals remaining. Small area (> 0.01% but < 0.1% of the total ecosystem type extent) of natural habitat of EN ecosystem type or large area (> 0.1%) of natural habitat of VU ecosystem type. Presence of Rare species. Globally significant populations of congregatory species (> 1% but < 10% of global population).
Medium	Confirmed or highly likely occurrence of populations of Near Threatened (NT) species, threatened species (CR, EN, VU) listed under Criterion A only and which have more than 10 locations or more than 10 000 mature individuals. Any area of natural habitat of threatened ecosystem type with status of VU. Presence of range-restricted species. > 50% of receptor contains natural habitat with potential to support SCC.
Low	No confirmed or highly likely populations of SCC. No confirmed or highly likely populations of range-restricted species. < 50% of receptor contains natural habitat with limited potential to support SCC.
Very Low	No confirmed and highly unlikely populations of SCC. No confirmed and highly unlikely populations of range-restricted species. No natural habitat remaining.

Table 2-2 Summary of Functional Integrity (FI) criteria

Functional Integrity	Fulfilling Criteria
Very High	Very large (> 100 ha) intact area for any conservation status of ecosystem type or > 5 ha for CR ecosystem types. High habitat connectivity serving as functional ecological corridors, limited road network between intact habitat patches. No or minimal current negative ecological impacts, with no signs of major past disturbance.
High	Large (> 20 ha but < 100 ha) intact area for any conservation status of ecosystem type or > 10 ha for EN ecosystem types. Good habitat connectivity, with potentially functional ecological corridors and a regularly used road network between intact habitat patches. Only minor current negative ecological impacts, with no signs of major past disturbance and good rehabilitation potential.
Medium	Medium (> 5 ha but < 20 ha) semi-intact area for any conservation status of ecosystem type or > 20 ha for VU ecosystem types. Only narrow corridors of good habitat connectivity or larger areas of poor habitat connectivity and a busy used road network between intact habitat patches. Mostly minor current negative ecological impacts, with some major impacts and a few signs of minor past disturbance. Moderate rehabilitation potential.
Low	Small (> 1 ha but < 5 ha) area. Almost no habitat connectivity but migrations still possible across some modified or degraded natural habitat and a very busy used road network surrounds the area. Low rehabilitation potential.

Solar Facility

Several minor and major current negative ecological impacts.

Very Low	Very small (< 1 ha) area. No habitat connectivity except for flying species or flora with wind-dispersed seeds. Several major current negative ecological impacts.
-----------------	--

BI can be derived from a simple matrix of CI and FI as provided in Table 2-3.

Table 2-3 Matrix used to derive Biodiversity Importance (BI) from Functional Integrity (FI) and Conservation Importance (CI)

Biodiversity Importance (BI)		Conservation Importance (CI)				
		Very high	High	Medium	Low	Very low
Functional Integrity (FI)	Very high	Very high	Very high	High	Medium	Low
	High	Very high	High	Medium	Medium	Low
	Medium	High	Medium	Medium	Low	Very low
	Low	Medium	Medium	Low	Low	Very low
	Very low	Medium	Low	Very low	Very low	Very low

The fulfilling criteria to evaluate RR are based on the estimated recovery time required to restore an appreciable portion of functionality to the receptor, as summarised in Table 2-4.

Table 2-4 Summary of Receptor Resilience (RR) criteria

Resilience	Fulfilling Criteria
Very High	Habitat that can recover rapidly (~ less than 5 years) to restore > 75% of the original species composition and functionality of the receptor functionality, or species that have a very high likelihood of: (i) remaining at a site even when a disturbance or impact is occurring, or (ii) returning to a site once the disturbance or impact has been removed.
High	Habitat that can recover relatively quickly (~ 5–10 years) to restore > 75% of the original species composition and functionality of the receptor functionality, or species that have a high likelihood of: (i) remaining at a site even when a disturbance or impact is occurring, or (ii) returning to a site once the disturbance or impact has been removed.
Medium	Will recover slowly (~ more than 10 years) to restore > 75% of the original species composition and functionality of the receptor functionality, or species that have a moderate likelihood of: (i) remaining at a site even when a disturbance or impact is occurring, or (ii) returning to a site once the disturbance or impact has been removed.
Low	Habitat that is unlikely to be able to recover fully after a relatively long period: > 15 years required to restore ~ less than 50% of the original species composition and functionality of the receptor functionality, or species that have a low likelihood of: (i) remaining at a site even when a disturbance or impact is occurring, or (ii) returning to a site once the disturbance or impact has been removed.
Very Low	Habitat that is unable to recover from major impacts, or species that are unlikely to: (i) remain at a site even when a disturbance or impact is occurring, or (ii) return to a site once the disturbance or impact has been removed.

Solar Facility

Subsequent to the determination of the BI and RR, the SEI can be ascertained using the matrix as provided in Table 2-5.

Table 2-5 Matrix used to derive Site Ecological Importance from Receptor Resilience (RR) and Biodiversity Importance (BI)

Site Ecological Importance		Biodiversity Importance (BI)				
		Very high	High	Medium	Low	Very low
Receptor Resilience (RR)	Very Low	Very high	Very high	High	Medium	Low
	Low	Very high	Very high	High	Medium	Very low
	Medium	Very high	High	Medium	Low	Very low
	High	High	Medium	Low	Very low	Very low
	Very High	Medium	Low	Very low	Very low	Very low

Interpretation of the SEI in the context of the proposed project is provided in Table 2-6.

Table 2-6 Guidelines for interpreting Site Ecological Importance in the context of the proposed development activities

Site Ecological Importance	Interpretation in relation to proposed development activities
Very High	Avoidance mitigation – no destructive development activities should be considered. Offset mitigation not acceptable/not possible (i.e., last remaining populations of species, last remaining good condition patches of ecosystems/unique species assemblages). Destructive impacts for species/ecosystems where persistence target remains.
High	Avoidance mitigation wherever possible. Minimisation mitigation – changes to project infrastructure design to limit the amount of habitat impacted, limited development activities of low impact acceptable. Offset mitigation may be required for high impact activities.
Medium	Minimisation and restoration mitigation – development activities of medium impact acceptable followed by appropriate restoration activities.
Low	Minimisation and restoration mitigation – development activities of medium to high impact acceptable followed by appropriate restoration activities.
Very Low	Minimisation mitigation – development activities of medium to high impact acceptable and restoration activities may not be required.

The SEI evaluated for each taxon can be combined into a single multi-taxon evaluation of SEI for the assessment area. Either a combination of the maximum SEI for each receptor should be applied, or the SEI may be evaluated only once per receptor but for all necessary taxa simultaneously. For the latter, justification of the SEI for each receptor is based on the criteria that conforms to the highest CI and FI, and the lowest RR across all taxa.

3 Results & Discussion

3.1 Desktop Baseline

3.1.1 Ecologically Important Landscape Features

The GIS analysis pertaining to the relevance of the proposed project to ecologically important landscape features is summarised in Table 3-1.

Table 3-1 *Summary of relevance of the proposed project to ecologically important landscape features*

Desktop Information Considered	Relevant/Irrelevant	Section
Ecosystem Threat Status	Relevant – Overlaps with a Least Concern ecosystem	3.1.1.1
Ecosystem Protection Level	Relevant – Overlaps with a Moderately Protected Ecosystem	3.1.1.2
Critical Biodiversity Area	Relevant – The project area overlaps with CBA2, ESA1, NNR and ONA classified areas	3.1.1.3
Protected Areas	Relevant – The project area overlaps with the Rustenburg Platinum Mines (Union Section) Private Nature Reserve	3.1.1.4
National Protected Areas Expansion Strategy	Relevant – The project area overlaps with a NPAES protected area	3.1.1.5
South African Inventory of Inland Aquatic Ecosystems	Relevant - The project area overlaps with two CR wetlands and is adjacent to 1 CR wetland	3.1.1.5
National Freshwater Priority Area	Relevant – The project area overlaps with an unclassified FEPA wetland and an unclassified FEPA river	3.1.1.6
Strategic Water Source Areas	Irrelevant- The project area is 57 km from the closest SWSA	-
Renewable Energy Development Zones	Irrelevant - The project area is 167 km for the closest REDZ	-
Powerline Corridor	Irrelevant- The project area falls 88km from the Northern Corridor	-

3.1.1.1 Ecosystem Threat Status

The Ecosystem Threat Status is an indicator of an ecosystem's wellbeing, based on the level of change in structure, function or composition. Ecosystem types are categorised as Critically Endangered (CR), Endangered (EN), Vulnerable (VU), Near Threatened (NT) or Least Concern (LC), based on the proportion of the original extent of each ecosystem type that remains in good ecological condition. According to the spatial dataset the proposed project overlaps with a LC ecosystem (Figure 3-1).




Figure 3-1 Map illustrating the ecosystem threat status associated with the project area.

3.1.1.2 Ecosystem Protection Level

This is an indicator of the extent to which ecosystems are adequately protected or under-protected. Ecosystem types are categorised as Well Protected (WP), Moderately Protected (MP), Poorly Protected (PP), or Not Protected (NP), based on the proportion of the biodiversity target for each ecosystem type that is included within one or more protected areas. NP, PP or MP ecosystem types are collectively referred to as under-protected ecosystems. The proposed project overlaps with a MP ecosystem (Figure 3-2).




Figure 3-2 Map illustrating the ecosystem protection level associated with the project area

3.1.1.3 Critical Biodiversity Areas and Ecological Support Areas

The conservation of CBAs is crucial, in that if these areas are not maintained in a natural or near-natural state, biodiversity conservation targets cannot be met. Maintaining an area in a natural state can include a variety of biodiversity compatible land uses and resource uses (SANBI-BGIS, 2017).

The provincial CBA spatial data for the North West province indicates that both feasibility areas don't traverse any CBA nor Ecological Support Areas (ESAs) and Other Natural Areas (ONAs). Based on the Limpopo Conservation Plan the SBPM feasibility area traverses ESA1 and NNR areas, whereas the SBPM feasibility area traverses ESA1, NNR and ONA area.

The purpose of the Limpopo C-Plan (2018) is to inform land-use planning and development on a provincial scale and to aid in natural resource management. One of the outputs is a map of Critical Biodiversity Areas (CBAs) and Ecological Support Areas (ESAs). These are classified into different categories, namely Protected Areas, CBA1 areas, CBA2 areas, ESA1 areas, ESA2 areas, Other Natural Areas (ONAs) and areas with No Natural Habitat Remaining (NNR) based on biodiversity characteristics, spatial configuration, and requirements for meeting targets for both biodiversity patterns and ecological processes.

Figure 3-3 shows the project area superimposed on the Terrestrial CBA maps. The project area overlaps with CBA2, ESA1, NNR and ONA classified areas. Development in these areas is feasible, but developments other than the preferred biodiversity-compatible land-uses should be investigated in detail and the mitigation hierarchy applied.




Figure 3-3 Map illustrating the locations of CBAs in the project area

3.1.1.4 Protected areas

According to the protected area spatial datasets from SAPAD (2021), the project area overlaps with the Rustenburg Platinum Mines (Union Section) Private Nature Reserve (Figure 3-4). From the imagery, and confirmed by the site visit, the portion of the reserve in which the project area is located is comprised of an old tailings dam in various stages of rehabilitation and is therefore not considered ecologically sensitive. Several additional private nature reserves are in close proximity to the project area. These are the Leopard Hills, Animalia, Youngs and Leeuwkopje private nature reserves. All of these reserves are within 5km of the project area which means that the project area is within the buffer zone of the nature reserves.




Figure 3-4 The project area in relation to the protected areas

3.1.1.5 National Protected Area Expansion Strategy

National Protected Area Expansion Strategy 2016 (NPAES) areas were identified through a systematic biodiversity planning process. They present the best opportunities for meeting the ecosystem-specific protected area targets set in the NPAES and were designed with a strong emphasis on climate change resilience and requirements for protecting freshwater ecosystems. These areas should not be seen as future boundaries of protected areas, as in many cases only a portion of a particular focus area would be required to meet the protected area targets set in the NPAES. They are also not a replacement for finescale planning which may identify a range of different priority sites based on local requirements, constraints and opportunities (NPAES, 2016). The project area overlaps with an NPAES protected area as can be seen in Figure 3-5. However as mentioned above the portion of the protected area in which the project area is located is comprised of an old tailings dam in various stages of rehabilitation and is therefore not considered ecologically sensitive.




Figure 3-5 The project area in relation to the National Protected Area Expansion Strategy

3.1.1.6 Hydrological Setting

The South African Inventory of Inland Aquatic Ecosystems (SAIIAE) was released with the NBA 2018. Ecosystem threat status (ETS) of river and wetland ecosystem types are based on the extent to which each river ecosystem type had been altered from its natural condition. Ecosystem types are categorised as CR, EN, VU or LT, with CR, EN and VU ecosystem types collectively referred to as ‘threatened’ (Van Deventer *et al.*, 2019; Skowno *et al.*, 2019). The project area borders on a CR wetland and CR NBA River (Figure 3-6). The reach of the Sefathlane River proximal to the area is regarded as CR.




Figure 3-6 Map illustrating ecosystem threat status of rivers and wetland ecosystems in the project area

3.1.1.7 National Freshwater Ecosystem Priority Area Status

In an attempt to better conserve aquatic ecosystems, South Africa has categorised its river systems according to set ecological criteria (i.e., ecosystem representation, water yield, connectivity, unique features, and threatened taxa) to identify Freshwater Ecosystem Priority Areas (FEPAs) (Driver *et al.*, 2011). The FEPAs are intended to be conservation support tools and envisioned to guide the effective implementation of measures to achieve the National Environment Management Biodiversity Act’s (NEM:BA) biodiversity goals (Nel *et al.*, 2011).

Figure 3-7 shows the project area overlaps with unclassified FEPA wetlands and unclassified FEPA rivers.




Figure 3-7 The project area in relation to the National Freshwater Ecosystem Priority Areas.

3.1.1.8 Renewable Energy Database

The Renewable Energy Database (<http://egis.environment.gov.za/>), shows that there are limited other projects in the near vicinity (Figure 3-8). This reduces the overall impact on the habitats in the area.

Solar Facility




Figure 3-8 The project area in relation to the renewable energy database projects in the area.

3.1.2 Flora Assessment

This section is divided into a description of the vegetation type expected to occur under natural conditions and the expected flora species.

3.1.2.1 Vegetation Type

The project area is situated in the Savanna biome. The savanna vegetation of South Africa represents the southernmost extension of the most widespread biome in Africa (Mucina & Rutherford, 2006). Major macroclimatic traits that characterise the Savanna biome include:

- a) Seasonal precipitation; and
- b) (Sub) tropical thermal regime with no or usually low incidence of frost (Mucina & Rutherford, 2006).

Most savanna vegetation communities are characterised by a herbaceous layer dominated by grasses and a discontinuous to sometimes very open tree layer (Mucina & Rutherford, 2006).

The savanna biome is the largest biome in South Africa, extending throughout the east and north-eastern areas of the country. Savannas are characterised by a dominant grass layer, over-topped by a discontinuous, but distinct woody plant layer. At a structural level, Africa’s savannas can be broadly categorised as either fine-leaved (microphyllous) savannas or broad-leaved savannas. Fine-leaved savannas typically occur on nutrient rich soils and are dominated by microphyllous woody plants of the Mimosaceae family (Common genera include *Acacia* and *Albizia*) and a generally dense herbaceous layer (Scholes & Walker, 1993).

On a fine-scale vegetation type, the project area overlaps with the Dwaalboom Thornveld vegetation type (Figure 3-9).




Figure 3-9 Map illustrating the vegetation type associated with the project area

3.1.2.1.1 Dwaalboom Thornveld

Dwaalboom Thornveld is restricted to, and is distributed in, the Limpopo and North-West Provinces within flats north of the Dwarsberge and associated ridges mainly west of the Crocodile River in the Dwaalboom area but including a patch around Sentrum. South of the ridges it extends eastwards from the Nietverdiend area, north of the Pilanesberg to the Northam area at an altitude range of between 900 and 1,200m AMSL. Its main vegetation and landscape features include plains with a layer of scattered, low to medium high, deciduous microphyllous trees and shrubs with a few broad-leaved tree species. There is almost a continuous herbaceous layer dominated by grass species.

Important Plant Taxa in Dwaalboom Thornveld

Based on Mucina and Rutherford’s (2006) vegetation classification, important plant taxa are those species that have a high abundance, a frequent occurrence (not being particularly abundant) or are prominent in the landscape within a particular vegetation type. They note the following species are important taxa in the Dwaalboom Thornveld vegetation type:

Trees: *Vachellia erioloba*, *Vachellia erubescens*, *Vachellia nilotica*, *Vachellia tortilis* subsp *heteracantha*, *Senegalia fleckii*, *Senegalia burkei*, *Searsia lancea* (Mucina & Rutherford, 2006).

Shrubs: *Diospyros lycioides* subsp. *lycioides*, *Grewia flava*, *Mystroxyton aethiopicum* subsp. *burkenum*, *Agathisanthemum bojeri* (Mucina & Rutherford, 2006).

Graminoids: *Aristida bipartite*, *Bothriochloa insculpta*, *Digitaria eriantha* subsp *eriantha*, *Ischaemum afrum*, *Panicum maximum* and *Cymbopogon pospischilii* (Mucina & Rutherford, 2006).

Conservation Status

According to Mucina and Rutherford (2006) Dwaalboom Thornveld is classified as Least Threatened. Although the target for conservation is 19%, only 6%of this vegetation type is currently under statutory

Solar Facility

conservation in reserves such as the Madikwe Game Reserve (approximately 150km west of the project area). Cultivation and to a lesser extent urbanisation have resulted in the transformation of approximately 14% of Dwaalboom Thornveld and exotic invasive plants are present. Incidences of erosion are low to very low (Mucina & Rutherford, 2006).

3.1.2.2 Expected Flora Species

The POSA database indicates that 428 species of indigenous plants are expected to occur within the project area (The full list of species will be provided in the final report). One (1) SCC based on their conservation status could be expected to occur within the project area and are provided in Table 3-2 below. It is believed that additional SCC will be recorded in the assessment. During the screening assessment a number of protected *Vachellia erioloba* (Camel thorn) trees were recorded within the SBPM feasibility area. This is a nationally protected tree.

Table 3-2 Threatened flora species that may occur within the project area.

Family	Taxon	Author	IUCN	Ecology
Apocynaceae	<i>Stenostelma umbelluliferum</i>	(Schltr.) Bester & Nicholas	NT	Indigenous; Endemic

3.1.3 Faunal Assessment

3.1.3.1 Amphibians

Based on the IUCN Red List Spatial Data and AmphibianMap, 30 amphibian species are expected to occur within the area (The full list will be provided in the final assessment). One (1) are regarded as threatened (Table 3-3).

Table 3-3 Threatened amphibian species that are expected to occur within the project area

Species	Common Name	Conservation Status		Likelihood of occurrence
		Regional (SANBI, 2016)	IUCN (2021)	
<i>Pyxicephalus adspersus</i>	Giant Bullfrog	NT	LC	Moderate

Giant Bull Frog (*Pyxicephalus adspersus*) is a species of conservation concern that will possibly occur in the project area, especially in the area with the wetlands. The Giant Bull Frog is listed as near threatened on a regional scale. It is a species of drier savannas where it is fossorial for most of the year, remaining buried in cocoons. They emerge at the start of the rains, and breed in shallow, temporary waters in pools, pans and ditches (IUCN, 2017).

3.1.3.2 Reptiles

Based on the IUCN Red List Spatial Data and the ReptileMAP database, 91 reptile species are expected to occur within the area (The full list will be provided in the final assessment). Three (3) are regarded as threatened, all having a low likelihood of occurrence (Table 3-4).

Table 3-4 Threatened reptile species that are expected to occur within the project area

Species	Common Name	Conservation Status		Likelihood of Occurrence
		Regional (SANBI, 2016)	IUCN (2022)	
<i>Crocodylus niloticus</i>	Nile Crocodile	VU	VU	Low
<i>Lygodactylus waterbergensis</i>	Waterberg Dwarf Gecko	NT	NT	Low
<i>Pseudocordylus transvaalensis</i>	Northern Crag Lizard	NT	NT	Low

3.1.3.3 Mammals

The IUCN Red List Spatial Data lists 85 mammal species that could be expected to occur within the area (The full list will be provided in the final assessment). This list includes large mammal species that are normally restricted to protected areas, as these were observed during the screening assessment. Twelve (12) (smaller non protected area restricted species) of these expected species are regarded as threatened

Solar Facility

(Table 3-5), five of these have a low likelihood of occurrence based on the lack of suitable habitat and food sources in the project area.

Table 3-5 Threatened mammal species that are expected to occur within the project area.

Species	Common Name	Conservation Status		Likelihood of occurrence
		Regional (SANBI, 2016)	IUCN (2022)	
<i>Aonyx capensis</i>	Cape Clawless Otter	NT	NT	Moderate
<i>Atelerix frontalis</i>	South Africa Hedgehog	NT	LC	Moderate
<i>Cloeotis percivali</i>	Short-eared Trident Bat	EN	LC	Moderate
<i>Crocidura mariquensis</i>	Swamp Musk Shrew	NT	LC	Moderate
<i>Crocuta crocuta</i>	Spotted Hyaena	NT	LC	Low
<i>Felis nigripes</i>	Black-footed Cat	VU	VU	Moderate
<i>Leptailurus serval</i>	Serval	NT	LC	Confirmed
<i>Panthera pardus</i>	Leopard	VU	VU	Low
<i>Parahyaena brunnea</i>	Brown Hyaena	NT	NT	Confirmed
<i>Poecilogale albinucha</i>	African Striped Weasel	NT	LC	Low
<i>Redunca fulvorufula</i>	Mountain Reedbuck	EN	LC	Low
<i>Smutsia temminckii</i>	Temminck's Ground Pangolin	VU	VU	Low

Aonyx capensis (Cape Clawless Otter) is the most widely distributed otter species in Africa (IUCN, 2017). This species is predominantly aquatic, and it is seldom found far from water. This species has a moderate likelihood of occurrence based on the rivers in the project area.

Atelerix frontalis (South African Hedgehog) has a tolerance to a degree for habitat modification and occurs in a wide variety of semi-arid and sub-temperate habitats (IUCN, 2017). Based on the Red List of Mammals of South Africa, Lesotho and Swaziland (2016), *A. frontalis* populations are decreasing due to the threats of electrocution, veld fires, road collisions, predation from domestic pets and illegal harvesting. Suitable grasslands occur in the project area, although somewhat disturbed, that can function as habitat for this species, as such the likelihood of occurrence is rated as moderate.

Cloeotis percivali (Short-eared Trident Bat) occurs in savanna areas where there is sufficient cover in the form of caves and mine tunnels for day roosting (IUCN, 2017). It feeds exclusively on moths and appears to be very sensitive to disturbance. Suitable habitat and roosting area can be found around the project area and therefore the likelihood of finding this species is rated as moderate.

Crocidura maquassiensis (Maquassie Musk Shrew) is listed as VU on a regional basis and is known to be found in rocky, mountain habitats. It may tolerate a wider range of habitats and individuals have been collected in Kwa-Zulu Natal from a garden, and in mixed bracken and grassland alongside a river at 1,500 m (IUCN, 2017). This species has a moderate likelihood of occurring based on the rocky habitat found in the project area.

Felis nigripes (Black-footed cat) is endemic to the arid regions of southern Africa. This species is naturally rare, has cryptic colouring, is small in size and is nocturnal. These factors have contributed to a lack of information on this species. The highest densities of this species have been recorded in the more arid Karoo region of South Africa. The habitat in the project area can be considered to be somewhat suitable for the species and the likelihood of occurrence is therefore rated as moderate.

Leptailurus serval (Serval) occurs widely through sub-Saharan Africa and is commonly recorded from most major national parks and reserves (IUCN, 2017). The Serval's status outside reserves is not certain, but they are inconspicuous and may be common in suitable habitat as they are tolerant of farming practices provided there is cover and food available. In sub-Saharan Africa they are found in habitat with well-watered savanna long-grass environments and are particularly associated with reedbeds and other

Solar Facility

riparian vegetation types. Suitable habitat is present for this species in the project area, as such the likelihood of occurrence is rated as high. This species was confirmed during the field assessment.

Parahyaena brunnea (Brown Hyaena) is endemic to southern Africa. This species occurs in dry areas, generally with annual rainfall less than 100 mm, particularly along the coast, semi-desert, open scrub and open woodland savanna. Given its known ability to persist outside of formally protected areas the likelihood of occurrence of this species in the project area is moderate to good. This species was confirmed during the screening assessment.

3.1.4 Literature Review

ENVASS conducted an updated biodiversity assessment for the Siyanda Bakgatla development that were conducted in 2020 is based on a 2006 baseline study conducted by Engelbrecht and Grosel (2006). It was reported that since 2006;

- Two Nationally Protected tree species were recorded, *Boscia albitrunca* (Shepard's Tree) and *Vachellia erioloba* (Camelthorn) with 4 Provincially Protected Species (NW);
- 35 mammal species were recorded in and around their project area specifically in the game farm and watercourse habitats. Of the 35 species that have been recorded, six (6) sighted are provincially protected within the North West and four (4) are considered SCC under the IUCN Red List, namely Cape Clawless Otter (*Aonyx capensis*) (NT), South Africa Hedgehog (*Atelerix frontalis*) (NT), Brown Hyaena (*Parahyaena brunnea*) (NT) as well as Plains Zebra (*Equus zebra*)(NT) however considered a captive species;
- Ten of the expected 20 reptile species was sighted within the area, three of these species said to be nationally protected, namely: *Python natalensis* (African Python), *Varanus albigularis* (Rock Monitor) and *Varanus niloticus* (Water Monitor); and
- Eleven of the expected 23 amphibian species was sighted within the area, with one considered SCC, namely *Pyxicephalus adspersus* (Giant Bullfrog) which is NT.

The Ecological Sensitivity Assessment was conducted using the same method as in section 2.3 of this report, it was determined that the habitats identified by the author (Mixed Bushveld, Watercourses and Vachellia Thornveld habitat unit units) were determined to be of a high sensitivity in the broader environment (Figure 3-10).




Figure 23: Illustration of the ecological sensitivity of the study area.

Figure 3-10 Map illustrating ecological sensitivity according to the literature review

3.2 Field Assessment

The following sections provide the results from the field survey for the proposed development that was undertaken from the 4th till the 7th of April 2022.

3.2.1 Flora Assessment

This section is divided into two sections:

- Indigenous flora; and
- Invasive Alien Plants (IAPs).

3.2.1.1 Indigenous flora

The vegetation assessment was conducted throughout the extent of the project area. A total of 84 tree, shrub and herbaceous plant species were recorded in the project area during the field assessment (Table 3-6). Plants listed as Category 1 alien or invasive species under the NEMBA appear in green text. Plants listed in Category 2 or as 'not indigenous' or 'naturalised' according to NEMBA, appear in blue text. Some of the plant species recorded can be seen in Figure 3-11.

Table 3-6 Trees, shrub and herbaceous plant species recorded in the project area.

Family	Scientific Name	Threat Status (SANBI, 2017)	SA Endemic	NEMBA Category
Acanthaceae	<i>Crabbea angustifolia</i>	LC	Yes	
Amaranthaceae	<i>Achyranthes aspera</i>			Naturalized exotic weed
Amaranthaceae	<i>Alternanthera pungens</i>	LC	Not Endemic	
Amaranthaceae	<i>Pupalia lappacea</i>	LC	No	
Amaranthaceae	<i>Hermibstaedtia odorata</i>	LC	No	
Amaryllidaceae	<i>Crinum crassicaule</i>	LC	No	
Amaryllidaceae	<i>Ammocharis coranica</i>	LC	Not Endemic	
Anacardiaceae	<i>Searsia lancea</i>	LC	No	
Anacardiaceae	<i>Searsia pyroides</i> var. <i>pyroides</i>	LC	No	
Apocynaceae	<i>Pentarrhinum insipidum</i>	LC	No	
Asparagaceae	<i>Asparagus cooperi</i>	LC	No	
Asparagaceae	<i>Asparagus laricinus</i>	LC	No	
Asteraceae	<i>Bidens pilosa</i>			Naturalized exotic weed
Asteraceae	<i>Conyza bonariensis</i>			Naturalized exotic weed
Asteraceae	<i>Flaveria bidentis</i>			NEMBA Category 1B
Asteraceae	<i>Hilliardiella elaeagnoides</i>	LC	Not Endemic	
Asteraceae	<i>Nidorella resedifolia</i>	LC	No	
Asteraceae	<i>Schkuhria pinnata</i>			Naturalized exotic weed
Asteraceae	<i>Tagetes minuta</i>			Naturalized exotic weed
Asteraceae	<i>Tithonia diversifolia</i>			NEMBA Category 1B
Asteraceae	<i>Xanthium spinosum</i>			NEMBA Category 1B
Asteraceae	<i>Zinnia peruviana</i>			Naturalized exotic weed
Asteraceae	<i>Argemone ochroleuca</i>			NEMBA Category 1b.
Asteraceae	<i>Cirsium vulgare</i>			NEMBA Category 1b.

Solar Facility

Asteraceae	<i>Verbena bonariensis</i>			NEMBA Category 1b.
Boraginaceae	<i>Ehretia rigida</i>	LC	Yes	
Boraginaceae	<i>Heliotropium amplexicaule</i>			Naturalized exotic weed
Cactaceae	<i>Opuntia ficus-indica</i>			NEMBA Category 1B
Celastraceae	<i>Gymnosporia buxifolia</i>	LC	No	
Combretaceae	<i>Combretum erythrophyllum</i>	LC	No	
Commelinaceae	<i>Commelina africana var. barberae</i>	LC	No	
Commelinaceae	<i>Commelina africana var. barberae</i>	LC	Not Endemic	
Commelinaceae	<i>Commelina erecta</i>	LC	No	
Commelinaceae	<i>Commelina livingstonii</i>	LC	Not Endemic	
Convolvulaceae	<i>Cuscuta campestris</i>			Naturalized exotic weed
Convolvulaceae	<i>Evolvulus alsinoides</i>	LC	Not Endemic	
Convolvulaceae	<i>Seddera capensis</i>	LC	No	
Cucurbitaceae	<i>Cucumis zeyheri</i>	LC	No	
Ebenaceae	<i>Diospyros lycioides subsp. lycioides</i>	LC	No	
Fabaceae	<i>Dichrostachys cinerea</i>	LC	No	
Fabaceae	<i>Elephantorrhiza elephantina</i>	LC	Not Endemic	
Fabaceae	<i>Peltophorum africanum</i>	LC	No	
Fabaceae	<i>Senegalia mellifera</i>	LC	No	
Fabaceae	<i>Sesbania bispinosa</i>			Naturalized exotic weed
Fabaceae	<i>Vachellia gerrardii subsp. gerrardii</i>	LC	No	
Fabaceae	<i>Vachellia hebeclada subsp. hebeclada</i>	LC	No	
Fabaceae	<i>Vachellia karoo</i>	LC	No	
Fabaceae	<i>Vachellia nilotica</i>	LC	No	
Fabaceae	<i>Vachellia tortilis</i>	LC	No	
Fabaceae	<i>Vachellia tenuispina</i>	LC	No	
Iridaceae	<i>Gladiolus elliotii</i>	LC	No	
Iridaceae	<i>Gladiolus sericeovillosus subsp. calvatus</i>	LC	Not Endemic	
Lamiaceae	<i>Leonotis schinzii</i>	LC	No	
Malvaceae	<i>Abutilon austro-africanum</i>	LC	No	
Malvaceae	<i>Grewia flava</i>	LC	No	
Malvaceae	<i>Grewia flavescens</i>	LC	No	
Malvaceae	<i>Hibiscus pusillus</i>	LC	Not Endemic	
Malvaceae	<i>Hibiscus trionum</i>			Naturalized exotic weed
Malvaceae	<i>Sida dregei</i>	LC	Not Endemic	
Nymphaeaceae	<i>Nymphaea lotus</i>	LC	Not Endemic	
Orobanchaceae	<i>Striga asiatica</i>	LC	No	
Pedaliaceae	<i>Ceratotheca triloba</i>	LC	No	
Pedaliaceae	<i>Dicercaryum senecioides</i>	LC	Not Endemic	

Solar Facility

Poaceae	<i>Arundo donax</i>			NEMBA Category 1B
Poaceae	<i>Bothriochloa insculpta</i>	LC	No	
Poaceae	<i>Brachiaria xantholeuca</i>	LC	Not Endemic	
Poaceae	<i>Cenchrus ciliaris</i>	LC	No	
Poaceae	<i>Chloris virgata</i>	LC	No	
Poaceae	<i>Cymbopogon caesius</i>	LC	No	
Poaceae	<i>Cynodon dactylon</i>	LC	No	
Poaceae	<i>Dichanthium annulatum var. papillosum</i>	LC	Not Endemic	
Poaceae	<i>Digitaria eriantha</i>	LC	No	
Poaceae	<i>Diheteropogon amplexans</i>	LC	No	
Poaceae	<i>Dinebra retroflexa var. condensata</i>	LC	Not Endemic	
Poaceae	<i>Enneapogon cenchroides</i>	LC	No	
Poaceae	<i>Eragrostis rigidior</i>	LC	No	
Poaceae	<i>Fingerhuthia africana</i>	LC	No	
Poaceae	<i>Heteropogon contortus</i>	LC	No	
Poaceae	<i>Ischaemum afrum</i>	LC	No	
Poaceae	<i>Melinis repens</i>	LC	No	
Poaceae	<i>Panicum maximum</i>	LC	No	
Poaceae	<i>Panicum maximum</i>	LC	Not Endemic	
Poaceae	<i>Perotis patens</i>	LC	Not Endemic	
Poaceae	<i>Phragmites australis</i>	LC	Not Endemic	
Poaceae	<i>Setaria sphacelata var. sphacelata</i>	LC	No	
Poaceae	<i>Setaria verticillata</i>	LC	No	
Poaceae	<i>Sorghum versicolor</i>	LC	No	
Poaceae	<i>Themeda triandra</i>	LC	Not Endemic	
Poaceae	<i>Tragus berteronianus</i>			Naturalized exotic weed
Poaceae	<i>Tragus berteronianus</i>			Naturalized exotic weed
Poaceae	<i>Urochloa mosambicensis</i>	LC	No	
Poaceae	<i>Hyparrhenia hirta</i>	LC	Not Endemic	
Rhamnaceae	<i>Ziziphus mucronata</i>	LC	No	
Rubiaceae	<i>Kohautia amatymbica</i>	LC	Not Endemic	
Sapindaceae	<i>Pappea capensis</i>	LC	No	
Solanaceae	<i>Datura ferox</i>			NEMBA Category 1B
Solanaceae	<i>Lycium horridum</i>	LC	No	
Solanaceae	<i>Solanum campylacanthum</i>	LC	No	
Zygophyllaceae	<i>Tribulus terrestris</i>	LC	No	




Figure 3-11 Photographs illustrating some of the flora recorded within the assessment area. A) *Hermbstaedtia odorata*, B) *Vachellia nilotica*, C) *Nymphaea lotus*, D) *Heliotropium amplexicaule* and E) *Hibiscus pusillus*

3.2.1.2 Invasive Alien Plants

Invasive Alien Plants (IAPs) tend to dominate or replace indigenous flora, thereby transforming the structure, composition and functioning of ecosystems. Therefore, it is important that these plants are controlled by means of an eradication and monitoring programme. Some invader plants may also degrade ecosystems through superior competitive capabilities to exclude native plant species.

NEMBA is the most recent legislation pertaining to alien invasive plant species. In August 2014, the list of Alien Invasive Species was published in terms of the NEMBA. The Alien and Invasive Species Regulations were published in the Government Gazette No. 43726, 18 September 2020. The legislation calls for the removal and / or control of AIP species (Category 1 species). In addition, unless authorised thereto in terms of the NWA, no land user shall allow Category 2 plants to occur within 30 meters of the 1:50 year flood line of a river, stream, spring, natural channel in which water flows regularly or intermittently, lake, dam or wetland. Category 3 plants are also prohibited from occurring within proximity to a watercourse. Below is a brief explanation of the three categories in terms of the NEMBA:

- *Category 1a:* Invasive species requiring compulsory control. Remove and destroy. Any specimens of Category 1a listed species need, by law, to be eradicated from the environment. No permits will be issued.
- *Category 1b:* Invasive species requiring compulsory control as part of an invasive species control programme. Remove and destroy. These plants are deemed to have such a high invasive potential that infestations can qualify to be placed under a government sponsored invasive species management programme. No permits will be issued.
- *Category 2:* Invasive species regulated by area. A demarcation permit is required to import, possess, grow, breed, move, sell, buy or accept as a gift any plants listed as Category 2 plants. No permits will be issued for Category 2 plants to exist in riparian zones.
- *Category 3:* Invasive species regulated by activity. An individual plant permit is required to undertake any of the following restricted activities (import, possess, grow, breed, move, sell, buy or accept as a gift) involving a Category 3 species. No permits will be issued for Category 3 plants to exist in riparian zones.

Note that according to the Alien and Invasive Species Regulations, a person who has under his or her control a category 1b listed invasive species must immediately:

- Notify the competent authority in writing
- Take steps to manage the listed invasive species in compliance with:
 - Section 75 of the NEMBA;
 - The relevant invasive species management programme developed in terms of regulation 4; and
 - Any directive issued in terms of section 73(3) of the NEMBA.

Four (4) IAP species were recorded within the project area. These species are listed under the Alien and Invasive Species List 2020, Government Gazette No. GN1003 as Category 1b. These IAP species must be controlled by implementing an IAP Management Programme, in compliance of section 75 of the NEMBA, as stated above.

3.2.2 Faunal Assessment

Herpetofauna and mammal observations and recordings fall under this section. A separate avifaunal report was compiled for this project.

3.2.2.1 Amphibians and Reptiles

Four (4) species of reptile and one species were recorded within the project area during the survey period (Table 3-7, Figure 3-12). However, there is the possibility of more species being present, as certain reptile species are secretive and require long-term surveys to ensure capture. None of the species recorded are regarded as threatened, albeit all are protected under provincial legislation.

Table 3-7 Summary of herpetofauna species recorded within the project area.

Species	Common Name	Conservation Status	
		Regional (SANBI, 2016)	Global (IUCN, 2022)
Reptiles			
<i>Acanthocercus atricollis</i>	Tree agama	LC	LC
<i>Lygodactylus capensis</i>	Cape dwarf gecko	LC	LC
<i>Stigmochelys pardalis</i>	Leopard Tortoise	LC	LC
<i>Trachylepis punctatissima</i>	Speckled Rock Skink	LC	Unlisted
<i>Trachylepis varia</i>	Variable Skink	LC	LC
<i>Varanus niloticus</i>	Water Monitor	LC	Unlisted
Amphibians			
<i>Cacosternum boettgeri</i>	Common Caco	LC	LC
<i>Tomopterna cryptotis</i>	Tremelo Sand Frog	LC	LC




Figure 3-12 Photographs illustrating the reptile species recorded within the assessment area associated with the project area during the survey period. A) *Tomopterna cryptotis*, B) *Lygodactylus capensis*, and C) *Varanus niloticus*, D) *Stigmochelys pardalis* and E) *Acanthocercus atricollis*.

3.2.2.2 Mammals

Twenty-one (21) mammal species were observed in total based on either direct observation or the presence of visual tracks and signs. Fourteen of these species could naturally occur outside of protected areas/game farms, while six species are considered mainly found restricted to protected areas/game farms as 'captive' species (Table 3-8) (Figure 3-13). Four SCC were observed, were two of the SCC are regarded as threatened species that could occur naturally (Figure 3-14).

Table 3-8 Summary of mammal species recorded within the project area. Mammal species are considered 'captive' species as these were only present within the game farm areas, marked in green text

Species	Common Name	Conservation Status	
		Regional (SANBI, 2016)	IUCN (2017)
<i>Aepyceros melampus</i>	Impala	LC	LC
<i>Alcelaphus buselaphus</i>	Hartebeest	LC	LC
<i>Canis mesomelas</i>	Black-backed Jackal	LC	LC
<i>Chlorocebus pygerythrus</i>	Vervet Monkey	LC	LC
<i>Connochaetes taurinus</i>	Blue Wildebeest	LC	LC
<i>Damaliscus pygargus phillipsi</i>	Blesbok	LC	LC
<i>Equus quagga</i>	Plains Zebra	LC	NT
<i>Genetta genetta</i>	Small-spotted Genet	LC	LC
<i>Giraffa camelopardalis</i>	Giraffe	LC	VU
<i>Herpestes sanguineus</i>	Slender Mongoose	LC	LC
<i>Hystrix africaeaustralis</i>	Cape Porcupine	LC	LC
<i>Kobus ellipsiprymnus</i>	Common Waterbuck	LC	LC
<i>Leptailurus serval</i>	Serval	NT	LC
<i>Lepus saxatilis</i>	Scrub Hare	LC	LC
<i>Mungos mungo</i>	Banded Mongoose	LC	LC
<i>Papio ursinus</i>	Chacma Baboon	LC	LC
<i>Paraxerus cepapi</i>	Tree Squirrel	LC	LC
<i>Parahyaena brunnea</i>	Brown Hyaena	NT	NT
<i>Phacochoerus africanus</i>	Common Warthog	LC	LC
<i>Raphicerus campestris</i>	Steenbok	LC	LC
<i>Tragelaphus strepsiceros</i>	Greater Kudu	LC	LC




Figure 3-13 Photographs illustrating the mammal species recorded within the project area during the survey period. A) *Equus quagga* and *Alcelaphus buselaphus*, B) *Mungos mungo*, C) *Damaliscus pygargus phillipsi*, D) *Chlorocebus pygerythrus* and E) *Giraffa camelopardalis*



Figure 3-14 Photographs illustrating a mammal SCC recorded within the project area during the survey period: *Leptailurus serval*

4 Habitat Assessment and Site Ecological Importance

4.1 Habitat Assessment

The main habitat types identified across the project area were initially identified largely based on aerial imagery. These main habitat types were refined based on the field coverage and data collected during the survey; the delineated habitats can be seen in Figure 4-1. Emphasis was placed on limiting timed meander searches within the natural habitats and therefore habitats with a higher potential of hosting SCC. Five habitats were identified in the project area, each of the habitats identified are discussed in the sub-sections below.




Figure 4-1 Habitats identified in the project area.

Degraded Bushveld

This habitat type is regarded as semi-natural bushveld, but slightly disturbed due adjacent land use, mismanagement (overgrazing) and also human infringement as it is being used as a game farm (Figure 4-2 and Figure 4-3). This habitat represents typical bushveld, with rocky extrusions in certain areas. The current ecological condition of this habitat regarding the main driving forces, are intact, which is evident in the amount and importance of the species recorded in the faunal assessment; and the high species diversity and number of plant species recorded. Current human infringement still occurs throughout, especially in areas close to roads. The difference between this habitat and the disturbed bushveld is the extent of the disturbance in the disturbed bushveld being more severe. Portions of this bushveld has been fragmented by linear infrastructure such as powerlines and fences and are conspired fragmented bushveld.

This habitat unit can be regarded as important, not only within the local landscape, but also regionally. The unit functions as remaining greenlands which supports viable plant species populations and is also used for foraging. The unit also serves as a movement corridor for fauna within a landscape fragmented. The habitat sensitivity is regarded as high sensitivity due to the role of this intact habitat to biodiversity within an area being more fragmented locally, which is supported by the various ecological datasets. This habitat functions as the CBA 2 it is classified as and portions classified as ESA, may be considered just as important.



Figure 4-2 A typical example of degraded Bushveld habitat from the project area.



Figure 4-3 A typical example of fragmented Bushveld habitat from the project area.

Disturbed Bushveld

This habitat is regarded as areas that have been impacted more by historic grazing, mismanagement and land use (Figure 4-4). Historical vegetation clearing for cultivation has led to localised bush encroachment and the presence of alien floral species within this habitat unit, with current grazing activities by livestock also taking place within this area. These habitats aren't entirely transformed but in a constant disturbed state, as they can't recover to a more natural state due to ongoing disturbances and impacts received from grazing and mismanagement. This habitat can be found in different conditions of disturbance, but in many cases has either been encroached on by *Dichrostachys cinerea*, *Vachellia tenuispina* or *Senegalia mellifera*. These areas are considered to have a low sensitivity, as they may be used as a movement corridor and in many cases form a barrier between the more degraded bushveld and the transformed areas.



Figure 4-4 A typical example of disturbed Bushveld habitat from the project area.

Transformed

This habitat unit represents all areas of urban area, agriculture, mining areas as well as the associated secondary roads (Figure 4-5 and Figure 4-6). The transformed areas have little to no remaining natural vegetation due to land transformation by urban and mining infrastructure, agriculture as well as informal homesteads. These habitats exist in a constant disturbed state as it cannot recover to a more natural state due to ongoing disturbances and impacts it receives.




Figure 4-5 Illustration of transformed habitat from the project area.



Figure 4-6 *Illustration of transformed habitat from the project area.*

Wetlands

Wetlands are identified in the wetland report (TBC, 2022). Even though somewhat disturbed, the ecological integrity, importance and functioning of these areas play a crucial role as a water resource system and an important habitat for various fauna and flora, including the SCC recorded (Figure 4-7 and Figure 4-8). The preservation of this system, which includes the CR Sefathlane River is the most important aspect to consider for the proposed project. This habitat needs to be protected and improved due to the role of this habitat as a water resource.



Figure 4-7 *Illustration of wetland habitat from the project area (Sefathlane River)*



Figure 4-8 *Illustration of wetland habitat from the project area*

4.2 Site Ecological Importance (SEI)

The biodiversity theme sensitivity, as indicated in the screening report, was derived to be Very High, mainly due to the project area being with an CBA 2 and ESA 1 (Figure 4-9), while the animal and plant species theme sensitivity shows that majority of the area is classified as medium and low sensitivity respectively.




Figure 4-9 Terrestrial Biodiversity Theme Sensitivity, TBC Screening Report

Four (4) different terrestrial habitat types were delineated within the project area, and one set of wetland habitats as a whole. Based on the criteria provided in Section 2.3 of this report, all habitats within the assessment area of the proposed project were allocated a sensitivity category Table 4-1. The sensitivities of the habitat types delineated are illustrated in Figure 4-10.

Table 4-1 Summary of habitat types delineated within the project area

Habitat (Area)	Conservation Importance	Functional Integrity	Biodiversity Importance	Receptor Resilience	Site Ecological Importance
Degraded Bushveld	Medium	High	Medium	Low	High
Wetlands	Medium	Medium	Medium	Low	High
Fragmented Bushveld	Medium	Low	Medium	Low	Medium

Disturbed Bushveld	Low	Low	Low	Medium	Low
Transformed	Very Low	Very Low	Very Low	Low	Very Low




Figure 4-10 Sensitivity of the project area

5 Impact Risk Assessment

5.1 Biodiversity: Risk Assessment Method

The assessment of the significance of direct, indirect and cumulative impacts was undertaken using the method as developed by Savannah. The assessment of the impact considers the following, the:

- Nature of the impact, which shall include a description of what causes the effect, what will be affected, and how it will be affected;
- Extent of the impact, indicating whether the impact will be local or regional;
- Duration of the impact, very short-term duration (0-1 year), short-term duration (2-5 years), medium-term (5-15 years), long-term (> 15 years) or permanent;
- Probability of the impact, describing the likelihood of the impact actually occurring, indicated as improbable, probable, highly probable or definite;
- Severity/beneficial scale, indicating whether the impact will be very severe/beneficial (a permanent change which cannot be mitigated/permanent and significant benefit with no real alternative to achieving this benefit); severe/beneficial (long-term impact that could be mitigated/long-term benefit); moderately severe/beneficial (medium- to long-term impact that could be mitigated/ medium- to long-term benefit); slight; or have no effect;
- Significance, which shall be determined through a synthesis of the characteristics described above and can be assessed as low medium or high;
- Status, which will be described as either positive, negative or neutral;
- Degree to which the impact can be reversed;
- Degree to which the impact may cause irreplaceable loss of resources; and
- Degree to which the impact can be mitigated.

5.1.1 Present Impacts to Biodiversity

Considering the anthropogenic activities and influences within the landscape, several negative impacts to biodiversity were observed within the project area. These include:

- Mining activities;
- Present energy distribution infrastructure, including powerlines;
- Wood harvesting;
- Historical cattle grazing land-use;
- Invasive species;
- Roads and associated vehicle traffic and road kills; and
- Fences.



Figure 5-1 *Photographs illustrating impacts to biodiversity A) Roads and fencing), B) Powerlines and associated servitude C) Wood harvesting D) Livestock, roads and railway and E) Livestock and litter*

5.1.2 Identification of Additional Potential Impacts

The potential impacts during the construction and operation phases of the project are presented in Table 5-1, and the infrastructure in relation to the SEI can be seen in Figure 5-2.

Table 5-1 Potential impacts to biodiversity associated with the proposed activity

Main Impact	Project activities that can cause loss/impacts to habitat (especially with regard to the proposed infrastructure areas):	Secondary impacts anticipated
1. Destruction, fragmentation and degradation of habitats and ecosystems	Physical removal of vegetation, including protected species.	Displacement/loss of flora & fauna (including possible SCC)
	Access roads and servitudes	Increased potential for soil erosion
	Soil dust precipitation	Habitat fragmentation
	Dumping of waste products	Increased potential for establishment of alien & invasive vegetation
	Random events such as fire (cooking fires or cigarettes)	Erosion
Main Impact	Project activities that can cause the spread and/or establishment of alien and/or invasive species	Secondary impacts anticipated
2. Spread and/or establishment of alien and/or invasive species	Vegetation removal	Habitat loss for native flora & fauna (including SCC)
	Vehicles potentially spreading seed	Spreading of potentially dangerous diseases due to invasive and pest species
	Unsanitary conditions surrounding infrastructure promoting the establishment of alien and/or invasive rodents	Alteration of fauna assemblages due to habitat modification
	Creation of infrastructure suitable for breeding activities of alien and/or invasive birds	
Main Impact	Project activities that can cause direct mortality of fauna	Secondary impacts anticipated
3. Direct mortality of fauna	Clearing of vegetation	Loss of habitat
		Loss of ecosystem services
	Roadkill due to vehicle collision	
	Pollution of water resources due to dust effects, chemical spills, etc.	Increase in rodent populations and associated disease risk
	Intentional killing of fauna for food (hunting)	
Main Impact	Project activities that can cause reduced dispersal/migration of fauna	Secondary impacts anticipated
4. Reduced dispersal/migration of fauna	Loss of landscape used as corridor	Reduced dispersal/migration of fauna
		Loss of ecosystem services
	Compacted roads	Reduced plant seed dispersal
	Removal of vegetation	
Main Impact	Project activities that can cause pollution in watercourses and the surrounding environment	Secondary impacts anticipated
5. Environmental pollution due to water runoff, spills from vehicles and erosion	Chemical (organic/inorganic) spills	Pollution in watercourses and the surrounding environment
		Faunal mortality (direct and indirectly)
	Erosion	Groundwater pollution
		Loss of ecosystem services
Main Impact	Project activities that can cause disruption/alteration of ecological life cycles due to sensory disturbance.	Secondary impacts anticipated
6. Disruption/alteration of ecological life cycles (breeding,	Operation of machinery (Large earth moving machinery, vehicles)	Disruption/alteration of ecological life cycles due to noise

<p>migration, feeding) due to noise, dust and light pollution.</p>	<p>Project activities that can cause disruption/alteration of ecological life cycles due to dust</p>	<p>Loss of ecosystem services Secondary impacts associated with disruption/alteration of ecological life cycles due to dust</p>
<p>Main Impact</p>	<p>Vehicles Project activities that can cause staff to interact directly with potentially dangerous fauna</p>	<p>Loss of ecosystem services Secondary impacts anticipated</p>
<p>8. Staff and others interacting directly with fauna (potentially dangerous) or poaching of animals</p>	<p>All unregulated/supervised activities outdoors</p>	<p>Loss of SCCs</p>




Figure 5-2 Proposed infrastructure layout overlaid with the SEI.

5.1.3 Assessment of Impact Significance

The assessment of impact significance was undertaken in accordance with the method developed by Savannah. The various identified impacts are assessed below for the different phases of the development. The impacts assessed may be re-assessed if an exact infrastructure layout has been provided

5.1.3.1 Construction Phase

The following potential main impacts on the biodiversity (based on the framework above) were considered for the construction phase of the proposed development. This phase refers to the period during construction when the proposed features are constructed; and is considered to have the largest direct impact on biodiversity. The actual footprint of the pole/pylon infrastructure has a small localised, impact. It is the clearance for the PV and substation areas as well as creation of access and service roads that is a more important aspect to consider and will be considered in relation to the powerlines. The following potential impacts to terrestrial biodiversity were considered:

- Destruction, further loss and fragmentation of the of habitats, ecosystems and vegetation community (Table 5-2),
- Introduction of alien species, especially plants (Table 5-3); and
-
- Displacement of faunal community due to habitat loss, direct mortalities and disturbance (road collisions, noise, dust, vibration and poaching) (Table 5-4).

Table 5-2 Impacts to biodiversity associated with the proposed construction phase.

Impact Nature: Loss of vegetation within development footprint		
Destruction, further loss and fragmentation of the of habitats, ecosystems and vegetation community		
	Without mitigation	With mitigation
Extent	High (4)	Moderate (3)
Duration	Permanent (5)	Moderate term (3)
Magnitude	Moderate (6)	Low (4)
Probability	Highly probable (4)	Probable (3)
Significance	Medium	Medium
Status (positive or negative)	Negative	Negative
Reversibility	Moderate	High
Irreplaceable loss of resources?	No	No
Can impacts be mitigated?	Yes, although this impact cannot be well mitigated as the loss of vegetation/habitat is unavoidable.	
Mitigation:		
See Biodiversity Management Outcomes		
Residual Impacts:		
The loss of currently intact vegetation is an unavoidable consequence of the project and cannot be entirely mitigated.		

Table 5-3 Impacts to biodiversity associated with the proposed construction phase.

Impact Nature: Introduction of alien species, especially plants		
Degradation and loss of surrounding natural vegetation		
	Without mitigation	With mitigation

Impact Nature: Introduction of alien species, especially plants		
Degradation and loss of surrounding natural vegetation		
Extent	High (4)	Low (2)
Duration	Long term (4)	Short term (2)
Magnitude	Moderate (6)	Minor (2)
Probability	Highly probable (4)	Improbable (2)
Significance	Medium (56)	Low (12)
Status (positive or negative)	Negative	Negative
Reversibility	Moderate	High
Irreplaceable loss of resources?	No	No
Can impacts be mitigated?	Yes	
Mitigation:		
See Biodiversity Management Outcomes		
Residual Impacts:		
Long-term broad scale. IAP infestation if not mitigated.		

Table 5-4 Impacts to biodiversity associated with the proposed construction phase.

Impact Nature: Displacement of faunal community due to habitat loss, direct mortalities and disturbance		
Construction activity will likely lead to direct mortality of fauna due to earthworks, vehicle collisions, accidental hazardous chemical spills and persecution. Disturbance due to dust and noise pollution and vibration may disrupt behaviour.		
	Without mitigation	With mitigation
Extent	High (4)	Low (2)
Duration	Moderate term (3)	Very short term (1)
Magnitude	Moderate (6)	Low (4)
Probability	Highly probable (4)	Improbable (2)
Significance	Medium	Low
Status (positive or negative)	Negative	Negative
Reversibility	Moderate	High
Irreplaceable loss of resources?	No	No
Can impacts be mitigated?	Yes, to some extent. Noise and disturbance cannot be well mitigated, impacts on fauna due to human presence, such as vehicle collisions, poaching, and persecution can be mitigated.	
Mitigation:		
See Biodiversity Management Outcomes		
Residual Impacts:		
It is probable that some individuals of susceptible species will be lost to construction-related activities despite mitigation. However, this is not likely to impact the viability of the local population of any fauna species.		

5.1.3.2 Operation Phase

The operational phase of the impact of daily activities is anticipated to further spread the IAP, as well as the deterioration of the habitats due to the increase of dust and edge effect impacts. Dust reduces the ability of plants to photosynthesize and thus leads to degradation/retrogression of the veld. Moving

maintenance and mining vehicles don't only cause sensory disturbances to fauna, affecting their life cycles and movement, but will lead to direct mortalities due to collisions.

The following potential impacts were considered:

- Continued fragmentation and degradation of habitats and ecosystems (Table 5-5);
- Spread of alien and/or invasive species (Table 5-6);
- Ongoing displacement and direct mortalities of faunal community (including SCC) due to disturbance (road collisions, collisions with substation, noise, light, dust, vibration) (Table 5-7).

Table 5-5 Impacts to biodiversity associated with the proposed operational phase

Impact Nature: Continued fragmentation and degradation of habitats and ecosystems		
Disturbance created during the construction phase will leave the project area vulnerable to erosion and IAP encroachment.		
	Without Mitigation	With Mitigation
Extent	Moderate (3)	Low (2)
Duration	Permanent (5)	Short term (2)
Magnitude	High (8)	Low (4)
Probability	Highly probable (4)	Improbable (2)
Significance	High	Low
Status (positive or negative)	Negative	Negative
Reversibility	Moderate	High
Irreplaceable loss of resources?	No	No
Can impacts be mitigated?	Yes, with proper management and avoidance, this impact can be mitigated to a low level.	
Mitigation:		
See Biodiversity Management Outcomes		
Residual Impacts		
There is still the potential some potential for erosion and IAP encroachment even with the implementation of control measures but would have a low impact.		

Table 5-6 Impacts to biodiversity associated with the proposed operational phase.

Impact Nature: Spread of alien and/or invasive species		
Degradation and loss of surrounding natural vegetation		
	Without mitigation	With mitigation
Extent	Moderate (3)	Low (2)
Duration	Long term (4)	Short term (2)
Magnitude	Moderate (6)	Minor (2)
Probability	Highly probable (4)	Improbable (2)
Significance	Medium	Low
Status (positive or negative)	Negative	Negative
Reversibility	Moderate	High
Irreplaceable loss of resources?	No	No
Can impacts be mitigated?	Yes	
Mitigation:		

Impact Nature: Spread of alien and/or invasive species
Degradation and loss of surrounding natural vegetation
See Biodiversity Management Outcomes
Residual Impacts:
Long term broad scale IAP infestation if not mitigated.

Table 5-7 Impacts to biodiversity associated with the proposed operational phase

Impact Nature: Ongoing displacement and direct mortalities of faunal community (including SCC) due to disturbance (road collisions, collisions with substation, noise, light, dust, vibration)		
The operation and maintenance of the proposed development may lead to disturbance or persecution of fauna in the vicinity of the development.		
	Without Mitigation	With Mitigation
Extent	Moderate (3)	Low (2)
Duration	Long term (4)	Short term (2)
Magnitude	High (8)	Low (4)
Probability	Highly probable (4)	Improbable (2)
Significance	Medium	Low
Status (positive or negative)	Negative	Negative
Reversibility	Moderate	High
Irreplaceable loss of resources?	No	No
Can impacts be mitigated?	Yes	
Mitigation:		
See Biodiversity Management Outcomes		
Residual Impacts		
Disturbance from maintenance activities will occur albeit at a low and infrequent level.		

5.1.3.3 Cumulative Impacts

Cumulative impacts are assessed in context of the extent of the proposed project area; other developments in the area; and general habitat loss and transformation resulting from other activities in the area.

Impact Nature: Cumulative habitat loss within the region		
The development of the proposed infrastructure will contribute to cumulative habitat loss within ESAs and thereby impact the ecological processes in the region.		
	Overall impact of the proposed development considered in isolation	Cumulative impact of the project and other projects in the area
Extent	Moderate (3)	High (4)
Duration	Moderate term (3)	Long term (4)
Magnitude	Moderate (6)	High (8)
Probability	Probable (3)	Probable (3)
Significance	Medium (15)	Medium (19)
Status (positive or negative)	Negative	Negative
Reversibility	High	High

Irreplaceable loss of resources?	No	No
Can impacts be mitigated	To some degree, but most of the impact results from the presence of the various facilities which cannot be well mitigated.	
Mitigation:		
Ensure that a rehabilitation plan and IAP management plan be compiled for each development and are effectively implemented.		

5.1.4 Biodiversity Management Outcomes

The purpose of the management outcomes is to allow for the mitigation measures associated with the impact assessment to be incorporated into the EMPr. These are provided in Table 5-8.

Table 5-8 Mitigation measures including requirements for timeframes, roles and responsibilities for this report

Impact Management Actions	Implementation		Monitoring	
	Phase	Responsible Party	Aspect	Frequency
Management outcome: Vegetation and Habitats				
Areas rated as High sensitivity in proximity to the development areas, should be declared as 'no-go' areas during the life of the project, and all efforts must be made to prevent access to this area from construction workers, machinery. The infrastructure should be realigned to prioritise development within very low/low sensitivity areas. Mitigated development in Moderate sensitivity areas is permissible. High sensitivity areas are to be avoided.	Life of operation	Project manager, Environmental Officer	Development footprint	Ongoing
Areas of indigenous vegetation, even secondary communities outside of the direct project footprint, should under no circumstances be fragmented or disturbed further than that proposed for the project. Clearing of vegetation should be minimized and avoided where possible.	Life of operation	Project manager, Environmental Officer	Areas of indigenous vegetation	Ongoing
Where possible, existing access routes and walking paths must be made use of.	Construction/Operational Phase	Environmental Officer & Design Engineer	Roads and paths used	Ongoing
All laydown, chemical toilets etc. should be restricted to very low/ low sensitivity areas. Any materials may not be stored for extended periods of time and must be removed from the project area once the construction/closure phase has been concluded. No storage of vehicles or equipment will be allowed outside of the designated project areas.	Construction/Operational Phase	Environmental Officer & Design Engineer	Laydown areas	Ongoing
Areas that are denuded during construction need to be re-vegetated with indigenous vegetation to prevent erosion during flood and wind events. This will also reduce the likelihood of encroachment by alien invasive plant species.	Operational phase	Environmental Officer & Contractor	Assess the state of rehabilitation and encroachment of alien vegetation	Quarterly for up to two years after the closure
Any woody material removed can be shredded and used in conjunction with the topsoil to augment soil moisture and prevent further erosion.	Operational and Decommissioning phase	Environmental Officer & Contractor	Woody material under powerline and in SS footprint	During Phase
A hydrocarbon spill management plan must be put in place, to ensure that should there be any chemical spill out or over that it does not run into the surrounding areas. The Contractor shall be in possession of an emergency spill kit that must always be complete and available on site. Drip trays or any form of oil absorbent material must be placed underneath vehicles/machinery and equipment when not in use. No servicing of equipment may occur on site, unless necessary. All contaminated soil / yard stone shall be treated in situ or removed and be placed in containers. Appropriately contain any generator diesel storage tanks, machinery spills (e.g. accidental spills of hydrocarbons oils, diesel etc.) in such a way as to prevent them leaking and entering the environment.	Life of operation	Environmental Officer & Contractor	Spill events, Vehicles dripping.	Ongoing

Storm Water run-off & Discharge Water Quality monitoring	Life of operation	Environmental Officer & Design Engineer	Water Quality and presence of erosion	Ongoing
It should be made an offence for any staff to take/ bring any plant species into/out of any portion of the project area. No plant species whether indigenous or exotic should be brought into/taken from the project area, to prevent the spread of exotic or invasive species or the illegal collection of plants.	Life of operation	Project manager, Environmental Officer	Any instances	Ongoing
A fire management plan needs to be compiled and implemented to restrict the impact fire might have on the surrounding areas.	Life of operation	Environmental Officer & Contractor	Fire Management	During Phase

Management outcome: Fauna

Impact Management Actions	Implementation		Monitoring	
	Phase	Responsible Party	Aspect	Frequency
The areas to be developed must be specifically demarcated to prevent movement of staff or any individual into the surrounding environments, <ul style="list-style-type: none"> Signs must be put up to enforce this 	Construction/Operational Phase	Project manager, Environmental Officer	Infringement into these areas	Ongoing
Noise must be kept to an absolute minimum during the evenings and at night, to minimize all possible disturbances to amphibian species and nocturnal mammals	Construction/Operational Phase	Environmental Officer	Noise levels	Ongoing
No trapping, killing, or poisoning of any wildlife is to be allowed. <ul style="list-style-type: none"> Signs must be put up to enforce this; 	Life of operation	Environmental Officer	Evidence of trapping etc	Ongoing
Outside lighting should be designed and limited to minimize impacts on fauna. All outside lighting should be directed away from highly sensitive areas. Fluorescent and mercury vapor lighting should be avoided and sodium vapor (green/red) lights should be used wherever possible.	Construction/Operational Phase	Project manager, Environmental Officer & Design Engineer	Light pollution and period of light.	Ongoing
All construction and maintenance motor vehicle operators should undergo an environmental induction that includes instruction on the need to comply with speed limits, to respect all forms of wildlife. Speed limits must still be enforced to ensure that road killings and erosion is limited.	Life of operation	Health and Safety Officer	Compliance to the training.	Ongoing
Schedule activities and operations during least sensitive periods, to avoid migration, nesting and breeding seasons.	Life of operation	Project manager, Environmental Officer & Design Engineer	Activities should take place during the day in the case.	Ongoing
Any excavations or holes must be conducted in a progressive manner. <ul style="list-style-type: none"> Should the holes/excavations stay open overnight they must be covered temporarily, to ensure no small fauna species fall in. 	Planning and construction	Environmental Officer & Contractor, Engineer	Presence of trapped animals and open holes	Ongoing
A qualified environmental control officer must be on site when construction begins. A site walk through is recommended by a suitably qualified ecologist prior to any construction activities, preferably during the wet season and any SSC should be noted. In situations where the threatened and protected plants must be removed, the proponent may only do so after the required permission/permits have been obtained in accordance with	Construction Phase	Environmental Officer, Contractor	Presence of any floral or faunal species.	During phase

Solar Facility

national and provincial legislation. In the abovementioned situation the development of a search, rescue and recovery program is suggested for the protection of these species. Should animals not move out of the area on their own relevant specialists must be contacted to advise on how the species can be relocated				
Heat generated from the substation must be monitored to ensure it does not negatively affect the local fauna	Life of operation	Environmental Officer & Contractor	Heat generated by substations	Ongoing
Once the development layout has been confirmed, the open areas must be fenced off appropriately pre-construction in order to allow animals to move or be moved into these areas before breaking ground activities occur. Construction activities must take place systemically, especially in relation to the game farm area.	Planning/Construction Phase	Environmental Officer & Design Engineer	Areas not to be developed and construction direction	Ongoing
Management outcome: Alien Vegetation and fauna				
Impact Management Actions	Implementation		Monitoring	
	Phase	Responsible Party	Aspect	Frequency
Compilation of and implementation of an alien vegetation management plan.	Life of operation	Project manager, Environmental Officer & Contractor	Assess presence and encroachment of alien vegetation	Twice a year
The footprint area of the construction should be kept to a minimum. The footprint area must be clearly demarcated to avoid unnecessary disturbances to adjacent areas	Construction/Operational Phase	Project manager, Environmental Officer & Contractor	Footprint Area	Life of operation
Waste management must be a priority and all waste must be collected and stored adequately. It is recommended that all waste be removed from site on a weekly basis to prevent rodents and pests entering the site	Life of operation	Environmental Officer & Health and Safety Officer	Presence of waste	Life of operation
A pest control plan must be put in place and implemented; it is imperative that poisons not be used due to the likely presence of SCCs	Life of operation	Environmental Officer & Health and Safety Officer	Evidence or presence of pests	Life of operation
Management outcome: Dust				
Impact Management Actions	Implementation		Monitoring	
	Phase	Responsible Party	Aspect	Frequency
Dust-reducing mitigation measures must be put in place and strictly adhered to. This includes wetting of exposed soft soil surfaces. <ul style="list-style-type: none"> No non environmentally friendly suppressants may be used, as this could result in pollution of water sources 	Life of operation	Contractor	Dustfall	Dust monitoring program.
Management outcome: Waste management				
Impact Management Actions	Implementation		Monitoring	
	Phase	Responsible Party	Aspect	Frequency

Solar Facility

<p>Waste management must be a priority and all waste must be collected and stored adequately. It is recommended that all waste be removed from site on a weekly basis to prevent rodents and pests entering the site.</p> <ul style="list-style-type: none"> • Refuse bins will be emptied and secured; • Temporary storage of domestic waste shall be in covered waste skips; and • Maximum domestic waste storage period will be 10 days. <p>Toilets at the recommended Health and Safety standards must be provided. These should be emptied twice a day, to prevent staff from using the surrounding vegetation.</p> <p>The Contractor should supply sealable and properly marked domestic waste collection bins and all solid waste collected shall be disposed of at a licensed disposal facility. Under no circumstances may domestic waste be burned on site</p> <p>Refuse bins will be emptied and secured. Temporary storage of domestic waste shall be in covered waste skips. Maximum domestic waste storage period will be 10 days.</p> <p>Suitable temporary solid waste facilities are to be incorporated into the design to prevent unsanitary conditions. These are to be cleared weekly and waste collected by the local waste management department. The residents must be encouraged to recycle.</p>	Construction Phase	Environmental Officer & Health and Safety Officer	Presence of waste	Life of operation
	Construction Phase	Environmental Officer & Health and Safety Officer	Number of toilets per staff member. Waste levels	Daily
	Construction Phase	Environmental Officer & Health and Safety Officer	Availability of bins and the collection of the waste.	Ongoing
	Construction Phase	Environmental Officer, Contractor & Health and Safety Officer	Management of bins and collection of waste	Ongoing
	Operational Phase	Project manager	Management of bins and collection of waste	Ongoing

Management outcome: Environmental awareness training

Impact Management Actions	Implementation		Monitoring	
	Phase	Responsible Party	Aspect	Frequency
All personnel and contractors to undergo Environmental Awareness Training. A signed register of attendance must be kept for proof. Discussions are required on sensitive environmental receptors within the project area to inform contractors and site staff of the presence of Red / Orange List species, their identification, conservation status and importance; and biology, habitat requirements and management requirements in the EA and EMPr. The avoidance and protection of the wetland areas must be included into a site induction. Contractors and employees must all undergo the induction and made aware of the "no-go" to be avoided.	Life of operation	Health and Safety Officer	Compliance to the training.	Ongoing

Management outcome: Erosion

Impact Management Actions	Implementation		Monitoring	
	Phase	Responsible Party	Aspect	Frequency
Speed limits must be put in place to reduce erosion.	Life of operation	Project manager, Environmental Officer	Water Runoff from road surfaces	Ongoing

Solar Facility

<ul style="list-style-type: none"> Reducing the dust generated by the listed activities above, especially the earthmoving machinery, through wetting the soil surface; putting up signs to enforce speed limit; and speed bumps built to force slow speeds; Signs must be put up to enforce this. 				
Where possible, existing access routes and walking paths must be made use of.	Life of operation	Project manager, Environmental Officer	Routes used within the area	Ongoing
Areas that are denuded during construction need to be re-vegetated with indigenous vegetation, to prevent erosion during flood events and strong winds.	Life of operation	Project manager, Environmental Officer	Re-establishment of indigenous vegetation	Progressively
A stormwater management plan must be compiled and implemented.	Life of operation	Project manager, Environmental Officer	Management plan	Before construction phase: Ongoing

6 Conclusion and Impact Statement

6.1 Conclusion

6.1.1 Terrestrial Biodiversity

The project area has been altered both currently and historically. The present land use had a direct impact on both the fauna and the flora in the area, which is evident in the disturbed and transformed habitats. Historically, Cultivation, overgrazing and mismanagement has led to the deterioration of most of the area to a disturbed Bushveld that is either encroached upon or invaded by exotic plant species.

However, the degraded Bushveld habitat and wetlands/watercourses in the wider project area can be regarded as important, not only within the local landscape, but also regionally; as they are used for habitat, foraging and movement corridors for fauna within a landscape fragmented by development.

The degraded Bushveld habitat and wetlands/watercourses in the project area have a **High ecological theme sensitivity**.

The habitat sensitivity of the degraded Bushveld and wetland/water resources is regarded as high, due to the species recorded and the role of this intact unique habitat to biodiversity within a very fragmented local landscape, not to mention the sensitivity according to various ecological datasets. The high sensitivity terrestrial areas still:

- Support nearby CBA/ESA's as per the LCP;
- and
- Support various organisms (Including SCC) and may play an important role in the ecosystem, if left to recover from the superficial impacts.

The ecological integrity, importance and functioning of these terrestrial biodiversity areas provide a variety of ecological services considered beneficial, with one key service being the maintenance of biodiversity. The preservation of these systems is the most important aspect to consider for the proposed project.

The results of the Ecological Sensitivity Assessment by (ENVASS,2020) and this report are in corroboration.

Any development in high sensitivity areas must be conducted with the provided mitigation measures.. The mitigation measures, management and associated monitoring regarding the expected impacts will be the most important factor of this project and must be considered by the issuing authority.

6.2 Impact Statement

The main expected impacts of the proposed infrastructure will include the following:

- Habitat loss and fragmentation;
- Degradation of surrounding habitat;
- Disturbance and displacement caused during the construction and maintenance phases; and
- Direct mortality during the construction phase.

Mitigation measures as described in this report can be implemented to reduce the significance of the risk but there is always possibility of unexpected impacts. Considering that this area that has been identified as being of significance for biodiversity maintenance and ecological processes, development may proceed but with caution and only with the implementation of mitigation measures.

Considering the above-mentioned information, no fatal flaws are evident for the proposed project. It is the opinion of the specialists that the project, may be favourably considered, on condition that all prescribed mitigation measures and supporting recommendations are implemented.

7 References

Bates, M.F., Branch, W.R., Bauer, A.M., Burger, M., Marais, J., Alexander, G.J & de Villiers, M.S. (Eds). 2014. Atlas and Red List of Reptiles of South Africa, Lesotho and Swaziland. Suricata 1. South African Biodiversity Institute, Pretoria.

BirdLife International. 2016a. *Afrotis afra*. The IUCN Red List of Threatened Species 2016: e.T22691975A93331501. <http://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T22691975A93331501.en>.

BGIS (Biodiversity GIS). (2017). <http://bgis.sanbi.org/>

BODATSA-POSA. (2021). Plants of South Africa - an online checklist. POSA ver. 3.0. <http://newposa.sanbi.org/>.

Boycott, R. and Bourquin, R. 2000. The Southern African Tortoise Book – A Guide to Southern African Tortoises, Terrapins and Turtles. Revised Edition. Hilton. 228 pages.

Branch, W.R. (1998). Field Guide to Snakes and Other Reptiles of Southern Africa. Struik, Cape Town.

Du Preez, L. & Carruthers, V. (2009) A Complete Guide to the Frogs of Southern Africa. Struik Nature, Cape Town.

Department of Water Affairs and Forestry (DWAF). 2005. A practical field procedure for identification and delineation of wetlands and riparian areas. Pretoria: Department of Water Affairs and Forestry.

EWT. (2016). Mammal Red List 2016. www.ewt.org.za

ENVASS (2020). Updated biodiversity assessment of the Siyanda Bakgatla (pty) ltd. Platinum mine area situated within the Thabazimbi and Moses Kotane local municipalities of the Limpopo province, South Africa. BIO-REP-250-19_20

Fish, L., Mashau, A.C., Moeaha, M.J. & Nembudani, M.T. (2015). Identification Guide to Southern African Grasses: An Identification Manual with Keys, Descriptions, and Distributions. SANBI, Pretoria.

IUCN. (2021). The IUCN Red List of Threatened Species. www.iucnredlist.org

Johnson, S. & Bytebier, B. (2015). Orchids of South Africa: A Field Guide. Struik publishers, Cape Town.

Kotze, D.C., Marneweck, G.C., Batchelor, A.L., Lindley, D.C. & Collins, N.B. (2009). A Technique for rapidly assessing ecosystem services supplied by wetlands. Mondi Wetland Project.

Land Type Survey Staff. (1972 - 2006). Land Types of South Africa: Digital Map (1:250 000 Scale) and Soil Inventory Databases. Pretoria: ARC-Institute for Soil, Climate, and Water.

Macfarlane DM and Bredin IP. 2017. Part 1: technical manual. Buffer zone guidelines for wetlands, rivers and estuaries

Macfarlane, D.M., Bredin, I.P., Adams, J.B., Zungu, M.M., Bate, G.C., Dickens, C.W.S. (2014). Preliminary guideline for the determination of buffer zones for rivers, wetlands and estuaries. Final Consolidated Report. WRC Report No TT 610/14, Water Research Commission, Pretoria.

Macfarlane, D.M., Dickens, J. & Von Hase, F. (2009). Development of a methodology to determine the appropriate buffer zone width and type for developments associated with wetlands, watercourses and estuaries Deliverable 1: Literature Review. INR Report No: 400/09.

Mucina, L. & Rutherford, M.C. (Eds.). 2006. The vegetation of South Africa, Lesotho and Swaziland. Strelizia 19. South African National Biodiversity Institute, Pretoria, South African.

Mucina, L., Rutherford, M.C. & Powrie, L.W. (Eds.). 2007. Vegetation map of South Africa, Lesotho and Swaziland. 1:1 000 000 scale sheet maps. 2nd ed. South African National Biodiversity Institute, Pretoria.

Nel JL, Murray KM, Maherry AM, Petersen CP, Roux DJ, Driver A, Hill L, Van Deventer H, Funke N, Swartz ER, Smith-Adao LB, Mbona N, Downsborough L and Nienaber S. 2011. Technical Report for the National Freshwater Ecosystem Priority Areas project. WRC Report No. K5/1801.

Ollis DJ, Snaddon CD, Job NM, and Mbona N. 2013. Classification System for Wetlands and other Aquatic Ecosystems in South Africa. User Manual: Inland Systems. SANBI Biodiversity Series 22. South African Biodiversity Institute, Pretoria.

Raimonde, D. (2009). Red list of South African Plants. SANBI, Pretoria.

Rountree, M.W. and Kotze, D.M. 2013. Manual for the Rapid Ecological Reserve Determination of Inland Wetlands (Version 2.0). Joint Department of Water Affairs/Water Research Commission Study. Report No 1788/1/12. Water Research Commission, Pretoria.

SADAP (South Africa Protected Areas Database) and SACAD (South Africa Conservation Areas Database) (2021). <http://egis.environment.gov.za>

SANBI. 2013. Grasslands Ecosystem Guidelines: landscape interpretation for planners and managers. Compiled by Cadman, M., de Villiers, C., Lechmere-Oertel, R. and D. McCulloch. South African National Biodiversity Institute, Pretoria. 139 pages.

SANBI-BGIS. 2017. Technical guidelines for CBA Maps: Guidelines for developing a map of Critical Biodiversity Areas & Ecological Support Areas using systematic biodiversity planning.

Skowno, A.L., Raimondo, D.C., Poole, C.J., Fizzotti, B. & Slingsby, J.A. (eds.). 2019. South African National Biodiversity Assessment 2018 Technical Report Volume 1: Terrestrial Realm. South African National Biodiversity Institute, Pretoria.

Smith, B. (2006). The Farming Handbook. Netherlands & South Africa: University of KwaZulu-Natal Press & CTA.

Soil Classification Working Group. (1991). Soil Classification A Taxonomic system for South Africa. Pretoria: The Department of Agricultural Development.

Soil Classification Working Group. (2018). Soil Classification A Taxonomic system for South Africa. Pretoria: The Department of Agricultural Development.

Van Deventer, H., Smith-Adao, L., Collins, N.B., Grenfell, M., Grundling, A., Grundling, P-L., Impson, D., Job, N., Lötter, M., Ollis, D., Petersen, C., Scherman, P., Sieben, E., Snaddon, K., Tererai, F. and Van der Colff D. 2019. *South African National Biodiversity Assessment 2018: Technical Report*. Volume 2b: Inland Aquatic (Freshwater) Realm. CSIR report number CSIR/NRE/ECOS/IR/2019/0004/A. South African National Biodiversity Institute, Pretoria. <http://hdl.handle.net/20.500.12143/6230>.

Van Deventer, H., Smith-Adao, L., Mbona, N., Petersen, C., Skowno, A., Collins, N.B., Grenfell, M., Job, N., Lötter, M., Ollis, D., Scherman, P., Sieben, E. & Snaddon, K. 2018. South African National Biodiversity Assessment 2018: Technical Report. Volume 2a: South African Inventory of Inland Aquatic Ecosystems (SAIIAE). Version 3, final released on 3 October 2019. Council for Scientific and Industrial Research (CSIR) and South African National Biodiversity Institute (SANBI): Pretoria, South Africa.

8 Appendix Items

8.1 Appendix A – Specialist Declaration of Independence

I, Lindi Steyn, declare that:

- I act as the independent specialist in this application;
- I will perform the work relating to the application in an objective manner, even if this results in views and findings that are not favourable to the applicant;
- I declare that there are no circumstances that may compromise my objectivity in performing such work;
- I have expertise in conducting the specialist report relevant to this application, including knowledge of the Act, regulations and any guidelines that have relevance to the proposed activity;
- I will comply with the Act, regulations, and all other applicable legislation;
- I have no, and will not engage in, conflicting interests in the undertaking of the activity;
- I undertake to disclose to the applicant and the competent authority all material information in my possession that reasonably has or may have the potential of influencing any decision to be taken with respect to the application by the competent authority; and the objectivity of any report, plan, or document to be prepared by myself for submission to the competent authority;
- All the particulars furnished by me in this form are true and correct; and
- I realise that a false declaration is an offence in terms of Regulation 71 and is punishable in terms of Section 24F of the Act.



Lindi Steyn

Biodiversity Specialist

The Biodiversity Company

February 2022