

11 ADDENDUM: PALAEOLOGICAL SPECIALIST STUDY



PALAEONTOLOGICAL SPECIALIST STUDY: COMBINED DESKTOP AND FIELD-BASED ASSESSMENTS

Proposed solar power generation facilities on the remaining extent of the farm Vetlaagte No. 4, De Aar, Northern Cape Province

John E. Almond PhD (Cantab.)
Natura Viva cc, PO Box 12410 Mill Street,
Cape Town 8010, RSA
naturaviva@universe.co.za

May 2012

1. SUMMARY

It is proposed to develop seven small (30-75 MW) photovoltaic solar power generation facilities with associated infrastructure within an area of c. 935 ha on the Farm Vetlaagte, located 7 km east of De Aar, Pixley ka Seme District Municipality, Northern Cape.

The potentially fossiliferous sediments of the Late Palaeozoic Karoo Supergroup (Ecca and Lower Beaufort Groups) that underlie the study area are almost entirely mantled in a thick layer of superficial deposits of probable Pleistocene to Recent age. These include various soils, gravels and – at least in some areas - a well-developed calcrete hardpan. The upper Ecca Group bedrocks in the northern portion of the study area contain locally abundant fossil wood (of palaeontological interest for dating and palaeoenvironmental studies), as well as low diversity non-marine trace fossil assemblages typical of the Waterford Formation, rather than the Tierberg Formation as mapped.

No vertebrate fossils and only scattered woody plant impressions of the Permian *Glossopteris* Flora were observed within the Lower Beaufort Group rocks that are very poorly exposed in the southern portion of the Vetlaagte study area. Trace fossils, silicified wood and rare vertebrate remains (therapsids, parareptiles) of the Middle Permian *Pristerognathus* Assemblage Zone have recently been recorded from this succession in the De Aar region (Almond 2010b). Extensive dolerite sills and dykes of the Early Jurassic Karoo Dolerite Suite intruding the Karoo Supergroup sediments are entirely unfossiliferous, as are rare intrusive kimberlite pipe rocks of Cretaceous age.

The diverse superficial deposits within the three study areas (e.g. soils, gravels, alluvium, calcrete hardpans) are of low palaeontological sensitivity as a whole. Abundant fragments of reworked fossil wood material of Ecca provenance occur widely within subsurface and surface gravels overlying the Ecca Group outcrop area.

The proposed solar power facility developments are inferred to be of LOW overall significance in terms of palaeontological heritage resource conservation because:

- The potentially fossiliferous Karoo Supergroup rocks within the development footprints (solar panel arrays, transmission lines, roads and other infrastructure) are generally buried beneath a thick mantle of fossil-poor superficial sediments (soils, gravels, calcretes);

- The Karoo Supergroup rocks are extensively disrupted by near-surface secondary calcrete formation. In many cases they have suffered baking during dolerite magma intrusion, further compromising their fossil heritage;
- The solar energy facilities each have a small footprint while extensive, deep bedrock excavations are not envisaged for this sort of alternative energy development.

The construction of new access roads and transmission lines in this region are likewise considered to be of low significance as far as fossil heritage is concerned.

Given the low overall palaeontological sensitivity of the region around De Aar, and the widespread occurrence elsewhere in the Great Karoo of the fossils so far recorded there, the development of several small PV sites on Vetlaagte near De Aar does not pose a significant cumulative impact on local fossil heritage.

In view of the overall low significance of the proposed development on palaeontological heritage resources, it is concluded that no further palaeontological heritage studies or specialist mitigation are required for these small PV projects, pending the exposure of any substantial fossil remains (e.g. vertebrate bones and teeth, large blocks of petrified wood) during the construction phase. The ECO responsible for these developments should be alerted to the possibility of fossil remains being found on the surface or exposed by fresh excavations during construction. Should substantial fossil remains be discovered during construction, these should be safeguarded (preferably *in situ*) and the ECO should alert SAHRA so that appropriate mitigation (e.g. recording, sampling or collection) can be taken by a professional palaeontologist.

The specialist involved would require a collection permit from SAHRA. Fossil material must be curated in an approved repository (e.g. museum or university collection) and all fieldwork and reports should meet the minimum standards for palaeontological impact studies developed by SAHRA.

These recommendations should be incorporated into the EMP for the Vetlaagte PV solar power developments.

2. INTRODUCTION & BRIEF

It is proposed to develop seven photovoltaic solar power generation facilities with associated infrastructure within an area of c. 935 ha on Portions A, B, C, D, E, F and G of the Farm Vetlaagte No. 4, located approximately 7 km east of De Aar, Pixley ka Seme District Municipality, Northern Cape (Fig. 1). Six of the proposed facilities will generate 75 MW each, while the seventh (on Portion G) will generate 30 MW. The main infrastructural components of the solar facilities include:

- Numerous photovoltaic (PV) solar panels and/or concentrated photovoltaic (CPV) panels and/or dish Stirling (DS) generating units and inverters, together with their foundations;
- A new power line with a capacity of 132 kV, linking the solar power generation facilities to the existing Eskom Hydra Substation (located on the southern border of the property);
- Seven transformers and small substations which will connect to the existing power lines or the proposed new 132 kV power line;
- Cabling between the PV panels and/or CPV panels and/or dish Stirling units and small substations;
- A short 132kV line connecting 2 of the transformers to the proposed power line;
- Internal access roads;
- A maintenance building and site office.

In accordance with the National Heritage Resources Act, 1999, a palaeontological heritage assessment is required as part of a Heritage Impact Assessment for these solar power projects since their development footprints overlie potentially fossiliferous Palaeozoic rocks of the Karoo Supergroup (Ecca and Beaufort Groups). The various categories of heritage resources recognised as part of the National Estate in Section 3 of the Heritage Resources Act include, among others:

- geological sites of scientific or cultural importance
- palaeontological sites
- palaeontological objects and material, meteorites and rare geological specimens

The present combined desktop and field-based palaeontological assessment for the entire Vetlaagte development area has been commissioned by AGES (Pty) Ltd, Pretoria (Contact details: Catherine da Camara. Ages (Pty) Ltd. Plot 356 Zwavelpoort, Zwavelpoort, Pretoria 0084. Tel: +27 12 751 2160; Fax:+27 86 607 2406; Cell: +27 83 922 6971; email: cdacamara@ages-group.com).




Fig. 1. Locality map for the seven proposed solar power facilities on Portions A to G of Farm Vetlaagte No. 4, located c. 7 km east of De Aar, Northern Cape. Note also proposed short new 132 kV line linking to the existing Eskom Hydra Substation on the southern border of the property (yellow line).




Fig. 2. Google earth© satellite image of the area east of De Aar, Northern Cape, showing the terrain within the Vetlaagte study area (*NB* North is towards the left of the image). Silty alluvium of the Brakrivier drainage system (pale buff) is seen in the east. The rusty-brown patches are underlain by dolerite intrusions. Grey areas in the north are Eccca Group mudrocks, while small areas of Lower Beaufort Group sediments are mapped between dolerite intrusions in the south.

2.1. Approach used for this specialist palaeontological study

This palaeontological report provides an assessment of the observed or inferred palaeontological heritage within the entire Vetlaagte study area, with recommendations for specialist palaeontological mitigation where this is considered necessary. The report is based on (1) a review of the relevant scientific literature, including recent palaeontological assessments for development projects in the De Aar region by the author and others (notably Almond 2012a); (2) published geological maps and accompanying sheet explanations, and (3) palaeontological field assessments carried out on 25th March 2012. Because the level of natural rock exposure within the flat-lying study areas was generally very poor, the far better exposed stratotype section of the main rock units involved (Tierberg and Waterford Formations) on the farm Swartkoppies, some 47 km north-east of De Aar, was also inspected for fossil remains associated with these formations. Data on fossil heritage within the Lower Beaufort Group near De Aar that was collected during the recent field assessment of two wind energy projects near De Aar (Almond 2012b) have also been referred to here.

In preparing a palaeontological desktop study the potentially fossiliferous rock units (groups, formations *etc*) represented within the study area are determined from geological maps. The known fossil heritage within each rock unit is inventoried from the published scientific literature, previous palaeontological impact studies in the same region, and the author's field experience (Consultation with professional colleagues as well as examination of institutional fossil collections may play a role here, or later following scoping during the compilation of the final report). This data is then used to assess the palaeontological sensitivity of each rock unit to development (Provisional tabulations of palaeontological sensitivity of all formations in the Western, Eastern and Northern Cape have already been compiled by J. Almond and colleagues; *e.g.* Almond & Pether 2008). The likely impact of the proposed development on local fossil heritage is then determined on the basis of (1) the palaeontological sensitivity of the rock units concerned and (2) the nature and scale of the development itself, most notably the extent of fresh bedrock excavation envisaged. When rock units of moderate to high palaeontological sensitivity are present within the development footprint, a field-based assessment by a professional palaeontologist is usually warranted.

The focus of the field-based assessment work is *not* simply to survey the development footprint or even the development area as a whole (*e.g.* farms or other parcels of land concerned in the development). Rather, the palaeontologist seeks to assess or predict the diversity, density and distribution of fossils within and beneath the study area, as well as their heritage or scientific interest. This is primarily achieved through a careful field examination of one or more representative exposures of all the sedimentary rock units present (*N.B.* Metamorphic and igneous rocks rarely contain fossils). The best rock exposures are generally those that are easily accessible, extensive, and fresh (*i.e.* unweathered) and include a large fraction of the stratigraphic unit concerned (*e.g.* formation). These exposures may be natural or artificial and include, for example, rocky outcrops in stream or river banks, cliffs, quarries, dams, dongas, open building excavations or road and railway cuttings. Uncemented superficial deposits, such as alluvium, scree or wind-blown sands, may occasionally contain fossils and should also be included in the scoping study where they are well-represented in the study area. It is normal practice for impact palaeontologists to collect representative, well-localized (*e.g.* GPS and stratigraphic data) samples of fossil material during Phase 1 studies; this should be supported by a collection permit from the responsible heritage management authority (in this case SAHRA, Cape Town). All fossil material collected must be properly curated within an approved repository (usually a museum or university collection).

Note that while fossil localities recorded during fieldwork within the study area itself are obviously highly relevant, most fossil heritage here is embedded within rocks beneath the land surface or obscured by surface deposits (soil, alluvium *etc*) and by vegetation cover. In many cases where levels of fresh (*i.e.* unweathered) bedrock exposure are low, the hidden fossil resources have to be *inferred* from palaeontological observations made from better exposures of the same formations elsewhere in the region but outside the immediate study area. Therefore a palaeontologist might

reasonably spend far *more* time examining road cuts and borrow pits close to, but outside, the study area than within the study area itself. Field data from localities even further afield (e.g. an adjacent province) may also be adduced to build up a realistic picture of the likely fossil heritage within the study area.

On the basis of the desktop and field assessment studies, the likely impact of the proposed development on local fossil heritage and any need for specialist mitigation are then determined. Adverse palaeontological impacts normally occur during the construction rather than the operational or decommissioning phase. Mitigation by a professional palaeontologist – normally involving the recording and sampling of fossil material and associated geological information (e.g. sedimentological data) – is usually most effective during the construction phase when fresh fossiliferous bedrock has been exposed by excavations. To carry out mitigation, the palaeontologist involved will need to apply for a palaeontological collection permit from the relevant heritage management authority (i.e. SAHRA, Cape Town). It should be emphasized that, *providing appropriate mitigation is carried out*, the majority of developments involving bedrock excavation can make a *positive* contribution to our understanding of local palaeontological heritage.

2.2. Assumptions & limitations

The accuracy and reliability of palaeontological specialist studies as components of heritage impact assessments are generally limited by the following constraints:

1. Inadequate database for fossil heritage for much of the RSA, given the large size of the country and the small number of professional palaeontologists carrying out fieldwork here. Most development study areas have never been surveyed by a palaeontologist.
2. Variable accuracy of geological maps which underpin these desktop studies. For large areas of terrain these maps are largely based on aerial photographs alone, without ground-truthing. The maps generally depict only significant (“mappable”) bedrock units as well as major areas of superficial “drift” deposits (alluvium, colluvium) but for most regions give little or no idea of the level of bedrock outcrop, depth of superficial cover (soil *etc*), degree of bedrock weathering or levels of small-scale tectonic deformation, such as cleavage. All of these factors may have a major influence on the impact significance of a given development on fossil heritage and can only be reliably assessed in the field.
3. Inadequate sheet explanations for geological maps, with little or no attention paid to palaeontological issues in many cases, including poor locality information.
4. The extensive relevant palaeontological “grey literature” - in the form of unpublished university theses, impact studies and other reports (e.g. of commercial mining companies) - that is not readily available for desktop studies.
5. Absence of a comprehensive computerized database of fossil collections in major RSA institutions which can be consulted for impact studies. A Karoo fossil vertebrate database is now accessible for impact study work.

In the case of palaeontological desktop studies without supporting Phase 1 field assessments these limitations may variously lead to either:

(a) *underestimation* of the palaeontological significance of a given study area due to ignorance of significant recorded or unrecorded fossils preserved there, or

(b) *overestimation* of the palaeontological sensitivity of a study area, for example when originally rich fossil assemblages inferred from geological maps have in fact been destroyed by tectonism or weathering, or are buried beneath a thick mantle of unfossiliferous “drift” (soil, alluvium *etc*).

Since most areas of the RSA have not been studied palaeontologically, a palaeontological desktop study usually entails *inferring* the presence of buried fossil heritage within the study area from relevant fossil data collected from similar or the same rock units elsewhere, sometimes at localities far away. Where substantial exposures of bedrocks or potentially fossiliferous superficial sediments are present in the study area, the reliability of a palaeontological impact assessment may be significantly enhanced through field assessment by a professional palaeontologist.

In the case of palaeontological field studies in the De Aar region, the main limitations are:

- Very extensive intrusion of the potentially fossiliferous Karoo Supergroup bedrocks by dolerite. Weathered dolerite colluvium (scree) and sheetwash blanket most of the hill slopes in the area, *i.e.* the very regions where fossiliferous bedrocks are usually exposed;
- High levels of bedrock cover by thick alluvial and colluvial soils as well as extensive calcrete hardpans;
- Conflicting views among geologists concerning the stratigraphic subdivision and palaeoenvironmental interpretation of the Ecca – Beaufort transition rocks in the De Aar / Philipstown area.

These limitations were in part addressed through palaeontological surveying of a much larger area beyond the boundaries of the present solar energy facility study area itself (*e.g.* as part of an impact study for nearby wind energy and solar power developments). Confidence levels in the conclusions presented here are in consequence moderately high.




Fig. 3. Geological map of the region east of De Aar, Northern Cape, showing approximate boundaries of the Vetlaagte study area east of De Aar (Abstracted from 1: 250 000 geology sheet 3024 Colesberg, Council for Geoscience, Pretoria).

The following rock units are mapped within the PV study areas:

grey (Pt) = Tierberg Formation (Ecca Group) (*NB* According to the author these sediments should rather be assigned to the Waterford Formation of the Ecca Group following recent fieldwork in the study area – see discussion in text)

pale green (Pa) = Adelaide Subgroup (Lower Beaufort Group)

pink (Jd) = intrusive dykes and sills of the Karoo Dolerite Suite

white = Quaternary to Recent superficial deposits (alluvium, colluvium etc)

small black diamond symbol = Kimberlite pipe

3. GEOLOGICAL BACKGROUND

The Farm Vetlaagte No. 4 on the eastern outskirts of De Aar is a largely flat-lying area lying at c. 1200-1300m amsl on the western side of a silted up intermittently flowing tributary of the Brakrivier drainage system (Fig. 2). The large Hydra substation is situated on its southeastern border. Low rocky dolerite *koppies* emerge from the *vlaktes* in the southern portion of the area. During summer the veld is mantled by abundant grasses. In general, levels of sedimentary bedrock exposure are very poor.

The geology of the Vetlaagte study area is outlined on the 1: 250 000 geology sheet 3024 Colesberg (Le Roux 1993) (Fig. 3). The area is underlain by Middle Permian sedimentary rocks of the Karoo Supergroup that are intruded by Early Jurassic dolerites (rusty-brown areas on satellite images, Fig. 2). According to the 1: 250 000 geological map the northern portion of the study area is underlain by sediments of the **Tierberg Formation** (Ecca Group) (Pt). However, in the author's opinion, as argued below in Section 3.1, these rocks rather belong to the **Waterford Formation** (previously known in this area as the Carnarvon Formation) at the top of the Ecca Group succession. Small outcrop areas of the **Adelaide Subgroup** (Lower Beaufort Group) (Pa) are mapped in the southern portion of Vetlaagte, but this region is largely dominated by dolerite intrusions of the **Karoo Dolerite Suite** (Jd). The great majority of the Ecca and Beaufort Group outcrop area is obscured by superficial sediments of probable Pleistocene to Holocene age, as well as by abundant karroid shrub and grassy vegetation (Figs. 7, 8). These superficial sediments include silty soils and alluvium related to the broader Brakrivier drainage system, doleritic colluvium (scree, sheetwash slope deposits), and downwasted surface gravels dominated by resistant clasts of hornfels, sandstone and dolerite that are modified by sheet wash processes.

The dolerite intrusions weather out at surface as low rocky ridges and *koppies* (Fig. 4) that show up in rusty-brown colours in satellite images (Fig. 2). They have baked (thermally metamorphosed) the adjacent Karoo Supergroup mudrocks to hornfels, and any sandstones to quartzites. Dolerite colluvial rubble extends well beyond the intrusions themselves to blanket adjacent slopes and *vlaktes*.

3.1. Upper Ecca Group

The **Tierberg Formation (Pt)** (Ecca Group, Karoo Supergroup) is a recessive-weathering, mudrock-dominated succession – predominantly consisting of dark, well-laminated, carbonaceous shales with subordinate thin, fine-grained sandstones (Prinsloo 1989, Le Roux 1993, Viljoen 2005, Johnson *et al.*, 2006). The Tierberg shales are Lower to Mid Permian in age and were deposited in a range of offshore, quiet water environments below wave base. These include basin plain, distal turbidite fan and distal prodelta settings in ascending order (Viljoen 2005, Almond 2008a). Thin coarsening-upwards cycles occur towards the top of the formation with local evidence of soft-sediment deformation, ripples and common calcareous concretions. A restricted, brackish water environment is reconstructed for the Ecca Basin at this time. Close to the contact with Karoo dolerite intrusions the Tierberg mudrocks are baked to a dark grey hornfels with a reddish-brown crust or patina (Prinsloo 1989).

It should be noted here that the stratigraphic as well as palaeoenvironmental interpretation of the Ecca / Beaufort boundary rocks in the De Aar – Philipstown area is more complex and unresolved than that suggested by the brief treatment in the Britstown sheet explanation by Le Roux (1993). For mapping purposes, the base of the first prominent-weathering sandstone within the Ecca / Beaufort boundary succession has been taken as the base of the Beaufort Group in this region (*ibid.*, p. 4, following Nel 1977). The marine / lacustrine, uppermost Ecca Group rocks here, though mapped as offshore / basinal Tierberg Formation, have in fact many features in common with the shallow shelf, storm-dominated, sandstone-rich facies seen at the top of the Ecca succession in the Carnarvon area to the west. These uppermost Ecca Group rocks were previously assigned to the **Carnarvon Formation** that has since been incorporated into the **Waterford Formation** (*e.g.* Johnson *et al.* 2006). They tend to be more sandstone-rich than the overlying Beaufort Group.

The “Carnarvon Facies” is characterised by upward-coarsening, yellowish-weathering, sandstone-rich successions containing storm-generated hummocky cross-stratification and wave ripples, large ferruginous carbonate concretions (*koffieklip*), ball-and-pillow load structures, and pervasive low intensity bioturbation by low diversity trace fossil assemblages. The latter have been assigned to the shallow marine *Cruziana* Ichnofacies as well as the marginal marine *Skolithos* and *Scoyenia* Ichnofacies (e.g. Siebrits 1987, Prinsloo 1989, Rust *et al.* 1991 and references therein). Petrified wood and other plant remains (e.g. leaf compressions) are locally abundant. The inshore shelf (shoreface) Carnarvon facies rocks have a gradational contact with the underlying offshore Tierberg mudrocks and are in turn conformably overlain by continental (subaerial), fluvial sediments of the Lower Beaufort Group. For the purpose of the present fossil heritage study, the upper Ecca Group sediments within the study area are assigned to the Waterford Formation, despite their attribution to the Tierberg Formation on the published 1: 250 000 geological map (Fig. 3) and the key SACS publication by Viljoen (2005).

Only small patchy exposures of grey-green Ecca mudrocks and sandstones (the latter sometimes showing heavy mineral lamination) are exposed from beneath superficial sediments in the northern portion of the study area. Large oblate ferruginous nodules as typically seen in the Carnarvon facies of the Waterford Formation occur locally (Fig. 6). Close to dolerite intrusions, the sandstones are ferruginised and baked. A band of thin-bedded, buff-weathering tempestite sandstones with wave rippled tops and abundant trace fossils is seen at Loc. 418 (Fig. 5).

3.2. Lower Beaufort Group (Adelaide Subgroup)

The **Adelaide Subgroup (Pa)** (Lower Beaufort Group, Karoo Supergroup) was deposited by large-scale meandering river systems flowing northwards from the youthful Cape Fold Belt across the extensive floodplains of the ancient Karoo Basin (Smith 1980, Rubidge 1995, Johnson *et al.* 2006). The sediments mainly comprise fine-grained overbank mudrocks with subordinate lenticular channel sandstones. These last commonly have a basal conglomeratic lag of rolled mudflake pellets and calcrete nodules, the latter reflecting the prevailing semi-arid climates in Middle to Late Permian times. Small, often transient playa lakes were also present on the floodplain. In the Britstown – Williston - Colesberg sheet areas the Lower Beaufort succession consists largely of blocky-weathering, blue-grey and reddish floodplain mudrocks, showing occasional mudcracks. There are also subordinate siltstones, fine-grained, lenticular, current cross-bedded channel sandstones, flat-laminated crevasse-splay sandstones, and occasional playa lake deposits (Prinsloo, 1989, Viljoen 1989, Le Roux 1993). Carbonate concretions, including ferruginous *koffieklip*, as well as calcrete nodules (pedogenic limestones) and silicified gypsum rosettes (“desert roses”) are common.

The precise stratigraphic assignment of the Lower Beaufort Group sediments east of De Aar is unresolved. According to the most recent fossil biozonation map of the Beaufort Group (Van der Walt *et al.* 2010) the sediments here are assigned to the *Pristerognathus* Assemblage Zone that characterises the uppermost Abrahamskraal Formation *plus* the Poortjie Member of the Teekloof Formation west of longitude 24° East, as well as the uppermost Koonap Formation and basal Middleton Formation to the east (Rubidge 1995). De Aar is situated on the (arbitrary) cut-off line between these two stratigraphic schemes. The lowermost Beaufort Group rocks in the region to the east of town contain numerous, closely-spaced sandstones with a yellowish hue, resembling in this respect the Poortjie Member recognised in the western part of the Karoo Basin. An assignment of these rocks to the **Poortjie Member** is supported (but not yet confirmed) by the sparse fossil vertebrate remains recently recorded here (Almond 2012b).

Compared with the underlying Abrahamskraal rocks, the Teekloof Formation has a generally higher proportion of sandstones while reddish mudrocks are more abundant here. Multi-storied sandstones are common in the basal arenaceous Poortjie Member. Thin, impersistent lenses of pinkish “cherts” are probably altered volcanic ashes (Johnson & Keyser 1979, Theron 1983, Smith & Keyser 1995, Rubidge *et al.* 2010). Several economically interesting uranium ore deposits occur within the Poortjie Member in association with brown-weathering, ferruginous channel sandstones

(“koffieklip”) and transported plant material. Interesting accounts of the sedimentology and palaeontology of the Poortjie Member are given by Stear (1978) as well as by Cole and Smith (2008). The Poortjie Member has a thickness of some 200m while the entire Teekloof succession is c. 1000m thick (Cole *et al.* 1990, Cole & Voster 1999). Recent, unpublished radiometric dating of zircons from tuff layers within the Poortjie Member gives an age of 261.3 Ma (Rubidge *et al.* 2010 and pers. comm. 2010), placing this stratigraphic unit within the Gaudalupian Epoch (late Middle Permian). Previously the Poortjie Member was considered to be earliest Late Permian or Lopingian in age (*cf* Smith & Keyser 1995, Rubidge 2005).

Apart from small gully exposures of baked, ferruginised sandstones, almost all the Beaufort Group rocks in the southern part of the study area are obscured by doleritic colluvium and reddish-brown soils. Occasional low rocky outcrops of baked quartzites, spotted with secondary iron minerals, also occur (Fig. 4).

3.3. Karoo Dolerites

The **Karoo Dolerite Suite (Jd)** is an extensive network of basic igneous bodies (dykes, sills) that were intruded into sediments of the Main Karoo Basin in the Early Jurassic Period, about 183 million years ago (Duncan & Marsh 2006). These dolerites form part of the Karoo Igneous Province of Southern Africa that developed in response to crustal doming and stretching preceding the break-up of Gondwana. Hard cappings of blocky, reddish-brown to rusty-weathering dolerite are a very typical feature of the flat-topped *koppies* in the Great Karoo region. As seen from geological maps (Fig. 3), extensive dolerite intrusion of both the upper Ecca Group as well as the Lower Beaufort Group rocks is observed in the De Aar region. The country rocks adjacent to the intrusions have often been extensively baked or thermally metamorphosed. Mudrocks are altered to flinty hornfels (“lydianite” of some authors), while sandstones are metamorphosed to resistant-weathering, siliceous quartzites. The Karoo rocks within the thermal aureole of the dolerite intrusions are also often chemically altered; they tend to be silicified, more brittle and contain numerous irregular *vugs* (cavities) lined or infilled with secondary minerals.

Blocky, well-jointed outcrops of Karoo dolerite are seen on the slopes and summits of low *koppies* in the southern study area (Fig. 4). Rounded corestones of dolerite occur embedded in the surrounding reddish colluvial deposits.

3.4. Kimberlite pipes

Numerous **kimberlite pipes** of Jurassic to Cretaceous age intrude the Karoo Supergroup rocks north of Victoria West, including several examples to the east of De Aar. They are variously assigned to the Victoria West and Group II Provinces (Skinner & Truswell 2006) and do not contain diamonds. According to Le Roux (1993) the ultramafic kimberlite pipe rocks in the Colesberg sheet area are highly weathered with no obvious surface expression. They can usually be located only on the basis of characteristic mineral assemblages (garnet, phlogopite mica) found in ant heaps, termite mounds and prospecting holes. A single pipe is mapped in the north-central portion of Vetlaage but was not relocated during the present field study (diamond symbol in map Fig. 3). Kimberlite rocks are unfossiliferous, although rich Cretaceous to Paleocene fossil assemblages may be found in associated crater lake facies (not present here).

3.5. Late Caenozoic superficial deposits

Quaternary to Recent superficial deposits (“drift”) cover all but the steepest slopes of the Karoo *koppies* as well as most of the plains at their feet, including dry river courses such as the Brakrivier in the broader De Aar study region. Various types of superficial deposits of geologically young, Late Caenozoic (Miocene / Pliocene to Recent) age (< 5 Ma) occur throughout the Great Karoo region (Prinsloo 1989, Le Roux 1993, with more extensive discussion in Holmes & Marker 1995,

Cole *et al.* 2004, Partridge *et al.* 2006). They include pedocretes (e.g. calcretes), colluvial slope deposits (dolerite, sandstone and hornfels scree *etc*), sandy, gravelly and bouldery river alluvium, as well as spring and pan sediments. These colluvial and alluvial deposits may be extensively calcretised (*i.e.* cemented with soil limestone), especially in the neighbourhood of dolerite intrusions.

Thin (usually < 1m) horizons of fine to coarse, angular gravels mantle the Palaeozoic bedrocks over much of the study area. Gravel clasts mostly consist of locally-derived Beaufort Group sandstones, hornfels and quartzite as well as weathered to fresh dolerite, including large rounded dolerite corestone boulders. Unconsolidated orange-brown to brown or buff surface soils, often overlying a thin calcrete hardpan, may be of alluvial, sheet wash or even in part aeolian origin. These superficial soils are probably Holocene in age. They contain, or are locally overlain by, downwasted surface gravels, concentrated and reworked by downwasting and sheetwash processes (Fig. 7). Thick silty alluvial deposits floor the intermittently flowing tributary of the Brakrivier in the eastern part of the study area (Fig. 8). Neither Karoo Supergroup bedrock exposures nor coarse gravels were seen here.



Fig. 4. View towards SW of low dolerite koppies in the southwestern portion of Vetlaagte No. 4. Sparse outcrop of baked and ferruginised sandstones of the Lower Beaufort Group in foreground.



Fig. 5. Thin-bedded tempestite sandstones of the Waterford Formation showing wave-rippled upper surfaces. These beds contain abundant trace fossils (See Figs. 10-11) (Loc. 418).



Fig. 6. Large spheroidal concretions, probably of ferruginous carbonate, within the Waterford Formation (Loc. 417) (Hammer = 27 cm). These concretions do not generally contain macrofossils.



Fig. 7. Typical view towards the northeast showing the flat-lying terrain in the northern part of Vetlaagte. Sparse fine-grained surface gravels exposed here contain abundant small pieces of petrified wood (Fig. 9).



Fig. 8. Reddish-brown fine silts associated with a tributary of the Brakrivier in the eastern part of Vetlaagte. Neither coarse alluvial gravels nor Karoo Supergroup bedrocks are exposed here.

4. PALAEOLOGICAL HERITAGE

Fossil biotas recorded from each of the main stratigraphic units mapped in the study area are briefly reviewed in this section. Bedding dips of the Karoo Supergroup sediments in the study region are generally horizontal to very shallow. Low levels of tectonic deformation and cleavage development are expected here, favouring good fossil preservation. However, extensive dolerite intrusion has compromised fossil heritage in the Karoo Supergroup sediments due to resulting thermal metamorphism. In addition, pervasive calcretisation of many near-surface bedrocks has further compromised their original fossil heritage.

4.1. Upper Ecca Group

The fossil record of the **Tierberg Formation** has been reviewed in detail by Almond (2008a). Rare body fossil records include disarticulated microvertebrates (eg fish teeth and scales) from calcareous concretions in the Koffiefontein sheet area (Zawada 1992) and allochthonous plant remains (drifted leaves, petrified wood). The latter become more abundant in the upper, more proximal (prodeltaic) facies of the Tierberg (eg Wickens 1984). Prinsloo (1989) records numerous plant impressions and unspecified “fragmentary vertebrate fossils” within fine-grained sandstones in the Britstown sheet area. Dark carbonaceous Ecca mudrocks are likely to contain palynomorphs (e.g. pollens, spores, acritarchs).

The commonest fossils by far in the Tierberg Formation are sparse to locally concentrated assemblages of trace fossils that are often found in association with thin event beds (eg distal turbidites, prodeltaic sandstones) within more heterolithic successions. A modest range of ten or so different ichnogenera have been recorded from the Tierberg Formation (e.g. Abel 1935, Anderson 1974, 1976, Wickens 1980, 1984, 1994, 1996, Prinsloo 1989, De Beer *et al.*, 2002, Viljoen 2005, Almond 2008a). These are mainly bedding parallel, epichnial and hypichnial traces, some preserved as undertracks. Penetrative, steep to subvertical burrows are rare, perhaps because the bottom sediments immediately beneath the sediment / water interface were anoxic. Most Tierberg ichnoassemblages display a low diversity and low to moderate density of traces. Apart from simple back-filled and / or lined horizontal burrows (*Planolites*, *Palaeophycus*) they include arthropod trackways (*Umfolozia*) and associated resting impressions (*Gluckstadtella*), undulose fish swimming trails (*Undichna*) that may have been generated by bottom-feeding palaeoniscoids, horizontal epichnial furrows (so-called *Scolicia*) often attributed to gastropods (these are also common in the co-eval Collingham Formation; Viljoen 1992, 1994), arcuate, finely striated feeding excavations of an unknown arthropod (*Vadoscavichnia*), beaded traces (“*Hormosiroidea*” or “*Neonereites*”), small sinusoidal surface traces (*Cochlichnus*), small star-shaped feeding burrows (*Stelloglyphus*) and zigzag horizontal burrows (*Beloraphe*), as well as possible narrow (<1cm) *Cruziana* scratch burrows. The symmetrical, four-pronged trace *Broomichnium* (= *Quadrispinichna* of Anderson, 1974 and later authors) often occurs in groups of identical size (c. 3.5cm wide) and similar orientation on the bedding plane. This trace has frequently been misinterpreted as a web-footed tetrapod or arthropod trackway (e.g. Van Dijk *et al.* 2002 and references therein). However, Braddy and Briggs (2002) present a convincing case that this is actually a current-orientated arthropod resting trace (cubichnion), probably made by small crustaceans that lived in schools of similar-sized individuals and orientated themselves on the seabed with respect to prevailing bottom currents. Distinctive broad (3-4cm), strap-shaped, horizontal burrows with blunt ends and a more-or-less pronounced transverse ribbing occur widely within the Tierberg mudrocks. They have been described as “fucoid structures” by earlier workers (e.g. Ryan 1967) by analogy with seaweeds, and erroneously assigned to the ichnogenera *Plagiogmus* by Anderson (1974) and *Lophoctenium* by Wickens (1980, 1984). Examples up to one metre long were found in Tierberg mudrocks near Calvinia in 1803 by H. Lichtenstein, who described them as “eel fish”. These are among the first historical records of fossils in South Africa (MacRae 1999). These as yet unnamed burrows are infilled with organized arrays of faecal pellets (Werner 2006). Sandstone sole surfaces with casts of complex networks of anastomosing (branching and fusing) tubular burrows have been attributed to the ichnogenus *Paleodictyon* (Prinsloo 1989) but may more appropriately assigned to *Megagraption* (Almond 1998). These so-called graphoglyptid burrows are associated with turbidite facies from the Ordovician to Recent

times and have been interpreted as gardening burrows or *agrichnia* (Seilacher, 2007). Microbial mat textures, such as *Kinneyia*, also occur in these offshore mudrocks but, like the delicate grazing traces with which they are often associated, are generally under-recorded.

As discussed previously (Section 3.1.) it is considered likely that the Ecca Group rocks in the study area belong to the **Waterford Formation** rather than the Tierberg Formation as mapped. Rare fragments of poorly-preserved tetrapod bone are recorded in channel lags within the upper Waterford Formation in the Williston sheet area (Viljoen 1989) and the southern Great Karoo. These probably belong to aquatic temnospondyl amphibians (“labyrinthodonts”) but large fish and terrestrial therapsids might also be represented. Scattered palaeoniscoid fish scales and fish coprolites are common in the Waterford Formation, and several genera of non-marine bivalves have been described from the southern Karoo (Bender *et al.* 1991, Cooper & Kensley 1984).

Upper delta platform facies of the Waterford Formation (including the Koedoesberg Formation of earlier authors) contain abundant, low diversity trace assemblages of the *Scoyenia* ichnofacies. They are dominated by the rope-like, horizontal and oblique burrows of the ichnogenus *Scoyenia* that has been attributed to small arthropods (possibly insects) and / or earthworms. These tubular, meniscate back-filled scratch burrows characterise intermittently moist, firm substrates such as channel and pond margins on the upper delta platform (Smith & Almond 1998, Buatois & Mángano 2004, 2007). Good examples, often associated with wave-rippled surfaces, are recorded from Waterford thin-bedded sandstones and siltstones in the Roggeveld Escarpment zone by Wickens (1984, 1996) and Viljoen (1989). Offshore delta platform facies of the Waterford Formation have very impoverished, poorly-preserved ichnofaunas due to rapid sedimentation rates with abundant soft-sediment deformation and perhaps also to fluctuating salinities.

Petrified wood and other plant material of the *Glossopteris* Flora (e.g. *Glossopteris*, *Phyllothea*) is also common in the Waterford Formation (Theron 1983, Anderson & Anderson 1985, Viljoen 1989, Wickens 1984, 1996, Rubidge *et al.* 2000). Leaves and stems of arthropytes (horsetails) such as *Schizoneura* have been observed in vertical life position. Substantial fossil logs (so-called “*Dadoxylon*”) showing clearly developed seasonal growth rings are mostly permineralised with silica but partially or completely calcified material is also known (Viljoen 1989). At least two different genera of gymnospermous woods, *Prototaxoxylon* and *Australoxylon*, have been identified so far (Bamford 1999, 2004).

The storm-dominated shelf sediments of the Carnarvon-type facies of the Waterford Formation, as seen near De Aar, are typically associated with pervasive low intensity bioturbation by low diversity trace fossil assemblages. The latter have been assigned to the shallow marine *Cruziana* Ichnofacies as well as the marginal marine *Skolithos* and *Scoyenia* Ichnofacies (e.g. Rust *et al.* 1991 and references therein). Good examples of these traces are illustrated by Siebrits (1987), Prinsloo (1989) and Rust *et al.* (1991). Prominent trace fossil taxa include cm-sized horizontal to oblique burrows with striated walls (*cf. Palaeophycus striatus*) and vertical spreiten burrows of the ichnogenus *Teichichnus*. Possible arthropod feeding traces of the ichnogenus *Cruziana* are also reported here. Petrified wood (“*Dadoxylon*”) showing well-developed seasonal growth lines and other plant remains (e.g. leaf compressions) are locally abundant.

Moderately diverse non-marine ichnoassemblages of the shallow water *Cruziana* ichnofacies occur in tempestite-dominated exposures of the Waterford Formation on Vetlaagte (Loc. 418) (Figs. 5 and 10). Among the various horizontal, oblique and vertical invertebrate burrows are distinctive arthropod scratch burrows of the ichnogenus *Rusophycus* that have only rarely been reported from Mid Permian Ecca Group rocks in the Main Karoo Basin (Fig. 11).

Reworked clasts of cherty silicified wood occur widely within downwasted surface or near-surface gravels on Vetlaagte, and are of widespread occurrence and locally common in the Ecca outcrop near De Aar (Figs. 7 and 9; e.g. Locs. 414-417 and many other areas). Elsewhere in the area to the east of De Aar larger petrified wood samples also occur within subsurface gravels overlying Ecca bedrocks where these are exposed at surface (Almond 2010a). The woods typically show well-developed seasonal growth rings and preservation of original the original woody

microstructure appears to be good; this would facilitate identification and possible dating of the samples.



Fig. 9. Typical examples of silicified wood fragments reworked from the Waterford Formation and commonly found scattered among surface gravels in the northern part of Vetlaagte. The examples seen here are up to 6 cm across.



Fig. 10. Float blocks of tempestite sandstones from the Waterford at Loc. 418 showing sparse but moderately diverse trace fossil assemblages (Scale = 16 cm).



Fig. 11. Detail of sole surface of block seen in previous figure showing small, transversely-striated scratch burrow (*Rusophycus*, arrowed) attributed to a burrowing non-marine arthropod (e.g. crustacean) (Scale in cm and mm).



Fig. 12. Baked mudrock (hornfels) of the Lower Beaufort Group showing scattered impressions of transported plant stems and twigs (Scale in cm and mm) (Loc. 419).

4.2. Adelaide Subgroup

The overall palaeontological sensitivity of the Lower Beaufort Group sediments is high (Rubidge 1995, Almond & Pether 2008). These fluvial and lacustrine sediments have yielded one of the richest fossil records of land-dwelling plants and animals of Permo-Triassic age anywhere in the world. Well-preserved tetrapod fossils, from isolated skulls and post-cranial bones to fully articulated skeletons, are mainly found in overbank mudrocks, often in association with pedogenic calcretes (palaeosol horizons). Disarticulated, water-worn bones occur in the channel lag conglomerates and sandstones (Smith 1980, 1993). Playa lake deposits may be associated with disarticulated amphibian bones and a range of trace fossils (e.g. *Scoyenia*). Fossils embedded within metamorphosed sediments (quartzites, hornfels) adjacent to dolerite intrusions may be well-preserved, but are very difficult to prepare out from the matrix and therefore usually of limited scientific value.

A chronological series of mappable fossil biozones or assemblage zones (AZ), defined mainly on their characteristic tetrapod faunas, has been established for the Main Karoo Basin of South Africa (Rubidge 1995). Maps showing the distribution of the Beaufort assemblage zones within the Main Karoo Basin have been provided by Kitching (1977), Keyser and Smith (1979) and Rubidge (1995). The first two articles do not specify an assemblage zone for the study area near De Aar. As mentioned earlier (Section 3.2) the sediments here are assigned to the *Pristerognathus* Assemblage Zone according to the most recent fossil biozonation map of the Beaufort Group published by Van der Walt *et al.* (2010). The paucity of fossil data for the Lower Beaufort succession in the Colesberg sheet explanation (Le Roux 1993) also suggests that this region is palaeontologically under-explored; any new fossil finds here are consequently of palaeontological significance. This is emphasized by the absence of fossil records from the De Aar area in the recent maps of Karoo vertebrate fossil sites produced by Nicolas (2007).

Fossils of the *Pristerognathus* Assemblage Zone characterize the arenaceous Poortjie Member as well as the uppermost beds of the underlying Abrahamskraal Formation in the western Main Karoo Basin as well as the laterally equivalent beds spanning the Koonap / Middleton Formation boundary in the eastern Karoo (Smith & Keyser 1995). This important terrestrial biota is dominated by various therapsids (“mammal-like reptiles”) such as the moderate-sized therocephalian carnivore *Pristerognathus* as well as several gorgonopsian predators / scavengers and herbivorous dicynodonts (Fig. 13). The commonest genus by far is the small burrowing dicynodont *Diictodon* (Keyser and Smith 1977-78, Smith & Keyser 1995b, MacRae 1999, Cole *et al.*, 2004, Rubidge 2005, Almond 2010, Nicolas 2007, Nicolas & Rubidge 2010). There are also large, rhino-sized herbivorous pareiasaur reptiles (*Bradysaurus* spp.), crocodile-like temnospondyl amphibians (*Rhinesuchus*), palaeoniscoid fish, vascular plant fossils of the *Glossopteris* Flora (fossil wood, leaves *etc*) and various trace fossils, including invertebrate and therapsid burrows as well as tetrapod trackways. The comparatively low number of specimens and major taxa represented in fossil collections from this biozone has been highlighted by Nicolas (2007). The fossil biota of the *Pristerognathus* AZ is of special interest because it possibly represents an impoverished post-extinction recovery fauna following a late Mid Permian extinction event that preceded the well-known end-Guadalupian biotic crisis (*cf* Benton 2003, Retallack *et al.*, 2006, Lucas 2009). During a recent palaeontological impact study several specimens of small dicynodonts – probably *Diictodon* and *Pristerodon* – as well as a rare specimen of the small tortoise-like parareptile *Eunotosaurus* but no dinocephalians were recorded to the east of De Aar (Almond 2012b). These new finds support correlation of the lowermost Beaufort Group beds here with the *Pristerognathus* Assemblage Zone.

Most fossils in the *Pristerognathus* Assemblage Zone are found in the softer-weathering mudrock facies (floodplain sediments) that are usually only exposed on steeper hill slopes and in stream gullies. Fossils here are often associated with pedogenic limestone nodules or calcretes (Smith 1993, Smith & Keyser 1995). The mudrocks lie between the more resistant-weathering channel sandstones, which in the Poortjie Member display a distinctive “golden yellow” tint. Fossil skeletal remains also occur in the lenticular channel sandstones, especially in intraformational lag conglomerates towards the base, but are usually very fragmentary and water-worn (“rolled bone”).




Fig. 13. Skulls of typical therapsids from the *Pristerognathus* Assemblage Zone: A. the dog-sized carnivorous therocephalian *Pristerognathus* and B. the small herbivorous dicynodont *Diictodon* (From Smith & Keyser 1995).

No vertebrate fossil remains were recorded from the very poor exposures of Lower Beaufort Group sediments on Vetlaagte. Local concentrations of unidentifiable reworked woody plant remains embedded within baked mudrocks (hornfels) were seen at Loc. 419 (Fig. 12).

4.3. Karoo Dolerite Suite

The dolerite outcrops in the De Aar PV study areas are in themselves of no palaeontological significance. These are high temperature igneous rocks emplaced at depth within the Earth's crust so they do not contain fossils. However, as a consequence of their proximity to large dolerite intrusions in the Great Escarpment zone, some of the Ecca and Beaufort Group sediments in the study area will have been thermally metamorphosed or "baked" (*ie.* recrystallised, impregnated with secondary minerals). Embedded fossil material of phosphatic composition, such as bones and teeth, is frequently altered by baking – bones may become blackened, for example - and can be very difficult to extract from the hard matrix by mechanical preparation (Smith & Keyser 1995). Thermal metamorphism by dolerite intrusions therefore tends to reduce the palaeontological heritage potential of Beaufort Group sediments. In some cases (*e.g.* fossil moulds of reptiles and palaeoniscoid fish) baking may enhance the quality of preservation of Ecca fossils while other fossil groups (*e.g.* carbonaceous remains of plants, organic-walled palynomorphs) are more likely to be compromised.

4.4. Quaternary to Recent superficial deposits

The central Karoo drift deposits have been comparatively neglected in palaeontological terms. However, they may occasionally contain important fossil biotas, notably the bones, teeth and horn cores of mammals as well as remains of reptiles like tortoises. Good examples are the Pleistocene mammal faunas at Florisbad, Cornelia and Erfkroon in the Free State and elsewhere (Wells & Cooke 1942, Cooke 1974, Skead 1980, Klein 1984, Brink, J.S. 1987, Bousman *et al.* 1988, Bender & Brink 1992, Brink *et al.* 1995, MacRae 1999, Meadows & Watkeys 1999, Churchill *et al.* 2000 Partridge & Scott 2000). Other late Caenozoic fossil biotas from these superficial deposits include non-marine molluscs (bivalves, gastropods), ostrich egg shells, tortoise remains, trace fossils (*e.g.* calcretised termitaria, coprolites), and plant material such as peats or palynomorphs (pollens) in organic-rich alluvial horizons (Scott 2000) and diatoms in pan sediments. In Quaternary deposits, fossil remains may be associated with human artefacts such as stone tools and are also of archaeological interest (*e.g.* Smith 1999 and refs. therein). Ancient solution hollows within extensive calcrete hardpans such as seen here may have acted as animal traps in the past. As with coastal and interior limestones, they might occasionally contain mammalian bones and teeth (perhaps associated with *hyaena dens*) or invertebrate remains such as snail shells.

The only fossil remains recorded within the Late Caenozoic superficial deposits on Vetlaagte were reworked fragments of Ecca Group silicified wood scattered among surface gravels in the northern part of the farm (See Section 4.1 and Figs. 7 and 9).

5. CONCLUSIONS & RECOMMENDATIONS

The seven small PV solar energy facilities proposed on the Farm Vetlaagte No. 4 near De Aar are located in an area that is underlain by potentially fossiliferous sedimentary rocks of Palaeozoic and younger, probably Quaternary age. The construction phase of these alternative energy developments will entail numerous, but mostly shallow, excavations into the superficial sediment cover and in some areas into the underlying bedrock as well. These include, for example, excavations for the solar panel foundations, underground cables, new electricity transmission lines, as well as new gravel access roads and the maintenance building. In addition, substantial areas of bedrock will be sealed-in or sterilized by infrastructure such as solar panel arrays and new gravel roads. All these developments may adversely affect fossil heritage within the development footprint by destroying, disturbing or permanently sealing-in fossils that are then no longer available for scientific research or other public good. The operational and decommissioning phases of the solar energy facilities will not involve further significant adverse or other impacts on palaeontological heritage.

The fossil record and inferred palaeontological sensitivity of fossil heritage within each of the main rock units represented in the study area near De Aar is summarized in Table 1 below (See also Almond & Pether 2008). The Ecca and Beaufort Group sediments here generally have a moderate to high palaeontological sensitivity. However, the potentially fossiliferous sediments of the Karoo Supergroup that underlie the study area are almost entirely mantled in a thick layer of superficial deposits of probable Pleistocene to Recent age. They include various soils, gravels and – at least in some areas - a well-developed calcrete hardpan. The upper Ecca Group bedrocks in the De Aar area contain locally abundant fossil wood (of palaeontological interest for dating and palaeoenvironmental studies) as well as low diversity trace fossil assemblages typical of the Waterford Formation, rather than the Tierberg Formation as mapped.

No vertebrate fossils were observed within the Lower Beaufort Group rocks that are very poorly exposed in the southern portion of the study area. However, various trace fossils, silicified woods and rare vertebrate remains (therapsids, parareptiles) of the Middle Permian *Pristerognathus* Assemblage Zone have recently been recorded elsewhere from these successions in the De Aar area (Almond 2010b). Jurassic dolerite sill and dyke rocks here are entirely unfossiliferous (igneous intrusions), as are rare kimberlite pipes of Cretaceous age.

The diverse superficial deposits in the study region are of low palaeontological sensitivity as a whole. Calcretized rhizoliths (root casts) and possible invertebrate burrows of probable Quaternary age were observed during recent field studies in the De Aar region (Almond 2010a). Good examples of silicified *Ecce* fossil woods are recorded from gravels directly overlying *Ecce* bedrock as well as reworked into surface gravels in the present study area. Fossil wood material probably occurs widely in similar settings throughout the *Ecce* Group outcrop area, albeit often buried beneath superficial deposits

Fossils exposed at the surface or underground may be damaged, disturbed or sealed-in during the construction phase of the proposed solar energy facilities on Vetlaagte near De Aar. However these developments are all inferred to be of LOW significance in terms of palaeontological heritage resource conservation because:

- The potentially fossiliferous Karoo Supergroup rocks within the development footprints (solar panel arrays, transmission lines, roads and other infrastructure) are generally buried beneath a mantle of fossil-poor superficial sediments (soils, gravels, calcretes);
- The Karoo Supergroup rocks are often extensively disrupted by near-surface secondary calcrete formation. Baking by dolerite intrusion has often further compromised their original fossil heritage;
- The solar energy facilities each have a small footprint while extensive, deep bedrock excavations are not envisaged for this sort of alternative energy development.

The construction of new access roads and transmission lines in this region are likewise considered to be of low significance as far as fossil heritage is concerned.

Given the low overall palaeontological sensitivity of the region around De Aar, and the widespread occurrence elsewhere in the Great Karoo of the fossils so far recorded there, the seven small PV solar power developments proposed here do not pose a significant cumulative impact on local fossil heritage.

In view of the overall low significance of the proposed development on palaeontological heritage resources, it is concluded that no further palaeontological heritage studies or specialist mitigation are required for these PV projects, pending the exposure of any substantial fossil remains (e.g. vertebrate bones and teeth, large blocks of petrified wood) during the construction phase. The ECO responsible for these developments should be alerted to the possibility of fossil remains being found on the surface or exposed by fresh excavations during construction. Should substantial fossil remains be discovered during construction, these should be safeguarded (preferably *in situ*) and the ECO should alert SAHRA so that appropriate mitigation (e.g. recording, sampling or collection) can be taken by a professional palaeontologist.

The specialist involved would require a collection permit from SAHRA. Fossil material must be curated in an approved repository (e.g. museum or university collection) and all fieldwork and reports should meet the minimum standards for palaeontological impact studies developed by SAHRA.

These recommendations should be incorporated into the EMP for the Vetlaagte PV solar power developments.

7. ACKNOWLEDGEMENTS

Ms Catherine da Camara of Ages (Pty) Ltd, Pretoria, is thanked for commissioning this study, for facilitating field work and for providing the necessary background information. I am very grateful to Ms Madelon Tusenius for her kind assistance and companionship in the field.

Table 1: Palaeontological record and sensitivity of rocks units represented in the broader De Aar study region

TABLE 1: FOSSIL HERITAGE IN THE DE AAR AREA				
GEOLOGICAL UNIT	ROCK TYPES & AGE	FOSSIL HERITAGE	PALAEONTOLOGICAL SENSITIVITY	RECOMMENDED MITIGATION
Superficial deposits (“drift”)	Alluvium, colluvium (scree), pan sediments, surface gravels, calcrete hardpans etc NEOGENE / QUATERNARY TO RECENT	Sparse remains of mammals (bones, teeth), reptiles, ostrich egg shells, molluscs shells, trace fossils (calcretized termitaria, rhizoliths), plant remains, palynomorphs, diatoms; reworked Karoo-age silicified wood clasts and stone artefacts in surface or subsurface gravels	LOW	Any substantial fossil finds to be reported by ECO to SAHRA
Kimberlite pipes (diamond symbol)	Ultramafic kimberlite CRETACEOUS	None within pipe itself	ZERO	None
Karoo Dolerite Suite (Jd)	Intrusive dolerite sills & dykes EARLY JURASSIC	NONE	ZERO	None
Adelaide Subgroup (Pa) BEAUFORT GROUP	Floodplain mudrocks with lenticular channel sandstones, tabular crevasse splay sandstones, minor playa lake sediments LATE MIDDLE PERMIAN	Important but low diversity terrestrial vertebrate fauna (esp. therapsids) of <i>Pristerognathus</i> Assemblage Zone, petrified wood, plant remains (incl. fossil wood, leaf & stem impressions), freshwater molluscs, trace fossils (trackways, burrows, coprolites)	HIGH	Any substantial fossil finds to be reported by ECO to SAHRA
Tierberg and Waterford Formations (Pt) ECCA GROUP	Dark basinal, prodelta and submarine fan mudrocks with minor sandstones (Tierberg Fm) OR Storm-influenced coastal sandstones and mudrocks (Carnarvon facies of Waterford Fm) EARLY TO MIDDLE PERMIAN	Locally abundant trace fossils, petrified wood, plant debris, microvertebrates	MEDIUM	Any substantial fossil finds to be reported by ECO to SAHRA

8. REFERENCES

- ABEL, O. 1935. Vorzeitliche Lebensspuren. xv+ 644 pp. Gustav Fischer, Jena.
- ALMOND, J.E. 1998. Non-marine trace fossils from the western outcrop area of the Permian Ecca Group, southern Africa. Tercera Reunión Argentina de Icnología, Mar del Plata, 1998, Abstracts p. 3.
- ALMOND, J.E. 2008a. Fossil record of the Loeriesfontein sheet area (1: 250 000 geological sheet 3018). Unpublished report for the Council for Geoscience, Pretoria, 32 pp. Natura Viva cc, Cape Town.
- ALMOND, J.E. 2008b. Palaeozoic fossil record of the Clanwilliam sheet area (1: 250 000 geological sheet 3218). Unpublished report for the Council for Geoscience, Pretoria, 49 pp. Natura Viva cc, Cape Town.
- ALMOND, J.E. 2009. Contributions to the palaeontology and stratigraphy of the Alexander Bay sheet area (1: 250 000 geological sheet 2816), 117 pp. Unpublished technical report prepared for the Council for Geoscience by Natura Viva cc, Cape Town.
- ALMOND, J.E. 2010. Eskom Gamma-Omega 765kV transmission line: Phase 2 palaeontological impact assessment. Sector 1, Tanqua Karoo to Omega Substation (Western and Northern Cape Provinces), 95 pp + appendix. Natura Viva cc, Cape Town.
- ALMOND, J.E. 2012a. Proposed Mulilo Renewable Energy PV2, PV3 and PV4 photovoltaic energy facilities on Farms Paarde Valley, Badenhorst Dam and Annex Du Plessis Dam near De Aar, Northern Cape Province. Palaeontological specialist study: combined desktop and field-based assessments, 45 pp. Natura Viva cc, Cape Town.
- ALMOND J.E. 2012b. Two wind energy facilities on the Eastern Plateau near De Aar, Northern Cape Province proposed by Mulilo Renewable Energy (Pty) Ltd. Palaeontological specialist study: combined desktop and field-based assessments, 55 pp. Natura Viva cc, Cape Town.
- ALMOND, J.E. & PETHER, J. 2008. Palaeontological heritage of the Northern Cape. Interim SAHRA technical report, 124 pp. Natura Viva cc., Cape Town.
- ANDERSON, A.M. 1974. Arthropod trackways and other trace fossils from the Early Permian lower Karoo Beds of South Africa. Unpublished PhD thesis, University of Witwatersrand, Johannesburg, 172 pp.
- ANDERSON, A.M. 1975. Turbidites and arthropod trackways in the Dwyka glacial deposits (Early Permian) of southern Africa. Transactions of the Geological Society of South Africa 78: 265-273.
- ANDERSON, A.M. 1976. Fish trails from the Early Permian of South Africa. Palaeontology 19: 397-409, pl. 54.
- ANDERSON, A.M. 1981. The *Umfolozia* arthropod trackways in the Permian Dwyka and Ecca Groups of South Africa. Journal of Paleontology 55: 84-108, pls. 1-4.
- ANDERSON, A.M. & MCLACHLAN, I.R. 1976. The plant record in the Dwyka and Ecca Series (Permian) of the south-western half of the Great Karoo Basin, South Africa. Palaeontologia africana 19: 31-42.
- ANDERSON, J.M. & ANDERSON, H.M. 1985. Palaeoflora of southern Africa. Prodrum of South African megafloras, Devonian to Lower Cretaceous, 423 pp. Botanical Research Institute, Pretoria & Balkema, Rotterdam.

- BENDER, P.A. 2004. Late Permian actinopterygian (palaeoniscid) fishes from the Beaufort Group, South Africa: biostratigraphic and biogeographic implications. Bulletin 135, 84pp. Council for Geoscience, Pretoria.
- BENDER, P.A., RUBIDGE, B.S., GARDINER, B.S., LOOCK, J.C. & BREMNER, A.T. 1991. The stratigraphic range of the palaeoniscoid fish *Namaichthys digitata* in rocks of the Karoo sequence and its palaeoenvironmental significance. South African Journal of Science 87: 468-469.
- BENDER, P.A. & BRINK, J.S. 1992. A preliminary report on new large mammal fossil finds from the Cornelia-Uitzoek site. South African Journal of Science 88: 512-515.
- BENTON, M.J. 2003. When life nearly died. The greatest mass extinction of them all, 336 pp. Thames & Hudson, London.
- BOUSMAN, C.B. *et al.* 1988. Palaeoenvironmental implications of Late Pleistocene and Holocene valley fills in Blydefontein Basin, Noupoort, C.P., South Africa. Palaeoecology of Africa 19: 43-67.
- BAMFORD, M. 1999. Permo-Triassic fossil woods from the South African Karoo Basin. Palaeontologia africana 35, 25-40.
- BAMFORD, M.K. 2004. Diversity of woody vegetation of Gondwanan southern Africa. Gondwana Research 7, 153-164.
- BRADY, S.J. & BRIGGS, D.E.G. 2002. New Lower Permian nonmarine arthropod trace fossils from New Mexico and South Africa. Journal of Paleontology 76: 546-557.
- BRINK, J.S. 1987. The archaeozoology of Florisbad, Orange Free State. Memoirs van die Nasionale Museum 24, 151 pp.
- BRINK, J.S. *et al.* 1995. A new find of *Megalotragus priscus* (Alcephalini, Bovidae) from the Central Karoo, South Africa. Palaeontologia africana 32: 17-22.
- BUATOIS, L. & MANGANO, M.G. 2004. Animal-substrate interactions in freshwater environments: applications of ichnology in facies and sequence stratigraphic analysis of fluvio-lacustrine successions. In: McIlroy, D. (Ed.) The application of ichnology to palaeoenvironmental and stratigraphic analysis. Geological Society, London, Special Publications 228, pp 311-333.
- BUATOIS, L.A. & MANGANO, M.G. 2007. Invertebrate ichnology of continental freshwater environments. In: Miller, W. III (Ed.) Trace fossils: concepts, problems, prospects, pp. 285-323. Elsevier, Amsterdam.
- CHURCHILL, S.E. *et al.* 2000. Erfkroon: a new Florisian fossil locality from fluvial contexts in the western Free State, South Africa. South African Journal of Science 96: 161-163.
- COLE, D.I., SMITH, R.M.H. & WICKENS, H. DE V. 1990. Basin-plain to fluvio-lacustrine deposits in the Permian Ecca and Lower Beaufort Groups of the Karoo Sequence. Guidebook Geocongress '90, Geological Society of South Africa, PO2, 1-83.
- COLE, D.I. & VORSTER, C.J. 1999. The metallogeny of the Sutherland area, 41 pp. Council for Geoscience, Pretoria.
- COLE, D.I., NEVELING, J., HATTINGH, J., CHEVALLIER, L.P., REDDERING, J.S.V. & BENDER, P.A. 2004. The geology of the Middelburg area. Explanation to 1: 250 000 geology Sheet 3124 Middelburg, 44 pp. Council for Geoscience, Pretoria.
- COLE, D. & SMITH, R. 2008. Fluvial architecture of the Late Permian Beaufort Group deposits, S.W. Karoo Basin: point bars, crevasse splays, palaeosols, vertebrate fossils and uranium. Field Excursion FT02 guidebook, AAPG International Conference, Cape Town October 2008, 110 pp.

COOKE, H.B.S. 1974. The fossil mammals of Cornelia, O.F.S., South Africa. In: Butzer, K.W., Clark, J.D. & Cooke, H.B.S. (Eds.) The geology, archaeology and fossil mammals of the Cornelia Beds, O.F.S. *Memoirs of the National Museum, Bloemfontein* 9: 63-84.

COOPER, M.R. & KENSLEY, B. 1984. Endemic South American Permian bivalve molluscs from the Ecca of South Africa. *Journal of Paleontology* 58: 1360-1363.

DE BEER, C.H., GRESSE, P.G., THERON, J.N. & ALMOND, J.E. 2002. The geology of the Calvinia area. Explanation to 1: 250 000 geology Sheet 3118 Calvinia. 92 pp. Council for Geoscience, Pretoria.

COLE, D.I., NEVELING, J., HATTINGH, J., CHEVALLIER, L.P., REDDERING, J.S.V. & BENDER, P.A. 2004. The geology of the Middelburg area. Explanation to 1: 250 000 geological sheet 3124 Middelburg, 43 pp. Council for Geoscience, Pretoria.

DUNCAN, A.R. & MARSH, J.S. 2006. The Karoo Igneous Province. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) *The geology of South Africa*, pp. 501-520. Geological Society of South Africa, Marshalltown.

HOLMES, P.J. & MARKER, M.E. 1995. Evidence for environmental change from Holocene valley fills from three central Karoo upland sites. *South African Journal of Science* 91: 617-620.

JOHNSON, M.R. & KEYSER, A.W. 1979. Die geologie van die gebied Beaufort-Wes. Explanation of geological Sheet 3222, 14 pp. Council for Geoscience, Pretoria.

JOHNSON, M.R., VAN VUUREN, C.J., VISSER, J.N.J., COLE, D.I., De V. WICKENS, H., CHRISTIE, A.D.M., ROBERTS, D.L. & BRANDL, G. 2006. Sedimentary rocks of the Karoo Supergroup. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) *The geology of South Africa*, pp. 461-499. Geological Society of South Africa, Marshalltown.

KEYSER, A.W. & SMITH, R.M.H. 1977-78. Vertebrate biozonation of the Beaufort Group with special reference to the Western Karoo Basin. *Annals of the Geological Survey of South Africa* 12: 1-36.

KITCHING, J.W. 1977. The distribution of the Karoo vertebrate fauna, with special reference to certain genera and the bearing of this distribution on the zoning of the Beaufort beds. *Memoirs of the Bernard Price Institute for Palaeontological Research, University of the Witwatersrand, No. 1*, 133 pp (incl. 15 pls).

KLEIN, R.G. 1984. The large mammals of southern Africa: Late Pliocene to Recent. In: Klein, R.G. (Ed.) *Southern African prehistory and paleoenvironments*, pp 107-146. Balkema, Rotterdam.

LE ROUX, F.G. 1993. Die geologie van die gebied Colesberg. Explanation to 1: 250 000 geology Sheet 3024, 12 pp. Council for Geoscience, Pretoria.

LE ROUX, F.G. & KEYSER, A.W. 1988. Die geologie van die gebied Victoria-Wes. Explanation to 1: 250 000 geology Sheet 3122, 31 pp. Council for Geoscience, Pretoria.

LUCAS, D.G. 2009. Global Middle Permian reptile mass extinction: the dinocephalian extinction event. *Geological Society of America Abstracts with Programs* 41, No. 7, p. 360.

MACRAE, C. 1999. Life etched in stone. *Fossils of South Africa*, 305 pp. The Geological Society of South Africa, Johannesburg.

MEADOWS, M.E. & WATKEYS, M.K. 1999. Palaeoenvironments. In: Dean, W.R.J. & Milton, S.J. (Eds.) *The karoo. Ecological patterns and processes*, pp. 27-41. Cambridge University Press, Cambridge.

NEL, L. 1977. Die geologie van die gebied suid van Hopetown. Unpublished PhD thesis, University of the Free State, 171 pp.

NICOLAS, M.V. 2007. Tetrapod diversity through the Permo-Triassic Beaufort Group (Karoo Supergroup) of South Africa. Unpublished PhD thesis, University of Witwatersrand, Johannesburg.

NICOLAS, M. & RUBIDGE, B.S. 2010. Changes in Permo-Triassic terrestrial tetrapod ecological representation in the Beaufort Group (Karoo Supergroup) of South Africa. *Lethaia* 43, 45-59.

PARTRIDGE, T.C. & SCOTT, L. 2000. Lakes and pans. In: Partridge, T.C. & Maud, R.R. (Eds.) *The Cenozoic of southern Africa*, pp.145-161. Oxford University Press, Oxford.

PARTRIDGE, T.C., BOTHA, G.A. & HADDON, I.G. 2006. Cenozoic deposits of the interior. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) *The geology of South Africa*, pp. 585-604. Geological Society of South Africa, Marshalltown.

PRINSLOO, M.C. 1989. Die geologie van die gebied Britstown. Explanation to 1: 250 000 geology Sheet 3022 Britstown, 40 pp. Council for Geoscience, Pretoria.

RETALLACK, G.J., METZGER, C.A., GREAVES, T., HOPE JAHREN, A., SMITH, R.M.H. & SHELDON, N.D. 2006. Middle – Late Permian mass extinction on land. *GSA Bulletin* 118, 1398-1411.

RUBIDGE, B.S. (Ed.) 1995. *Biostratigraphy of the Beaufort Group (Karoo Supergroup)*. South African Committee for Biostratigraphy, Biostratigraphic Series No. 1., 46 pp. Council for Geoscience, Pretoria.

RUBIDGE, B.S. 2005. Re-uniting lost continents – fossil reptiles from the ancient Karoo and their wanderlust. 27th Du Toit Memorial Lecture. *South African Journal of Geology* 108, 135-172.

RUBIDGE, B.S., HANCOX, P.J. & CATUNEANU, O. 2000. Sequence analysis of the Ecca-Beaufort contact in the southern Karoo of South Africa. *South African Journal of Geology* 103, 81-96.

RUBIDGE, B.S., ERWIN, D.H., RAMEZANI, J., BOWRING, S.A. & DE KLERK, W.J. 2010. The first radiometric dates for the Beaufort Group, Karoo Supergroup of South Africa. *Proceedings of the 16th conference of the Palaeontological Society of Southern Africa*, Howick, August 5-8, 2010, pp. 82-83.

RUST, I.C., SHONE, R.W. & SIEBRITS, L.B. 1991. Carnarvon Formasie: golf-oorheesde sedimentasie in 'n vlak Karoo-see. *South African Journal of Science* 87, 198-202.

RYAN, P.J. 1967. Stratigraphic and palaeocurrent analysis of the Ecca Series and lowermost Beaufort Beds in the Karoo Basin of South Africa. Unpublished PhD thesis, University of the Witwatersrand, Johannesburg, 210 pp.

SCOTT, L. 2000. Pollen. In: Partridge, T.C. & Maud, R.R. (Eds.) *The Cenozoic of southern Africa*, pp.339-35. Oxford University Press, Oxford.

SEILACHER, A. 2007. *Trace fossil analysis*, xiii + 226pp. Springer Verlag, Berlin.

SIEBRITS, L.B. 1987. Die sedimentologie van die Formasie Carnarvon in die omgewing van Carnarvon. Unpublished MSc thesis, University of Port Elizabeth, v + 92 pp.

- SIEBRITS, L.B. 1989. Die geologie van die gebied Sakrivier. Explanation of 1: 250 000 geology sheet 3020, 19 pp. Council for Geoscience, Pretoria.
- SKEAD, C.J. 1980. Historical mammal incidence in the Cape Province. Volume 1: The Western and Northern Cape. 903pp. Department of Nature and Environmental Conservation, Cape Town.
- SKINNER, E.M.W. & TRUSWELL, J.F. 2006. Kimberlites. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) The geology of South Africa, pp. 651-659. Geological Society of South Africa, Marshalltown.
- SMITH, A.M. & ZAWADA, P.K. 1988. The Ecca-Beaufort transition zone near Philipstown, Cape Province: a marine shelf sequence. South African Journal of Geology 91, 75-82.
- SMITH, A.M. & ZAWADA, P.K. 1989. Permian storm current-produced offshore bars from an ancient shelf sequence: northwestern Karoo basin, Republic of South Africa. Journal of African Earth Sciences 9, 363-370.
- SMITH, A.B. 1999. Hunters and herders in the Karoo landscape. Chapter 15 in Dean, W.R.J. & Milton, S.J. (Eds.) The Karoo; ecological patterns and processes, pp. 243-256. Cambridge University Press, Cambridge.
- SMITH, R.M.H. 1980. The lithology, sedimentology and taphonomy of flood-plain deposits of the Lower Beaufort (Adelaide Subgroup) strata near Beaufort West. Transactions of the Geological Society of South Africa 83, 399-413.
- SMITH, R.M.H. 1993. Sedimentology and ichnology of floodplain paleosurfaces in the Beaufort Group (Late Permian), Karoo Sequence, South Africa. Palaios 8, 339-357.
- SMITH, R.M.H. & KEYSER, A.W. 1995. Biostratigraphy of the *Tapinocephalus* Assemblage Zone. Pp. 8-12 in Rubidge, B.S. (ed.) Biostratigraphy of the Beaufort Group (Karoo Supergroup). South African Committee for Stratigraphy, Biostratigraphic Series No. 1. Council for Geoscience, Pretoria.
- SMITH, R.M.H. & ALMOND, J.E. 1998. Late Permian continental trace assemblages from the Lower Beaufort Group (Karoo Supergroup), South Africa. Abstracts, Tercera Reunión Argentina de Icnología, Mar del Plata, 1998, p. 29.
- STEAR, W.M. 1978. Sedimentary structures related to fluctuating hydrodynamic conditions in flood plain deposits of the Beaufort Group near Beaufort West, Cape. Transactions of the Geological Society of South Africa 81, 393-399.
- THERON, J.N. 1983. Die geologie van die gebied Sutherland. Explanation of 1: 250 000 geological Sheet 3220, 29 pp. Council for Geoscience, Pretoria.
- VAN DER WALT M., DAY M., RUBIDGE B., COOPER A.K. & NETTERBERG I. 2010. A new GIS-based biozone map of the Beaufort Group (Karoo Supergroup), South Africa. Palaeontologia Africana, 45, 5pp.
- VAN DIJK, D.E., CHANNING, A. & VAN DEN HEEVER, J.A. 2002. Permian trace fossils attributed to tetrapods (Tierberg Formation, Karoo Basin, South Africa). Palaeontologia africana 38: 49-56.
- VILJOEN, J.H.A. 1989. Die geologie van die gebied Williston. Explanation to geology sheet 3120 Williston, 30 pp. Council for Geoscience, Pretoria.

- VILJOEN, J.H.A. 1992. Lithostratigraphy of the Collingham Formation (Ecca Group), including the Zoute Kloof, Buffels River and Wilgehout River Members and the Matjiesfontein Chert Bed. South African Committee for Stratigraphy, Lithostratigraphic Series No. 22, 10 pp.
- VILJOEN, J.H.A. 1994. Sedimentology of the Collingham Formation, Karoo Supergroup. South African Journal of Geology 97: 167-183.
- VILJOEN, J.H.A. 2005. Tierberg Formation. SA Committee for Stratigraphy, Catalogue of South African Lithostratigraphic Units 8: 37-40.
- VISSER, J.N.J. & LOOCK, J.C. 1974. The nature of the Ecca-Beaufort transition in the western and central Orange Free State. Transactions of the Geological Society of South Africa 77, 371-372.
- WELLS, L.H. & COOKE, H.B.S. 1942. The associated fauna and culture of Vlakkraal thermal springs, O.F.S.; III, the faunal remains. Transactions of the Royal Society of South Africa 29: 214-232.
- WERNER, M. 2006. The stratigraphy, sedimentology and age of the Late Palaeozoic Mesosaurus Inland Sea, SW-Gondwana: new implications from studies on sediments and altered pyroclastic layers of the Dwyka and Ecca Group (lower Karoo Supergroup) in southern Namibia. Dr rer. nat. thesis, University of Würzburg, 428 pp, 167 figs, 1 table.
- WICKENS, H. DE V. 1980. Verslag oor kartering in die Calvinia gebied. Unpublished report, Council for Geoscience, Pretoria, 19 pp.
- WICKENS, H. DE V. 1984. Die stratigraphie en sedimentologie van die Group Ecca wes van Sutherland. Unpublished MSc thesis, University of Port Elizabeth, viii + 86 pp.
- WICKENS, H. DE V. 1992. Submarine fans of the Permian Ecca Group in the SW Karoo Basin, their origin and reflection on the tectonic evolution of the basin and its source areas. In: De Wit, M.J. & Ransome, I.G.D. (Eds.) Inversion tectonics of the Cape Fold Belt, Karoo and Cretaceous Basins of southern Africa, pp. 117-126. Balkema, Rotterdam.
- WICKENS, H. DE V. 1994. Submarine fans of the Ecca Group. Unpublished PhD thesis, University of Port Elizabeth. 350 pp.
- WICKENS, H. DE V. 1996. Die stratigraphie en sedimentologie van die Ecca Groep wes van Sutherland. Council for Geosciences, Pretoria Bulletin 107, 49pp.
- ZAWADA, P.K. 1992. The geology of the Koffiefontein area. Explanation of 1: 250 000 geology sheet 2924, 30 pp. Council for Geoscience, Pretoria.

APPENDIX: VETLAAGTE, DE AAR - GPS LOCALITY DATA

All GPS readings were taken in the field using a hand-held Garmin GPSmap 60CSx instrument. The datum used is WGS 84.

LOC. NO.	GPS DATA	COMMENTS
414	S30 39.216 E24 05.479 1198 m	Small fragments of petrified wood in surface gravels
415	S30 39.873 E24 05.813 1254 m	As above
416	S30 40.141 E24 05.961 1256 m	As above. Calcrete hard pan beneath surface soils.
417	S30 40.422 E24 05.960 1260 m	Large ferruginous carbonate nodules within Ecca sandstones, plus fossil wood fragments
418	S30 40.750 E24 05.479 1271 m	Thin tempestite sandstones of Waterford Formation with moderately diverse trace fossil assemblages
419	S30 40.985 E24 05.332 1283 m	Baked Beaufort Group mudrocks with transported fossil plant remains

QUALIFICATIONS & EXPERIENCE OF THE AUTHOR

Dr John Almond has an Honours Degree in Natural Sciences (Zoology) as well as a PhD in Palaeontology from the University of Cambridge, UK. He has been awarded post-doctoral research fellowships at Cambridge University and in Germany, and has carried out palaeontological research in Europe, North America, the Middle East as well as North and South Africa. For eight years he was a scientific officer (palaeontologist) for the Geological Survey / Council for Geoscience in the RSA. His current palaeontological research focuses on fossil record of the Precambrian - Cambrian boundary and the Cape Supergroup of South Africa. He has recently written palaeontological reviews for several 1: 250 000 geological maps published by the Council for Geoscience and has contributed educational material on fossils and evolution for new school textbooks in the RSA.

Since 2002 Dr Almond has also carried out palaeontological impact assessments for developments and conservation areas in the Western, Eastern and Northern Cape under the aegis of his Cape Town-based company *Natura Viva* cc. He is a long-standing member of the Archaeology, Palaeontology and Meteorites Committee for Heritage Western Cape (HWC) and an advisor on palaeontological conservation and management issues for the Palaeontological Society of South Africa (PSSA), HWC and SAHRA. He is currently compiling technical reports on the provincial palaeontological heritage of Western, Northern and Eastern Cape for SAHRA and HWC. Dr Almond is an accredited member of PSSA and APHP (Association of Professional Heritage Practitioners – Western Cape).

Declaration of Independence

I, John E. Almond, declare that I am an independent consultant and have no business, financial, personal or other interest in the proposed solar energy projects, application or appeal in respect of which I was appointed other than fair remuneration for work performed in connection with the activity, application or appeal. There are no circumstances that compromise the objectivity of my performing such work.



Dr John E. Almond
Palaeontologist
***Natura Viva* cc**