

**PALAEONTOLOGICAL ASSESSMENT
(DESKTOP STUDY)
PROSPECTING RIGHT APPLICATION FOR SEA CONCESSION 12C
CONTINENTAL SHELF OFF OLIFANTRIVIER MOUTH, WESTERN CAPE**

By

**John Pether, M.Sc., Pr. Sci. Nat. (Earth Sci.)
Geological and Palaeontological Consultant**

P. O. Box 48318, Kommetjie, 7976

Tel./Fax (021) 7833023

Cellphone 083 744 6295

jpether@iafrica.com

Prepared at the Request of

Vanessa Maitland, Maritime Archaeologist

African Centre for Heritage Activities (ACHA)

Cell: +27 82 490 4066

Email: vanessa@cocojams.co.za

Client

Aqua Marine Diamonds 12 (Pty) Ltd

18 October 2022

SUMMARY

Site Name

Diamond Sea Concession 12C off southern Namaqualand – Prospecting Right Application by Aqua Marine Diamonds 12 (Pty) Ltd

Location

Sea Concession 12C (221245 ha) is located offshore just north of the Olifantsrivier (Figure 1) and extends northwards from off the Olifantsrivier mouth for ~34 km opposite the coastal farms De Punt, Geelwal Karoo and Klipvley Karoo Kop. The inshore boundary is ~5 km offshore where water depths are 70-80 m below sea level (bsl.) (-70 to -80 m elevation). The offshore boundary is 65 to 90 km offshore in depths of 190 to 230 m bsl. (Figure 2).

Proposed Activities

The Applicant, Aqua Marine Diamonds 12 (Pty) Ltd., proposes to carry out geological exploration for diamonds in Sea Concession 12C off southern Namaqualand (Figure 1).

The proposed prospecting is intended to be accomplished in phases:

- Phase 1 - Geophysical Survey – Multibeam Swath Bathymetry to derive a high-resolution map of the topography of the seabed and shallow seismics to acquire sub-bottom profiles.
- Phase 2a – Vibracoring and Grab Sampling – In depths shallower than ~100 m a vibracorer will be deployed at 100-200 sites to acquire cores of ~10 cm in diameter and up to ~3 m in length. Surface sediments may be sampled at 20-50 sites using a Van Veen grab sampler.
- Phase 2b – Drillship Sampling – To be undertaken by a chartered drillship with a capability of drilling a cased hole 2.5 m in diameter (5 m²) and up to 12 m deep in weakly-lithified sediments.
- Phase 3 - Feasibility Studies - Involves consideration of all factors and potential costs of defining and exploiting the diamond resource.

Impacted Palaeontological Resources

Cretaceous fossil wood occurs primarily in the gravels on the flat middle shelf which directly overlie the source Cretaceous formations.

The unstudied Cenozoic shelly macrofauna comprises phosphatic shell casts and more rare intact shells of Eocene, Miocene and Pliocene ages.

Fossil bones and teeth include the phosphatized/petrified bones and teeth of sharks and other fishes, the skulls of extinct whale species and the occasional bones and teeth of land-living animals that roamed the ice-age exposed shelf.

Shells from the Last Transgression Sequence occur in the sediments accumulated on the shelf during the last 20 thousand years as it was submerged following the low sea level of the Last Ice Age.

The discoveries of the Cretaceous fossil woods, the Miocene petrified whale fossils and the Subantarctic, Algoa and West African molluscan taxa in the Last Transgression Sequence, are all published in the international scientific literature.

Recommendations

The Environmental Management Plans for the prospecting and mining rights areas must therefore include provisions for the collection of representative examples of the fossils that occur therein. As part of Environmental Awareness Training, geological staff involved in logging must be informed of the need to watch for fossil material and rescue such from the vibracores, grab samples and the drillship gravel oversize screen. The prospecting/mining company must apply to SAHRA for a general permit to destroy, damage, excavate, disturb and collect fossils identified during sampling and mining, as per the NHRA.

Fossils may be found during the processing of the vibracores and grab samples. All material of potential interest must have the details of context recorded and be kept for identification by an appropriate specialist and if significant, to be deposited in a curatorial institution such as the IZIKO SA Museum. The best outcome for a set of cores from this poorly-known area is that they are the subject of a detailed study, such as for a B.Sc. Honours or M.Sc. project, with radiocarbon dates. It is possible that a core or two might intersect rarely preserved lagoonal deposits which are important for providing points on the sea-level curve applicable to the West Coast (Runds et al., 2018).

During sampling by a drillship, as part of the normal sampling and mining process, the material crossing the oversize screen (Figure 6) must be monitored for the occurrence of the various fossil types. Potential fossil material should be collected for later identification and evaluation, with labelling and recording of the find data.

Collected samples are to be stored by the company. When a collection of fossil material has been accumulated, the appointed palaeontologist should undertake the identification and evaluation of the fossil material and compile the report for submission to SAHRA. A selection of material could be removed for further study. The Environmental Manager/Officer is to liaise with the appointed palaeontologist on the progress of the fossil collection and the scheduling of the evaluation.

During all operations, personnel can send queries and images by email to an appointed palaeontologist for evaluation and prompt feedback.

CONTENTS

1	INTRODUCTION	1
2	LOCATION.....	1
3	PROPOSED ACTIVITIES.....	2
4	GEOLOGICAL SETTING.....	3
5	PALAEONTOLOGICAL HERITAGE.....	4
6	ANTICIPATED IMPACTS	6
6.1	Vibracores and Grabs	6
6.2	Drillship Sampling	7
6.2.1	Extents	7
6.2.2	Duration.....	7
6.2.3	Intensity	7
	Cretaceous Fossil Wood.....	7
	Cenozoic Shelly Macrofauna	8
	Fossil Bones and Teeth	8
	Shells from the Last Transgression Sequence	8
7	RECOMMENDATIONS.....	9
7.1	Vibracores and Grab Samples.....	9
7.2	Collection of Fossil Material during Prospecting and Mining	9
8	REFERENCES	10
9	APPENDIX 1. METHODOLOGY FOR ASSESSING THE SIGNIFICANCE OF IMPACTS	12
10	APPENDIX 2. PALAEONTOLOGICAL SENSITIVITY RATING	13

SPECIALIST DETAILS, EXPERTISE AND DECLARATION

CURRICULUM VITAE

John Pether, M.Sc., Pr. Sci. Nat. (Earth Sci.)

Independent Consultant/Researcher recognized as an authority with 37 years' experience in the field of coastal-plain and continental-shelf palaeoenvironments, fossils and stratigraphy, mainly involving the West Coast/Shelf of southern Africa. Has been previously employed in academia (South African Museum) and industry (Trans Hex, De Beers Marine). At present an important involvement is in Palaeontological Impact Assessments (PIAs) and mitigation projects in terms of the National Heritage Resources Act 25 (1999) (~350 PIA reports to date) and is an accredited member of the Association of Professional Heritage Practitioners (APHP). Continues to be involved as consultant to offshore and onshore marine diamond exploration ventures. Expertise includes:

- Coastal plain and shelf stratigraphy (interpretation of open-pit exposures, on/offshore cores and exploration drilling).
- Sedimentology and palaeoenvironmental interpretation of shallow marine, aeolian and other terrestrial surficial deposits.
- Marine macrofossil taxonomy (molluscs, barnacles, brachiopods) and biostratigraphy.
- Marine macrofossil taphonomy.
- Sedimentological and palaeontological field techniques in open-cast mines (including finding and excavation of vertebrate fossils (bones).

Membership of Professional Bodies

- South African Council of Natural Scientific Professions. Earth Science. Reg. No. 400094/95.
- Geological Society of South Africa.
- Palaeontological Society of Southern Africa.
- Southern African Society for Quaternary Research.
- Association of Professional Heritage Practitioners (APHP), Western Cape. Accredited Member No. 48.

Past Clients Palaeontological Assessments

AECOM SA (Pty) Ltd.	Guillaume Nel Environmental Management Consultants.
Agency for Cultural Resource Management (ACRM).	Klomp Group.
AMATHEMBA Environmental.	Megan Anderson, Landscape Architect.
Anél Blignaut Environmental Consultants.	Ninham Shand (Pty) Ltd.
Arcus Gibb (Pty) Ltd.	PD Naidoo & Associates (Pty) Ltd.
ASHA Consulting (Pty) Ltd.	Perception Environmental Planning.
Aurecon SA (Pty) Ltd.	PHS Consulting.
BKS (Pty) Ltd. Engineering and Management.	Resource Management Services.
Bridgette O'Donoghue Heritage Consultant.	Robin Ellis, Heritage Impact Assessor.
Cape Archaeology, Dr Mary Patrick.	Savannah Environmental (Pty) Ltd.
Cape EAPrac (Cape Environmental Assessment Practitioners).	Sharples Environmental Services cc
CCA Environmental (Pty) Ltd.	Site Plan Consulting (Pty) Ltd.
Centre for Heritage & Archaeological Resource Management (CHARM).	SRK Consulting (South Africa) (Pty) Ltd.
Chand Environmental Consultants.	Strategic Environmental Focus (Pty) Ltd.
CK Rumboll & Partners.	UCT Archaeology Contracts Office (ACO).
CNdV Africa	UCT Environmental Evaluation Unit
CSIR - Environmental Management Services.	Urban Dynamics.
Digby Wells & Associates (Pty) Ltd.	Van Zyl Environmental Consultants
Enviro Logic	Western Cape Environmental Consultants (Pty) Ltd, t/a ENVIRO DINAMIK.
Environmental Resources Management SA (ERM).	Wethu Investment Group Ltd.
Greenmined Environmental	Withers Environmental Consultants.

Stratigraphic consulting including palaeontology

Afri-Can Marine Minerals Corp	Council for Geoscience
De Beers Marine (SA) Pty Ltd.	De Beers Namaqualand Mines.
Geological Survey Namibia	IZIKO South African Museum.
Namakwa Sands (Pty) Ltd	NAMDEB

DECLARATION OF INDEPENDENCE

BRIEF PALAEOLOGICAL ASSESSMENT (DESKTOP STUDY)

PROSPECTING RIGHT APPLICATION FOR SEA CONCESSION 12C

CONTINENTAL SHELF OFF OLIFANTSRIVIER MOUTH, WESTERN CAPE


Terms of Reference

This assessment forms part of the Heritage Assessment and it assesses the overall palaeontological (fossil) sensitivities of formations underlying the Project Area in terms of the proposed development.

Declaration

I ...**John Pether**....., as the appointed independent specialist hereby declare that I:

- » act/ed as the independent specialist in the compilation of the above report;
- » regard the information contained in this report as it relates to my specialist input/study to be true and correct, and
- » do not have and will not have any financial interest in the undertaking of the activity, other than remuneration for work performed in terms of the NEMA, the Environmental Impact Assessment Regulations, 2014 and any specific environmental management Act;
- » have and will not have any vested interest in the proposed activity proceeding;
- » have disclosed to the EAP any material information that has or may have the potential to influence the decision of the competent authority or the objectivity of any report, plan or document required in terms of the NEMA, the Environmental Impact Assessment Regulations, 2014 and any specific environmental management act;
- » have provided the EAP with access to all information at my disposal regarding the application, whether such information is favourable to the applicant or not; and
- » am aware that a false declaration is an offence in terms of regulation 48 of the 2014 NEMA EIA Regulations.



Signature of the specialist.

Date: 18 October 2021.

GEOLOGICAL TIME SCALE TERMS

For more detail see www.stratigraphy.org.

ka: Thousand years or kilo-annum (10^3 years). Implicitly means “ka ago” *i.e.* duration from the present, but “ago” is omitted. The “Present” refers to 1950 AD. Not used for durations not extending from the Present. For a duration only “kyr” is used.

Ma: Millions years, mega-annum (10^6 years). Implicitly means “Ma ago” *i.e.* duration from the present, but “ago” is omitted. The “Present” refers to 1950 AD. Not used for durations not extending from the Present. For a duration only “Myr” is used.

Tertiary Period: Refers to the Palaeogene and Neogene Periods together. Supposed to be obsolete, but still widely in use.

Mesozoic and Cenozoic Chronostratigraphy
From: International Commission on Stratigraphy.
Chronostratigraphic Chart 2016-12.pdf

Eonothem / Eon Erathem / Era System / Period		Series / Epoch	Stage / Age	GSSP	numerical age (Ma)	
Phanerozoic	Cenozoic	Quaternary	Holocene		present	
			Upper		0.0117	
			Middle		0.126	
		Pleistocene	Calabrian		0.781	
			Gelasian		1.80	
			Zanclean		2.58	
		Pliocene	Piacenzian		3.600	
			Messinian		5.333	
		Neogene	Miocene	Tortonian		7.246
				Serravallian		11.63
				Langhian		13.82
	Burdigalian				15.97	
	Aquitanian				20.44	
	Chatthian				23.03	
	Rupelian				28.1	
	Paleogene	Oligocene	Priabonian		33.9	
			Bartonian		37.8	
			Lutetian		41.2	
		Eocene	Ypresian		47.8	
			Thanetian		56.0	
			Selandian		59.2	
			Danian		61.6	
		Paleocene	Maastrichtian		66.0	
Campanian				72.1 ± 0.2		
Santonian				83.6 ± 0.2		
Mesozoic	Upper Cretaceous	Coniacian		86.3 ± 0.5		
		Turonian		89.8 ± 0.3		
		Cenomanian		93.9		
		Albian		100.5		
	Lower Cretaceous	Aptian		~ 113.0		
		Barremian		~ 125.0		
		Hauterivian		~ 129.4		
		Valanginian		~ 132.9		
Berriasian		~ 139.8				
					~ 145.0	

ICS-approved 2009 Quaternary (SQS/INQUA) proposal

ERA	PERIOD	EPOCH & SUBEPOCH	AGE	AGE (Ma)	GSSP
CENOZOIC	QUATERNARY	HOLOCENE			
			'Tarantian'	0.012	
			'Ionian'	0.126	
	PLIOCENE	Early	Calabrian	0.781	
			Gelasian	1.806	
		Late	Piacenzian	2.588	
			Zanclean	3.600	
	Ng	PLIOCENE			
			Zanclean	5.332	

Vrica, Calabria
Monte San Nicola, Sicily

Holocene: The most recent geological epoch commencing 11.7 ka till the present.

Pleistocene: Epoch from 2.6 Ma to 11.7 ka.
Late Pleistocene 11.7–126 ka.
Middle Pleistocene 135–781 ka.
Early Pleistocene 781–2588 ka.

Quaternary: The current Period, from 2.6 Ma to the present, in the Cenozoic Era.
The Quaternary includes both the Pleistocene and Holocene epochs. As used herein, early and middle Quaternary correspond with the Pleistocene divisions, but late Quaternary includes the Late Pleistocene and the Holocene.

1 INTRODUCTION

The Applicant, Aqua Marine Diamonds 12 (Pty) Ltd., proposes to carry out geological exploration for diamonds in Sea Concession 12C off southern Namaqualand (Figure 1). Consultant Philip le Roux has been appointed to facilitate the Prospecting Right Application process for the proposed exploration activities. Maritime Archaeologist Vanessa Maitland has been appointed to undertake the Heritage Study to inform the application process. This report is the contribution to the Heritage Impact Assessment and it describes the potential palaeontological heritage which may be affected by the proposed activities.

2 LOCATION




Figure 1. Location of Sea Concession 12C and regional geological context.

Sea Concession 12C (221245 ha) is located offshore just north of the Olifantsrivier (Figure 1) and extends northwards from the Olifantsrivier mouth for ~34 km opposite the coastal farms De Punt, Geelwal Karoo and Klipvley Karoo Kop. The inshore boundary is ~5 km offshore where water depths

are 70-80 m below sea level (bsl.) (-70 to -80 m elevation). The offshore boundary is 65 to 90 km offshore in depths of 190 to 230 m bsl. (Figure 2).

3 PROPOSED ACTIVITIES

The proposed prospecting is intended to be accomplished in phases:

- **Phase 1 - Geophysical Survey** – A geophysical survey vessel will undertake acquisition of Multibeam Swath Bathymetry to derive a high-resolution map of the topography of the seabed (DEM). Sub-bottom profiles are to be acquired using a CHIRP shallow seismic system to reveal the sedimentary units and their geometry within local sediment basins. The data will be interpreted and targets for sampling will be defined.
- **Phase 2a – Vibracoring and Grab Sampling** – In depths shallower than ~100 m a vibracorer will be deployed at 100-200 sites to acquire cores of ~10 cm in diameter and up to ~3 m in length. Surface sediments may be sampled at 20-50 sites using a Van Veen grab sampler.
- **Phase 2b – Drillship Sampling** – To be undertaken by a chartered drillship with a capability of drilling a cased hole 2.5 m in diameter (5 m²) and up to 12 m deep in weakly-lithified sediments.
- **Phase 3 - Feasibility Studies** - Involves consideration of all factors and potential costs of exploiting the mineral resource including mining vessel costs, financial modelling, market risks, taxes, environmental impacts, mitigation *etc.*, in order to ascertain the portion of the mineral resource which is a “bankable” mineral reserve which is profitable to mine.




Figure 2. Bathymetry of Sea Concession 12C, surficial shelf sediments and coastal bedrock geology.

4 GEOLOGICAL SETTING

The seabed geology of the continental shelf in the surrounding area is depicted in Figure 1 which shows overall successively younger formations seawards. The Cretaceous and Paleogene units comprise the main bulk of the Olifants Delta and are succeeded by cappings of Miocene and Pliocene units.

Sea Concession 12C is on the flat middle shelf, seaward of the steep slope of the inner-shelf Pre-Cretaceous bedrock. Cretaceous formations (Figure 1). Extrapolating from the seashore exposures, the shallowest margin of 12C inner-shelf bedrock is expected to be mainly comprised of highly deformed, metasedimentary schists, quartzites and limestones of the Gifberg Group which were originally deposited in a rifted ocean basin between 780-760 million years ago (Ma) during the late Precambrian Eon. Older crustal basement gneisses occur in the north.

The Cretaceous, Paleogene and Mio-Pliocene formations comprise the “footwall” to overlying, much younger late Quaternary to present-day deposits which for the most part are a veneer over the continental shelf, but are much thicker where they have accumulated in the accommodation provided by the slope break between the steeper inner shelf and the nearly flat middle shelf (Figure 3) where they constitute the “Holocene Mudbelt” with sediment thicknesses in excess of 10 m.




Figure 3. Schematic geology of the shallow continental shelf based on geophysical seismic profiles, illustrating the “mudbelt” sediment wedge overlying the inner to midshelf transition. Adapted from Birch et al., 1991.

In general the C-concession middle shelf the sediments seaward of the Holocene Mudbelt are a thin veneer usually not exceeding ~1 m thick. However, in the area of 12C the Quaternary sediments are thicker due to sediment delivery by the Olifants River and the terrigenous sediment cover extends farther offshore across the top of the Olifants Delta than to the north (Figure 2). Beneath the fine-grained sediment cover the marine gravels are expected to be thicker in bedrock depressions eroded in the Cretaceous and Cenozoic shelf formations bedrock aka “footwall”. The oldest deposits are also patchily preserved in deeper, local bedrock depressions and palaeochannels in the shelf formations, beneath the latest Quaternary basal gravels. Based on personal observations there are several permutations for what type of deposits might be preserved, largely dependent on the space (depth) within the depressions and position on the shelf. These may include earlier-Quaternary marine conglomerates and sandstones as remnants that escaped erosion during the latest transgression from the Last Ice Age (Last Glacial Maximum) low sea level (Figure 4).

An idealized sequence in a deeper bedrock depression may include all or part of the following (refer to Figure 4);

1. A gravel deposited during the MIS5/6 transgression ~135 ka.
2. A succeeding muddy shelf sand deposited during the LIG highstand.
3. A shelly, variously-cemented sandstone deposited during the subsequent regression (MISs 5a & 4), constituting the basal unit of the **Last Regression Sequence (LRS)**.
4. Subsequent terrestrial deposits that may include colluvium, pan deposits and dune sands, with land snails, possible animal remains and perhaps archaeological material, capped by a soil and/or incipient calcrete (MISs 3 & 2). The terrestrial deposits will only occur inshore of the LGM lowstand.
5. An erosion surface overlain by the basal gravel of the **Last Transgression Sequence (LTS)** (early MIS 1).
6. Succeeding shelly sand fining up to muddy shelf sand (current interglacial highstand, later MIS 1).

However, for the most part the shelf bedrock is overlain only by the LTS basal gravel unit (5) and the shelf upward-fining unit (6).




Figure 4. Sea level history and palaeoclimatic nomenclature for the last 200 ka.

5 PALAEOLOGICAL HERITAGE

Cretaceous fossil wood occurs primarily in the gravels on the flat middle shelf which directly overlie the source Cretaceous formations. Petrified wood is common and includes areas where petrified logs litter the seabed, the "Fossil Forests". Specimens obtained via diamond exploration are providing valuable insights into the palaeo climates of the Cretaceous West Coast, when wide, well-watered coastal plains were covered by forests of primitive yellow wood (podocarp) trees (Bamford & Corbett, 1994; Bamford & Stevenson, 2002; Stevenson & Bamford, 2003).

The **Cenozoic shelly macrofauna** comprises phosphatic shell casts and more rare intact shells of various ages. During later Neogene and Quaternary times the shelf was dominated by upwelling processes, with high organic productivity and authigenic mineralization of seabed rocks, clays and biogenic particles by phosphatization and glauconization. Extensive cemented crusts or “hardgrounds” formed on formations exposed at the seabed. Sea level oscillated repeatedly, dropping to ice-age palaeoshorelines as much as 140 m below present sea level. The hardgrounds were eroded during the ice-age/glacial shallowing episodes, releasing these fossils for incorporation into the LTS gravels.




Figure 5. Sea-level curve for the last deglaciation, showing contexts of dated shells from the West Coast Shelf.

Fossil bones and teeth include the bones and teeth of sharks and other fishes, the skulls of extinct whale species and the occasional remains of land-living animals that roamed the ice-age exposed shelf are also phosphatized and reworked into the latest, loose Last Transgression Sequence sediments on the seabed. A sample of this reworked material turns up in bottom-trawl fishnets, scientific dredging and during diamond-mining operations and the specimens which have been donated to scientific institutions have been invaluable contributions (e.g. Bianucci, Lambert & Post, 2007; Bianucci, Post & Lambert, 2007). All such material should be collected.

Shells from the Last Transgression Sequence refers to the “sub-fossil” shells that occur abundantly in the sediments accumulated on the shelf during the last 20 thousand years as it was submerged to increasing depths. The marine shell fossils which occur in the LTS are predominantly the species expected on the West Coast Shelf, in a deepening-water faunal succession with littoral epifaunal species in the basal gravels, succeeded by infaunal bivalves in clean sands, succeeded by bivalves adapted to dwelling in the capping sulphidic muds. However, unexpected species and “extralimitals” (species beyond their normal home range) are actually quite common.

For instance, the Last Ice Age palaeoshoreline gravels are dominated by a “Venus shell” clam, *Tawera philomela* (Figure 5). This cold-water species, along with others, reached the Cape coast from the mid-Atlantic islands of Tristan da Cunha and Gough, apparently thrived here and then became extinct locally during the last deglaciation (Pether, 1993). During the subsequent deglaciation/warming, several warm-water species from the south and east coasts “invaded” the western shelf temporarily (Figure 5).

This shows a more marked influence of Agulhas water rounding the Cape and affecting the Benguela System during the global-warming steps of the last deglaciation (Pether, 1994). These Agulhas extralimitals have mainly been found during diamond sampling/mining off northern Namaqualand – one may expect them to be more abundant further south such as in 12C, as well as more species occurring. It is important to obtain a comprehensive sample of these occurrences for future study. In addition to dating the incursions on Agulhas influence, the individual shells are snapshot archives of the palaeoceanographic conditions at the time, as revealed by incremental analyses of stable isotopes and trace elements by “spot trails” across the growth lines of shells.

6 ANTICIPATED IMPACTS

Fossils are rare objects, often preserved due to unusual circumstances. This is particularly applicable to vertebrate fossils (bones), which tend to be sporadically preserved and have high value with respect to palaeoecological and biostratigraphic (dating) information. Such fossils are non-renewable resources. Provided that no subsurface disturbance occurs, the fossils remain sequestered. The absence of management and operator mitigatory actions to be alert for fossils and retrieve them will result in their loss. This loss of the opportunity to recover fossils and record their contexts when exposed at a particular site is a negative, irreversible impact.

If mitigatory efforts are made to watch out for and rescue the fossils then the impact is positive for palaeontology. However, there remains a medium to high risk of valuable fossils being lost in spite of management actions to mitigate such loss. The fossils may simply not be noticed or not recognized. Even the most diligent attempts at mitigation can only hope to acquire some fraction of the fossils. This is particularly the case if the fossils are sparsely distributed in the deposits, which is generally the case for scientifically-valuable fossil bone material. A misperception exists that if fossils are sparse in a deposit then the intensity of impact will be low. This is not the case as it is the valuable fossils that are usually scarce, such as fossil bones. The very scarcity of such fossils makes for the added importance of watching for them.

6.1 VIBRACORES AND GRABS

The palaeontological impact of the vibracoring and grab sampling in Concession 12C is considered to be negligible, in view of the minimal volumes of sediment affected.

In the vibracores the small volumes involved greatly reduce the likelihood of capturing the sparse fossils reworked from the older, pre- late Quaternary formations and the “extralimitals” in the Last Transgression Sequence. The grab samples are purposed for obtaining the upper, modern fauna and are unlikely to capture fossils which are usually lower down. Notwithstanding, the grabs may capture poorly-known modern species.

However, should extralimital Subantarctic and Agulhas species occur in the cores they are more important specimens than those selected from the loose, mixed shells crossing the oversize screens on sampling/mining vessels, as they have context in the geological and faunal succession in the core and the expense to have specimens radiocarbon dated is more worthwhile.

6.2 DRILLSHIP SAMPLING

The target areas and number of drillsite sample sites are yet to be determined on the basis of interpretation of the geophysical survey. For each metre drilled a 5 m² drill footprint delivers ~5 m³ of material to the gravel screening plant on the vessel. For the most part the excavated material is the Last Transgression Sequence deposits with expected “subfossil” extant shell species in the “overburden” of fine sands and silts. Scientifically important extralimital species and rare reworked old fossil shells, bones and teeth occur in the gravels and sands of the lowermost metre of drill holes. Ignoring the variable thickness of “overburden”, each drill hole includes a basal, potentially fossiliferous 5 m³ of material.

6.2.1 Extents

In sampling and mining the fossil content of a prescribed volume of deposit is destroyed which is the physical extent of the impact. *i.e.* the extent is local. On the other hand, fossils uncovered during sampling and mining of the coastal plain and offshore are often of sufficient note to publish about them, which is a scientific impact on a national to international scale. For example, the discoveries of the Cretaceous fossil woods, the Miocene petrified whale fossils and the Subantarctic, Algoa and West African molluscan taxa in the Last Transgression Sequence, are all published in the international scientific literature.

6.2.2 Duration

The impact of both the finding or the loss of fossils is permanent. Destroyed fossils are lost to posterity. The found fossils must be preserved for posterity.

6.2.3 Intensity

The intensity of the potential impact of sampling and mining on fossil resources is determined by the palaeontological sensitivity of the affected formations - the potential scientific value of the fossils which are included in it, together with the volume of the formation which is excavated. Overall, the palaeontological sensitivity of marine deposits is HIGH (Almond & Pether, 2009) due to a few, crucial fossil bone finds of high scientific importance that provided the age constraints for the formations. However, there are complications as marine formations usually contain more than one type of fossil of differing importance, *e.g.* common shells and rare bones. Quaternary fossil shell assemblages consist mainly of well-known, usual taxa and it is the unexpected, out of range or unknown, new shell species which are important to distinguish from the expected, common species.

Cretaceous Fossil Wood

Large well-preserved chunks of various types of petrified fossil wood are directly derived from the Cretaceous formations exposed on the middle shelf, as well as transported, abraded cobbles and pebbles in the marine gravels. The impact of the rescue or loss of valuable specimens is rated of MEDIUM intensity.

CRITERIA	WITHOUT MITIGATION	WITH MITIGATION
Extent	National 4	National 4
Duration	Permanent 5	Permanent 5
Intensity	Medium 6	Medium 6
Probability	Probable 3	Probable 3
Confidence	High	High
Significance	Medium negative 45	Medium positive 45
Reversibility	Irreversible	
Mitigation potential	Medium	

Cenozoic Shelly Macrofauna

The fossil shells eroded from the Eocene, Miocene and Pliocene “footwall” formations and incorporated into the overlying LTS basal gravels are usually poorly preserved shell casts and moulds. However, in places on the middle shelf these formations contain beds with well-preserved whole shells which are brought up by the drill. These fossils which are directly derived from the Cenozoic formations are scientifically valuable and practically unstudied. The impact of the rescue or loss of valuable specimens is rated of HIGH intensity.

CRITERIA	WITHOUT MITIGATION	WITH MITIGATION
Extent	National 4	National 4
Duration	Permanent 5	Permanent 5
Intensity	High 8	High 8
Probability	Probable 3	Probable 3
Confidence	High	High
Significance	Medium-high - negative 51	Medium-high - positive 51
Reversibility	Irreversible	
Mitigation potential	Medium	

Fossil Bones and Teeth

This category includes fossil bones and teeth of any origin as there is no purpose for distinctions when they must be captured before going overboard. Recent fresh bones such as those of fish and seals are common and are excluded. The fossil material is phosphatized (petrified) to various degrees and worn by transport and/or pitted by boring organisms. This material is scarce, but the large volumes involved increase the probability that some will be encountered and could be of high scientific value.

CRITERIA	WITHOUT MITIGATION	WITH MITIGATION
Extent	National 4	National 4
Duration	Permanent 5	Permanent 5
Intensity	High 8	High 8
Probability	Probable 3	Probable 3
Confidence	High	High
Significance	Medium-high - negative 51	Medium-high - positive 51
Reversibility	Irreversible	
Mitigation potential	Medium	

Shells from the Last Transgression Sequence

The concern here are shell species which are not typical of the normal faunal assemblages of the Namaqua shelf, are unusual and are often sparse. The table below refers to such occurrences.

CRITERIA	WITHOUT MITIGATION	WITH MITIGATION
Extent	National 4	National 4
Duration	Permanent 5	Permanent 5
Intensity	Medium 6	Medium 6
Probability	Probable 3	Probable 3
Confidence	High	High
Significance	Medium - negative 45	Medium - positive 45
Reversibility	Irreversible	
Mitigation potential	Medium	

7 RECOMMENDATIONS

The exploration and mining for diamonds in the marine environment is a once-off, never to be repeated opportunity to obtain fossils from various areas of the continental shelf, from deposits of various ages. It is cutting-edge, commercially-driven exploration at a scale and detail unaffordable by the state. In order to not overlook such opportunity to advance science co-operatively, the ambit of contemporary environmental management includes such concerns. The additional input from fossil information will be of benefit for the geological interpretation of the deposits. A find of an important fossil can generate favourable publicity. In the longer term, the offshore fossil heritage should also be made available in more permanent exhibitions at an appropriate facility.

The EMPs for the prospecting and mining rights areas must therefore include provisions for the collection of representative examples of the fossils that occur therein. As part of Environmental Awareness Training, geological staff involved in logging must be informed of the need to watch for fossil material and rescue such from the vibracores, grab samples and the drillship gravel oversize screen.

The prospecting/mining company must apply to SAHRA for a general permit to destroy, damage, excavate, disturb and collect fossils identified during sampling and mining, as per the NHRA.

7.1 VIBRACORES AND GRAB SAMPLES

Fossils may be found during the processing of the vibracores and grab samples. These may be obvious, such as petrified bone and teeth and shell casts, usually phosphatic. All material of potential interest must have the details of context recorded and be kept for identification by an appropriate specialist and if significant, to be deposited in a curatorial institution such as the IZIKO SA Museum.

The identification of extralimital, Agulhas “sub-fossil” shell species in the loose shells of the Last Transgression Sequence requires a level of seashell knowledge. The best outcome for a set of cores from this poorly-known area is that they are the subject of a detailed study, such as for a B.Sc. Honours or M.Sc. project, with radiocarbon dates. It is possible that a core or two might intersect rarely preserved lagoonal deposits which are important for providing points on the sea-level curve applicable to the West Coast (Runds *et al.*, 2018).

7.2 COLLECTION OF FOSSIL MATERIAL DURING PROSPECTING AND MINING

As part of the normal sampling and mining process the material crossing the oversize screen (Figure 6) must be monitored for the occurrence of the various fossil types. Potential fossil material should be collected for later identification and evaluation.

For overall monitoring purposes it is suggested that a few small bulk samples of shells (~5 litres) be collected on occasion. The idea is to sample the typical assemblage at a few points in the sampling/mining area. It is possible that an uncommon assemblage may be encountered, such as a shallow-water fauna or a lagoonal fauna, in which case it should also be sampled.

Data to be recorded during fossil collection includes:

- Date
- Company name
- Sample no.
- Collector's name
- Position (co-ordinates)
- Water depth
- Sample subsurface depth
- Vessel
- Brief description and photographs

- A copy of the graphic log of the sample drill hole or mining face showing the vertical sequence of units and the estimated location of the fossil in the sequence.
- A map of the fossil finds in the particular sampling/mining area, such as a contoured multibeam bathymetric image showing the context of samples in relation to the bedrock topography and sediment bodies.



Figure 6. The gravel oversize screen on a typical diamond mining vessel where the geological personnel monitor the material being dredged and where fossil collection takes place.

Collected samples are to be temporarily stored by the company.

When a collection of fossil material has been accumulated, the appointed palaeontologist should undertake the identification and evaluation of the fossil material and compile the report for submission to SAHRA. A selection of material could be removed for further study. The Environmental Manager/Officer is to liaise with the appointed palaeontologist on the progress of the fossil collection and the scheduling of the evaluation.

During all operations, personnel can send queries and images by email to an appointed palaeontologist for evaluation and prompt feedback.

8 REFERENCES

- Almond, J.E. & Pether, J. 2009. Palaeontological Heritage of the Northern Cape. SAHRA Palaeotechnical Report, Natura Viva cc., Cape Town.
- Bamford M.K. & Stevenson I.R. 2002. A submerged Late Cretaceous podocarpaceous forest, west coast, South Africa. *S. Afr. J. Sci.* 98, 181–185.
- Bamford, M.K. & Corbett, I.B. 1994. Fossil wood of Cretaceous age from the Namaqualand continental shelf, South Africa. *Palaeontologia Africana* 31: 83-95.

- Bianucci, G., Lambert, O. & Post, K. 2007. A high diversity in fossil beaked whales (Odontoceti, Ziphiidae) recovered by trawling from the sea floor off South Africa. *Geodiversitas* 29(4); 5–62.
- Bianucci, G., Post, K. & Lambert, O. 2007. Beaked whale mysteries revealed by seafloor fossils trawled off South Africa. *South African Journal of Science* 104; 140-142.
- Birch, G.F., Day, R.W. & Du Plessis, A. 1991. Nearshore Quaternary sediments on the West Coast of southern Africa. *Geological Survey of South Africa Bulletin* 101. 14 pp.
- De Wet, W. M. 2012. Bathymetry of the South African continental shelf. M.Sc thesis, Department of Geological Sciences, University of Cape Town. 255 pp.
- Dingle, R.V. & Siesser, W.G. 1977. Geology of the continental margin between Walvis Bay and Ponto do Ouro. Map - Geological Survey of South Africa Marine Geoscience Series.
- Pether, J. 1993. Relict shells of Subantarctic Mollusca from the Orange Shelf, Benguela Region, off southwestern Africa. *The Veliger* 36 (3): 276-284.
- Pether, J. 1994. Molluscan evidence for enhanced deglacial advection of Agulhas water in the Benguela Current, off southwestern Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* 111: 99-117.
- Pether, J. 1995. *Belichnus* new ichnogenus, a ballistic trace on mollusc shells from the Holocene of the Benguela region, South Africa. *Journal of Paleontology* 69: 171-181.
- Runds, M.J., Bordy, E.M. & Pether, J. 2019. Late Quaternary sedimentological history of a submerged gravel barrier beach complex, southern Namibia. *Geo-Marine Letters* 39: 469-491.
- Siddall, M., Chappell, J. & Potter, E.-K. 2007. Eustatic sea level during past interglacials. *The Climate of Past Interglacials: Developments in Quaternary Science* 7: 75-92.
- Stevenson I.R. & Bamford, M. 2003. Submersible-based observations of in situ fossil tree trunks in Late Cretaceous seafloor outcrops, Orange Basin, western offshore, South Africa. *South African Journal of Geology* 106: 315-326.

9 APPENDIX 1. METHODOLOGY FOR ASSESSING THE SIGNIFICANCE OF IMPACTS

EFFECT	Extents/Spatial Scale		E
	Localized	At localized scale and a few hectares in extent.	1
	Study area	The proposed site and its immediate environs.	2
	Regional	District and Provincial level.	3
	National	Country.	4
	International	Internationally.	5
	Duration/Temporal Scale		D
	Very short	Less than 1 year.	1
	Short term	Between 2 to 5 years.	2
	Medium term	Between 5 and 15 years.	3
	Long term	Exceeding 15 years and from a human perspective almost permanent.	4
	Permanent	Resulting in a permanent and lasting change.	5
	Magnitude/Intensity (Palaeontological Sensitivity)		M
	No potential	Formations entirely lacking fossils such as igneous rocks.	0
	Marginal	Limited probability for producing fossils from certain contexts at localized outcrops.	2
	Low	Depositional environment where fossils are unlikely to be preserved, or are judged unlikely to produce unique fossil remains.	4
	Medium	Strong potential to yield fossil remains based on stratigraphy and/or geomorphologic setting.	6
	High	Formations known to contain palaeontological resources that include rare, well-preserved fossil materials.	8
	Very high	Formations/sites known or likely to include vertebrate fossils pertinent to human ancestry and palaeoenvironments and which are of international significance.	10
	Probability/Likelihood		P
	Very improbable	Probably will not happen.	1
	Improbable	Some possibility, but low likelihood.	2
	Probable	Distinct possibility of these impacts occurring.	3
	Highly probable	The impact is most likely to occur.	4
Definite	The impact will definitely occur regardless of prevention measures.	5	

SIGNIFICANCE = (E+D+M)P		
< 30	LOW	The impact would not have a direct influence on the decision to develop in the area
30-60	MEDIUM	The impact could influence the decision to develop in the area unless it is effectively mitigated
>60	HIGH	The impact must have an influence on the decision process to develop in the area

10 APPENDIX 2. PALAEOONTOLOGICAL SENSITIVITY RATING

Palaeontological Sensitivity refers to the likelihood of finding significant fossils within a geologic unit.

VERY HIGH: Formations/sites known or likely to include vertebrate fossils pertinent to human ancestry and palaeoenvironments and which are of international significance.

HIGH: Assigned to geological formations known to contain palaeontological resources that include rare, well-preserved fossil materials important to on-going palaeoclimatic, palaeobiological and/or evolutionary studies. Fossils of land-dwelling vertebrates are typically considered significant. Such formations have the potential to produce, or have produced, vertebrate remains that are the particular research focus of palaeontologists and can represent important educational resources as well.

MODERATE: Formations known to contain palaeontological localities and that have yielded fossils that are common elsewhere, and/or that are stratigraphically long-ranging, would be assigned a moderate rating. This evaluation can also be applied to strata that have an unproven, but strong potential to yield fossil remains based on its stratigraphy and/or geomorphologic setting.

LOW: Formations that are relatively recent or that represent a high-energy subaerial depositional environment where fossils are unlikely to be preserved, or are judged unlikely to produce unique fossil remains. A low abundance of invertebrate fossil remains can occur, but the palaeontological sensitivity would remain low due to their being relatively common and their lack of potential to serve as significant scientific resources. However, when fossils are found in these formations, they are often very significant additions to our geologic understanding of the area. Other examples include decalcified marine deposits that preserve casts of shells and marine trace fossils, and fossil soils with terrestrial trace fossils and plant remains (burrows and root fossils)

MARGINAL: Formations that are composed either of volcanoclastic or metasedimentary rocks, but that nevertheless have a limited probability for producing fossils from certain contexts at localized outcrops. Volcanoclastic rock can contain organisms that were fossilized by being covered by ash, dust, mud, or other debris from volcanoes. Sedimentary rocks that have been metamorphosed by the heat and pressure of deep burial are called metasedimentary. If the meta sedimentary rocks had fossils within them, they may have survived the metamorphism and still be identifiable. However, since the probability of this occurring is limited, these formations are considered marginally sensitive.

NO POTENTIAL: Assigned to geologic formations that are composed entirely of volcanic or plutonic igneous rock, such as basalt or granite, and therefore do not have any potential for producing fossil remains. These formations have no palaeontological resource potential.

Adapted from Society of Vertebrate Paleontology. 1995. Assessment and Mitigation of Adverse Impacts to Nonrenewable Paleontologic Resources - Standard Guidelines. News Bulletin, Vol. 163, p. 22-27.