

SiVEST SA (PTY) LTD

**PROPOSED CONSTRUCTION OF THE KOUP 2 WIND
ENERGY FACILITY AND ASSOCIATED GRID
INFRASTRUCTURE, NEAR BEAUFORT WEST,
WESTERN CAPE PROVINCE, SOUTH AFRICA**

Palaeontological Heritage Report

DEFF Reference: 14/12/16/3/3/2/2121
Report Prepared by: Dr John E. Almond
Natura Viva cc
PO Box 12410 Mill Street
CAPE TOWN 8010, RSA

Issue Date: 4 April 2022
Version No.: Draft2

SIVEST SA (PTY) LTD

PROPOSED CONSTRUCTION OF THE KOUP 2 WIND ENERGY FACILITY AND ASSOCIATED GRID INFRASTRUCTURE, NEAR BEAUFORT WEST, WESTERN CAPE PROVINCE, SOUTH AFRICA

PALAEONTOLOGICAL HERITAGE REPORT

EXECUTIVE SUMMARY

Genesis Enertrag Koup 2 Wind (Pty) Ltd is proposing to construct the Koup 2 WEF, comprising thirty-two wind turbines with a maximum total energy generation capacity of up to 140MW, with a 132kV overhead power line connection to the national grid. A Battery Energy Storage System (BESS) will be located next to the onsite 33/132kV substation. The WEF and grid project areas are located in the Great Karoo region some 60 km south of Beaufort West, falling within the Beaufort West Local Municipality (Central Karoo District Municipality) of the Western Cape Province.

The Koup 2 WEF and grid connection project area is underlain by continental (fluvial / lacustrine) sediments of the Abrahamskraal and Teekloof Formations (Lower Beaufort Group, Karoo Supergroup) which are of Middle to Late combined Permian age and are provisionally assigned a Very High sensitivity on the SAHRIS palaeosensitivity map as well as the DFFE screening map. These bedrocks contain sparse, unpredictable to locally concentrated vertebrate fossils as well as rare trace fossils (e.g. tetrapod burrows) and plant material that are of scientific and conservation value. A significant number of new fossil vertebrate sites (cranial and post-cranial material of large-bodied dinocephalians, small dicynodonts, rare tetrapod burrow casts) have been recorded within the combined WEF / grid connection project areas during a 5-day site visit, while several fossil sites have previously been mapped shortly outside its margins. These palaeontological sites, together with their sedimentological context, provide important data for on-going research into the pattern and causes of the Middle Permian Mass Extinction Event on land around 260 million years ago.

Scientifically-valuable and legally-protected fossil heritage resources, preserved at or beneath the ground surface, within the project footprint are potentially threatened by surface clearance and bedrock excavations during the construction phase of the WEF and grid connection (e.g. for access roads, wind turbine foundations). The majority of the recorded fossil sites lie outside the project footprint but most of the WEF and grid connection footprint has yet to be palaeontologically surveyed on foot. A significant number of unrecorded sites likely to exist within or very close to the project footprint.

No Very High Sensitivity or No-Go palaeontological sites or areas have been identified within the WEF and grid connection project areas. Since all known fossil sites can be readily mitigated through professional recording and collection of fossil material in the pre-construction phase, no recommendations for micro-siting of infrastructure such as wind turbine, pylon positions or access roads are therefore made here. There are no preferences on palaeontological heritage grounds for specific site options for the Koup 2 WEF on-site substation and construction laydown area. There is no preference for Grid Option 1 (now the preferred corridor) or 2 (either alternative) on

palaeontological heritage grounds. Since they are of similar length and traverse bedrocks of comparable palaeosensitivity, they are likely to have similar potential impacts on fossil sites.

The proposed Koup 2 WEF and associated grid connection developments are assigned a similar overall impact significance rating (Construction Phase) of NEGATIVE MEDIUM without mitigation and NEGATIVE LOW following mitigation. No significant further impacts on fossil heritage resources are anticipated in the planning, operational and decommissioning phases. The No-Go Option may have a NEGATIVE LOW impact significance; fossils will continue to be exposed and destroyed by natural weathering processes while the positive benefits of professional mitigation (*viz.* improved palaeontological database) will be lost. Anticipated cumulative impacts in the context of several planned or authorized renewable energy projects in the region are assessed as NEGATIVE MEDIUM before mitigation and NEGATIVE LOW after mitigation.

Recommended mitigation comprises:

- (1) A specialist palaeontological walk-down of the final WEF and grid connection project area in the pre-construction phase,
- (2) Implementation of a Chance Fossil Finds Protocol (See Appendix 4) by the ECO / ESO during the construction phase. The specialist palaeontologist responsible will be required to submit a Work Plan for approval by Heritage Western Cape.

Conclusion

No fatal flaws were identified and anticipated impacts can be substantially reduced through mitigation during the pre-construction and construction phases. On condition that the recommended mitigation measures are included within the relevant EMPs and implemented in full, there are no objections on palaeontological heritage grounds to the authorization of the proposed Koup 2 WEF and the associated grid connection.

NATIONAL ENVIRONMENTAL MANAGEMENT ACT, 1998 (ACT NO. 107 OF 1998) AND ENVIRONMENTAL IMPACT REGULATIONS, 2014 (AS AMENDED) - REQUIREMENTS FOR SPECIALIST REPORTS (APPENDIX 6)

Regulation GNR 326 of 4 December 2014, as amended 7 April 2017, Appendix 6	Section of Report
1. (1) A specialist report prepared in terms of these Regulations must contain- a) details of- i. the specialist who prepared the report; and ii. the expertise of that specialist to compile a specialist report including a curriculum vitae;	1.2 & Appendix 1
b) a declaration that the specialist is independent in a form as may be specified by the competent authority;	7.7 & Addendum
c) an indication of the scope of, and the purpose for which, the report was prepared;	1.1 & Appendix 4
(cA) an indication of the quality and age of base data used for the specialist report;	1.3.1.
(cB) a description of existing impacts on the site, cumulative impacts of the proposed development and levels of acceptable change;	5
d) the date and season of the site investigation and the relevance of the season to the outcome of the assessment;	1.3.1.
e) a description of the methodology adopted in preparing the report or carrying out the specialised process inclusive of equipment and modelling used;	1.3.1.
f) details of an assessment of the specific identified sensitivity of the site related to the proposed activity or activities and its associated structures and infrastructure, inclusive of a site plan identifying site alternatives;	3.2 & 6
g) an identification of any areas to be avoided, including buffers;	6.2.
h) a map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	Figure 61
i) a description of any assumptions made and any uncertainties or gaps in knowledge;	2
j) a description of the findings and potential implications of such findings on the impact of the proposed activity, (including identified	5,6, & 7

alternatives on the environment) or activities;	
k) any mitigation measures for inclusion in the EMPr;	8 & Appendix 4
l) any conditions for inclusion in the environmental authorisation;	8
m) any monitoring requirements for inclusion in the EMPr or environmental authorisation;	8 & Appendix 4
n) a reasoned opinion- <ul style="list-style-type: none"> i. (as to) whether the proposed activity, activities or portions thereof should be authorised; <ul style="list-style-type: none"> (iA) regarding the acceptability of the proposed activity or activities; and ii. if the opinion is that the proposed activity, activities or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan; 	9
o) a description of any consultation process that was undertaken during the course of preparing the specialist report;	n/a
p) a summary and copies of any comments received during any consultation process and where applicable all responses thereto; and	n/a
q) any other information requested by the competent authority.	
2) Where a government notice <i>gazetted</i> by the Minister provides for any protocol or minimum information requirement to be applied to a specialist report, the requirements as indicated in such notice will apply.	

SiVEST SA (PTY) LTD

PROPOSED CONSTRUCTION OF THE KOUP 2 WIND ENERGY FACILITY AND ASSOCIATED GRID INFRASTRUCTURE, NEAR BEAUFORT WEST, WESTERN CAPE PROVINCE, SOUTH AFRICA

Contents

1.	INTRODUCTION	12
1.1	Terms of Reference	12
1.2	Specialist Credentials	13
1.3	Assessment Methodology	13
1.3.1	Information sources	13
1.3.2	Study approach.....	14
1.3.3	Legislative context	15
2.	ASSUMPTIONS AND LIMITATIONS	16
3.	TECHNICAL DESCRIPTION	18
3.1	Project Location	18
3.1.1	WEF	18
3.1.2	Grid Connection.....	19
3.2	Project Description.....	21
3.2.1	Wind Farm Components.....	21
3.2.2	Grid Components.....	22
3.3	Layout alternatives.....	22
3.3.1	Wind Energy Facility.....	22
3.3.2	Grid Components.....	24
3.3.3	No-go Alternative.....	24
4.	LEGAL REQUIREMENT AND GUIDELINES	24
4.1	Legislative context	24
5.	DESCRIPTION OF THE RECEIVING ENVIRONMENT	26
5.1	Geological context	26
5.2	Palaeontological heritage.....	37
6.	IDENTIFICATION AND ASSESSMENT OF IMPACTS	60

6.1	Palaeontological sensitivity of the project area.....	60
6.2	Results of the Palaeontological Desktop and Field Study	61
6.2.1	WEF project area.....	61
6.2.2	Grid connection project area	62
6.3	Identification of Potential Impacts.....	66
6.4	Assessment of WEF and grid connection project impacts.....	66
6.4.1	Construction Phase: Disturbance, damage or destruction of fossils.....	67
6.4.2	No-Go Option impacts	67
6.5	Cumulative impacts.....	67
6.6	Overall Impact Rating	73
7.	COMPARATIVE ASSESSMENT OF ALTERNATIVES.....	75
7.1	Koup 2 WEF	75
7.2	Koup 2 grid connection	75
8.	PROPOSED MONITORING AND MITIGATION: INPUT TO EMPR.....	76
9.	SUMMARY & CONCLUSIONS.....	77
9.1	Summary of Findings	77
9.2	Conclusions and Impact Statement.....	78
10.	REFERENCES.....	78
11.	ACKNOWLEDGEMENTS.....	83

List of Tables

Table 1: Renewable energy developments proposed within a 35km radius of the Koup 2 WEF application site (See also text for Heuweltjies WEF, Kraaltjies WEF, Kwagga 1-3 WEFs not listed here).....	70
Table 2: Assessment of paleontological heritage impacts for the proposed Koup 2 Wind Energy Facility (Construction Phase).....	71
Table 3: Assessment of paleontological heritage impacts for the proposed Koup 2 Wind Energy Facility grid connection (Construction Phase).....	71
Table 4: Assessment of impacts for the No Go Option.....	72
Table 5: Assessment of cumulative impacts for the Koup 1 & 2 WEFs and other renewable energy developments in the region.	72
Table 6: Overall impact rating for the Koup 2 WEF project.....	73
Table 7: Overall impact rating for the Koup 2 WEF grid connection project	74
Table 8: Overall cumulative impact rating for the Koup 2 WEF and grid connection project	74
Table 9: Comparative assessment of WEF layout options.....	75
Table 10: Comparative assessment of WEF grid connection options	75

List of Figures

Figure 1: Regional Context Map	18
Figure 2: Koup 2 WEF Site Locality	19
Figure 3: Proposed 132kV Power Line Route Alignment.....	20
Figure 4: Alternatives proposed as part of the Koup 2 WEF.....	23
Figure 5: Extract from 1: 250 000 geology sheet 3222 Beaufort West showing the boundaries of the combined Koup 1 and Koup 2 WEF project area to the south of Beaufort West (yellow polygons). Note numerous W-E trending fold axes in the region which falls within the northern margins of the Cape Fold Belt. Pa (pale green) = Abrahamskraal Formation (Adelaide Subgroup, Lower Beaufort Group). Pt (dark green) = Teekloof Formation (Adelaide Subgroup, Lower Beaufort Group). Yellow = Late Caenozoic / Quaternary superficial sediments, including alluvium, sheet wash, colluvium, soils, locally cemented by pedocretes such as calcrete. To the west of the N12 and largely outside the WEF project area triangular symbols indicate fossil localities within the Pristerognathus Assemblage Zone (N.B. This fossil biozone data is now outdated – see updated stratigraphic chart presented below). A single fossil site for the underlying Tapinocephalus Assemblage zone (star symbol) is indicated c. 4 km to the NW of the WEF project area.	27
Figure 6: Stratigraphic subdivision of the Karoo Supergroup with the rock units and fossil biozones most relevant to the present PIA study outlined in green (Modified from Smith et al. 2020). In the Koup WEF project area fossil assemblages within the uppermost Abrahamskraal Formation (Karelskraal Member) and lower part of the Poortjie Member of the Teekloof Formation are now assigned to the Diictodon-Styracocephalus Assemblage Zone.....	28
Figure 7: View towards a low ridge of Poortjie Member sandstones along the southern margins of the Koup 2 WEF project area (Farm 11/380). The lower terrain in the foreground is underlain by the upper Abrahamskraal Formation.....	31
Figure 8: Typical dissected, hilly terrain the central portion of the Koup 2 WEF project area (Farm 11/380). Dark grey areas are exposures of Beaufort group mudrock while the central yellowish to orange patch features weathered tuffite (mixed volcanic ash and terrigenous sediment) gravels.....	31
Figure 9: Intensely gullied lower hillslopes in the Koup 2 WEF project area (Farm 1/380) showing extensive good Karelskraal Member overbank mudrock exposure.	32
Figure 10: Low, flat-topped koppies in the central portion of the Koup 2 WEF (Farm 1/380, looking to the NW) showing stepped slopes of Abrahamskraal Formation sandstones and thin sandstones and capped by thicker Poortjie Member sandstones.....	32
Figure 11: Outstanding exposures of multi-hued Karelskraal Member mudrocks with intercalated thin, pale crevasse splay sandstones capped by Poortjie Member sandstones, Koup 2 WEF project area (Farm 1/380).	33
Figure 12: Horizon of large (boulder-sized), spheroidal concretions of rusty-brown ferruginous carbonate within the Karelskraal Member in the Koup 2 WEF project area (Farm 1/380).	33
Figure 13: Well-developed palaeosol horizon (fossil soil) marked by pale brown, ferruginous calcrete concretions in the Karelskraal Member, Koup 2 WEF project area (Farm 1/380). These horizons are apparently only rarely fossiliferous.....	34
Figure 14: Thin palaeosol horizon (arrowed) marked by cobble-sized, sphaeroidal calcrete concretions with a micritic composition, Karelskraal Member, Koup 2 WEF project area (Farm 1/380). Many fossil vertebrate specimens are associated with these fossil semi-arid climate soils.	34
Figure 15: Well-developed lens of koffieklip (ferruginous carbonate), here associated with an underlying pale yellowish-green tuffite bed of possible lacustrine origin, Karelskraal Member, Koup 2 WEF project area (Farm 8/380).	35
Figure 16: Boudinage or soft-sediment loading of thin crevasse-splay sandstones overlying dark, water-rich mudrocks of probable lacustrine origin, Karelskraal Member, Koup 2 WEF project area (Farm 8/380) (Hammer = 30 cm).	35
Figure 17: One of several thin, blocky-weathering tuffite horizons within the Karelskraal – Poortjie stratigraphic interval Koup 2 WEF project area (Farm 8/380) (Hammer = 30 cm). These waterlain volcanic ash layers can be accurately dated using radiometric methods.....	36
Figure 18: Thick, massive to subtly bedded, silty to finely-gravelly Late Caenozoic alluvial deposits overlying coarser basal gravels, stream bank exposure in the Koup 2 WEF project area (Farm 1/380) (Hammer = 30 cm).	36

Figure 19: Typical orange-hued, well-rounded gravels generated by corestone weathering of fine-grained Karelskraal Member sandstones, Koup 2 WEF project area (Farm 8/380). Flaked stone artefacts manufactured from corestones are frequently found among such deposits. ...37

Figure 20: Distribution of recorded vertebrate fossil sites within the southern portion of the Main Karoo Basin (modified from Nicolas 2007). The approximate location of the Koup WEF project area to the south of Beaufort West (BW) is indicated by the red rectangle. The high density of recorded fossil sites along the N12 here as well as further to the north is notable. .38

Figure 21: Vertebrate fossil localities within the Lower Beaufort Group in the region south of Beaufort West where the boundary between the Abrahamskraal and Teekloof Formations is highly folded (dotted line). Tapinocephalus Assemblage Zone specimens are found in the far south (small open circles) while Pristerognathus Assemblage Zone fossils (black dots) are associated with outcrops of the Poortjie Member (lowermost Teekloof Formation) (Map abstracted from Keyser & Smith 1977-78). The paucity of fossil records east of the N12 tar road reflects the generally poor bedrock exposure in the area (“min dagsome”) compared with more dissected terrain to the west of the road.39

Figure 22: Chart showing the ranges of known terrestrial tetrapod genera from the Middle to Late Permian of the Main Karoo Basin (From Day et al. 2015b). The boundary between the Abrahamskraal and Teekloof Formations is associated with a catastrophic extinction event at the end of the Capitanian Stage (c. 260 Ma) that has been dated here on the basis of a tuff horizon close to the contact of the Karelskraal and Poortjie Members (yellow star). Key victims of the extinction event were almost all the large-bodied dinocephalians and pareiasaur parareptiles as well as many (but not all) dicynodonts and therocephalians.39

Figure 23: Skulls of two key fossil therapsid tetrapods from the upper part of the Tapinocephalus Assemblage Zone – the small-bodied dicynodont Diictodon (top) and the large-bodied dinocephalian Styracocephalus. Note the very thick cranial roof in the latter, a possible adaptation for head-butting shown by many tapinocephalid dinocephalians.40

Figure 24: Skeleton of the tapinocephalid (thick-skulled) dinocephalian Moschops, a rhino-sized herbivorous therapsid that reached lengths of 2.5 to 3 m and may have lived in small herds. Postcranial remains (and much rarer fossil skull material) of several dinocephalians have been recorded within and just outside the Koup WEF project area.41

Figure 25: Artist’s reconstructions of the indescribably ugly late Middle Permian dinocephalians Styracocephalus (left) and Criocephalosaurus (right). One or both these taxa may occur within or close to the WEF project area.41

Figure 26: Large proximal limb bone of a large-bodied tetrapod (probably dinocephalian) embedded within dark Karelskraal Member mudrocks (Loc. 665) (Scale is 15 cm long).44

Figure 27: Weathered-out large vertebrae collected from stream gravels downslope of the in situ specimen illustrated above and probably belonging to the same animal (Loc. 665) (Top RHS block 15 cm across).44

Figure 28: Partial skull roof of a tapinocephalid dinocephalian – probably Criocephalosaurus (Loc. 724) found on the Farm Bloemendal, just N of the Koup WEF project area (Scale is 15 cm long). The specimen probably comes from the lower Poortjie and, if so, would count among the last surviving members of the Dinocephalia known (See also following figure).45

Figure 29: Oblique ventral view of the tapinocephalid specimen illustrate above. The skull is c. 25 cm wide and the skull roof as seen here is over 15 cm thick.45

Figure 30: In situ, ferruginized partial postcranial skeleton of a dinocephalian therapsid embedded within Karelskraal Member mudrocks (Loc. 603) (Scale = 15 cm). Pyrite pseudomorph crystals within some of the bones as well as rippled sandstones in the vicinity suggest preservation of a corpse along waterlogged lake margin.46

Figure 31: Heap of fragmentary, disarticulated postcranial and cranial material collected in the immediate vicinity of the in situ dinocephalian skeleton illustrated above (Loc. 603) (Hammer = 30 cm).46

Figure 32: Weathered jaw fragment of a dinocephalian with deep-rooted teeth associated with the skeleton in Figure 30 above (Loc. 603) (Block is c. 10 cm across).47

Figure 33: Probable fragmentary cranial and other remains associated with the skeleton in Figure 30 (Loc. 603) (Scale = 15 cm).47

Figure 34: Fragments of limb bones from the same individual (Loc. 603) (Scale = 15 cm).48

Figure 35: Further fragments of limb bones from the same individual (Loc. 603) (Scale = 15 cm).48

Figure 36: Surface scatter of white, weathered bone material of an unidentified large-bodied tetrapod found as surface float overlying the upper Karelskraal Member (Loc. 660) (Scale = 15 cm).	49
Figure 37: Close up of one of the sandstone and bone blocks illustrated above showing highly fragmentary preservation, possibly due to sun-cracking during pre-burial exposure on the ancient Karoo floodplain (Loc. 660) (Scale in cm).	49
Figure 38: Typical sparse scatter of fragmentary postcranial bones of an unidentified large-bodied tetrapod – pareiasaur or dinocephalian - from the Karelskraal Member, found as surface float (Loc. 623) (Largest block is c. 15.5 cm across). Such poorly-preserved postcranial material is mostly of limited scientific value.	50
Figure 39: Thin channel sandstone containing dispersed, reworked post-cranial remains of a large-bodied tetrapod (arrowed) in the upper Karelskraal Member (Loc. 675) (Scale = 15 cm). This site is associated with a downslope trail of weathered-out bones within sandstone blocks.	50
Figure 40: Pedogenic calcrete nodule in float containing several articulated vertebrae, possibly from the tail, of a medium- to large-bodied tetrapod, upper Karelskraal Member (Loc. 613) (Block is c. 9 cm across).	51
Figure 41: Ferruginized calcrete concretion within cleaved overbank mudrocks of the upper Karelskraal Member containing a row of ribs of a large-bodied tetrapod (Loc. 668) (Concretion is c. 30 cm across).	51
Figure 42: Articulated complete skull of a small dicynodont therapsid almost entirely enclosed in a pedogenic calcrete concretion, found among float overlying the Karelskraal Member (Loc. 577) (Scale in cm and mm). The specimen shows several post-canine teeth (not illustrated here) and so does not belong to the common genus Diictodon.	52
Figure 43: Skull of a small dicynodont from the Karelskraal Member (Loc. 631) showing a broad skull table, i.e. not Diictodon (Specimen is 7 cm long).	52
Figure 44: Poorly preserved dicynodont skull within a pedogenic calcrete concretion, Karelskraal Member (Loc. 693) (Specimen is c. 10 cm long).	53
Figure 45: Possible (but equivocal) large, inclined vertebrate burrow cast within cleaved mudrocks of the Karelskraal Member (Loc. 594) (Hammer = 30 cm).	53
Figure 46: Possible (but equivocal) large, inclined vertebrate burrow cast in the Karelskraal Member (Loc. 595), recorded within a few meters of the previous specimen (Hammer = 30 cm).	54
Figure 47: Skull of a small-bodied dicynodont – probably Diictodon - from a mudrock interval within the Poortjie Member, seen in dorsal view (Loc. 641) (Specimen is c. 6.5 cm long).	54
Figure 48: Postcranial remains of a small-bodied therapsid, probably dicynodont, preserved within a pedogenic calcrete concretion from Poortjie Member mudrocks (Loc. 641) (Block is c. 11 cm across).	55
Figure 49: Well-preserved skull with articulated lower jaw and clear canine tusk preserved in situ within Poortjie Member mudrocks, oblique dorso-lateral view (Loc. 649) (Specimen is c. 8.5 cm long).	55
Figure 50: Small pedogenic calcrete concretion from the Poortjie Member enclosing the skull of a small-bodied dicynodont (Loc. 699) (Scale in cm).	56
Figure 51: In situ skull of a small dicynodont within cleaved siltstones of the Poortjie Member (Loc. 706) (Scale in cm and mm).	56
Figure 52: Post-crania, including limbs and rib cage, of a small-bodied dicynodont preserved inside a calcrete concretion embedded within cleaved Poortjie Member mudrocks (Loc. 687) (Scale = 15 cm).	57
Figure 53: Skull of small dicynodont within cleaved Poortjie Member mudrocks (Loc. 689) (Scale in cm).	57
Figure 54: Skull of small dicynodont with a broad skull table and clear pineal foramen weathering out of Poortjie Member mudrocks (Loc. 696).	58
Figure 55: Convex sandstone cast of a small vertebrate burrow weathering out of cleaved Poortjie Member mudrocks (Loc. 688) (Scale in cm and mm).	58
Figure 56: Cast of small subhorizontal burrow, either tetrapod or perhaps invertebrate, within Poortjie Member mudrocks (Loc. 647) (Scale in cm).	59
Figure 57: Sandstone cast of small, inclined burrow (sloping down to the LHS) with a smooth central zone, possibly constructed by a small vertebrate (Loc. 648) (Scale = 15 cm).	59

Figure 58: Paleontological sensitivity map for the Koup 2 WEF project area. Due to the scarcity of well-preserved, scientifically important fossils over much of this region, based on desktop studies and fieldwork, it is inferred that large parts of the project area are in practice of LOW palaeontologically sensitivity. However, there is also a substantial number of, dispersed and unpredictable fossil sites here of HIGH to VERY HIGH sensitivity. Applying the precautionary principle, the Koup 2 WEF and associated grid connection project area are best assigned an overall High Sensitivity as far as palaeontological heritage is concerned.....61

Figure 59: Google Earth© satellite image of the adjoining Koup 1 WEF (orange polygon) and Koup 2 WEF (yellow polygon) project areas, showing the numbered new fossil sites recorded during the site visit. No No-Go palaeontological heritage sites or areas have been identified. Most recorded fossil sites lie outside the project footprint and all can be mitigated through pre-construction collection. Most of the WEF and grid connection footprint has not been palaeontologically surveyed on foot and is likely to contain additional fossil sites at surface that are of palaeontological and conservation value. A pre-construction palaeontological walkdown of the final Koup 1 WEF and grid connection footprints is therefore recommended here. 63

Figure 60: Map showing project areas for authorized and proposed renewable energy projects within a 35 km radius of the Koup 1 and Koup 2 WEF and grid connection project areas (Image provided by SiVEST). Additional unmapped WEFs proposed to the east and southeast of the Koup project area (Heuweltjies WEF, Kraaltjies WEF, Kwagga 1-3 WEFs) have also been taken into consideration here..... 69

List of Appendices

- Appendix 1: Short CV of the author.
- Appendix 2: GPS data and description of recorded fossil sites.
- Appendix 3: Site Sensitivity Verification Report.
- Appendix 4: Chance Fossil Finds Protocol
- Appendix 5: Terms of Reference

List of Abbreviations

amsl	above mean sea level
DFFE	Department of Forestry, Fisheries and the Environment
ECO	Environmental Control Officer
EMPr	Environmental Management Programme
ESO	Environmental Site Officer
HWC	Heritage Western Cape
Ma	millions of years ago
PIA	palaeontological heritage impact assessment
SAHRA	South African Heritage Resources Agency
SAHRIS	South African Heritage Resources Information System

SiVEST SA (PTY) LTD

PROPOSED CONSTRUCTION OF THE KOUP 2 WIND ENERGY FACILITY AND ASSOCIATED GRID INFRASTRUCTURE, NEAR BEAUFORT WEST, WESTERN CAPE PROVINCE, SOUTH AFRICA

PALAEONTOLOGICAL HERITAGE REPORT

1. INTRODUCTION

Genesis Enertrag Koup 2 Wind (Pty) Ltd (hereafter referred to as “Genesis”), has appointed SiVEST Environmental (hereafter referred to as “SiVEST”) to undertake the required EIA / BA Processes for the proposed construction of the Koup 2 Wind Energy Facility (WEF) and associated grid connection infrastructure near Beaufort West in the Western Cape Province of South Africa.

The overall objective of the development is to generate electricity by means of renewable energy technology capturing wind energy to feed into the National Grid.

It is anticipated that the proposed Koup 2 WEF will comprise thirty-two (32) wind turbines with a maximum total energy generation capacity of up to approximately 140MW. The electricity generated by the proposed WEF development will be fed into the national grid via a 132kV overhead power line. A Battery Energy Storage System (BESS) will be located next to the onsite 33/132kV substation. The storage capacity and type of technology would be determined at a later stage during the development phase, but most likely will comprise an array of containers, outdoor cabinets and/or storage tanks.

In terms of the Environmental Impact Assessment (EIA) Regulations, which were published on 04 December 2014 [GNR 982, 983, 984 and 985] and amended on 07 April 2017 [promulgated in Government Gazette 40772 and Government Notice (GN) R326, R327, R325 and R324 on 7 April 2017], various aspects of the proposed development are considered listed activities under GNR 327 and GNR 324 which may have an impact on the environment and therefore require authorisation from the National Competent Authority (CA), namely the Department of Environment, Forestry and Fisheries (DEFF), prior to the commencement of such activities. Specialist studies have been commissioned to assess and verify the project under the new Gazetted specialist protocols.

1.1 Terms of Reference

The present combined desktop and field-based PIA report assesses potential impacts to palaeontological heritage resources that may result from the proposed Koup 2 WEF and its associated grid connection. It will contribute to the over-arching Heritage Impact Assessments, co-ordinated by PGS Heritage and SiVEST

Environmental Division, as part of the Basic Assessment / Environmental Impact Assessment processes for these developments as well as to the relevant EMPs.

Please see Appendix 4 for the SiVEST Terms of Reference applicable to this report.

1.2 Specialist Credentials

The author, Dr John Almond, is a specialist palaeontologist who has over 40 years of experience in palaeontological research and teaching in Europe, South Africa and elsewhere. He also has more than 20 years of experience in the palaeontological heritage impact assessment world in the RSA and has been involved with numerous PIAs in the Karoo region and elsewhere (Please see Appendix 1 for a short Specialist CV).

1.3 Assessment Methodology

1.3.1 Information sources

The desktop and field-based palaeontological heritage study of the Koup 2 WEF and grid connection project areas was based on the following information resources:

1. A detailed project outline, kmz files, screening report and maps provided by SiVEST Environmental Division and PGS Heritage;

2. A desktop review of:

(a) the relevant 1:50 000 scale topographic map (3222 CD Daskop) and the 1:250 000 scale topographic map 3222 Beaufort West,

(b) Google Earth© satellite imagery,

(c) published geological and palaeontological literature, including 1:250 000 geological maps (3222 Beaufort West) and relevant sheet explanations (Johnson & Keyser 1979) as well as

(d) several previous and on-going fossil heritage (PIA) assessments in the Great Karoo region to the south of Beaufort West by the author (e.g. Almond 2018, 2020a, 2020b, 2021a, 2021b & 2021c *plus* various earlier studies on renewable energy projects in the region listed in the References as well as on-going field-based palaeontological heritage studies for the proposed Kwagga 1, Kwagga 2 & Kwagga 3 WEFs and the proposed Heuweltjies and Kraaltjies WEFs, all of which are situated in the southern Great Karoo shortly to the east and southeast of the present project area);

3. The author's field experience with the formations concerned and their palaeontological heritage (*cf* Almond & Pether 2008 and PIA reports listed in the References); and

4. A five-day field assessment of the combined Koup 1 and Koup 2 WEF project area, including portions of all land parcels involved, by the author and an experienced field assistant (Ms Madelon Tusenius, *Natura Viva* cc), during the period 14 to 19 November 2020. A local farm lessee, Ms Marietjie Mostert (Farm Bloemendal), kindly shared her knowledge of local fossil occurrences. Accessible sectors of the Grid

Connection project area within the combined WEF project areas were surveyed in part, but *not* those sectors lying *outside* the WEF project area itself.

The season in which the site visit took place has no critical bearing on the palaeontological study, although palaeontological fieldwork in the Karoo winter was somewhat hampered by shorter days, occasional rain and low-angle light, making fossils more difficult to discern and to photograph effectively.

1.3.2 Study approach

In preparing a palaeontological desktop study the potentially fossiliferous rock units (groups, formations, members *etc.*) represented within the study area are determined from geological maps and satellite images. The known fossil heritage within each rock unit is inventoried from the published scientific literature, previous palaeontological impact studies in the same region, and the author's field experience (consultation with professional colleagues as well as examination of institutional fossil collections may play a role here, or later following scoping during the compilation of the final report). This data is then used to assess the palaeontological sensitivity of each rock unit to development (provisional tabulations of palaeontological sensitivity of all formations in the Western Cape have already been compiled by J. Almond and colleagues; *e.g.* Almond & Pether 2008) and are shown on the palaeosensitivity map on the SAHRIS (South African Heritage Resources Information System) website. The likely impact of the development on local fossil heritage is then determined on the basis of (1) the palaeontological sensitivity of the rock units concerned and (2) the nature and scale of the development itself, most notably the extent of fresh bedrock excavation and ground clearance envisaged. When rock units of moderate to high palaeontological sensitivity are present within the development footprint, a field assessment study by a professional palaeontologist is usually warranted.

The focus of palaeontological field assessment is *not* simply to survey the development footprint or even the development area as a whole (*e.g.* farms or other parcels of land concerned in the development). Rather, the palaeontologist seeks to assess or predict the diversity, density and distribution of fossils within and beneath the study area, as well as their heritage or scientific interest. This is primarily achieved through a careful field examination of one or more *representative* exposures of all the sedimentary rock units present (*N.B.* Metamorphic and igneous rocks rarely contain fossils). The best rock exposures are generally those that are easily accessible, extensive, fresh (*i.e.* unweathered) and include a large fraction of the stratigraphic unit concerned (*e.g.* formation). These exposures may be natural or artificial and include, for example, rocky outcrops in stream or river banks, cliffs, quarries, dams, dongas, open building excavations or road and railway cuttings. Consolidated as well as uncemented superficial deposits, such as alluvium, scree or wind-blown sands, may occasionally contain fossils and should also be included in the field study where they are well-represented in the study area. It is occasional practice for impact palaeontologists to collect representative, well-localised (*e.g.* GPS and stratigraphic data) samples of fossil material during field assessment studies. In order to do so, a fossil collection permit from Heritage Western Cape (HWC) is required and all fossil material collected must be properly curated within an approved repository (usually a museum or university collection).

Note that while fossil localities recorded during field work within the study area itself are obviously highly relevant, most fossil heritage here is embedded within rocks beneath the land surface or obscured by surface deposits (soil, alluvium, *etc.*) and by vegetation cover. In many cases where levels of fresh (*i.e.*

unweathered) bedrock exposure are low, the hidden fossil resources have to be *inferred* from palaeontological observations made from better exposures of the same formations elsewhere in the region but outside the immediate study area. Therefore a palaeontologist might reasonably spend far *more* time examining road cuts and borrow pits close to, but outside, the study area / project footprint than within the study area / project footprint itself. Field data from localities even further afield (e.g. an adjacent province) may also be adduced to build up a realistic picture of the likely fossil heritage within the study area.

Given 1) the large project areas concerned with the Koup 2 WEF project (approx. 2477.408 ha) and (2) the extensive bedrock exposure in this region of the Great Karoo, the palaeontological heritage field study largely entailed the examination of selected potentially fossiliferous sites with good Beaufort Group mudrock exposure – especially along drainage lines as well as gentler hillslopes and erosion gullies. Since previous field experience shows that in the lower part of the Beaufort Group outcrop area important fossil sites may also occur in association with crevasse splay and channel sandstones, a representative selection of such sites as well as good sections through Late Caenozoic alluvial deposits were also examined. It is emphasised that it is simply *not* practicable to record all, or even a major portion, of fossil sites within such a large area within the course of a few days' fieldwork, and that the occurrence of fossils at surface in the Great Karoo has a large element of unpredictability. Several fossil sites were discovered simply by chance. It is therefore inevitable that the recent site visit can only hope to locate a *representative subsample* of surface fossil sites present within the WEF project areas. The absence of recorded sites within an area does *not* therefore mean that palaeontologically significant material is not present there, either on or beneath the ground surface.

1.3.3 Legislative context

The present combined desktop and field-based palaeontological heritage report falls under Sections 35 and 38 (Heritage Resources Management) of the South African Heritage Resources Act (Act No. 25 of 1999), and it will also inform the EMPr for this project.

The various categories of heritage resources recognised as part of the National Estate in Section 3 of the National Heritage Resources Act include, among others:

- geological sites of scientific or cultural importance;
- palaeontological sites;
- palaeontological objects and material, meteorites and rare geological specimens.

According to Section 35 of the National Heritage Resources Act, dealing with archaeology, palaeontology and meteorites:

(1) The protection of archaeological and palaeontological sites and material and meteorites is the responsibility of a provincial heritage resources authority.

(2) All archaeological objects, palaeontological material and meteorites are the property of the State.

(3) Any person who discovers archaeological or palaeontological objects or material or a meteorite in the course of development or agricultural activity must immediately report the find to the responsible heritage resources authority, or to the nearest local authority offices or museum, which must immediately notify such heritage resources authority.

(4) No person may, without a permit issued by the responsible heritage resources authority—

- (a) destroy, damage, excavate, alter, deface or otherwise disturb any archaeological or palaeontological site or any meteorite;
 - (b) destroy, damage, excavate, remove from its original position, collect or own any archaeological or palaeontological material or object or any meteorite;
 - (c) trade in, sell for private gain, export or attempt to export from the Republic any category of archaeological or palaeontological material or object, or any meteorite; or
 - (d) bring onto or use at an archaeological or palaeontological site any excavation equipment or any equipment which assist in the detection or recovery of metals or archaeological and palaeontological material or objects, or use such equipment for the recovery of meteorites.
- (5) When the responsible heritage resources authority has reasonable cause to believe that any activity or development which will destroy, damage or alter any archaeological or palaeontological site is under way, and where no application for a permit has been submitted and no heritage resources management procedure in terms of section 38 has been followed, it may—
- (a) serve on the owner or occupier of the site or on the person undertaking such development an order for the development to cease immediately for such period as is specified in the order;
 - (b) carry out an investigation for the purpose of obtaining information on whether or not an archaeological or palaeontological site exists and whether mitigation is necessary;
 - (c) if mitigation is deemed by the heritage resources authority to be necessary, assist the person on whom the order has been served under paragraph (a) to apply for a permit as required in subsection (4); and
 - (d) recover the costs of such investigation from the owner or occupier of the land on which it is believed an archaeological or palaeontological site is located or from the person proposing to undertake the development if no application for a permit is received within two weeks of the order being served.

Minimum standards for the palaeontological component of heritage impact assessment reports (PIAs) have been published by SAHRA (2013) and by Heritage Western Cape (2021).

2. ASSUMPTIONS AND LIMITATIONS

The accuracy and reliability of palaeontological specialist studies as components of heritage impact assessments are generally limited by the following constraints:

1. Inadequate database for fossil heritage for much of the RSA, given the large size of the country and the small number of professional palaeontologists carrying out fieldwork here. Most development study areas have never been surveyed by a palaeontologist.
2. Variable accuracy of geological maps which underpin these desktop studies. For large areas of terrain these maps are largely based on aerial photographs alone, without ground-truthing. The maps generally depict only significant (“mappable”) bedrock units as well as major areas of superficial “drift” deposits (alluvium, colluvium) but for most regions give little or no idea of the level of bedrock outcrop, depth of superficial cover (soil *etc*), degree of bedrock weathering or levels of small-scale tectonic deformation, such as cleavage. All of these factors may have a major influence on the impact significance of a given development on fossil heritage and can only be reliably assessed in the field.

3. Inadequate sheet explanations for geological maps, with little or no attention paid to palaeontological issues in many cases, including poor locality information.
4. The extensive relevant palaeontological “grey literature” - in the form of unpublished university theses, impact studies and other reports (e.g. of commercial mining companies) - that is not readily available for desktop studies.
5. Absence of a comprehensive computerised database of fossil collections in major RSA institutions which can be consulted for impact studies.

In the case of palaeontological desktop studies without supporting Phase 1 field assessments these limitations may variously lead to either:

- a) *underestimation* of the palaeontological significance of a given study area due to ignorance of significant recorded or unrecorded fossils preserved there, or
- b) *overestimation* of the palaeontological sensitivity of a study area, for example when originally rich fossil assemblages inferred from geological maps have in fact been destroyed by tectonism or weathering, or are buried beneath a thick mantle of unfossiliferous “drift” (soil, alluvium etc).

Since most areas of the RSA have not been studied palaeontologically, a palaeontological desktop study usually entails *inferring* the presence of buried fossil heritage within the study area from relevant fossil data collected from similar or the same rock units elsewhere, sometimes at localities far away. Where substantial exposures of bedrocks or potentially fossiliferous superficial sediments are present in the study area, the reliability of a palaeontological impact assessment may be significantly enhanced through field assessment by a professional palaeontologist, as in the case of the present study.

In the case of the combined Koup 1 WEF and Koup 2 WEF project area bedrock exposure is often remarkably good in highly-dissected, hilly regions but is highly constrained by extensive superficial deposits in areas of low relief (e.g. NE sector of Koup 1 WEF project area), as well as, to a lesser extent, by shrubby vegetation. The project area is very extensive (> 4000 ha) and with remarkably few access roads, probably because much of the area is not currently being farmed at present. Unavoidably, only a small fraction of the entire project area could be surveyed on foot within the time available (5 days). Short days, low angle light and occasional rainy weather in winter further constrained the field survey.

Nevertheless, sufficient (c. 150-200) bedrock exposures – including many of excellent quality - were examined during the course of the five-day field study to assess the palaeontological heritage sensitivity of the main rock units represented within the combined Koup 1 and Koup 2 WEF and grid connection study area (See satellite image Fig. A2.1 as well as palaeontological data table in Appendix 2). Since access permission for sectors of the grid connection project area lying outside the combined WEF project area was not available at the time of the palaeontological field survey, these sectors are only treated at a desktop level in the present report.

Comparatively few academic palaeontological studies or palaeontological impact assessments have been carried-out hitherto in this region of the Great Karoo, so any new data from impact studies here are of

scientific interest. Confidence levels for this impact assessment are rated as *medium*, despite the unavoidable constraints of limited time and access in the project area.

3. TECHNICAL DESCRIPTION

3.1 Project Location

The proposed WEF and associated grid connection infrastructure is located approximately 55km south of Beaufort West in the Western Cape Province and is within the Beaufort West and Prince Albert Local Municipalities, in the Central Karoo District Municipality (**Figure 1**).




Figure 1: Regional Context Map for the proposed Koup 2 WEF south of Beaufort West.

3.1.1 WEF

The WEF application site as shown on the locality map below (**Figure 2**) is approximately 2477.408 hectares (ha) in extent and incorporates the following farm portions:

- Portion 1 of the farm Kaffirs Kraal No 380
- Portion 8 of the Kaffirs Kraal No 380

A smaller buildable area (1575.2.114ha) has however been identified as a result of a preliminary suitability assessment undertaken by Genesis and this area is likely to be further refined with the exclusion of sensitive areas determined through various specialist studies being conducted as part of the EIA process.




Figure 2: Koups 2 WEF Site Locality

3.1.2 Grid Connection

At this stage, it is proposed that the 132kV power lines will connect the Koups 2 WEF on-site substation to the national grid via the proposed Koups 1 collector substation, located on the Koups 1 WEF project site. Two grid connection route options were originally assessed (Figure 3a) of which Grid Option 1 has now been chosen (Figure 3b).




Figure 3a: Two 132kV Power Line Route Alignments originally assessed to connect the Koups 2 WEF on-site substation to the national grid via the proposed Koups 1 collector substation.




Figure 4b: The chosen 132kV Power Line Route Alignment connecting the Koups 2 WEF on-site substation to the national grid via the proposed Koups 1 collector substation.

3.2 Project Description

It is anticipated that the proposed Koup 2 WEF will comprise thirty-two (32) wind turbines with a maximum total energy generation capacity of up to approximately 140MW. The electricity generated by the proposed WEF development will be fed into the national grid via a 132kV overhead power line. The 132kV overhead power line will however require a separate EA and is subject to a Basic Assessment (BA) process, which is currently being undertaken in parallel to the EIA process. The proposed Koup 2 WEF will include the following components:

3.2.1 Wind Farm Components

- Up to 32 wind turbines, each between 5.6MW and 6.6MW, with a maximum export capacity of approximately 140MW. This will be subject to allowable limits in terms of the Renewable Energy Independent Power Producer Procurement Programme (REIPPPP). The final number of turbines and layout of the WEF will, however, be dependent on the outcome of the Specialist Studies conducted during the EIA process;
- Each wind turbine will have a hub height and rotor diameter of up to approximately 200m;
- Permanent compacted hardstanding areas / platforms (also known as crane pads) of approximately 90m x 50m (total footprint of approx. 4 500m²) per turbine during construction and for on-going maintenance purposes for the lifetime of the proposed development;
- Each wind turbine will consist of a foundation of up to approximately 15m x 15m in diameter. In addition, the foundations will be up to approximately 3m in depth;
- Electrical transformers adjacent to each wind turbine (typical footprint of up to approximately 2m x 2m) to step up the voltage to 33kV;
- One (1) new 33/132kV on-site substation and/or combined collector substation, occupying an area of approximately 1.5 ha . The proposed substation will be a step-up substation and will include an Eskom portion and an IPP portion, hence the substation has been included in the WEF EIA and in the grid infrastructure BA (substation and 132kV overhead power line) to allow for handover to Eskom. Following construction, the substation will be owned and managed by Eskom. The current applicant will retain control of the low voltage components (*i.e.* 33kV components) of the substation, while the high voltage components (*i.e.* 132kV components) of this substation will likely be ceded to Eskom shortly after the completion of construction.
- The wind turbines will be connected to the proposed substation via medium voltage (33kV) cables. Cables will be buried along access roads wherever technically feasible.
- A Battery Energy Storage System (BESS) will be located next to the onsite 33/132kV substation. The storage capacity and type of technology would be determined at a later stage during the development phase, but most likely will comprise an array of containers, outdoor cabinets and/or storage tanks;
- Internal roads with a width of between 8m and 10m will provide access to each wind turbine. Existing site roads will be used wherever possible, although new site roads will be constructed where necessary. Turns will have a radius of up to 50m for abnormal loads (especially turbine blades) to access the

various wind turbine positions. It should be noted that the proposed application site will be accessed via an existing gravel road from the N12 National Route;

- One (1) construction laydown / staging area of up to approximately 2.2ha. It should be noted that no construction camps will be required in order to house workers overnight as all workers will be accommodated in the nearby town;
- One (1) permanent Operation and Maintenance (O&M) building, including an on-site spares storage building, a workshop and an operations building to be located on the site identified for the construction laydown area.
- A wind measuring lattice (approximately 120m in height) mast has already been strategically placed within the wind farm application site in order to collect data on wind conditions;
- No new fencing is envisaged at this stage. Current fencing is standard farm fence approximately 1-1.5m in height. Fencing might be upgraded (if required) to be up to approximately 2m in height; and
- Water will either be sourced from existing boreholes located within the application site or will be trucked in, should the boreholes located within the application site be limited.

3.2.2 Grid Components

The proposed grid connection infrastructure to serve the Koup 2 WEF will include the following components:

- One (1) new 33/132kV on-site substation and/or collector substation, occupying an area of up to approximately 1ha. The proposed substation will be a step-up substation and will include an Eskom portion and an IPP portion, hence the substation has been included in both the EIA for the WEF and in the BA for the grid infrastructure to allow for handover to Eskom. The applicant will remain in control of the low voltage components (*i.e.* 33kV components) of the substation, while the high voltage components (*i.e.* 132kV components) of this substation will likely be ceded to Eskom shortly after the completion of construction; and
- One (1) new 132kV overhead power line connecting the on-site and/or collector substation *via* the proposed Koup 1 collector substation and thereby feeding the electricity into the national grid. Power line towers being considered for this development include self-supporting suspension monopole structures for relatively straight sections of the line and angle strain towers where the route alignment bends to a significant degree. Maximum tower height is expected to be approximately 25m.

3.3 Layout alternatives

3.3.1 Wind Energy Facility

Design and layout alternatives have been considered and assessed as part of the EIA. These include alternatives for the Substation locations and also for the construction / laydown area. The proposed site alternatives are shown in **Figure 4a** below while the final proposed layout is shown in **Figure 4b**.




Figure 5a: Alternatives originally proposed as part of the Koups 2 WEF




Figure 6b: Final layout proposed for the Koups 2 WEF

3.3.2 Grid Components

The grid connection infrastructure proposals originally included two (2) switching and collector substation site alternatives and two (2) power line route alignment alternatives (**Figure 3a**). These alternatives have since been considered and assessed as part of the BA process and will be amended or refined to avoid identified environmental sensitivities.

All two (2) power line route alignments were assessed within 600m and 300m wide assessment corridor (150m on either side of power line). These alternatives are described below:

- Power Line Corridor Option 1 is approximately 12km in length, linking either substation / collector Option 1 or Option 2 to the proposed Koup 1 Collector Option 1 or Option 2. This route alignment will be assessed within a 600m wide corridor (300m on either side of the power line).
- Power Line Corridor Option 2 is approximately 13.2km in length, linking either substation / collector Option 1 or Option 2 to the proposed Koup 1 Collector Option 1 or Option 2. This route alignment will be assessed within a 300m wide corridor (150m on either side of the power line).

As shown in **Figure 3b**, Power Line Corridor Option 1 has now been chosen.

3.3.3 No-go Alternative

The 'no-go' alternative is the option of not undertaking the proposed WEF and / or grid connection infrastructure projects. Hence, if the 'no-go' option is implemented, there would be no development. This alternative would result in no environmental impacts from the proposed project on the site or surrounding local area. It provides the baseline against which other alternatives are compared and will be considered throughout the report.

4. LEGAL REQUIREMENT AND GUIDELINES

4.1 Legislative context

The present combined desktop and field-based palaeontological heritage report falls under Sections 35 and 38 (Heritage Resources Management) of the South African Heritage Resources Act (Act No. 25 of 1999), and it will also inform the EMP for this project.

The various categories of heritage resources recognised as part of the National Estate in Section 3 of the National Heritage Resources Act include, among others:

- geological sites of scientific or cultural importance;
- palaeontological sites;
- palaeontological objects and material, meteorites and rare geological specimens.

According to Section 35 of the National Heritage Resources Act, dealing with archaeology, palaeontology and meteorites:

- (1) The protection of archaeological and palaeontological sites and material and meteorites is the responsibility of a provincial heritage resources authority.
- (2) All archaeological objects, palaeontological material and meteorites are the property of the State.
- (3) Any person who discovers archaeological or palaeontological objects or material or a meteorite in the course of development or agricultural activity must immediately report the find to the responsible heritage resources authority, or to the nearest local authority offices or museum, which must immediately notify such heritage resources authority.
- (4) No person may, without a permit issued by the responsible heritage resources authority—
 - (e) destroy, damage, excavate, alter, deface or otherwise disturb any archaeological or palaeontological site or any meteorite;
 - (f) destroy, damage, excavate, remove from its original position, collect or own any archaeological or palaeontological material or object or any meteorite;
 - (g) trade in, sell for private gain, export or attempt to export from the Republic any category of archaeological or palaeontological material or object, or any meteorite; or
 - (h) bring onto or use at an archaeological or palaeontological site any excavation equipment or any equipment which assist in the detection or recovery of metals or archaeological and palaeontological material or objects, or use such equipment for the recovery of meteorites.
- (5) When the responsible heritage resources authority has reasonable cause to believe that any activity or development which will destroy, damage or alter any archaeological or palaeontological site is under way, and where no application for a permit has been submitted and no heritage resources management procedure in terms of section 38 has been followed, it may—
 - (e) serve on the owner or occupier of the site or on the person undertaking such development an order for the development to cease immediately for such period as is specified in the order;
 - (f) carry out an investigation for the purpose of obtaining information on whether or not an archaeological or palaeontological site exists and whether mitigation is necessary;
 - (g) if mitigation is deemed by the heritage resources authority to be necessary, assist the person on whom the order has been served under paragraph (a) to apply for a permit as required in subsection (4); and
 - (h) recover the costs of such investigation from the owner or occupier of the land on which it is believed an archaeological or palaeontological site is located or from the person proposing to undertake the development if no application for a permit is received within two weeks of the order being served.

Where Preconstruction of Construction Phase mitigation, comprising palaeontological recording and collection of fossil material and associated geological data, is required as a condition of Environmental Authorization (as here), this must be carried out by a suitably qualified professional palaeontologist under a Fossil Collection Permit issued by the relevant Heritage Resources Management Agency (in the present case, Heritage Western Cape, Cape Town). The fossil material collected must be curated in an approved repository (museum / university collection). Standards for palaeontological reporting and mitigation in the RSA have been established by Heritage Western Cape (2016, 2021) and SAHRA (2013). A tabulated Chance Fossil Finds Protocol which must be implemented throughout the Construction Phase of the WEF projects is provided in Appendix 4 to this report.

5. DESCRIPTION OF THE RECEIVING ENVIRONMENT

This section of the PIA report presents a short, illustrated overview of the geology and palaeontological heritage encountered within the Koup 2 WEF project area, including the associated grid connection project area. It also draws on geological and palaeontological observations from the adjoining Koup 1 WEF project area.

5.1 Geological context

A short outline of the geology of the Koup 2 WEF project area (including the associated grid connection project areas) is provided in this section of the report as context for the palaeontological heritage data discussed in the following subsection.

The Koup 2 WEF project area lies within Beaufort West District of the Western Cape (Fig. 1). The project area is situated on the western side of the N12 trunk road between Beaufort West and Oudtshoorn within semi-arid, low-relief to highly-dissected hilly terrain towards the southern margins of the Great Karoo region *sensu stricto*. The eastern margins of the combined Koup 1 and Koup 2 project area (including much of grid line project area) lie on the margins of the Aberdeen *Vlakte*, an ancient peneplanated land surface of possible Miocene age (Partridge & Maud 1987). Relief here is generally low, with gentle hillslopes largely mantled with colluvium (scree, hillwash). Elevations are around 1000 to 1100 m amsl. in this region which forms a watershed between west- and east-flowing drainage systems. Bedrock exposure here is localized and often very poor due to the pervasive mantle of Late Caenozoic superficial deposits such as alluvium, eluvium, sheetwash deposits and skeletal soils.

Further towards the west (Koup 2 WEF project area) the terrain is more dissected, hillslopes are steeper and bedrock exposure levels are much higher, occasionally superb by southern Karoo standards (See Frontispiece, **Figure 9** & **Figure 10**). This applies even to the more readily weathered and eroded Beaufort Group mudrock facies. Several of the ridges and peaks here are named and, given their concordance, most of them are probably erosional outliers of the Aberdeen *Vlakte* surface which is more extensively preserved further east. The highest elevation is Wolwekop (1022 m amsl) on southern edge of the combined WEF project area with isolated lower peaks to the north such as Turksvykop (950 m), Gouwermentskop (984), Daskop (962), Platkop (908 m), and Syfeerfonteinkop (910 m). Drainage is largely *via* intermittent-flowing (non-perennial) water courses. It flows mainly to the N and NW into the major, ancient Gamka River drainage system *via* small tributaries of the Veldmansrivier such as the Kareerivier, Platdoringrivier, Pieterskraalrivier, Diepkloof and Houtbosrivier but there are also minor streams running to the west into the Gatsrivier.

As is clear from satellite images (e.g. **Figure 61**) as well as the geological map (**Figure 7**), the WEF project area is characterized by broadly W-E trending ridges and intervening narrow rocky *vlaktes*. The topography here strongly reflects the pronounced folding of the Beaufort Group bedrocks along E-W fold axes within the northern margins of the Permo-Triassic Cape Fold Belt (CFB). Bedding dips are nevertheless moderate to low for the most part, reaching up to 45° on some fold limbs (dips up to 20° are marked on the geological map). Numerous small-scale faults cut the Beaufort Group bedrocks in this region, as best seen in road cuttings along the N12. Low-angle reverse / thrust faults as well as some steeper normal faults are marked by well-developed quartz mineral lineation, thick, laterally-persistent quartz veins, milky quartz rubble and slickensides. A well-developed, almost pervasive axial-planar cleavage affects most of the Beaufort Group bedrocks, expressed as a spaced cleavage in sandstone facies and many siltstones as well as fine slaty or pencil cleavage in the darker, finest-grain mudrocks. The CFB folds in this sector of the southern Karoo

appear to be narrower and more tightly spaced compared with regions immediately to the N and S; hence the higher levels of tectonic deformation encountered here.

The geology of the combined Koup 1 and Koup 2 WEF project areas is covered by 1: 250 000 geology sheet 3222 Beaufort West (Council for Geoscience, Pretoria; Johnson & Keyser 1979) (**Figure 7**). The greater part of the lower-lying terrain here is underlain by Middle Permian continental sediments of the **Abrahamskraal Formation** (Lower Beaufort Group / Adelaide Subgroup, Karoo Supergroup) (Pa, pale green in **Figure 7**) (Johnson & Keyser 1979, Johnson *et al.* 2006). It is likely the majority of the bedrocks here can be largely or entirely assigned to the mudrock-dominated **Karelskraal Member** situated at the top of the very thick Abrahamskraal Formation succession but this requires confirmation from detailed field mapping that is beyond the scope of the present PIA study. The broadly west-east trending ridges and associated *koppies* located within the WEF project area, especially towards its southern and northern margins, are built of the conformably overlying, sandstone-rich **Poortjie Member** which lies at the base of the **Teekloof Formation** (Adelaide Subgroup) (Pt, dark green in **Figure 7**). The sedimentology of the Abrahamskraal – Teekloof transition has been addressed recently by Paiva (2015). Early Jurassic intrusions of the **Karoo Dolerite Suite** are not mapped within the project area but do occur closer to the Great Escarpment at Beaufort West.




Figure 7: Extract from 1: 250 000 geology sheet 3222 Beaufort West showing the boundaries of the combined Koup 1 and Koup 2 WEF project area to the south of Beaufort West (yellow polygons). Note numerous W-E trending fold axes in the region which falls within the northern margins of the Cape Fold Belt. Pa (pale green) = Abrahamskraal Formation (Adelaide Subgroup, Lower Beaufort Group). Pt (dark green) = Teekloof Formation (Adelaide Subgroup, Lower Beaufort Group). Yellow = Late Caenozoic / Quaternary superficial sediments, including alluvium, sheet wash, colluvium, soils, locally cemented by pedocretes such as calcrete. To the west of the N12 and largely outside the WEF

project area triangular symbols indicate fossil localities within the *Pristerognathus* Assemblage Zone (*N.B.* This fossil biozone data is now outdated – see updated stratigraphic chart presented below). A single fossil site for the underlying *Tapinocephalus* Assemblage zone (star symbol) is indicated c. 4 km to the NW of the combined WEF project area.




Figure 8: Stratigraphic subdivision of the Karoo Supergroup with the rock units and fossil biozones most relevant to the present PIA study outlined in green (Modified from Smith et al. 2020). In the combined Koup WEF project area fossil assemblages within the uppermost Abrahamskraal Formation (Karelskraal Member) and lower part of the Poortjie Member of the Teekloof Formation are now assigned to the *Diictodon-Styracocephalus* Assemblage Zone.

The **Abrahamskraal Formation** (Pa in Figure 7) is a very thick (c. 2.4 km) succession of fluvial deposits laid down in the Main Karoo Basin by meandering rivers on an extensive, low-relief floodplain during the Middle Permian Period, some 268-261 million years ago (Rossouw & De Villiers 1952, Johnson & Keyser 1979, Turner 1981, Theron 1983, Smith 1979, 1980, 1990, 1993a, 1993b, Smith & Keyser 1995a, Looek *et al.*, 1994, McCarthy & Rubidge 2005, Johnson *et al.*, 2006, Wilson *et al.* 2014, Cole *et al.* 2016). These sediments include (a) lenticular to sheet-like channel sandstones, often associated with thin, impersistent intraformational breccio-conglomerates (larger clasts mainly of reworked mudflakes, calcrete nodules, *plus* sparse rolled bones, teeth, petrified wood), (b) well-bedded to laminated, grey-green to purple-brown floodplain mudrocks with common pedocrete horizons (calcrete nodules formed in ancient soils), (c) thin, sheet-like crevasse-splay sandstones, as well as more (d) localized playa lake deposits (e.g. wave-rippled sandstones, laminated mudrocks, limestones, evaporites) (Figure 11 to Figure 17. Figure 18 & Figure 19). A number of yellowish-green to reddish-weathering, silica-rich “chert” horizons are also found. Some of these appear to be secondarily silicified mudrocks or limestones of possible lacustrine origin but at least some contain high levels of reworked volcanic ash (tuffs and tuffites). A wide range of sedimentological and

palaeontological observations point to deposition under seasonally arid climates. These include, for example, the abundance of calcretes and evaporites (silicified gypsum pseudomorphs or “desert roses” *cf* Keyser 1968), reddened mudrocks, sun-cracked muds, “flashy” river systems, sun-cracked fossil bones, well-developed seasonal growth rings in fossil wood, rarity of fauna, and little evidence for substantial bioturbation or vegetation cover (*e.g.* root casts) on floodplains away from the river banks.

The Karelskraal Member is generally well-exposed in many sectors of the Koup WEF project area. This applies especially to the mudrock facies which elsewhere are usually poorly-exposed due to sandstone colluvium from the overlying Poortjie Member. Lower down the Karelskraal succession includes several thin channel sandstone / wacke packages of variable lithofacies. They tend to be thin, tabular, thin-bedded, fine-grained, well-sorted, and greyish to grey-green or speckled but occasional friable “golden yellow” sandstones more typical of the Poortjie member are found, especially within the upper part of the Karelskraal succession. Breccio-conglomerates are generally not well developed but locally thin mudflake intraclast breccias do occur along erosional basal contacts. Intervening mudrock packages vary from thick and massive to thick-bedded or thin-bedded. The majority are grey to grey-green but subordinate purple-brown or mottled mudrocks occur as well. They are intercalated with thin crevasse-splay sandstones with current ripple cross-laminated tops and flat to locally loaded bases.

Well-developed calcrete palaeosol (pedocrete) horizons are common, especially lower down in the Karelskraal succession. They are often a few dm thick, pale brown, cobbly to lenticular, and blocky-weathering. The more fossiliferous palaeosol horizons tend to be thinner, laterally persistent with greyish-green to rusty-brown, sphaeroidal to irregular-shaped concretions and are pale grey (micritic) to brownish internally. Prominent-weathering, laterally-extensive rusty-brown, ferruginous carbonate lenses or cobble- to boulder-sized sphaeroidal concretions occur at intervals, perhaps reflecting intervals of high water tables. Well-developed lenses and horizons of dark brown *koffieklip* up to several dm thick are also encountered, in at least one case directly overlying a tuffite bed.

Mudrock-dominated distal floodplain to lacustrine packages appear to predominate within the uppermost Karelskraal Member succession. The mudrocks are usually dark grey to purple-grey or blue-grey, massive to thin bedded or occasionally laminated and locally pencil cleaved. Sedimentological features suggesting periods of high water tables alternating with intermittent aridification within this interval include horizons with large ferruginous carbonate concretions, loading and boudinage of thin crevasse-splay sandstones, locally abundant (and often large large) stellate gypsum pseudomorphs as well as thin, blocky-weathering, greenish tuffites, occasionally showing ripple cross-lamination and wave rippled bed tops. Rectilinear, cross-cutting networks of desiccation cracks infilled with gypsiferous sandstone admixed with mudflakes are common within darker, fine-grained mudrocks (Possible neptunian dykes or dewatering features should also be considered here). However, these beds are typically fossil-poor (See following subsection).

Compared with the underlying rocks, the **Teekloof Formation** (Pt in **Figure 7**) has a generally higher proportion of sandstones and reddish mudrocks are generally more abundant here. Multi-storied sandstones are common in the basal arenaceous **Poortjie Member**, as are thin, impersistent lenses or persistent layers of pinkish “cherts” that are probably altered volcanic ashes or lacustrine limestones (Johnson & Keyser 1979, Smith & Keyser 1995b). Several economically interesting uranium ore deposits occur within the Poortjie Member in association with brown-weathering, ferruginous channel sandstones (*koffieklip*) and transported plant material. Interesting accounts of the sedimentology and palaeontology of the Poortjie Member at the farm Putfontein, some 25 km to the north of the present WEF study area, are given by Stear (1978) as well as by Cole and Smith (2008). The base of the Poortjie Member has recently been dated to 260 Ma on the basis of a white tuff unit 3.5 m above the basal sandstone (Day *et al.* 2015). Several laterally-persistent tuff or tuffitic units are represented within the upper Abrahamskraal – Poortjie Member interval, as well seen in satellite images of the region as well as in the field, in the Koup 1 and 2 WEF project area. They may be of use in future in constraining the end-Guadalupian mass extinction event (*cf* Day *et al.* 2015, McKay *et al.* 2015, Almond 2018, Day & Rubidge 2021) (See palaeontological discussion below).

Within the combined Koup 1 & 2 WEF project area a succession of several (3-4 at least), closely-spaced, moderately thick (c. 5-10 m) packages of channel sandstone build the higher ground along the crests of E-W trending ridges and cap several isolated *koppies*. The Poortjie sandstones are typically “golden-yellow” weathering, flat- to cross-bedded and medium-grained with a high feldspathic component, giving them a friable or biscuit-like texture. However, some well-sorted, tabular bedded, fine-grained packages occur here too. Thin basal breccias with reworked calcrete glaebules occur locally but the lower contacts of the sandstone packages are usually mantled by scree. Thin packages of grey to grey-green, massive to well-bedded silty mudrocks (often highly cleaved) are interleaved between the channel sandstones. They contain horizons of small to large pedogenic calcrete concretions (often ferruginised) and small to large radial gypsum pseudomorphs.

In areas of low topographic relief, such as gentler hillslopes, extensive gravelly to sandy *vlaktes* (*i.e.* plains) as well as in the vicinity of larger water courses, the Beaufort Group bedrocks are mantled by various **Late Caenozoic superficial sediments (Figure 20 & Figure 21)**. For the most part these comprise downwasted or eluvial surface gravels, rubbly sandstone colluvium, silty, sandy and gravelly alluvium (pale yellow areas mapped in **Figure 7**) as well as skeletal soils, with local development along drainage lines of spring deposits such as calcrete. Most of these superficial deposits are unconsolidated and probably of Late Pleistocene to Holocene age (*i.e.* last 2.5 million years) but some alluvium is well-calcretised and might be somewhat older. High Level gravel terraces are not well-developed in the region, implying low levels of stream incision.

The greyish to pale brown, predominantly sandy to fine-gravelly alluvial deposits may reach thicknesses of several meters with finer-grained, massive to well-bedded or locally cross-bedded alluvium overlying basal gravels dominated by angular wacke clasts. Some of these deposits are probably catastrophic flood inundites. Alluvial patches are mapped in the eastern sector of the combined WEF project area in association with the Platdoringrivier and Kaatjies se Loop (**Figure 7**). In the extensive alluvial *vlaktes* in the E and S portions of the project area (N of Kareerivier and E of Platdoorns as well as on Oskloof) shallow streams expose rubbly, cobbly to pebbly alluvial gravels of wacke, tuff, vein quartz and pedogenic calcrete concretions. Finer, often subrounded, downwasted eluvial surface gravels of the same lithologies and modified by sheetwash processes occur at surface. Yellow-hued, blocky sandstone colluvium forms aprons below low *kranzes* of Poortjie sandstone while the crests of higher-lying ridges and crests are usually mantled with rubbly, weathered relictual sandstone deposits and sandy soils. The finer-grained, better-cemented wackes within the Abrahamskraal Formation show well-developed corestone weathering, generating distinctive rounded, orange-patinated wacke cobbles that are often anthropogenically flaked. Sheets of yellow-green to bright orange weathering, blocky gravels of tuffite are also characteristic of the Karelskraal – Poortjie stratigraphic interval and, like the milky vein quartz gravels associated with faults lines, can often be picked out on satellite imagery.



Figure 9: View towards a low ridge of Poortjie Member sandstones along the southern margins of the Koup 2 WEF project area (Farm 11/380). The lower terrain in the foreground is underlain by the upper Abrahamskraal Formation.



Figure 10: Typical dissected, hilly terrain the central portion of the Koup 2 WEF project area (Farm 11/380). Dark grey areas are exposures of Beaufort Group mudrock while the central yellowish to orange patch features weathered tuffite (mixed volcanic ash and terrigenous sediment) gravels.



Figure 11: Intensely gullied lower hillslopes in the Koup 2 WEF project area (Farm 1/380) showing extensive good Karelskraal Member overbank mudrock exposure.



Figure 12: Low, flat-topped koppies in the central portion of the Koup 2 WEF (Farm 1/380, looking to the NW) showing stepped slopes of Abrahamskraal Formation sandstones and thin sandstones and capped by thicker Poortjie Member sandstones.




Figure 13: Outstanding exposures of multi-hued Karelskraal Member mudrocks with intercalated thin, pale crevasse splay sandstones capped by Poortjie Member sandstones, Koup 2 WEF project area (Farm 1/380).




Figure 14: Horizon of large (boulder-sized), spheroidal concretions of rusty-brown ferruginous carbonate within the Karelskraal Member in the Koup 2 WEF project area (Farm 1/380).



Figure 15: Well-developed palaeosol horizon (fossil soil) marked by pale brown, ferruginous calcrete concretions in the Karelskraal Member, Koup 2 WEF project area (Farm 1/380). These horizons are apparently only rarely fossiliferous.



Figure 16: Thin palaeosol horizon (arrowed) marked by cobble-sized, sphaeroidal calcrete concretions with a micritic composition, Karelskraal Member, Koup 2 WEF project area (Farm 1/380). Many fossil vertebrate specimens are associated with these fossil semi-arid climate soils.



Figure 17: Well-developed lens of *koffieklip* (ferruginous carbonate), here associated with an underlying pale yellowish-green tuffite bed of possible lacustrine origin, Karelskraal Member, Koup 2 WEF project area (Farm 8/380).




Figure 18: Boudinage or soft-sediment loading of thin crevasse-splay sandstones overlying dark, water-rich mudrocks of probable lacustrine origin, Karelskraal Member, Koup 2 WEF project area (Farm 8/380) (Hammer = 30 cm).



Figure 19: One of several thin, blocky-weathering tuffite horizons within the Karelskraal – Poortjie stratigraphic interval Koup 2 WEF project area (Farm 8/380) (Hammer = 30 cm). These waterlain volcanic ash layers can be accurately dated using radiometric methods.



Figure 20: Thick, massive to subtly bedded, silty to finely-gravelly Late Caenozoic alluvial deposits overlying coarser basal gravels, stream bank exposure in the Koup 2 WEF project area (Farm 1/380) (Hammer = 30 cm).



Figure 21: Typical orange-hued, well-rounded gravels generated by corestone weathering of fine-grained Karelskraal Member sandstones, Koup 2 WEF project area (Farm 8/380). Flaked stone artefacts manufactured from corestones are frequently found among such deposits.

5.2 Palaeontological heritage

Continental (terrestrial / lacustrine / fluvial) fossil biotas within the upper part of the Abrahamskraal Formation (Moordenaars and Karelskraal Members) as well as within the lowermost portion of the Poortjie Member of the Teekloof Formation are now assigned to the ***Diictodon – Styracocephalus* Subzone** of the revised ***Tapinocephalus* Assemblage Zone** (AZ) that is of Middle Permian age (Late Capitanian, c. 262-260 Ma) (Day & Rubidge 2020) (See stratigraphic column in **Error! Reference source not found.**). These biotas are of special palaeobiological interest in that they reflect the major Late Capitanian or Guadalupian (Late Middle Permian) Mass Extinction Event on land (See biostratigraphic chart in **Figure 24**). The highly impoverished, post-extinction vertebrate fauna represented in the uppermost part of the *Diictodon – Styracocephalus* Subzone (upper Karelskraal Member - lowermost Poortjie Member) includes – or is inferred to include – only a few representatives of several tetrapod subgroups including temnospondyl amphibians, parareptiles (pareiasaurs, *Eunotosaurus*), dinocephalians (e.g. *Criocephalosaurus*, perhaps also *Anteosaurus*, *Titanosuchus*), dicynodonts (e.g. *Diictodon*), therocephalians (e.g. *Pristerognathus*) and gorgonopsians (Retallack *et al* 2006, Smith *et al.* 2012, Day *et al.* 2015a, 2015b, Day & Rubidge 2020, Day & Rubidge 2021, Marchetti *et al.* 2020) (**Figure 24** to **Figure 27**).




Figure 22: Distribution of recorded vertebrate fossil sites within the southern portion of the Main Karoo Basin (modified from Nicolas 2007). The approximate location of the combined Koup WEF project area to the south of Beaufort West (BW) is indicated by the red rectangle. The high density of recorded fossil sites along the N12 here as well as further to the north is notable.




Figure 23: Vertebrate fossil localities within the Lower Beaufort Group in the region south of Beaufort West where the boundary between the Abrahamskraal and Teekloof Formations is highly folded (dotted line). *Tapinocephalus* Assemblage Zone specimens are found in the far south (small open circles) while *Pristerocephalus* Assemblage Zone fossils (black dots) are associated with outcrops of the Poortjie Member (lowermost Teekloof Formation) (Map abstracted from Keyser & Smith 1977-78). The paucity of fossil records east of the N12 tar road reflects the generally poor bedrock exposure in the area (“min dagsome”) compared with more dissected terrain to the west of the road.




Figure 24: Chart showing the ranges of known terrestrial tetrapod genera from the Middle to Late Permian of the Main Karoo Basin (From Day *et al.* 2015b). The boundary between the Abrahamskraal and Teekloof Formations is associated with a catastrophic extinction event at the end of the Capitanian Stage (c. 260 Ma) that has been dated here on the basis of a tuff horizon close to the contact of the Karelskraal and Poortjie Members (yellow star). Key victims of the extinction event were almost all the large-bodied dinocephalians and pareiasaur parareptiles as well as many (but not all) dicynodonts and therocephalians.




Figure 25: Skulls of two key fossil therapsid tetrapods from the upper part of the *Tapinocephalus* Assemblage Zone – the small-bodied dicynodont *Diictodon* (top) and the large-bodied dinocephalian *Styraecocephalus*. Note the very thick cranial roof in the latter, a possible adaptation for head-butting shown by many tapinocephalid dinocephalians.




Figure 26: Skeleton of the tapinocephalid (thick-skulled) dinocephalian *Moschops*, a rhino-sized herbivorous therapsid that reached lengths of 2.5 to 3 m and may have lived in small herds. Postcranial remains (and much rarer fossil skull material) of several dinocephalians have been recorded within and just outside the Koup WEF project area.



Figure 27: Artist's reconstructions of the indescribably ugly late Middle Permian dinocephalians *Styracocephalus* (left) and *Criocephalosaurus* (right). One or both these taxa may occur within or close to the WEF project area.

A number of vertebrate fossil sites of the (now outdated) *Pristerognathus* Assemblage Zone have been within the Poortjie Member both within and (mainly) outside the Koup WEF project area by Keyser and Smith (1977-1978) (**Figure 23**). Many of these are indicated on the published 1: 250 000 scale geological map (triangular symbols on map **Error! Reference source not found.**) while only a few sites for the underlying *Tapinocephalus* Assemblage Zone are recorded here (star-shaped symbols in map **Error! Reference source not found.** 5). The more recent map of Karoo fossil vertebrates published by Nicolas (2007) (**Figure**

22) shows several fossil sites along the N12 but none just to the west where the Koup 1 & 2 WEF project area is located. A number of dinocephalian, dicynodont and therocephalian fossil sites have been found on Farm 380 Bloemendal, situated just north of the WEF project area (Ms Marietjie Mostert, pers. comm., June 2021) One of these sites features substantial cranial fragments of several thick-skulled tapinocephalid dinocephalians (**Figure 30** & **Figure 31**). These fossils are of particular interest in that they are mapped within the lower part of the Poortjie Member. If correct (this requires confirmation), they would then represent very rare survivors of end Middle Permian Mass Extinction Event during which almost all members of the Dinocephalia went extinct (Day *et al.* 2015a, 2015b, 2020, Day & Rubidge 2021). A wide range of vertebrate and other fossil remains have been recorded from the critical upper Abrahamskraal – lower Poortjie Member stratigraphic interval during several recent PIA field studies to the east of the N12 adjacent or close to the present project area (Almond 2018, Almond in prep. 2021).

The extensive (> 6750 ha) combined Koup 1 & 2 WEF project area contains large areas of dissected hilly terrain with good to excellent exposure of both mudrock and sandstone facies of the potentially fossiliferous Lower Beaufort Group (Section 5.1). Sizeable portions of the combined WEF project area, especially in the NE, show low relief and here the bedrocks are mantled by Late Caenozoic superficial sediments (*e.g.* alluvium, soils, surface gravels) of low palaeontological sensitivity. During the 5-day palaeontological field survey by the author and an experienced assistant, numerous (over 50) new vertebrate fossil sites were recorded within the more accessible portions of the combined WEF and grid connection study area (See satellite map **Figure 61** and A2.2). GPS locality data together with a short description of each site as well as recommended mitigation (if any) is tabulated in Appendix 2 of this report while selected fossil specimens from the combined WEF project area are illustrated in **Figure 28** to **Figure 59** below with explanatory figure legends.

The main categories of fossils found here, associated with both sandstone and mudrock facies as well as downwasted surface gravels, include:

- Surface scatters and rare concentrations of disarticulated to semi-articulated skull and post-cranial skeletal elements of large-bodied tetrapods within the lower parts of the Karelskraal Member and apparently also within the lowermost part of the Poortjie Member. Most or all of these specimens are probably referable to one or (possibly) more members of **tapinocephalid Dinocephalia** (“horrible heads”) but diagnostic cranial or dental material is very rare (Alternatively, some of the material might be **pareisaur reptile** in origin but the presence of this group remains unconfirmed). Much of the bony material is fragmentary, secondarily mineralised (*e.g.* by pyrite), weathered (*e.g.* suncracked) and probably unidentifiable, so is of limited research or conservation interest. Some bones are encased within ferruginous carbonate pedoconcrete concretions or sandstone. However, a number of specimens are of scientific value and, if threatened by the proposed development, should be formally collected for storage in a museum collection (*e.g.* Iziko Museums, Cape Town). These include a concentration of several cranial fragments of very thick-skulled tapinocephalids (probably *Criocephalosaurus*) from the lower Poortjie Member on the Farm Bloemendal (Loc. 724), just N of and outside the WEF project area. These specimens may represent the remains of some of the last dinocephalians that survived the end-Middle Permian Extinction Event; a few other examples of this genus are recorded in the Beaufort West area (*cf* Day *et al.* 2015a, Almond 2020a).

Small-bodied herbivorous dicynodonts are the commonest fossils found in both the lower Karelskraal Member and the Poortjie Member. Many of these are referable to the common dassie-sized genus Diictodon (

- **Figure 25).** A number of specimens showing a broader skull table or (rarely) post-canine teeth belong to other dicynodont genera and are of higher palaeontological interest. Dense local concentrations of small *Diictodon* and other dicynodont skulls are found within mudrock intervals of the Poortjie Member at several sites.
- Sparse records of **trace fossils** include several (mostly equivocal) sandstone burrow casts of tetrapods, possible smaller-scale invertebrate burrows as well as fine horizontal burrows associated with subaqueous or pond margin microbial mats. No lungfish burrows were seen.
- Small carnivorous **therocephalians**, often preserved within fine-grained sandstones or pedoconcrete concretions, are known from several sites on the margins of the WEF project area but were not recorded during the recent field survey.
- Likewise, poorly preserved, ferruginised, **petrified wood** which is recorded weathering out of the Poortjie Member just to the east of the present study area as well as closer to Beaufort West (Almond 2020a) was not encountered here.

The Lower Beaufort Group fossil sites lie within both the uppermost Abrahamskraal Formation (Karelskraal Member) as well as the lower part of the overlying Poortjie Member of the Teekloof Formation. They therefore span the critical end-Guadalupian Mass Extinction event of c. 260 million years ago (*cf* **Figure 24**) and are of special biostratigraphic and palaeobiological interest for this reason. It is notable that the thick packages of silty mudrocks within the uppermost Karelskraal Member are generally largely unfossiliferous, even where they are very well exposed within the WEF project area. As mentioned in Section 5.1, the sedimentology of these mudrock intervals suggests strongly fluctuating climates with more pluvial, lacustrine intervals with high water tables alternating with periods of intense floodplain aridification. These highly unstable environmental conditions may be related to the end Middle Permian extinction episode. As noted previously, several thin tuff (volcanic ash) or tuffite horizons recorded within the Abrahamskraal – Poortjie contact zone are datable and could potentially constrain the age of the mass extinction event. The possible role of contemporary extra-basinal volcanism in the Karoo vertebrate extinctions has yet to be explored.

The diverse Late Caenozoic superficial deposits within the South African interior, including the Great Karoo region, have been comparatively neglected in palaeontological terms. However, sediments associated with ancient drainage systems, springs and pans in particular may occasionally contain important fossil biotas, notably the bones, teeth and horn cores of mammals as well as remains of reptiles like tortoises (*e.g.* Skead 1980, Klein 1984b, Brink, J.S. 1987, Bousman *et al.* 1988, Bender & Brink 1992, Brink *et al.* 1995, MacRae 1999, Meadows & Watkeys 1999, Churchill *et al.* 2000, Partridge & Scott 2000, Brink & Rossouw 2000, Rossouw 2006, De Ruyter *et al.* 2010, Backwell *et al.* 2017). Other late Caenozoic fossil biotas that may occur within these superficial deposits include non-marine molluscs (bivalves, gastropods), ostrich egg shells, trace fossils (*e.g.* calcretised termitaria, coprolites, invertebrate burrows, rhizocretions), and plant material such as peats or palynomorphs (pollens) in organic-rich alluvial horizons (Scott 2000) and diatoms in pan sediments. In Quaternary deposits, fossil remains may be associated with human artefacts such as stone tools and are also of archaeological interest (*e.g.* Smith 1999 and refs. therein). Ancient solution hollows within extensive calcrete hardpans may have acted as animal traps in the past. As with coastal and interior limestones, they might occasionally contain mammalian bones and teeth (perhaps associated with hyaena dens) or invertebrate remains such as snail shells. Although a number of Late Caenozoic deposits

(including thicker alluvium, spring-related calcretes) were examined during the palaeontological survey, no fossil remains were recorded within them.



Figure 28: Large proximal limb bone of a large-bodied tetrapod (probably dinocephalian) embedded within dark Karelskraal Member mudrocks (Loc. 665) (Scale is 15 cm long).



Figure 29: Weathered-out large vertebrae collected from stream gravels downslope of the *in situ* specimen illustrated above and probably belonging to the same animal (Loc. 665) (Top RHS block 15 cm across).



Figure 30: Partial skull roof of a tapinocephalid dinocephalian – probably *Criocephalosaurus* (Loc. 724) found on the Farm Bloemendal, just N of the Koup WEF project area (Scale is 15 cm long). The specimen probably comes from the lower Poortjie Member and, if so, would count among the last surviving members of the Dinocephalia known (See also following figure).



Figure 31: Oblique ventral view of the tapinocephalid specimen illustrate above. The skull is c. 25 cm wide and the skull roof as seen here is over 15 cm thick.



Figure 32: *In situ*, ferruginized partial postcranial skeleton of a dinocephalian therapsid embedded within Karelskraal Member mudrocks (Loc. 603) (Scale = 15 cm). Pyrite pseudomorph crystals within some of the bones as well as rippled sandstones in the vicinity suggest preservation of a corpse along a waterlogged lake margin.



Figure 33: Heap of fragmentary, disarticulated postcranial and cranial material collected in the immediate vicinity of the *in situ* dinocephalian skeleton illustrated above (Loc. 603) (Hammer = 30 cm).



Figure 34: Weathered jaw fragment of a dinocephalian with deep-rooted teeth associated with the skeleton in Figure 32 above (Loc. 603) (Block is c. 10 cm across).



Figure 35: Probable fragmentary cranial and other remains associated with the skeleton in Figure 32 (Loc. 603) (Scale = 15 cm).



Figure 36: Fragments of limb bones from the same individual (Loc. 603) (Scale = 15 cm). These are potentially of taxonomic value.



Figure 37: Further fragments of limb bones from the same individual (Loc. 603) (Scale = 15 cm).



Figure 38: Surface scatter of white, weathered bone material of an unidentified large-bodied tetrapod found as surface float overlying the upper Karelskraal Member (Loc. 660) (Scale = 15 cm). Such material is usually unidentifiable and accordingly of limited scientific value.




Figure 39: Close up of one of the sandstone and bone blocks illustrated above showing highly fragmentary preservation, possibly due to sun-cracking during pre-burial exposure on the ancient Karoo floodplain (Loc. 660) (Scale in cm).



Figure 40: Typical sparse scatter of fragmentary postcranial bones of an unidentified large-bodied tetrapod – pareiasaur or dinocephalian - from the Karelskraal Member, found as surface float (Loc. 623) (Largest block is c. 15.5 cm across). Such poorly-preserved post-cranial material is mostly of limited scientific value.



Figure 41: Thin channel sandstone containing dispersed, reworked post-cranial remains of a large-bodied tetrapod (arrowed) in the upper Karelskraal Member (Loc. 675) (Scale = 15 cm). This site is associated with a downslope trail of weathered-out bones within sandstone blocks.



Figure 42: Pedogenic calcrete nodule in float containing several articulated vertebrae, possibly from the tail, of a medium- to large-bodied tetrapod, upper Karelskraal Member (Loc. 613) (Block is c. 9 cm across).



Figure 43: Ferruginized calcrete concretion within cleaved overbank mudrocks of the upper Karelskraal Member containing a row of ribs of a large-bodied tetrapod (Loc. 668) (Concretion is c. 30 cm across).



Figure 44: Articulated complete skull of a small dicynodont therapsid almost entirely enclosed in a pedogenic calcrete concretion, found among float overlying the Karelskraal Member (Loc. 577) (Scale in cm and mm). The specimen shows several post-canine teeth (not illustrated here) and so does not belong to the common dicynodont genus *Diictodon*.



Figure 45: Skull of a small dicynodont from the Karelskraal Member (Loc. 631) showing a broad skull table, *i.e.* not *Diictodon* (Specimen is 7 cm long).



Figure 46: Poorly preserved dicynodont skull within a pedogenic calcrete concretion, Karelskraal Member) (Loc. 693) (Specimen is c. 10 cm long).



Figure 47: Possible (but equivocal) large, inclined vertebrate burrow cast within cleaved mudrocks of the Karelskraal Member (Loc. 594) (Hammer = 30 cm).



Figure 48: Possible (but equivocal) large, inclined vertebrate burrow cast in the Karelskraal Member (Loc. 595), recorded within a few meters of the previous specimen (Hammer = 30 cm).



Figure 49: Skull of a small-bodied dicynodont – probably *Diictodon* - from a mudrock interval within the Poortjie Member, seen in dorsal view (Loc. 641) (Specimen is c. 6.5 cm long).



Figure 50: Postcranial remains of a small-bodied therapsid, probably dicynodont, preserved within a pedogenic calcrete concretion from Poortjie Member mudrocks (Loc. 641) (Block is c. 11 cm across).



Figure 51: Well-preserved skull with articulated lower jaw and clear canine tusk preserved in situ within Poortjie Member mudrocks, oblique dorso-lateral view (Loc. 649) (Specimen is c. 8.5 cm long).



Figure 52: Small pedogenic calcrete concretion from the Poortjie Member enclosing the skull of a small-bodied dicynodont (Loc. 699) (Scale in cm).



Figure 53: *In situ* skull of a small dicynodont within cleaved siltstones of the Poortjie Member (Loc. 706) (Scale in cm and mm).



Figure 54: Post-crania, including limbs and rib cage, of a small-bodied dicynodont preserved inside a calcrete concretion embedded within cleaved Poortjie Member mudrocks (Loc. 687) (Scale = 15 cm).



Figure 55: Skull of small dicynodont within cleaved Poortjie Member mudrocks (Loc. 689) (Scale in cm).



Figure 56: Skull of small dicynodont with a broad skull table and clear pineal foramen weathering out of Poortjie Member mudrocks (Loc. 696).




Figure 57: Convex sandstone cast of a small vertebrate burrow weathering out of cleaved Poortjie Member mudrocks (Loc. 688) (Scale in cm and mm).



Figure 58: Cast of small subhorizontal burrow, either a small tetrapod or perhaps an invertebrate, within Poortjie Member mudrocks (Loc. 647) (Scale in cm).



Figure 59: Sandstone cast of small, inclined burrow (sloping down to the LHS) with a smooth central zone, possibly constructed by a small vertebrate (Loc. 648) (Scale = 15 cm).

6. IDENTIFICATION AND ASSESSMENT OF IMPACTS

The potential impact of the proposed Koup 2 WEF development and the associated grid connection on legally-protected local fossil heritage resources is evaluated in this section of the report and summarized in Tables 2 to 8 below. This assessment applies only to the *construction phase* of the developments since further significant impacts on fossil heritage during the planning, operational and decommissioning phases of the facility are not anticipated. The first assessment (Table 2) applies to all the key infrastructure described in Section 3 that will be situated within the main WEF project area (*i.e.* wind turbine foundations, access roads, on-site substation, pylons, underground cables, as well as the construction camp, laydown areas and operational and maintenance buildings, BESS *etc.*). Impacts of the grid connection options under consideration (See Fig. 3) are separately assessed in Table 3. Potential impacts here refer mainly to any associated new access roads, which may entail substantial surface disturbance or clearance, since bedrock excavations for the pylon footings are generally small.

6.1 Palaeontological sensitivity of the project area

The proposed Koup 2 WEF and grid connection project area is located in a region of the Great Karoo that is underlain by potentially fossiliferous sedimentary rocks of Late Palaeozoic and younger, Late Tertiary or Quaternary, age. In particular, these include (1) Middle to Late Permian continental sediments of the Abrahamskraal and Teekloof Formations (Lower Beaufort Group, Karoo Supergroup) which contain scientifically important fossils of vertebrates, trace fossils and terrestrial plants, as well as (2) Late Caenozoic alluvium that may contain important mammalian remains such as teeth and bones (These rock units and their fossils are described in more detail in Section 5 of this report).

The generally high palaeontological heritage sensitivity of the Lower Beaufort Group bedrocks in the Great Karoo is emphasized on the SAHRIS palaeosensitivity map maintained by SAHRA. The palaeontological heritage Site Sensitivity Map prepared by SiVEST Environmental for the Koup 2 WEF project area identifies areas underlain by the Lower Beaufort Group as being of a Very High Sensitivity (**Figure 60**). However, both desktop and field studies within this and neighbouring WEF project areas (*e.g.* Trakas, Beaufort West, Heuweltjies, Kraaltjies, Kwaggas 1-3 WEFs) demonstrate that, while a significant number of scientifically valuable, well-preserved fossils do indeed occur here, sometimes in high concentrations, in practice they are often scarce in this region and their distribution is to a large extent unpredictable. However, as a consequence of the considerably higher topographic relief – and hence bedrock exposure levels – encountered within the combined Koup 1 and 2 WEF and grid connection project areas, a significantly higher concentration of fossil sites is recorded here than within WEF project areas towards the east.

It is concluded that, applying the precautionary principle, the Koup 2 WEF and grid connection project areas are best assigned an overall High Sensitivity as far as palaeontological heritage is concerned, while recognising that, in practice, fossil sites are scarce at surface throughout much of the area (*N.B.* Additional fossils are preserved in the subsurface and may be impacted by excavations during the construction phase).




Figure 60: Paleontological sensitivity map for the Koup 2 WEF and grid connection project area. Due to the scarcity of well-preserved, scientifically important fossils over much of this region, based on desktop studies and fieldwork, it is inferred that large parts of the project area are in practice of LOW palaeontologically sensitivity. However, there is also a substantial number of, dispersed and unpredictable fossil sites here of HIGH to VERY HIGH sensitivity. Applying the precautionary principle, the Koup 2 WEF and associated grid connection project area are best assigned an overall High Sensitivity as far as palaeontological heritage is concerned. Areas underlain by thick alluvial sediments are generally of LOW sensitivity, although important concentrations of Cenozoic mammal remains might occur here.

6.2 Results of the Palaeontological Desktop and Field Study

6.2.1 WEF project area

A desktop review of the palaeontology of the Middle to Late Permian Lower Beaufort Group in the southern Great Karoo region, including palaeontological heritage impact assessments for a number of other authorized or proposed renewable energy developments in the region, shows that well-preserved fossil remains are generally scarce in this sector of the Great Karoo. However, a substantial number of scientifically important occurrences of vertebrate fossils are recorded here, and the vertebrate fossils may occur locally in high concentrations along the ridges as well as in low-lying terrain (Section 5). The 5-day palaeontological heritage survey of numerous exposures of Karoo Supergroup bedrocks as well as Late Cenozoic superficial sediments within the combined Koup 1 and Koup 2 WEF project area (See Appendix 2 and fossil locality maps, **Figure 61A-C** and A2.2) indicates that well-preserved, scientifically valuable fossils are present, for the most part sparsely distributed but occasionally in dense concentrations. Most of the

fossils recorded here lie outside the project footprint and are assigned a moderate to low provisional field rating (Appendix 2). The occurrence of important fossil remains in the subsurface obviously cannot be excluded and only a small subsample of all surface fossil sites have been detected by the reconnaissance-level field survey. Broadly comparable palaeontological findings have previously been obtained for the adjoining Trakas and Beaufort West WEF project areas (Almond 2018) as well as further WEF PIA work in the region (Almond, in prep.). However, as a result of the much higher bedrock exposure levels in the Koup 1 and Koup 2 WEF project areas, the total number and density of fossil sites recorded and expected here is considerably higher.

The potentially fossiliferous Permian bedrocks within the WEF project area are mostly mantled with Late Cenozoic colluvial and alluvial deposits as well as surface gravels and gravelly soils, none of which is palaeontologically sensitive in general. Few of the recorded fossil sites lies within or very close to (< 10 m) the proposed footprint of the WEF (**Figure 61A-C**) and therefore most of them should not be directly threatened by the proposed development. Several of the recorded fossil sites are associated with areas of good bedrock exposure that tend to occur along drainage lines and that are therefore generally protected by standard environmental buffer zones for water courses. No palaeontological Very High Sensitivity or No-Go areas have been identified within the project area. However, given the very incomplete coverage of the project footprint during the short field survey (see tracks in Fig. A2.1) - driven largely by access roads and the distribution of good bedrock exposure - many additional, unrecorded fossil sites must lie within or close to the footprint. The great majority (and probably all) of these unrecorded sites can be effectively mitigated through (1) judicious professional recording and collection during the recommended pre-construction specialist palaeontological walkdown of the project footprint complemented by (2) consistent application of a Chance Fossils Finds Procedure during the construction phase itself, as outlined in Section 8 and Appendix 4.

6.2.2 *Grid connection project area*

Several fossil sites are recorded within the grid connection project area for the Koup 2 WEF (**Figure 61B-C**). The corridor terrain traversing the eastern sector of the WEF project area in part comprises topographically subdued *vlaktes* mantled by palaeontologically insensitive alluvial deposits and surface gravels. As for the Koup 2 WEF development, a pre-construction specialist palaeontological walkdown of the finally selected grid connection footprint complemented by consistent application of a Chance Fossils Finds Procedure during the construction phase (Appendix 4) is recommended here.




Figure 61A: Google Earth© satellite image of the adjoining Koup 1 WEF (orange polygon) and Koup 2 WEF (yellow polygon) project areas, showing the numbered new fossil sites recorded during the site visit. No No-Go palaeontological heritage sites or areas have been identified. Most recorded fossil sites lie outside the project footprint and all can be mitigated through pre-construction collection. Most of the WEF and grid connection footprint has not been palaeontologically surveyed on foot and is likely to contain additional fossil sites at surface that are of palaeontological and conservation value. A pre-construction palaeontological walkdown of the final Koup 2 WEF and grid connection footprints is therefore recommended here.




Figure 59B: Google Earth© satellite image showing new (red) and previously recorded (yellow) fossil sites within the Lower Beaufort Group in the context of the Koup 2 WEF and originally assessed grid connection project areas (Please see following satellite map for more detail).




Figure 62C: Google Earth© satellite image showing new fossil sites (numbered in red) in the context of the Koup 2 WEF and originally assessed grid connection project areas and the provisional WEF layout (access roads and turbine locations shown in white). No No-Go palaeontological heritage sites or areas have been identified. Most recorded fossil sites lie outside the project footprint and all can be mitigated through pre-construction collection. Most of the WEF and chosen grid connection footprint (Option 1, blue corridor) has not been palaeontologically surveyed on foot and is likely to contain additional fossil sites at surface that are of palaeontological and conservation value. A pre-construction palaeontological walkdown of the final Koup 2 WEF and grid connection footprints is therefore recommended here.

6.3 Identification of Potential Impacts

The construction phase of the proposed WEF will entail extensive surface clearance as well as excavations into the superficial sediment cover and underlying bedrock (e.g. for widened or new access roads, wind turbine foundations, hardstanding areas, on-site substation, underground cables, construction laydown area, O&M building and BESS). Construction of the facility may adversely affect potential fossil heritage within the development footprint by damaging, destroying, disturbing or permanently sealing-in fossils preserved at or beneath the surface of the ground that are then no longer available for scientific research or other public good. The planning, operational and de-commissioning phases of the facility are unlikely to involve further adverse impacts on local palaeontological heritage and are therefore not separately assessed in this report. The potential palaeontological heritage resource impacts identified during the PIA assessment can be briefly summarized as follows:

- **Planning / Pre-construction Phase**

No significant impacts on palaeontological heritage anticipated.

- **Construction Phase**

Potential Impact 1: Disturbance, damage or destruction of fossil heritage resources preserved at or below the ground due to surface clearance and excavations (especially into sedimentary bedrock).

- **Operational Phase**

No significant impacts on palaeontological heritage anticipated.

- **Decommissioning Phase**

No significant impacts on palaeontological heritage anticipated

- **Cumulative impacts**

Potential loss of a significant fraction of scientifically important fossil heritage – especially fossil vertebrates - preserved within the Abrahamskraal and Teekloof Formations of the southern Great Karoo south of Beaufort West through multiple renewable energy developments in the region.

6.4 Assessment of WEF and grid connection project impacts

Potential impacts of the construction phase of the proposed Koup 2 WEF and associated grid connection on local fossil heritage resources, with and without mitigation, are assessed below in Tables 2 and 3 respectively, according to the Environmental Impact Assessment (EIA) Methodology developed by SiVEST. Further significant impacts on fossil heritage during the planning, operational and decommissioning phases of the facility are not anticipated.

Given the closely comparable geology of the WEF and grid connection project areas, the inferred impact ratings are the same in both cases.

6.4.1 Construction Phase: Disturbance, damage or destruction of fossils

The destruction, damage or disturbance out of context of legally-protected, scientifically-important fossils preserved at the ground surface or below ground that may occur during construction of the WEF / grid connection entail *direct negative* impacts to palaeontological heritage resources that are confined to the development footprint (*site*). These impacts can often be mitigated but cannot be fully rectified (*i.e.* they are *irreversible*). All the sedimentary formations represented within the study area contain fossils of *some* sort, and bedrock exposure levels within the development footprint are good, so impacts at some level on fossil heritage are *definite*. While most (but *not* all) of the fossils concerned are probably of widespread occurrence elsewhere within the outcrop areas of the formations concerned, *some* unique, well-preserved, scientifically-important fossils are known to occur in this region of the Great Karoo. The potential losses of irreplaceable fossil resources without mitigation is therefore conservatively rated as *significant*. Such impacts are of *permanent* duration. Their intensity / magnitude during the construction phase is rated as *medium* without mitigation as a precautionary measure since most of the project footprint has not been surveyed on foot. Without mitigation, a NEGATIVE MEDIUM impact significance is accordingly inferred for both the WEF and grid connection projects.

Potential negative impacts can be substantially reduced through implementation of the proposed mitigation measures, *viz.* a pre-construction palaeontological specialist walkdown of the final footprint (with recording and judicious collection of fossil material) backed up by a Chance Fossil Finds Procedure during the construction phase. With mitigation, the impact significance of the proposed WEF / grid connection project falls to NEGATIVE LOW.

Due to the reconnaissance level of the field survey of the extensive study area, confidence levels for this palaeontological heritage assessment are only moderate (*medium*). These conclusions are supported, however, by several previous palaeontological field assessments undertaken in the broader southern Karoo region by the author (See References and discussion on cumulative impacts below).

6.4.2 No-Go Option impacts

The No-Go Option, as assessed by the SiVEST system, is rated as NEGATIVE LOW (Table 4) in so far as even without development fossils will still be destroyed by natural weathering and erosion. In the case of the No-Go Alternative (*i.e.* no WEF / grid development), the possible loss of local heritage resources through construction activities (negative impact) would be avoided while potential improvements in palaeontological understanding through professional mitigation - *i.e.* recording and collection of palaeontological material and data (positive impacts) - would be lost. The slow destruction of fossils exposed at the surface through natural weathering and erosion would continue, but at the same time new fossils are revealed for scientific study. On balance, it is concluded that No-Go alternative would have a *neutral* impact on palaeontological heritage.

6.5 Cumulative impacts

Cumulative impacts addressed here principally concern the *potential* loss of a significant fraction of scientifically valuable and legally-protected fossil heritage preserved within the upper Abrahamskraal and

lower Teekloof Formations of the southern Karoo margins through multiple alternative energy developments in the region to the south of Beaufort West (**Figure 63** and Table 1). The cumulative impacts analysis shown in Table 5 is based on the Environmental Impact Assessment (EIA) Methodology developed by SiVEST.

Relevant renewable energy projects within a 35 km radius of the combined Koup 1 and 2 WEF and grid connection project areas are mapped in **Figure 63** below (No data is available for any other large-scale industrial developments in the region). PIA data for the proposed Leeu Gamka Solar Power Plant is not yet available. PIA studies for the authorized Mainstream Trakas and Beaufort West WEFs as well as the Lombardskraal Renewable Energy Facility have been undertaken by the present author (Almond 2018, 2020a). In addition, there are the proposed Mainstream Heuweltjies WEF and Kraaltjies WEF to the SE and E of the Koup WEF project area as well as for the proposed ABO Kwagga 1 to Kwagga 3 WEFs further to the east, for all of which palaeontological heritage impact assessments have been or are being conducted by the present author (Almond in prep., Almond 2021a-c). A number of further wind and solar renewable energy projects have been proposed on the southern outskirts of Beaufort West but these largely lie outside the 35 km cut-off radius applied here. Relevant published palaeontological literature for the region has also been considered (e.g. Day & Rubidge 2014, Rubidge & Day 2020, Day & Rubidge 2020). This cumulative impact assessment applies only to the construction phases of the renewable energy developments, since significant additional impacts on palaeontological heritage during the planning, operational and decommissioning phases are not anticipated.

It should be emphasized that, in the case of palaeontological heritage, it only makes sense to consider cumulative impacts on *comparable fossil assemblages* present in the same rock units (groups, formations, members *etc*) that are represented in the present study area as well as in the broader study region. For example, impacts on Mid-Palaeozoic aquatic fossil invertebrates in the Cape Supergroup that crops out in the Cape Fold Mountains to the south of the present study area are not directly relevant to - or cannot be reasonably weighed against - impacts on Middle Permian fossil assemblages of terrestrial vertebrates in the Lower Beaufort Group that is represented in the present study area. The analysis in Table 5 is therefore restricted to considering cumulative impacts on fossil heritage preserved within rock units and fossil assemblages that are represented in the Koup 1 and 2 WEF project areas as well as in nearby WEF and solar project areas – specifically the uppermost Abrahamskraal Formation and lowermost Teekloof Formation (upper *Tapinocephalus* Assemblage Zone). Since potentially fossiliferous, consolidated Late Caenozoic alluvial deposits will normally not be impacted in WEF developments because they usually lie along well-buffered drainage lines they are not considered for the purpose of this analysis.

In all the strictly *relevant* field-based Karoo palaeontological studies listed above the palaeontological sensitivity of the project area and the palaeontological heritage impact significance for the developments concerned has been rated as *low*. In all cases it was concluded by the author that, despite the undoubted occurrence of scientifically-important fossil remains (notably fossil vertebrates, petrified wood), the overall impact significance of the proposed developments was low because the probability of significant impacts on *scientifically important, unique or rare fossils* was slight. While fossils do indeed occur within most of the formations present, they tend to be sparse – especially as far as fossil vertebrates are concerned - while the great majority represent common forms that occur widely within the outcrop areas of the rock units concerned. Important exceptions include rare, semi-articulated skeletal remains of therapsids and pareiasaur reptiles as well as well-preserved dicynodont skulls of biostratigraphic significance from the *Tapinocephalus* Assemblage Zone.

Anticipated cumulative impacts of the known renewable energy projects proposed or authorised for the margins of the Great Karoo region to the south of Beaufort West – including the proposed adjoining Koup 1 and Koup 2 Wind Energy Facilities - are assessed as *NEGATIVE MEDIUM* without mitigation. Overall impact significance may fall to *NEGATIVE LOW* with full mitigation since impacts will then occur at a lower intensity and will be partially offset by valuable new scientific data. The analysis only applies *provided that* all the proposed monitoring and mitigation recommendations made for all these various projects are followed through (*N.B.* This is inherently unpredictable, and, sadly, unlikely). Unavoidable residual negative impacts may be partially offset by the improved understanding of Karoo palaeontology resulting from appropriate professional mitigation. This is regarded as a *positive* impact for Karoo palaeontological heritage.

In conclusion, the cumulative impacts on local fossil heritage anticipated for the various renewable energy projects in the southern Great Karoo margins region due south of Beaufort West – including the proposed Koup 1 and Koup 2 Wind Energy Facilities - are acceptable, *provided that* all recommended mitigation recommendations for these projects are followed through.




Figure 63: Map showing project areas for authorized and proposed renewable energy projects within a 35 km radius of the Koup 1 and Koup 2 WEF and grid connection project areas (Image provided by SiVEST). Additional unmapped WEFs proposed to the east and southeast of the Koup project area (Heuweltjies WEF, Kraaltjies WEF, Kwagga 1-3 WEFs) have also been taken into consideration here.

Table 1: Renewable energy developments proposed within a 35km radius of the Koup 2 WEF application site (See also text for Heuweltjies WEF, Kraaltjies WEF, Kwagga 1-3 WEFs not listed here).

Project	DEA Reference No	Technology	Capacity	Status of Application / Development
Proposed Beaufort West Wind Farm	12/12/20/1784/1	Wind	140MW	Approved
Proposed Trakas Wind Farm	12/12/20/1784/2	Wind	140MW	Approved
Proposed Wind and Solar Facility on the Farm Lombardskraal 330	14/12/16/3/3/2/406	Solar	20MW	EIA in Process
Proposed Leeu Gamka Solar Power Plant	12/12/20/2296	Solar	-	EIA in Process
Proposed Koup 1 WEF	TBA	Wind	140MW	EIA in Process

Table 2: Assessment of paleontological heritage impacts for the proposed Koup 2 Wind Energy Facility (Construction Phase)

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION								
		E	P	R	L	D	I / M	TOTAL STATUS S(+)	S	E		P	R	L	D	I / M	TOTAL STATUS S(+)	S		
Construction Phase																				
Fossil heritage resources	Disturbance, damage or destruction of fossils at or beneath the ground surface due to surface clearance and bedrock excavations	1	4	4	3	4	2	32	-	Medium	Pre-construction walkdown (with fossil recording / collection) of final footprint by specialist palaeontologist. Chance Fossil Finds Procedure during construction phase.	1	2	4	2	4	1	13	-	Low

Table 3: Assessment of paleontological heritage impacts for the proposed Koup 2 Wind Energy Facility grid connection (Construction Phase)

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION								
		E	P	R	L	D	I / M	TOTAL STATUS S(+)	S	E		P	R	L	D	I / M	TOTAL STATUS S(+)	S		
Construction Phase																				
Fossil heritage resources	Disturbance, damage or destruction of fossils at or beneath the ground surface due to surface clearance and bedrock excavations	1	4	4	3	4	2	32	-	Medium	Pre-construction walkdown (with fossil recording / collection) of final footprint by specialist palaeontologist. Chance Fossil Finds Procedure during construction phase.	1	2	4	2	4	1	13	-	Low

Table 4: Assessment of impacts for the No Go Option

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION										
		E	P	R	L	D	I / M	TOTAL STATUS S(+)	S	E		P	R	L	D	I / M	TOTAL STATUS S(+)	S				
Construction Phase																						
Fossil heritage resources	Disturbance, damage or destruction of fossils at or beneath the ground surface due to natural weathering and erosion	1	4	4	2	4	1	15	-	Low	N/A											N/A

Table 5: Assessment of cumulative impacts for the Koup 1 & 2 WEFs and other renewable energy developments in the region.

ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION									RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION								
		E	P	R	L	D	I / M	TOTAL STATUS S(+)	S	E		P	R	L	D	I / M	TOTAL STATUS S(+)	S		
CumulativePhase																				
Fossil heritage resources	Disturbance, damage or destruction of fossils at or beneath the ground surface due to surface clearance and bedrock excavations	1	4	4	3	4	2	32	-	Medium	(N.B. Vary between projects) Pre-construction walkdown (with fossil recording / collection) of final footprint by specialist palaeontologist. Chance Fossil Finds Procedure during construction phase.	1	2	4	2	4	1	13	-	Low

6.6 Overall Impact Rating

Overall impact ratings for the Koup 2 WEF and associated grid connection projects are provided in Tables 6 & 7 below. The significance of relevant cumulative impacts is assessed in Table 8. Recommended monitoring and mitigation measures for these developments are outlined in more detail in Section 8 of this report.

Table 6: Overall impact rating for the Koup 2 WEF project

KOU P 2 WEF																				
ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION							RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION										
		E	P	R	L	D	I/M	TOTAL STATUS (+ OR -)		S	E	P	R	L	D	I/M	TOTAL STATUS (+ OR -)	S		
Construction Phase																				
Fossil heritage resources	Disturbance, damage or destruction of fossils at or beneath the ground surface due to surface clearance and bedrock excavations	1	4	4	3	4	2	32	-	M	Pre-construction walkdown (with fossil recording / collection) of final footprint by specialist palaeontologist. Chance Fossil Finds Procedure during construction phase	1	2	4	2	4	1	13	-	L

Table 7: Overall impact rating for the Koup 2 WEF grid connection project

KOU P 2 WEF GRID CONNECTION																				
ENVIRONMENTAL PARAMETER	ISSUE / IMPACT / ENVIRONMENTAL EFFECT/ NATURE	ENVIRONMENTAL SIGNIFICANCE BEFORE MITIGATION								RECOMMENDED MITIGATION MEASURES	ENVIRONMENTAL SIGNIFICANCE AFTER MITIGATION									
		E	P	R	L	D	I/M	TOTAL	STATUS (+ OR -)		S	E	P	R	L	D	I/M	TOTAL	STATUS (+ OR -)	S
Construction Phase																				
Fossil heritage resources	Disturbance, damage or destruction of fossils at or beneath the ground surface due to surface clearance and bedrock excavations	1	4	4	3	4	2	32	-	M	Pre-construction walkdown (with fossil recording / collection) of final footprint by specialist palaeontologist. Chance Fossil Finds Procedure during construction phase	1	2	4	2	4	1	13	-	L

Table 8: Overall cumulative impact rating for the Koup 2 WEF and grid connection project

Cumulative																				
Fossil heritage resources	Disturbance, damage or destruction of fossils at or beneath the ground surface due to surface clearance and bedrock excavations	1	4	4	3	4	2	32	-	M	(N.B. Vary between projects) Pre-construction walkdown (with fossil recording / collection) of final footprint by specialist palaeontologist. Chance Fossil Finds Procedure during	1	2	4	2	4	1	13	-	L

7. COMPARATIVE ASSESSMENT OF ALTERNATIVES

7.1 Koup 2 WEF

A comparable NEGATIVE MEDIUM impact significance (without mitigation), as assessed in Table 2 applies equally to all project infrastructure alternatives and layout options under consideration that are outlined in Section 3.3 of this report. This includes the various site options for the on-site substation and construction laydown area. Given their very similar geological - and hence palaeontological - contexts, there are no preferences on palaeontological heritage grounds for any layout among the various options under consideration. The proposed final layout is shown in **Figure 4b**.

Key

PREFERRED	The alternative will result in a low impact / reduce the impact / result in a positive impact
FAVOURABLE	The impact will be relatively insignificant
LEAST PREFERRED	The alternative will result in a high impact / increase the impact
NO PREFERENCE	The alternative will result in equal impacts

Table 9: Comparative assessment of WEF layout options

Alternative	Preference	Reasons (incl. potential issues)
SUBSTATION SITE ALTERNATIVES		
Substation Option 1	None	Similar geological / palaeontological context
Substation Option 2	None	
CONSTRUCTION LAYDOWN AREA SITE ALTERNATIVES		
Construction Laydown Area Option 1	None	Similar geological / palaeontological context
Construction Laydown Area Option 2	None	

7.2 Koup 2 grid connection

There is no preference for the originally proposed Grid Option 1 or 2 (either alternative) on palaeontological heritage grounds. Since they are of similar length and traverse bedrocks of comparable palaeosensitivity, they are likely to have similar potential impacts on fossil sites. Grid Option 1 has now been chosen as the preferred alternative (**Figure 3b**).

Table 10: Comparative assessment of WEF grid connection options (Option 1 has now been chosen)

Alternative	Preference	Reasons (incl. potential issues)
GRID CONNECTION ALTERNATIVES		
Grid Option 1 (Sub1)	None	All options have similar impact significance due to similar geological/ palaeontological context.
Grid Option 1 (Sub2)		
Grid Option 2 (Sub1)		

Alternative	Preference	Reasons (incl. potential issues)
Grid Option 2 (Sub2)		

8. PROPOSED MONITORING AND MITIGATION: INPUT TO EMPR

A significant number of new fossil sites are recorded within the combined Koup 1 and Koup 2 WEF project areas (Section 5, Appendix 2) as well as in the vicinity of some of the grid connection options under consideration. However, (1) the majority of these fossil sites lie well away from the proposed infrastructure footprints (see satellite images **Figure 61A-C**), (2) many of them are rated as being of low scientific or conservation significance (See Appendix 2) while (3) all of them can be mitigated, if necessary, through professional palaeontological collection during the construction phase. ***The distribution of fossil sites need therefore have no influence on the proposed layout of the WEFs or associated grid connections.***

A pre-construction palaeontological heritage walkdown of the final WEF and grid connection layout by a suitably qualified palaeontologist is recommended here. This motivated by:

- the very incomplete coverage of the WEF and grid connection project areas (including provisional infrastructure footprints) during the short, reconnaissance-level palaeontological site visit;
- the highly unpredictable distribution of important fossil sites, which may occur within both mudrock and sandstone facies;
- the documented presence of a substantial number of scientifically-valuable fossil remains in the region based on desktop and field studies, favoured by extensive areas of good bedrock exposure.

The recommended palaeontological walkdown should involve the recording and judicious collection of valuable fossil material as well as relevant geological data (e.g. on stratigraphic context, preservation style / taphonomy) within or close to (within ~10 m) the project footprint. This mitigation phase is *essential* because all fossil heritage resources in the RSA are protected by law and it is illegal to disturb, damage or destroy fossils here without a permit from the relevant provincial heritage resources agency (South African Heritage Resources Act, Act No. 25 of 1999). The palaeontological heritage mitigation report would then make recommendations for further studies and mitigation (if any are necessary) during the construction phase of the renewable energy project. Since mitigation through recording and collection is almost invariably feasible, late-stage modifications to the final WEF / grid infrastructure layout (e.g. micro-siting changes to access roads, turbine or pylon locations) are not anticipated here.

The palaeontologist responsible for the mitigation work will be required to submit a Work Plan for approval by Heritage Western Cape (HWC) and a Mitigation Report must be submitted to HWC for consideration. All fieldwork and reporting should meet the standards of international best practice as well as those developed for PIA reports by SAHRA (2013) and Heritage Western Cape (2021). Fossil material collected must be safeguarded and curated within an approved palaeontological repository (e.g. museum or university collection) with full collection data.

During the construction phase the Chance Fossil Finds Protocol summarized in Appendix 4 should be fully implemented.

The Environmental Control Officer (ECO) / Environmental Site Officer (ESO) responsible for the development should be made aware of the possibility of important fossil remains (vertebrate bones, teeth, petrified wood, plant-rich horizons etc.) being found or unearthed during the construction phase of the development. Monitoring for fossil material of all major surface clearance and deeper (>1m) excavations by the Environmental Site Officer on an on-going basis during the construction phase is therefore recommended. Significant fossil finds should be safeguarded and reported at the earliest opportunity to Heritage Western Cape for recording and sampling by a professional palaeontologist (Contact details: Heritage Western Cape, 3rd Floor Protea Assurance Building, 142 Longmarket Street, Green Market Square, Cape Town 8000. Private Bag X9067, Cape Town 8001. Tel: 021 483 5959 Email: ceoheritage@westerncape.gov.za).

These recommendations must be included within the EMPr for the Koup 2 WEF and associated grid connection development.

9. SUMMARY & CONCLUSIONS

9.1 Summary of Findings

The Koup 2 WEF and grid connection project area is underlain by continental (fluvial / lacustrine) sediments of the Abrahamskraal and Teekloof Formations (Lower Beaufort Group, Karoo Supergroup) which are of Middle to Late Permian age. These bedrocks contain sparse, unpredictable to locally concentrated vertebrate fossils as well as rare trace fossils (e.g. tetrapod burrows) and plant material of scientific and conservation value. A substantial number of new fossil vertebrate sites (cranial and post-cranial material of large-bodied dinocephalians, small dicynodonts, rare tetrapod burrow casts) have been recorded during within the WEF project area during the short site visit, while several more sites have previously been mapped shortly outside its margins. These palaeontological sites, together with their sedimentological context, provide important data for on-going research into the pattern and causes of the Middle Permian Mass Extinction Event on land around 260 million years ago.

Scientifically-valuable and legally-protected fossil heritage resources preserved at or beneath the ground surface within the project footprint are potentially threatened by clearance and bedrock excavations during the construction phase of the WEF and grid connection (e.g. for access roads, wind turbine foundations). The majority of the recorded fossil sites lie outside the project footprint but most of the WEF and grid connection footprint has yet to be palaeontologically surveyed on foot. A significant number of unrecorded sites almost undoubtedly lies within of very close to the project footprint.

No Very High Sensitivity or No-Go palaeontological sites or areas have been identified within the WEF or grid connection project areas. Since all known fossil sites can be readily mitigated through professional recording and collection of fossil material in the pre-construction phase, no recommendations for micro-siting of infrastructure such as wind turbine, pylon positions or access roads are therefore made here. There are no preferences on palaeontological heritage grounds for specific site options for the Koup 2 WEF on-site substation and construction laydown area. Grid Option 1 and 2 (either alternative) are equally supported for the grid connection since they have a similar potential impacts on fossil sites. There are no objections on palaeontological heritage grounds to authorization of the proposed final layout.

The proposed Koup 2 WEF and associated grid connection developments are assigned a similar overall impact significance rating (Construction Phase) of NEGATIVE MEDIUM without mitigation and NEGATIVE LOW following mitigation. No significant further impacts on fossil heritage resources are anticipated in the planning, operational and decommissioning phases. The No-Go Option might have a NEGATIVE LOW or perhaps neutral impact significance; fossils will continue to be exposed and destroyed by natural weathering processes while the positive benefits of professional mitigation (*viz.* improved palaeontological database) will be lost. Anticipated cumulative impacts in the context of several planned or authorized renewable energy projects in the region are assessed as NEGATIVE MEDIUM before mitigation and NEGATIVE LOW after mitigation.

Recommended mitigation comprises (1) a specialist palaeontological walk-down of the final WEF and grid connection project areas in the pre-construction phase and (2) implementation of a Chance Fossil Finds Protocol (See Appendix 4) by the ECO / ESO during the construction phase. The specialist palaeontologist responsible will be required to submit a Work Plan for approval by Heritage Western Cape.

The proposed WEF and grid connection developments are not fatally flawed and, on condition that the recommended mitigation measures are included within the relevant EMPs and implemented in full, there are no objections on palaeontological heritage grounds to their authorization.

9.2 Conclusions and Impact Statement

In terms of palaeontological heritage resources, the proposed Koup 2 WEF and associated grid connection developments are assigned a similar overall impact significance rating (Construction Phase) of NEGATIVE MEDIUM without mitigation and NEGATIVE MEDIUM following mitigation. No significant further impacts on fossil heritage resources are anticipated in the planning, operational and decommissioning phases. The No-Go Option might have a NEGATIVE LOW impact significance. Anticipated cumulative impacts in the context of several planned or authorized renewable energy projects in the region are assessed as NEGATIVE MEDIUM without mitigation and NEGATIVE LOW after mitigation.

The proposed WEF and grid connection developments are not fatally flawed and, on condition that the recommended mitigation measures are included within the EMP and implemented in full, there are no objections on palaeontological heritage grounds to their authorization.

10. REFERENCES

ALMOND, J.E. 2010a. Eskom Gamma-Omega 765kV transmission line: Phase 2 palaeontological impact assessment. Sector 1, Tanqua Karoo to Omega Substation (Western and Northern Cape Provinces), 95 pp + appendix. Natura Viva cc, Cape Town.

ALMOND, J.E. 2010b. Palaeontological impact assessment: pre-scoping desktop study. Proposed Mainstream wind farm to the south of Beaufort West, Western Cape, 19 pp. Natura Viva cc., Cape Town.

ALMOND, J.E. 2010c. Areas proposed for low-cost housing, Beaufort West, Western Cape Province. Palaeontological impact assessment: combined desktop & scoping study, 19 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2011a. Proposed Photovoltaic Power Facility, Farm Steenrotsfontein 168, Beaufort West Municipality, Western Cape Province. Palaeontological impact assessment: desktop study, 23 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2011b. Proposed windfarm development, Beaufort West Municipality, Western Cape. Palaeontological specialist study: combined desktop & field-based assessment, 27 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2014. Proposed Droërivier Solar Facility, Portion 55 of Farm 168 Steenrotsfontein and a portion of Portion 10 of Farm 170 Weltevreden, Beaufort West Municipality, Western Cape. Palaeontological impact assessment: combined desktop & field-based study, 53 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2015g. Proposed Gunstfontein Wind Energy Facility near Sutherland, Karoo Hoogland Local Municipality, Northern Cape Province. Palaeontological heritage assessment: combined desktop & field-based study, 62 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2018. Proposed Trakas and Beaufort West 140 MW Wind Farms and associated electrical infrastructure near Beaufort West, Central Karoo District, Western Cape. Combined desktop and field-based study, 61 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2020a. Lombardskraal Renewable Energy Facility project area, Beaufort West, Western Cape. Palaeontological heritage site sensitivity assessment: combined desktop & field-based study, 22 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2020b. Grid connection for the proposed Redcap Nuweveld Wind Farms, Beaufort West Local Municipality, Central Karoo District Municipality, Western Cape. Palaeontological heritage assessment: desktop and field-based report, 101 pp. Natura Viva cc Cape Town.

ALMOND, J.E. 2021a. Proposed Development of the Kwagga 1 Wind Energy Facility near Beaufort West in the Central Karoo District, Western Cape. Palaeontological heritage: combined desktop & field-based screening study & site sensitivity verification, 18 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2021b. Proposed Development of the Kwagga 2 Wind Energy Facility near Beaufort West in the Central Karoo District, Western Cape. Palaeontological heritage: combined desktop & field-based screening study & site sensitivity verification, 17 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2021c. Proposed Development of the Kwagga 3 Wind Energy Facility near Beaufort West in the Central Karoo District, Western Cape. Palaeontological heritage: combined desktop & field-based screening study & site sensitivity verification, 18 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. & PETHER, J. 2008. Palaeontological heritage of the Western Cape. Interim SAHRA technical report, 20 pp. Natura Viva cc., Cape Town.

BAMFORD, M. 1999. Permo-Triassic fossil woods from the South African Karoo Basin. *Palaeontologia africana* 35, 25-40.

BENDER, P.A. 2004. Late Permian actinopterygian (palaeoniscid) fishes from the Beaufort Group, South Africa: biostratigraphic and biogeographic implications. *Council for Geoscience Bulletin* 135, 84 pp.

BENDER, P.A. & BRINK, J.S. 1992. A preliminary report on new large mammal fossil finds from the Cornelia-Uitzoek site. *South African Journal of Science* 88: 512-515.

BLACKWELL, L., STEININGER, C., NEVELING, J. ABDALA, F., PEREIRA, L., MAYER, E., ROSSOUW, L., DE LA PEÑA P. & BRINK, J. 2017. Holocene large mammal mass death assemblage from South Africa. *Quaternary International* xxx (2017), p1-15.

BOONSTRA, L.D. 1969. The fauna of the *Tapinocephalus* Zone (Beaufort Beds of the Karoo). *Annals of the South African Museum* 56: 1-73.

BOUSMAN, C.B. *et al.* 1988. Palaeoenvironmental implications of Late Pleistocene and Holocene valley fills in Blydefontein Basin, Noupoot, C.P., South Africa. *Palaeoecology of Africa* 19: 43-67.

BRINK, J.S. 1987. The archaeozoology of Florisbad, Orange Free State. *Memoirs van die Nasionale Museum* 24, 151 pp.

BRINK, J.S. *et al.* 1995. A new find of *Megalotragus priscus* (Alcephalini, Bovidae) from the Central Karoo, South Africa. *Palaeontologia africana* 32: 17-22.

BRINK, J.S. & ROSSOUW, L. 2000. New trial excavations at the Cornelia-Uitzoek type locality. *Navorsing van die Nasionale Museum Bloemfontein* 16, 141-156.

CHURCHILL, S.E. *et al.* 2000. Erfkroon: a new Florisian fossil locality from fluvial contexts in the western Free State, South Africa. *South African Journal of Science* 96: 161-163.

COLE, D.I. & VORSTER, C.J. 1999. The metallogeny of the Sutherland area, 41 pp. Council for Geoscience, Pretoria.

COLE, D.I. AND WIPPLINGER, P.E. 2001, Sedimentology and molybdenum potential of the Beaufort Group in the main Karoo Basin, South Africa, Council for Geoscience Memoir, South Africa 80, 225 pp.

COLE, D.I., JOHNSON, M.R. & DAY, M.O. 2016. Lithostratigraphy of the Abrahamskraal Formation (Karoo Supergroup), South Africa. *South African Journal of Geology* 119.2, 415-424.

DAY, M.O. & RUBIDGE, B.S. 2010. Biostratigraphy of the Tapinocephalus Assemblage Zone (Beaufort Group, Karoo Supergroup), South Africa. *South African Journal of Geology* 123, 149-164.

DAY, M.O. AND RUBIDGE, B.S., 2014. A brief lithostratigraphic review of the Abrahamskraal and Koonap Formations of the Beaufort Group, South Africa: Towards a basin-wide stratigraphic scheme for the Middle Permian Karoo. *Journal of African Earth Sciences* 100, 227-242.

DAY, M.O., GUVEN, S., ABDALA, F., JIRAH, S., RUBIDGE, B.S. AND ALMOND, J. 2015a. Youngest dinocephalian fossils extend the Tapinocephalus Zone, Karoo Basin, South Africa. *South African Journal of Science* 111, 78-82.

DAY, M.O., RAMEZANI, J., BOWRING, S.A., SADLER, P.M., ERWIN, D.H., ABDALA, F. & RUBIDGE, B.S. 2015b. When and how did the terrestrial mid-Permian mass extinction occur? Evidence from the tetrapod record of the Karoo Basin, South Africa. *Proc. R. Soc. B* 282: 20150834. <http://dx.doi.org/10.1098/rspb.2015.0834>

DAY, M.O. & RUBIDGE, B.S.. 2020. Biostratigraphy of the *Tapinocephalus* Assemblage Zone (Beaufort Group, Karoo Supergroup), South Africa. *South African Journal of Geology* 123, 149 - 164.

DAY, M.O. & SMITH, R.M.S. 2020. Biostratigraphy of the *Endothiodon* Assemblage Zone (Beaufort Group, Karoo Supergroup), South Africa. *South African Journal of Geology* 123, 164 - 180.

DAY, M.O. & RUBIDGE, B.S. 2021. The Late Capitanian mass extinction of terrestrial vertebrates in the Karoo Basin of South Africa. *Frontiers in Earth Science* 9, article 631198, 15 pp.

DE RUITER, D.J., BROPHY, J.K., LEWIS, P.J., KENNEDY, A.M., STIDHAM, T.A., CARLSON, K.B. & HANCOX, P.J. 2010. Preliminary investigation of Matjhabeng, a Pliocene fossil locality in the Free State of South Africa. *Palaeontologia Africana* 45, 11-22.

HERITAGE WESTERN CAPE 2016. Guide for minimum standards for archaeology and palaeontology reports submitted to Heritage Western Cape, 4 pp.

HERITAGE WESTERN CAPE 2021. Guide for minimum standards for archaeology and palaeontology reports submitted to Heritage Western Cape - June 2021, 6 pp.

JOHNSON, M.R. & KEYSER, A.W. 1979. The geology of the Beaufort West area. Explanation of geological Sheet 3222, 14 pp. Council for Geoscience, Pretoria.

JOHNSON, M.R., VAN VUUREN, C.J., VISSER, J.N.J., COLE, D.I., WICKENS, H. DE V., CHRISTIE, A.D.M., ROBERTS, D.L. & BRANDL, G. 2006. Sedimentary rocks of the Karoo Supergroup. In: Johnson. M.R., Anhaeusser, C.R. & Thomas, R.J. (eds.) *The geology of South Africa*, pp. 461-499. Geological Society of South Africa, Johannesburg & the Council for Geoscience, Pretoria.

KEYSER, A. W. 1966. Some indications of arid climate during the deposition of the Beaufort Series. *Annals of the Geological Survey of South Africa* 5,77–79.

KEYSER, A.W. & SMITH, R.M.H. 1977-78. Vertebrate biozonation of the Beaufort Group with special reference to the Western Karoo Basin. *Annals of the Geological Survey of South Africa* 12: 1-36.

KITCHING, J.W. 1977. The distribution of the Karoo vertebrate fauna, with special reference to certain genera and the bearing of this distribution on the zoning of the Beaufort beds. *Memoirs of the Bernard Price Institute for Palaeontological Research, University of the Witwatersrand*, No. 1, 133 pp (incl. 15 pls).

KLEIN, R.G. 1984. The large mammals of southern Africa: Late Pliocene to Recent. In: Klein, R.G. (Ed.) *Southern African prehistory and paleoenvironments*, pp 107-146. Balkema, Rotterdam.

MACRAE, C. 1999. *Life etched in stone. Fossils of South Africa*, 305 pp. The Geological Society of South Africa, Johannesburg.

MARCHETTI L. et al. 2019. Permian-Triassic vertebrate footprints from South Africa: Ichnotaxonomy, producers and biostratigraphy through two major faunal crises. *Gondwana Research* 72, 139-168.

McCARTHY, T. & RUBIDGE, B. 2005. *The story of Earth and life: a southern African perspective on a 4.6-billion-year journey*. 334pp. Struik, Cape Town.

MEADOWS, M.E. & WATKEYS, M.K. 1999. Palaeoenvironments. In: Dean, W.R.J. & Milton, S.J. (Eds.) *The karoo. Ecological patterns and processes*, pp. 27-41. Cambridge University Press, Cambridge.

NEUMANN, S. 2020. Taxonomic revision of the tapinocephalid dinocephalian subfamilies Moschopinae, Tapinocephalinae and Reibeckosaurinae – the key to understanding Middle Permian tetrapod biodiversity. Unpublished PhD thesis, University of the Witwatersrand, Johannesburg, 408pp.

NICOLAS, M.V. 2007. Tetrapod diversity through the Permo-Triassic Beaufort Group (Karoo Supergroup) of South Africa. Unpublished PhD thesis, University of Witwatersrand, Johannesburg.

PAIVA, F., 2015. Fluvial facies architecture and provenance history of the Abrahamskraal-Teekloof Formation transition (Lower Beaufort Group) in the main Karoo Basin. Unpublished M.Sc. dissertation, University of Cape Town, Cape Town, 98pp.

PARTRIDGE, T.C. & MAUD, R.R. 1987. Geomorphic evolution of southern Africa since the Mesozoic. *South African Journal of Geology* 90: 179-208.

PARTRIDGE, T.C., BOTHA, G.A. & HADDON, I.G. 2006. Cenozoic deposits of the interior. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) *The geology of South Africa*, pp. 585-604. Geological Society of South Africa, Marshalltown.

PARTRIDGE, T.C. & SCOTT, L. 2000. Lakes and pans. In: Partridge, T.C. & Maud, R.R. (Eds.) *The Cenozoic of southern Africa*, pp.145-161. Oxford University Press, Oxford.

RETALLACK, G.J., METZGER, C.A., GREAVER, T., HOPE JAHREN, A., SMITH, R.M.H. & SHELDON, N.D. 2006. Middle – Late Permian mass extinction on land. *GSA Bulletin* 118, 1398-1411.

ROSSOUW, L. 2006. Florisian mammal fossils from erosional gullies along the Modder River at Mitasrust Farm, Central Free State, South Africa. *Navorsinge van die Nasionale Museum Bloemfontein* 22, 145-162.

SAHRA 2013. Minimum standards: palaeontological component of heritage impact assessment reports, 15 pp. South African Heritage Resources Agency, Cape Town.

SCOTT, L. 2000. Pollen. In: Partridge, T.C. & Maud, R.R. (Eds.) The Cenozoic of southern Africa, pp.339-35. Oxford University Press, Oxford.

SKEAD, C.J. 1980. Historical mammal incidence in the Cape Province. Volume 1: The Western and Northern Cape, 903pp. Department of Nature and Environmental Conservation, Cape Town.

SMITH, R.M.H. & KEYSER, A.W. 1995a. Biostratigraphy of the Tapinocephalus Assemblage Zone. Pp. 8-12 in Rubidge, B.S. (ed.) Biostratigraphy of the Beaufort Group (Karoo Supergroup). South African Committee for Stratigraphy, Biostratigraphic Series No. 1. Council for Geoscience, Pretoria.

SMITH, R.M.H. & KEYSER, A.W. 1995b. Biostratigraphy of the Pristerognathus Assemblage Zone. Pp. 13-17 in Rubidge, B.S. (ed.) Biostratigraphy of the Beaufort Group (Karoo Supergroup). South African Committee for Stratigraphy, Biostratigraphic Series No. 1. Council for Geoscience, Pretoria.

SMITH, R., RUBIDGE, B. & VAN DER WALT, M. 2012. Therapsid biodiversity patterns and paleoenvironments of the Karoo Basin, South Africa. Chapter 2 pp. 30-62 in Chinsamy-Turan, A. (Ed.) Forerunners of mammals. Radiation, histology, biology. xv + 330 pp. Indiana University Press, Bloomington & Indianapolis.

SMITH, R.M.S. et al. 2020. Introduction to the tetrapod biozonation of the Karoo Supergroup. South African Journal of Geology 123, 131-140 • doi:10.25131/sajg.123.0009.

VAN DER WALT, J. 2019. Phase 2 Strategic Environmental Assessment for Wind and Solar Photovoltaic Energy Development in South Africa. Appendix A3. Heritage Scoping Assessment Report, 65 pp. CSIR.

11. ACKNOWLEDGEMENTS

I am grateful to Ms Michelle Guy of the SiVEST Environmental Division as well as Mnr Wouter Fourie of PGS Heritage for commissioning this study as well as for providing the necessary project information. Ms Niki Mann of PGS is heartily thanked for helpful editorial input. Ms Marietjie Mostert of the farm Bloemendal very kindly shared her impressive knowledge of local Karoo vertebrate fossils. The landowners concerned with the WEF study area are thanked for access permission to their property. Finally, the companionship in the field, substantial palaeontological contributions as well as logistical backup kindly provided by Ms Madelon Tusenius are all highly appreciated, as always.

Declaration of Independence

I, John E. Almond, declare that I am an independent consultant and have no business, financial, personal or other interest in the proposed development, application or appeal in respect of which I was appointed other than fair remuneration for work performed in connection with the activity, application or appeal. There are no circumstances that compromise the objectivity of my performing such work.



Dr John E. Almond
Palaeontologist
Natura Viva cc

APPENDIX 1: JOHN ALMOND SHORT CV

Dr John Almond has an Honours Degree in Natural Sciences (Zoology) as well as a PhD in Palaeontology from the University of Cambridge, UK. He has been awarded post-doctoral research fellowships at Cambridge University and the University of Tübingen in Germany, and has carried out palaeontological research in Europe, North America, the Middle East as well as North and South Africa and Madagascar. For eight years he was a scientific officer (palaeontologist) for the Geological Survey / Council for Geoscience in the RSA. His current palaeontological research focuses on fossil record of the Precambrian - Cambrian boundary and the Cape Supergroup of South Africa. He has recently written palaeontological reviews for several 1: 250 000 geological maps published by the Council for Geoscience and has contributed educational material on fossils and evolution for new school textbooks in the RSA.

Since 2002 Dr Almond has also carried out numerous palaeontological impact assessments for developments and conservation areas in the Western, Eastern and Northern Cape, Limpopo, Northwest Province, Mpumalanga, Gauteng, KwaZulu-Natal and the Free State under the aegis of his Cape Town-based company *Natura Viva* cc. He has served as a member of the Archaeology, Palaeontology and Meteorites Committee for Heritage Western Cape (HWC) and an advisor on palaeontological conservation and management issues for the Palaeontological Society of South Africa (PSSA), HWC and SAHRA. He is currently compiling technical reports on the provincial palaeontological heritage of Western, Northern and Eastern Cape for SAHRA and HWC. Dr Almond is an accredited member of PSSA and APHP (Association of Professional Heritage Practitioners – Western Cape).

APPENDIX 2: KOUP 1 & 2 WEFS FOSSIL SITE DATA – JUNE 2021

All GPS readings were taken in the field using a hand-held Garmin GPSmap 64s instrument. The datum used is WGS 84.

Please note that:

- Locality data for South African fossil sites in *not* for public release, due to conservation concerns.
- The table does *not* represent all potential fossil sites within the project area but only those sites recorded during the 5-day field survey. The absence of recorded fossil sites in any area therefore does *not* mean that no fossils are present there.
- The detailed stratigraphic data for each site is provisional and has yet to be confirmed.

Loc.	GPS data	Comments
574	S32° 50' 42.7" E22° 26' 07.3"	Farm 1/380. Karelskraal Member. Isolated, small fragment of bone within cleaved massive grey mudrocks. Proposed Field Rating IIIC Local Resource. No mitigation required.
576	S32° 50' 43.3" E22° 26' 09.5"	Farm 1/380. Karelskraal Member. Thick, grey, sandy to finely gravelly semi-consolidated alluvium exposed in banks of stream showing range of subfossil insect burrows (probably termite) as well as plant root moulds. Proposed Field Rating IIIC Local Resource. No mitigation required.
577	S32° 50' 31.2" E22° 25' 48.8"	Farm 1/380. Karelskraal Member. Articulated skull and lower jaw of small dicynodont entirely enclosed within pedogenic calcrete concretion in float. Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
587	S32° 50' 02.3" E22° 24' 21.6"	Farm 8/380. Karelskraal Member. Float block of sandstone containing unidentifiable white postcranial bones of medium-sized tetrapod. Proposed Field Rating IIIC Local Resource. No mitigation required.
594	S32° 49' 52.9" E22° 23' 13.3"	Farm 8/380. Karelskraal Member. Shallow stream bed exposures of hackly, cleaved blue-grey mudrock with ferruginous carbonate concretions and <i>equivocal</i> , gently-inclined vertebrate burrow cast (c. 30 cm wide) in sandstone (requires confirmation). Proposed Field Rating IIIB Local Resource. Specimen to be recorded in more detail if it falls within or close to project footprint.
595	S32° 49' 53.4" E22° 23' 13.3"	As above. <i>Possible</i> second example of a large vertebrate burrow cast (c. 50 cm wide) (requires confirmation). Proposed Field Rating IIIB Local Resource. Specimen to be recorded in more detail if falls within or close to project footprint.
603	S32° 50' 56.8" E22° 25' 09.0"	Farm 1/380. Karelskraal Member. Grey-green mudrocks exposed alongside shallow stream bed with concentration of partially weathered-out, ferruginized, semi-articulated postcranial bones of a dinocephalian therapsid with trail of bone material extending into adjacent stream gravels. Majority of material is post-cranial (limb bones, vertebrae, ribs, girdle fragments <i>etc</i>) but several pieces are probably of cranial origin, including a badly-weathered jaw fragment with series of dinocephalian-type teeth. Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
606	S32° 50' 55.7" E22° 24' 57.0"	Farm 1/380. Karelskraal Member. Surface scatter of white, postcranial bone fragments of large-bodied tetrapod preserved within pedogenic concretions overlying cleaved grey-green mudrocks. Proposed Field Rating IIIC Local Resource. No mitigation required.
613	S32° 51' 09.8" E22° 24' 36.8"	Farm 1/380. Karelskraal Member. Pedogenic calcrete concretion in float containing series of vertebrae of a medium-sized tetrapod. Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
622	S32° 51' 00.2" E22° 23' 03.4"	Farm 1/380. Karelskraal Member. <i>Possible</i> horizontal, convex-upwards invertebrate burrow with grey-green hackly mudrocks. Proposed Field Rating IIIC Local Resource. No mitigation required.
623	S32° 51' 06.2" E22° 22' 53.1"	Farm 1/380. Karelskraal Member. Fragmentary postcranial remains, including end of large limb bone, of large-bodied tetrapod in float, probably weathered out of

		sandstone. Some elements show evidence of sun-cracking. Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
625	S32° 51' 06.1" E22° 22' 52.7"	Farm 1/380. Karelskraal Member. Further post-cranial fragments – probably of same animal as at Loc. 623. Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
630	S32° 51' 48.9" E22° 25' 56.3"	Farm 5/380. Karelskraal Member. Fragmentary postcranial remains, including vertebra and probably fragmentary limb bone, of large-bodied tetrapod in float. Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
631	S32° 51' 49.6" E22° 25' 59.9"	Farm 5/380. Karelskraal Member. Skull of small dicynodont (broad skull table – not <i>Diictodon</i>) preserved within pedogenic calcrete concretion, in float. Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
632	S32° 51' 57.8" E22° 25' 56.2"	Farm 5/380. Karelskraal Member. Fragment of highly weathered bone of medium / large tetrapod in float. Proposed Field Rating IIIC Local Resource. No mitigation required.
638	S32° 52' 19.1" E22° 25' 48.6"	Farm 11/380. Poortjie Member mudrock. Skull with lower jaw of small dicynodont preserved within pedogenic calcrete nodule in float (broad skull table – not <i>Diictodon</i>). Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
639	S32° 52' 20.6" E22° 25' 52.8"	Farm 11/380. Poortjie Member mudrock. Partially preserved skull of small dicynodont <i>in situ</i> , facing side-upwards with articulated lower jaw. Calcretised, flattened-cylindrical horizontal burrow cast (c. 4 cm wide) of possible invertebrate origin. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
640	S32° 52' 20.3" E22° 25' 51.9"	Farm 11/380. Poortjie Member mudrock. Incomplete snout of small dicynodont preserved in float. Proposed Field Rating IIIC Local Resource. No mitigation required.
641	S32° 52' 19.9" E22° 25' 49.7"	Farm 11/380. Poortjie Member mudrock. Weathered-out pedogenic calcrete concretions containing skull of small dicynodont (probably <i>Diictodon</i>) as well as incomplete postcrania of similar animal. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
642	S32° 52' 19.4" E22° 25' 48.1"	Farm 11/380. Poortjie Member mudrock. Weathered-out, ferruginized pedogenic calcrete concretions containing skull of small dicynodont (probably <i>Diictodon</i>) as well as postcranial bones of similar animal. Proposed Field Rating IIIC Local Resource. No mitigation required.
643	S32° 52' 19.3" E22° 25' 47.6"	Farm 11/380. Poortjie Member mudrock. Several poorly-preserved skulls and postcranial bones of small dicynodonts preserved within pedogenic calcrete concretions. Proposed Field Rating IIIC Local Resource. No mitigation required.
644	S32° 52' 19.8" E22° 25' 46.6"	Farm 11/380. Poortjie Member mudrock. Float skull specimen of small dicynodont with articulated lower jaw. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
645	S32° 52' 20.8" E22° 25' 47.9"	Farm 11/380. Poortjie Member mudrock. Poorly-preserved, ferruginized skull specimen of small dicynodont with articulated lower jaw in float. Proposed Field Rating IIIC Local Resource. No mitigation required.
646	S32° 52' 21.2" E22° 25' 47.7"	Farm 11/380. Poortjie Member mudrock. Small bone fragment of medium-sized or larger tetrapod in float. Field Rating IIIC Local Resource. No mitigation required.
647	S32° 52' 23.2" E22° 25' 55.4"	Farm 5/380. Poortjie Member mudrock. Calcretised sandstone cast (c. 5 cm wide) of small, gently inclined burrow – possibly of small tetrapod or invertebrate – preserved within grey mudrock. Proposed Field Rating IIIB Local Resource. Specimen to be collected / recorded if falls within or close to project footprint.
648	S32° 52' 23.4" E22° 25' 56.0"	Farm 11/380. Poortjie Member mudrock. Gently inclined sandstone cast of small tetrapod or invertebrate burrow c. 8 cm wide with smooth, possibly bioturbated interior. Proposed Field Rating IIIB Local Resource. Specimen to be collected / recorded if falls within or close to project footprint.
649	S32° 52' 19.9" E22° 25' 55.6"	Farm 11/380. Poortjie Member mudrock. Well-preserved <i>in situ</i> skull of small dicynodont (broad skull table – not <i>Diictodon</i>) preserved within pedogenic calcrete concretion in grey-green mudrocks. Proposed Field Rating IIIB Local Resource. Specimen to be collected / recorded if falls within or close to project footprint
650	S32° 52' 02.6" E22° 25' 40.8"	Farm 11/380. Karelskraal Member. Poorly-preserved, incomplete snout of small dicynodont within sandstone float block. Proposed Field Rating IIIC Local Resource. No mitigation required.
653	S32° 52' 06.0"	Farm 10/380. Karelskraal Member. Poorly-preserved skull of small dicynodont

	E22° 24' 20.9"	within pedogenic calcrete concretion in float. Proposed Field Rating IIIC Local Resource. No mitigation required.
657	S32° 51' 35.5" E22° 27' 53.7"	Farm 5/380. Karelskraal Member. Sandstone float block containing white, weathered indeterminate bone of large tetrapod. Probable separate rib fragment enclosed within calcrete concretion. Proposed Field Rating IIIC Local Resource. No mitigation required.
660	S32° 51' 56.0" E22° 27' 27.4"	Farm 5/380. Karelskraal Member. Surface scatter of numerous white fragments of postcranial bone of large-bodied tetrapod, some showing extensive sun-cracking. Proposed Field Rating IIIC Local Resource. No mitigation required.
665	S32° 51' 42.1" E22° 28' 17.3"	Farm 5/380. Karelskraal Member. Dispersed postcranial bones of large tetrapod (dinocephalian / pareiasaur) embedded <i>in situ</i> within grey-green mudrocks as well as weathered-out into neighbouring stream gravels. Material includes several large limb bones and vertebrae. Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
668	S32° 51' 46.1" E22° 28' 24.9"	Farm 5/380. Karelskraal Member. Ferruginous carbonate concretion within grey-green mudrocks containing several ribs of medium-sized tetrapod. Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
670	S32° 51' 31.9" E22° 28' 25.5"	Farm 5/380. Karelskraal Member. Float block of pedogenic calcrete concretion containing bone fragments of large tetrapod (possibly ribs or vertebral neural spines). Proposed Field Rating IIIC Local Resource. No mitigation required.
675	S32° 51' 58.1" E22° 29' 18.3"	Farm 5/380. Karelskraal Member. Thin channel sandstone containing concentration of disarticulated postcranial bones of large-bodied tetrapod, with numerous fragments of white bone in sandstone within colluvial gravels downslope. Probably unidentifiable. Proposed Field Rating IIIC Local Resource. No mitigation required.
684	S32° 51' 55.2" E22° 31' 46.8"	Farm 11/374. Poortjie Member. Small dicynodont skull within pedogenic calcrete concretion. Proposed Field Rating IIIB Local Resource. Specimen to be collected if falls within or close to project footprint.
687	S32° 51' 56.2" E22° 29' 55.7"	Farm 11/374. Poortjie Member. Ferruginous pedogenic calcrete concretion within cleaved grey-green mudrocks containing post-crania (limb bones, ribs <i>etc</i>) of small tetrapod. V. close by is <i>in situ</i> skull of small dicynodont as well as float block of ferruginous carbonate (2 parts) with incomplete skull of larger dicynodont. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
688	S32° 51' 58.0" E22° 29' 55.0"	Farm 11/374. Poortjie Member. Sandstone cast of small tetrapod burrow (possibly dicynodont), c. 10 cm wide, preserved within cleaved grey-green mudrock. Proposed Field Rating IIIC Local Resource. No mitigation required.
689	S32° 51' 57.6" E22° 29' 54.7"	Farm 11/374. Poortjie Member. <i>In situ</i> skull of small dicynodont within cleaved grey-green mudrocks. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
690	S32° 51' 57.6" E22° 29' 54.7"	Farm 11/374. Poortjie Member. <i>In situ</i> skull of small dicynodont within cleaved grey-green mudrocks. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
691	S32° 50' 32.0" E22° 26' 23.0"	Farm 6/380 (just outside and N of WEF project area). Karelskraal or Poortjie Member. <i>In situ</i> postcranial remains of a large tetrapod preserved within channel sandstone (Marietjie Mostert., pers. comm., June 2021). No mitigation requires (outside project area).
693	S32° 51' 16.0" E22° 28' 46.7"	Farm 5/ 380. Karelskraal Member. 4 skulls of small dicynodonts preserved within pedogenic calcrete concretions weathered out into gully gravels. Possibly not all <i>Diictodon</i> . Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
696	S32° 50' 41.8" E22° 29' 20.5"	Farm 5/ 380. Poortjie Member. <i>In situ</i> small dicynodont skull preserved <i>in situ</i> , dorsal side-up, within pedogenic calcrete concretion (broad skull table – not <i>Diictodon</i>). Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
697	S32° 50' 41.6" E22° 29' 21.4"	Farm 5/ 380. Poortjie Member. Float pedogenic calcrete concretions with small dicynodont skull, postcrania of similar animal. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
698	S32° 50' 41.6" E22° 29' 21.6"	Farm 5/ 380. Poortjie Member. Cluster of several pedogenic calcrete concretions in float with skulls and postcrania of small dicynodonts. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
699	S32° 50' 41.7" E22° 29' 21.8"	Farm 5/ 380. Poortjie Member. Cluster of several pedogenic calcrete concretions in float with skulls, postcrania of small dicynodonts (some with broad skull tables). Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within

		or close to project footprint.
700	S32° 50' 41.4" E22° 29' 22.3"	Farm 5/ 380. Poortjie Member. Cluster of several pedogenic calcrete concretions in float with skulls, postcrania of small dicynodonts (some with broad skull tables). Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
701	S32° 50' 41.3" E22° 29' 22.4"	Farm 5/ 380. Poortjie Member. Pedogenic calcrete concretions in float with poorly-preserved skulls of small dicynodonts (some with broad skull tables). Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
702	S32° 50' 41.5" E22° 29' 22.5"	Farm 5/ 380. Poortjie Member. Pedogenic calcrete concretion with postcranial bones of small tetrapod. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
703	S32° 50' 41.2" E22° 29' 22.9"	Farm 5/ 380. Poortjie Member. Poorly-preserved bones of small tetrapod(s) within pedogenic calcrete concretions. Proposed Field Rating IIIC Local Resource. No mitigation required.
704	S32° 50' 41.8" E22° 29' 24.8"	Farm 5/ 380. Poortjie Member. Skulls and postcrania of small tetrapods / dicynodonts within pedogenic calcrete concretions. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
705	S32° 50' 42.2" E22° 29' 25.5"	Farm 5/ 380. Poortjie Member. <i>In situ</i> skull of small dicynodont within pedogenic calcrete concretion. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
706	S32° 50' 42.2" E22° 29' 25.5"	Farm 5/ 380. Poortjie Member. <i>Several in situ</i> skulls of small dicynodonts. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
713	S32° 50' 46.7" E22° 29' 50.8"	Farm 5/ 380. Poortjie Member. Pedogenic calcrete concretion in float with small dicynodont skull. Proposed Field Rating IIIB Local Resource. Specimens to be collected if fall within or close to project footprint.
724	S32° 50' 09.8" E22° 25' 22.0"	Farm 380 Bloemendal (outside and just N of WEF project area). Mudrock sandstone interval within Poortjie Member (as mapped, but requires checking). Several very thick-boned skull fragments as well as fragmentary postcranial remains of tapinocephalid dinocephalians (probably several individuals, perhaps including juveniles), probably all belonging to the genus <i>Criocephalosaurus</i> . Proposed Field Rating IIIA Local Resource. No mitigation recommended because specimens lie <i>outside</i> project footprint.




Figure A2.1: Google Earth© satellite image of the adjoining Koup 1 WEF (orange polygon) and Koup 2 WEF (yellow polygon) project areas to the south of Beaufort West showing the tracks (blue lines) and numbered waypoints of localities visited during the 5-day palaeontological site visit. Due to a GPS technical error, no waypoints were recorded for fieldwork in the north-eastern sector of the area (Fortunately this area is largely of low palaeosensitivity). The sites visited were largely determined by access and the presence of good bedrock exposure. Substantial portions of the project area (including most of the provisional WEF and grid connection footprints) have not yet been surveyed.




Figure A2.2: Google Earth© satellite image of the adjoining Koup 1 WEF (orange polygon) and Koup 2 WEF (yellow polygon) project areas to the south of Beaufort West showing the numbered new fossil sites recorded during the site visit (Please see Appendix 1 for GPS locality details and brief description). These are necessarily only a small subset of the fossil sites within the project areas; blank areas may well also contain important fossil remains.

APPENDIX 3: SITE SENSITIVITY VERIFICATION REPORT (IN TERMS OF PART A OF THE ASSESSMENT PROTOCOLS PUBLISHED IN GN 320 ON 20 MARCH 2020)

1. Introduction

Genesis Enertrag Koup 2 Wind (Pty) Ltd is proposing to construct the Koup 2 WEF, comprising thirty-two wind turbines with a maximum total energy generation capacity of up to 140MW, with a 132kV overhead power line connection to the national grid. A Battery Energy Storage System (BESS) will be located next to the onsite 33/132kV substation. The WEF and grid project areas are located in the Great Karoo region some 60 km south of Beaufort West, falling within the Beaufort West Local Municipality (Central Karoo District Municipality) of the Western Cape Province.

In accordance with Appendix 6 of the National Environmental Management Act (Act 107 of 1998, as amended) (NEMA) Environmental Impact Assessment (EIA) Regulations of 2014, a site sensitivity verification has been undertaken in order to confirm the current land use and environmental sensitivity of the proposed project area as identified by the National Web-Based Environmental Screening Tool (Screening Tool).

2. Site sensitivity verification

The site sensitivity verification of the proposed Koup 2 WEF and associated grid connection is based on:

- A desktop review of (a) the relevant 1:50 000 scale topographic maps and the 1:250 000 scale topographic map 3222 Beaufort West, (b) Google Earth© satellite imagery, (c) published geological and palaeontological literature, including 1:250 000 geological maps (3222 Beaufort West) and relevant sheet explanations (Johnson & Keyser 1979) as well as (d) several previous and on-going fossil heritage (PIA) assessments in the Great Karoo region to the south of Beaufort West by the author .
- A five-day field assessment of the combined Koup 1 and Koup 2 WEF project area by the author and an experienced field assistant during the period 14 to 19 November 2020. Accessible sectors of the Grid Connection project area within the combined WEF project areas were surveyed in part, but *not* those sectors lying outside the WEF project area itself.

3. Outcome of site sensitivity verification

The proposed Koup 2 WEF and grid connection project area is located in a region of the Great Karoo that is underlain by potentially fossiliferous sedimentary rocks of Late Palaeozoic and younger, Late Tertiary or Quaternary, age. In particular, these include (1) Middle to Late Permian continental sediments of the Abrahamskraal and Teekloof Formations (Lower Beaufort Group, Karoo Supergroup) which contain scientifically important fossils of vertebrates, trace fossils and terrestrial plants, as well as (2) Late Caenozoic alluvium that may contain important mammalian remains such as teeth and bones (These rock units and their fossils are described in more detail in Section 5 of the PIA report).

The generally high palaeontological heritage sensitivity of the Beaufort Group bedrocks in the Great Karoo is emphasized on the SAHRIS palaeosensitivity map maintained by SAHRA. Both desktop and field studies within this and neighbouring WEF project areas demonstrate that, while a significant number of scientifically valuable, well-preserved fossils do indeed occur here, in practice they are often scarce in this region and their distribution is to a large extent unpredictable. As a consequence of the considerably higher topographic relief – and hence bedrock exposure levels - encountered within the combined Koup 2 WEF and grid connection project area, a higher concentration of fossil sites is recorded here than within WEF project areas towards the east.

It is concluded that, applying the precautionary principle, the Koup 2 WEF and grid connection project areas are best assigned an overall High Sensitivity as far as palaeontological heritage is concerned, while recognising that, in practice, fossils are scarce at surface throughout much of the area (*N.B.* Additional fossils are preserved in the subsurface and may be impacted by excavations during the construction phase).

4. National Environmental Screening Tool

The palaeontological heritage Site Sensitivity Map for the Koup 2 WEF project area prepared using the DFFE screening tool identifies areas underlain by the Lower Beaufort Group as being of a High Sensitivity while those mantled by substantial alluvial deposits are assigned a Low Sensitivity (Fig. A3.1).

Due to the scarcity of well-preserved, scientifically important fossils in this region, based on desktop studies and fieldwork, it is inferred herein that the much of project area is in fact largely of Low palaeontologically sensitivity but with a substantial number of dispersed and unpredictable fossil sites of High to Very High sensitivity. Applying the precautionary principle, the Koup 2 WEF and grid connection project areas are best assigned an overall High Sensitivity as far as palaeontological heritage is concerned. The DFFE screening tool sensitivity map in Figure A3.1 is therefore supported in essence.

5. Conclusion

The palaeontological heritage site sensitivity of the combined Koup 2 WEF and associated grid connection project areas has been verified on the basis of desktop studies as well as a 5-day site visit. Applying the Precautionary Principle, an overall High Palaeontological Sensitivity is inferred for the WEF and grid connection project areas.




Figure A3.1: Paleontological sensitivity map for the Koup 2 WEF project area. Due to the scarcity of well-preserved, scientifically important fossils in this region, based on desktop studies and fieldwork, it is inferred herein that much of the project area is in fact largely of LOW palaeontological sensitivity but with a substantial number of dispersed and unpredictable fossil sites of HIGH to VERY HIGH sensitivity. Applying the precautionary principle, the Koup 2 WEF and grid connection project area are best assigned an overall High Sensitivity as far as palaeontological heritage is concerned.

APPENDIX 4: CHANCE FOSSIL FINDS PROTOCOL

KOUP 1 & 2 WIND ENERGY FACILITIES and GRID CONNECTIONS south of Beaufort West	
Province & region:	Western Cape (Central Karoo District): Beaufort West and Prince Albert Local Municipalities
Responsible Heritage Resources Agency	Heritage Western Cape (Contact details: Heritage Western Cape. 3 rd Floor Protea Assurance Building, 142 Longmarket Street, Green Market Square, Cape Town 8000. Private Bag X9067, Cape Town 8001. Tel: 021 483 5959 Email: ceoheritage@westerncape.gov.za)
Rock unit(s)	Abrahamskraal & Teekloof Formations (Lower Beaufort Group), Late Caenozoic alluvium
Potential fossils	Fossil vertebrate bones, teeth, trace fossils, trackways, petrified wood, plant-rich beds in the Lower Beaufort Group bedrocks. Fossil mammal bones, teeth, horn cores, freshwater molluscs, plant material in Late Caenozoic alluvium.
ECO protocol	1. Once alerted to fossil occurrence(s): alert site foreman, stop work in area immediately (<i>N.B.</i> safety first!), safeguard site with security tape / fence / sand bags if necessary.
	2. Record key data while fossil remains are still <i>in situ</i> : <ul style="list-style-type: none"> • Accurate geographic location – describe and mark on site map / 1: 50 000 map / satellite image / aerial photo • Context – describe position of fossils within stratigraphy (rock layering), depth below surface • Photograph fossil(s) <i>in situ</i> with scale, from different angles, including images showing context (<i>e.g.</i> rock layering)
	3. If feasible to leave fossils <i>in situ</i> : <ul style="list-style-type: none"> • Alert Heritage Resources Agency and project palaeontologist (if any) who will advise on any necessary mitigation • Ensure fossil site remains safeguarded until clearance is given by the Heritage Resources Agency for work to resume
	3. If <i>not</i> feasible to leave fossils <i>in situ</i> (emergency procedure only): <ul style="list-style-type: none"> • <i>Carefully</i> remove fossils, as far as possible still enclosed within the original sedimentary matrix (<i>e.g.</i> entire block of fossiliferous rock) • Photograph fossils against a plain, level background, with scale • Carefully wrap fossils in several layers of newspaper / tissue paper / plastic bags • Safeguard fossils together with locality and collection data (including collector and date) in a box in a safe place for examination by a palaeontologist • Alert Heritage Resources Agency and project palaeontologist (if any) who will advise on any necessary mitigation
	4. If required by Heritage Resources Agency, ensure that a suitably-qualified specialist palaeontologist is appointed as soon as possible by the developer.
	5. Implement any further mitigation measures proposed by the palaeontologist and Heritage Resources Agency
Specialist palaeontologist	Record, describe and judiciously sample fossil remains together with relevant contextual data (stratigraphy / sedimentology / taphonomy). Ensure that fossils are curated in an approved repository (<i>e.g.</i> museum / university / Council for Geoscience collection) together with full collection data. Submit Palaeontological Mitigation report to Heritage Resources Agency. Adhere to best international practice for palaeontological fieldwork and Heritage Resources Agency minimum standards.

APPENDIX 5: SiVEST SPECIALIST TERMS OF REFERENCE

Specialist Assessment Reports / Compliance Statements

Specialists are requested to provide *one (1)* scoping phase report and / or compliance statement that provides an assessment of the proposed Koup 2 WEF *and* the associated grid connection infrastructure (132kV overhead power line on-site switching / collector substation). The report should however include separate assessment and impact rating chapters/sections for the WEF and the grid connection proposals respectively.

During the EIA phase, specialists will be required to update the scoping phase specialist report to provide a review of their findings in accordance with revised site layouts and to address any comments or concerns arising from the public participation process.

The specialist assessment reports and / or compliance statements should include the following sections:

8.2.1 Project Description

The specialist report must include the project description as provided by SiVEST.

8.2.2 Terms of Reference

The specialist report must include an explanation of the terms of reference (TOR) applicable to the specialist study. The gazetted Environmental Assessment Protocols of the NEMA EIA Regulations (2014, as amended), prescribes Procedures for the Assessment and Minimum Criteria for Reporting on the Identified Environmental Themes in terms of Sections 24(5)(A) and (H) and 44 of the National Environmental Management Act, 1998. These procedures must be considered.

Where a specialist assessment is required and no specific environmental theme protocol has been prescribed, the required level of assessment must be based on the findings of the site sensitivity verification and must comply with Appendix 6 of the EIA Regulations; and any relevant legislation and guidelines deemed necessary

Where relevant, a table must be provided at the beginning of the specialist report, listing the requirements for specialist reports in accordance with Appendix 6 of the EIA Regulations, 2014 (as amended) and cross referencing these requirements with the relevant sections in the report.

8.2.3 Legal Requirements and Guidelines

The specialist report must include a thorough overview of all applicable best practice guidelines, relevant legislation, prescribed Assessment Protocols and authority requirements.

8.2.4 Methodology

The report must include a description of the methodology applied in carrying out the specialist assessment.

8.2.5 Specialist Findings / Identification of Impacts

The report must present the findings of the specialist studies and explain the implications of these findings for the proposed development (e.g. permits, licenses etc.). This section of the report should also identify any sensitive and/or 'no-go' areas on the development site or within the power line assessment corridors. These areas must be mapped clearly with a supporting explanation provided.

This section of the report should also specify if any further assessment will be required.

8.2.6 Environmental Impact Assessment

The impacts (both direct and indirect) of the proposed WEF and the proposed grid connection infrastructure (during the Construction, Operation and Decommissioning phases) are to be assessed and rated *separately* according to the methodology developed by SiVEST. Specialists will be required to make use of the impact rating matrix provided (in Excel format) for this purpose, and *separate tables* must be provided for the WEF and for the grid connection infrastructure respectively. Please note that the significance of Cumulative Impacts should also be rated in this section. Both the methodology and the rating matrix will be provided by SiVEST.

Please be advised that this section must include mitigation measures aimed at minimising the impact of the proposed development.

8.2.7 Input To The Environmental Management Programme (EMPr)

The report must include a description of the key monitoring recommendations for each applicable mitigation measure identified for each phase of the project for inclusion in the Environmental Management Programme (EMPr) or Environmental Authorisation (EA).

Please make use of the Impact Rating Table (in Excel format) for each of the phases i.e. Design, Construction, Operation and Decommissioning.

8.2.8 Cumulative Impact Assessment

Cumulative impact assessments must be undertaken for the proposed WEF and associated grid connection infrastructure to determine the cumulative impact that will materialise if other Renewable Energy Facilities (REFs) and large scale industrial developments are constructed within 35kms of the proposed development.

The cumulative impact assessment must contain the following:

- A cumulative environmental impact statement noting whether the overall impact is acceptable; and
- A review of the specialist reports undertaken for other REFs and an indication of how the recommendations, mitigation measures and conclusion of the studies have been considered.

In order to assist the specialists in this regard, SiVEST will provide the following documentation/data:

- A summary table listing all REFs identified within 35kms of the proposed WEF;

It should be noted that it is the specialist's responsibility to source the relevant EIA / BA reports that are available in the public domain. SiVEST will assist, where possible.

8.2.9 No Go Alternative

Consideration must be given to the “no-go” option in the EIA process. The “no-go” option assumes that the site remains in its current state, *i.e.* there is no construction of a WEF and associated infrastructure in the proposed project area and the status quo would be preserved.

8.2.10 Comparative Assessment Of Alternatives

As mentioned, alternatives for the Substation location, construction / laydown area and power line route alignment have been identified. These alternatives are being considered as part of the EIA / BA processes and as such specialists are required to undertake a comparative assessment of the alternatives mentioned above as per the latest table provided by SiVEST.

8.2.11 Conclusion / Impact Statement

The conclusion section of the specialist report must include an Impact Statement, indicating whether any fatal flaws have been identified and ultimately whether the proposed development can be authorised or not (*i.e.* whether EA should be granted / issued or not).

8.2.12 Executive Summary

Specialists must provide an Executive Summary summarising the findings of their report to allow for easy inclusion in the EIA / BA reports.

8.2.13 Specialist Declaration of Independence

A copy of the Specialist Declaration of Interest (DoI) form, containing original signatures, must be appended to all Draft and Final Reports. This form will be provided to the specialists. *Please note that the undertaking / affirmation under oath section of the report must be signed by a Commissioner of Oaths.*