

**PALAEONTOLOGICAL DESKTOP
ASSESSMENT FOR THE KWAGGA NORTH
PROJECT: AN EXTENSION OF THE OPTIMUM
COLLIERY NEAR THE TOWN OF ARNOT,
STEVE TSHWETE LOCAL MUNICIPALITY,
NKANGALA DISTRICT MUNICIPALITY,
MPUMALANGA**

For:

HIA CONSULTANTS



DATE: 7 JULY 2013

By

GIDEON GROENEWALD

EXECUTIVE SUMMARY

Gideon Groenewald was appointed by PSG Heritage to undertake a desktop survey, assessing the potential palaeontological impact of the proposed Kwagga North project at Optimum Coal Mine outside the town of Arnot, Steve Tshwete Local Municipality, Nkangala District Municipality, Mpumalanga Province.

This report forms part of the Environmental Impact Assessment and complies with the requirements of the South African National Heritage Resource Act No 25 of 1999. In accordance with Section 38 (Heritage Resources Management), a Heritage Impact Assessment (HIA) is required to assess any potential impacts to palaeontological heritage within the footprint of the development.

The Kwagga North Section is an extension of Optimum Colliery into reserves previously owned and mined by AngloCoal. The proposed mining layout of the Kwagga North Project region can be seen in Figure 1.6. Arnot Colliery, mined by AngloCoal and the neighbouring Kwagga North Section, was contracted to supply bituminous coal to Eskom's 2100 MW Arnot Power station at a rate of up to 6,18 million tons per annum.

The study area is mainly underlain by sedimentary rocks of the Permian Ecca Group, Karoo Supergroup. The Northern section of the study area is underlain by Vaalian-aged volcanic rocks of the Damwal Formation of the Rooiberg Group.

The Vryheid Formation is well-known for the occurrence of coal beds that resulted from the accumulation of plant material over long periods of time. Plant fossils described by Bamford (2011) from the Vryheid Formation are; *Azaniodendron fertile*, *Cyclodendron leslii*, *Sphenophyllum hammanskraalensis*, *Annularia sp.*, *Raniganjia sp.*, *Asterotheca spp.*, *Liknopetalon enigmata*, *Glossopteris > 20 species*, *Hirsutum 4 spp.*, *Scutum 4 spp.*, *Ottokaria 3 spp.*, *Estcourtia sp.*, *Arberia 4 spp.*, *Lidgettonia sp.*, *Noeggerathiopsis sp.* and *Podocarpidites sp.* Although no vertebrate fossils have been recorded from the Vryheid Formation, invertebrate trace fossils have been described in some detail. Due to its age and igneous character the Damwal Formation will not contain fossils.

The sedimentary rocks of the Vryheid Formation will contain fossils. Although these rock sequences will have a high palaeontological sensitivity, the chances of finding fossils before actual mining of the rocks takes place is very low. For this reason a medium palaeontological sensitivity is allocated to the areas underlain by the Vryheid Formation. Due to the igneous nature of the Damwal Formation, the rocks will not contain fossils and a low palaeontological sensitivity is allocated to these rocks.

It is recommended that:

1. The developer and the ECO of the mining project are made aware of the fact that coal mining is by definition the mining of fossil plant material.
2. That the developer applies for a destruction permit for plant fossils encountered during the mining operation.
3. That the developer should look out for exceptionally well preserved fossils and appoint a palaeontologist to collect representative samples of these fossils for further study at an appropriate institute, such as the Bernard Price Institute for Palaeontology at WITS University.

TABLE OF CONTENT

1. INTRODUCTION.....	1
1.1. Background.....	1
1.2. Aims and Methodology.....	1
1.3. Scope and Limitations of the Desktop Study.....	2
2. DESCRIPTION OF THE PROPOSED DEVELOPMENT.....	3
3. GEOLOGY.....	3
3.1. Ecca Group:.....	3
3.2. Damwal Formation:.....	4
4. PALAEOLOGY OF THE AREA.....	4
4.1. Ecca Group:.....	4
4.1.1. Vryheid Formation.....	4
4.2. Damwal Formation:.....	5
5. PALAEOLOGICAL SENSITIVITY.....	6
6. CONCLUSION AND RECOMMENDATIONS.....	7
7. REFERENCES.....	7
8. QUALIFICATIONS AND EXPERIENCE OF THE AUTHOR.....	8
9. DECLARATION OF INDEPENDENCE.....	8

LIST OF FIGURES

Figure 2.1 Location of the Kwagga North Project, Optimum Coal Mine.....	3
Figure 3.1 Map showing the geology of the study area.....	4
Figure 5.1 Palaeontological sensitivity of the study area.....	6

LIST OF TABLES

Table 1.1 Palaeontological Sensitivity Analysis Outcome Classification.....	2
---	---

1. INTRODUCTION

1.1. Background

Gideon Groenewald was appointed by PSG Heritage to undertake a desktop survey, assessing the potential palaeontological impact of the proposed Kwagga North project, Optimum Coal Mine, outside the town of Arnot, Steve Tshwete Local Municipality, Nkangala District Municipality, Mpumalanga Province.

This report forms part of the Environmental Impact Assessment and complies with the requirements of the South African National Heritage Resource Act No 25 of 1999. In accordance with Section 38 (Heritage Resources Management), a Heritage Impact Assessment (HIA) is required to assess any potential impacts to palaeontological heritage within the footprint of the development.

Categories of heritage resources recognised as part of the National Estate in Section 3 of the Heritage Resources Act, and which therefore fall under its protection, include:

- geological sites of scientific or cultural importance;
- objects recovered from the soil or waters of South Africa, including archaeological and palaeontological objects and material, meteorites and rare geological specimens;
- objects with the potential to yield information that will contribute to an understanding of South Africa's natural or cultural heritage.

1.2. Aims and Methodology

Following the *"SAHRA APM Guidelines: Minimum Standards for the Archaeological & Palaeontological Components of Impact Assessment Reports"*, the aims of the palaeontological impact assessment are:

- to identify exposed and subsurface rock formations that are considered to be palaeontologically significant;
- to assess the level of palaeontological significance of these formations;
- to comment on the impact of the development on these exposed and/or potential fossil resources; and
- to make recommendations as to how the developer should conserve or mitigate damage to these resources.

In preparing a palaeontological desktop study the potential fossiliferous rock units (groups, formations, etc) represented within the study area are determined from geological maps. The known fossil heritage within each rock unit is inventoried from the published scientific literature and previous palaeontological impact studies in the same region.

The likely impact of the proposed development on local fossil heritage is determined on the basis of the palaeontological sensitivity of the rock units concerned and the nature and scale of the development itself, most notably the extent of fresh bedrock excavation envisaged. The different sensitivity classes used are explained in Table 1.1 below.

Table 1.1 Palaeontological Sensitivity Analysis Outcome Classification

Sensitivity	Description
Low Sensitivity	Areas where a negligible impact on the fossil heritage is likely. This category is reserved largely for areas underlain by igneous rocks. However, development in fossil bearing strata with shallow excavations or with deep soils or weathered bedrock can also form part of this category.
Moderate Sensitivity	Areas where fossil bearing rock units are present, but fossil finds are localised or within thin or scattered sub-units. Pending the nature and scale of the proposed development, the chances of finding fossils are moderate. A field-based assessment by a professional palaeontologist is usually warranted.
High Sensitivity	Areas where fossil bearing rock units are present with a very high possibility of finding fossils of a specific assemblage zone. Fossils will most probably be present in all outcrops and the chances of finding fossils during a field-based assessment by a professional palaeontologist are very high. Palaeontological mitigation measures need to be incorporated into the Environmental Management Plan.

1.3. Scope and Limitations of the Desktop Study

The study will include: i) an analysis of the area’s stratigraphy, age and depositional setting of fossil-bearing units; ii) a review of all relevant palaeontological and geological literature, including geological maps, and previous palaeontological impact reports; iii) data on the proposed development provided by the developer (e.g. location of footprint, depth and volume of bedrock excavation envisaged); and iv) where feasible, location and examination of any fossil collections from the study area (e.g. museums).

The key assumption for this scoping study is that the existing geological maps and datasets used to assess site sensitivity are correct and reliable. However, the geological maps used were not intended for fine scale planning work and are largely based on aerial photographs alone, without ground-truthing. There is also an inadequate database for fossil heritage for much of the RSA, due to the small number of professional palaeontologists carrying out fieldwork in the RSA. Most development study areas have never been surveyed by a palaeontologist.

These factors may have a major influence on the assessment of the fossil heritage significance of a given development and, without supporting field assessments, may lead to either:

- an underestimation of the palaeontological significance of a given study area, due to ignorance of significant recorded or unrecorded fossils preserved there, or
- an overestimation of the palaeontological sensitivity of a study area, for example when originally rich fossil assemblages inferred from geological maps have in fact been destroyed by tectonism or weathering, or are buried beneath a thick mantle of unfossiliferous “drift” (soil, alluvium etc.).

2. DESCRIPTION OF THE PROPOSED DEVELOPMENT

The Kwagga North Section is an extension of Optimum Colliery into reserves previously owned and mined by AngloCoal. The proposed mining layout of the Kwagga North Project region can be seen in Figure 1.6. Arnot Colliery, mined by AngloCoal and neighbouring the Kwagga North Section, was contracted to supply bituminous coal to Eskom's 2100 MW Arnot Power station at a rate of up to 6,18 million tons per annum.

The mining method used is opencast, using draglines combined with truck and shovel for boxcut areas where necessary.



Figure 2.1 Location of the Kwagga North Project, Optimum Coal Mine.

3. GEOLOGY

The study area is mainly underlain by sedimentary rocks of the Permian Ecca Group, Karoo Supergroup. The Northern section of the study area is underlain by Vaalian-aged volcanic rocks of the Damwal Formation of the Rooiberg Group (Figure 3.1).

3.1. Ecca Group:

The study area is underlain by sedimentary rocks of the Ecca Group (Pe, Figure 3.1). Although not mapped out on the geological map (Figure 3.1), the Ecca Group consists of a lower Pietermaritzburg Formation (shale), middle Vryheid Formation (Grey sandstone with prominent coal beds) and an upper Volksrust Formation (shale). Due to the fact that most of the study area refers to a potential

mining area, it can be concluded that the site is underlain by rocks of the Vryheid Formation, consisting mainly of sandstone, grit, sandstone, conglomerate, shale and coal.

3.2. Damwal Formation:

Rocks of the Damwal Formation consist of glassy amygdaloidal pseudospherulitic and porphyritic black rhyolite, red granophyric rhyolite, black rhyolite and leptite.




Figure 3.1 Map showing the geology of the study area

4. PALAEOLOGY OF THE AREA

4.1. Ecca Group:

4.1.1. Vryheid Formation

The Vryheid Formation is well-known for the occurrence of coal beds that resulted from the accumulation of plant material over long periods of time. Plant fossils described by Bamford (2011) from the Vryheid Formation are; *Azaniodendron fertile*, *Cyclodendron leslii*, *Sphenophyllum hammanskraalensis*, *Annularia sp.*, *Raniganja sp.*, *Asterotheca spp.*, *Liknopetalon enigmata*,

Glossopteris > 20 species, *Hirsutum* 4 spp., *Scutum* 4 spp., *Ottokaria* 3 spp., *Estcourtia* sp., *Arberia* 4 spp., *Lidgettonia* sp., *Noeggerathiopsis* sp. and *Podocarpidites* sp.

According to Bamford (2011), little data has been published on these potentially fossiliferous deposits. Around the coalmines there is most likely to be good material and yet in other areas the exposures may be too poor to be of interest. When they do occur, fossil plants are usually abundant and it would not be feasible to preserve and maintain all the sites, however, in the interests of heritage and science such sites should be well recorded, sampled and the fossils kept in a suitable institution.

Although no vertebrate fossils have been recorded from the Vryheid Formation, invertebrate trace fossils have been described in some detail by Mason and Christie (1986). It should be noted, however, that the aquatic reptile, *Mesosaurus*, which is the earliest known reptile from the Karoo Basin, as well as fish (*Palaeoniscus capensis*), have been recorded in equivalent-aged strata in the Whitehill Formation in the southern part of the basin (MacRae, 1999). Indications are that the Whitehill Formation in the main basin might be correlated with the mid-Vryheid Formation. If this assumption proves correct, there is a possibility that *Mesosaurus* could be found in the Vryheid Formation.

The late Carboniferous to early Jurassic Karoo Supergroup of South Africa includes economically important coal deposits within the Vryheid Formation of Natal. The Karoo sediments are almost entirely lacking in body fossils but ichnofossils (trace fossils) are locally abundant. Modern sedimentological and ichnofaunal studies suggest that the north-eastern part of the Karoo basin was marine. In KwaZulu-Natal a shallow basin margin accommodated a prograding fluviodeltaic complex forming a broad sandy platform on which coal-bearing sediments were deposited. Ichnofossils include U-burrows (formerly *Corophioides*) which are assigned to ichnogenus *Diplocraterion* (Mason and Christie, 1986).

4.2. Damwal Formation:

Due to the age and igneous nature of the rocks, there will be no fossils in this formation.

5. PALAEOLOGICAL SENSITIVITY




Figure 5.1 Palaeontological sensitivity of the study area

The sedimentary rocks of the Vryheid Formation will contain fossils. Although these rock sequences will have a high palaeontological sensitivity, the chances of finding fossils before actual mining of the rocks takes place is very low. For this reason, a medium palaeontological sensitivity is allocated to the areas underlain by the Vryheid Formation (Figure 5.1). The mining of coal will by definition be the mining of fossil plant material and it is recommended that the developer apply for a destruction permit for all fossil material encountered during the mining operations. If well preserved examples of plant fossils are recorded, it would be of scientific value if these examples can be recorded by a palaeontologist and representative samples provided to a registered institution such as the Bernard Price Institute for Palaeontology at WITS University, for further studies.

Due to the age and igneous nature of the Damwal Formation, the rocks will not contain fossils and a low palaeontological sensitivity is allocated to these rocks.

6. CONCLUSION AND RECOMMENDATIONS

The study area at the Kwagga North project of the Optimum Coal Mine is mainly underlain by Permian aged sedimentary rocks of the Vryheid Formation, Ecca Group of the Karoo Supergroup, with a small section of the area underlain by Vaalium aged rocks of the Damwal Formation, Rooiberg Group. The Vryheid Formation is known for containing an abundant assemblage of plant fossils and the mining of coal is by definition the mining of fossil plant material. The Damwal Formation will not contain fossils.

Due to the fact that the Vryheid Formation sediments and coal beds will only be exposed during the mining operations, it is unlikely that fossils will be observed before the mining takes place. For this reason a medium palaeontological sensitivity is allocated to the larger part of the study area.

It is recommended that:

1. The developer and the ECO of the mining project are made aware of the fact that coal mining is by definition the mining of fossil plant material.
2. That the developer applies for a destruction permit for plant fossils encountered during the mining operation.
3. That the developer should look out for exceptionally well preserved fossils and appoint a palaeontologist to collect representative samples of these fossils for further study at an appropriate institute such as the Bernard Price Institute for Palaeontology at WITS University.

7. REFERENCES

Bamford M. 2011. Desktop study Palaeontology Ermelo to Empangeni – Eskom powerline. Internal report Bernard Price Institute for Palaeontological Research. University of the Witwatersrand.

Cairncross, B, Beukes, NJ, Coetzee, LL and Rehfeld, U. 2005. The Bivalve *Megadesmus* from the Permian Volksrust Shale Formation (Karoo Supergroup), north-eastern Karoo Basin, South Africa: implications for late Permian Basin development. *South African Journal of Geology* 108: 547-556

Johnson MR, Anhausser CR and Thomas RJ. 2009. The Geology of South Africa. Geological Society of South Africa.

MacRae C. 1999. Life Etched in Stone. Geological Society of South Africa.

Mason TR and Christie ADM. 1986. Palaeoenvironmental significance of ichnogenus *Diplocraterion* torell from the Permian Vryheid Formation of the Karoo Supergroup, South Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* 53(3-4):249-265.

8. QUALIFICATIONS AND EXPERIENCE OF THE AUTHOR

Dr Gideon Groenewald has a PhD in Geology from the University of Port Elizabeth (Nelson Mandela Metropolitan University) (1996) and the National Diploma in Nature Conservation from Technicon RSA (the University of South Africa) (1989). He specialises in research on South African Permian and Triassic sedimentology and macrofossils with an interest in biostratigraphy, and palaeoecological aspects. He has extensive experience in the locating of fossil material in the Karoo Supergroup and has more than 20 years of experience in locating, collecting and curating fossils, including exploration field trips in search of new localities in the southern, western, eastern and north-eastern parts of the country. His publication record includes multiple articles in internationally recognized journals. Dr Groenewald is accredited by the Palaeontological Society of Southern Africa (society member for 25 years).

9. DECLARATION OF INDEPENDENCE

I, Gideon Groenewald, declare that I am an independent specialist consultant and have no financial, personal or other interest in the proposed development, nor the developers or any of their subsidiaries, apart from fair remuneration for work performed in the delivery of palaeontological heritage assessment services. There are no circumstances that compromise the objectivity of my performing such work.



Dr Gideon Groenewald
Geologist