

PALAEONTOLOGICAL IMPACT ASSESSMENT:

Environmental Impact Assessment for the Proposed Development of a 75 MW Solar Photovoltaic Facility (GEMSBOK SOLAR PV5) on Portion 8 of Gemsbok Bult Farm 120, north-east of Kenhardt, Northern Cape Province

Report prepared for:

CSIR – Environmental Management Services

P O Box 17001

Congella, Durban, 4013

South Africa

Report prepared by:

Dr John Almond - Natura Viva cc

P.O. Box 12410

Mill Street, Cape Town, 8010

South Africa

January 2016

SPECIALIST EXPERTISE

Dr John Almond has an Honours Degree in Natural Sciences (Zoology) as well as a PhD in Palaeontology from the University of Cambridge, UK. He has been awarded post-doctoral research fellowships at Cambridge University and in Germany, and has carried out palaeontological research in Europe, North America, the Middle East as well as North and South Africa and Madagascar. For eight years he was a scientific officer (palaeontologist) for the Geological Survey / Council for Geoscience in the RSA. His current palaeontological research focuses on fossil record of the Precambrian - Cambrian boundary and the Cape Supergroup of South Africa. He has recently written palaeontological reviews for several 1:250 000 geological maps published by the Council for Geoscience and has contributed educational material on fossils and evolution for new school textbooks in the RSA.

Since 2002 Dr Almond has also carried out numerous palaeontological impact assessments for developments and conservation areas in the Western, Eastern and Northern Cape, Free State, Northwest, Mpumalanga and Gauteng under the aegis of his Cape Town-based company *Natura Viva* cc. He was a long-standing member of the Archaeology, Palaeontology and Meteorites Committee for Heritage Western Cape (HWC) and an advisor on palaeontological conservation and management issues for the Palaeontological Society of South Africa (PSSA), HWC and the South African Heritage Resources Agency (SAHRA). He is currently compiling technical reports on the provincial palaeontological heritage of Western, Northern and Eastern Cape for SAHRA and HWC. Dr Almond is an accredited member of PSSA and the Association of Professional Heritage Assessment Practitioners – Western Cape (APHAP).

SPECIALIST DECLARATION

I, Dr John Edward Almond, as the appointed independent specialist, in terms of the 2014 EIA Regulations, hereby declare that I:

- I act as the independent specialist in this application;
- I perform the work relating to the application in an objective manner, even if this results in views and findings that are not favourable to the applicant;
- regard the information contained in this report as it relates to my specialist input/study to be true and correct, and do not have and will not have any financial interest in the undertaking of the activity, other than remuneration for work performed in terms of the NEMA, the Environmental Impact Assessment Regulations, 2014 and any specific environmental management Act;
- I declare that there are no circumstances that may compromise my objectivity in performing such work;
- I have expertise in conducting the specialist report relevant to this application, including knowledge of the Act, Regulations and any guidelines that have relevance to the proposed activity;
- I will comply with the Act, Regulations and all other applicable legislation;
- I have no, and will not engage in, conflicting interests in the undertaking of the activity;
- I have no vested interest in the proposed activity proceeding;
- I undertake to disclose to the applicant and the competent authority all material information in my possession that reasonably has or may have the potential of influencing - any decision to be taken with respect to the application by the competent authority; and - the objectivity of any report, plan or document to be prepared by myself for submission to the competent authority;
- I have ensured that information containing all relevant facts in respect of the specialist input/study was distributed or made available to interested and affected parties and the public and that participation by interested and affected parties was facilitated in such a manner that all interested and affected parties were provided with a reasonable opportunity to participate and to provide comments on the specialist input/study;
- I have ensured that the comments of all interested and affected parties on the specialist input/study were considered, recorded and submitted to the competent authority in respect of the application;
- all the particulars furnished by me in this specialist input/study are true and correct; and
- I realize that a false declaration is an offence in terms of regulation 48 and is punishable in terms of section 24F of the Act.

John E. Almond

Signature of the specialist:

Name of Specialist: Dr John Edward Almond

Date: 26 January 2016

EXECUTIVE SUMMARY

Gemsbok Solar PV5 (Pty) Ltd (a wholly owned Subsidiary of Mulilo Renewable Project Developments (PTY) LTD (“Mulilo”)) is proposing to develop the Gemsbok Solar PV5 75 MW Solar Photovoltaic (PV) Facility on Portion 8 of Gemsbok Bult Farm 120, situated c. 36 km north-east of Kenhardt, Northern Cape Province. The study area (including preferred and alternative sites *plus* transmission lines) for the proposed PV facility is underlain at depth by Precambrian basement rocks (c. 1-2 billion years old) assigned to the Namaqua-Natal Province. These ancient igneous and high-grade metamorphic rocks - mainly granites and gneisses of the Keimoes Suite (granitoids) and Korannaland Supergroup (high grade metasediments) - crop out at surface in small areas and are entirely unfossiliferous. A large proportion of the basement rocks is mantled by a range of superficial sediments of Late Cenozoic age that may contain sparse fossil remains. These predominantly thin, unconsolidated deposits include small patches of calcretes, gravelly to sandy river alluvium, pan sediments, surface gravels, colluvium (scree) as well as Pleistocene to Recent wind-blown sands of the Gordonia Formation (Kalahari Group). Most of these younger rock units are of widespread occurrence and low palaeontological sensitivity. Scientifically important vertebrate fossil remains (e.g. Pleistocene mammalian bones and teeth) have been recorded within older stratified pan and river sediments elsewhere in the Bushmanland region where they are often associated with stone artefacts, while a limited range of trace fossils (e.g. plant root casts, termitaria and other invertebrate burrows) may be found within calcrete horizons.

No previously recorded areas or sites of exceptional fossil heritage sensitivity or significance have been identified within the Nieuwehoop Solar Development project area near Kenhardt as a whole. Due to the inferred scarcity of scientifically important fossil remains within the Gemsbok Solar PV5 study area, the overall impact significance of the construction phase of the proposed solar energy project is assessed as VERY LOW (before and after mitigation). No significant impacts on fossil heritage are anticipated during the operational and decommissioning phases of the proposed solar energy facility. The potentially fossiliferous sedimentary rock units represented within the study area (e.g. Gordonia sands, calcrete, alluvium) are of widespread occurrence and this is also likely to apply to most of the fossils they contain. It is concluded that the cumulative impacts on fossil heritage resource posed by the known alternative energy and other infrastructural developments in the region (as explained in Chapter 4 of the EIA Report) is very low. There are no fatal flaws in the proposed Gemsbok Solar PV5 75 MW Solar PV Facility development, nor are there objections to its authorisation as far as fossil heritage conservation is concerned, since significant impacts on scientifically valuable fossils or fossil sites are not anticipated here. The no-go option (no solar developments) will have a neutral impact on local palaeontological heritage resources. The only proposed condition to accompany the Environmental Authorisation is that the recommendations for monitoring and mitigation included in the Environmental Management Programme (EMPr) are fully complied with.

Given the low palaeontological sensitivity of the eastern Bushmanland region, as determined from desktop and field-based studies, as well as the inferred very low impact significance of the Gemsbok Solar PV5 75 MW Solar PV Facility for fossil heritage conservation, no specialist palaeontological monitoring or mitigation is recommended here, pending the potential discovery of substantial new fossil remains during construction. During the construction phase all substantial bedrock excavations should be monitored for fossil material by the responsible Environmental Control Officer. Should significant fossil remains - such as vertebrate bones and teeth, plant-rich fossil lenses, petrified wood or dense fossil burrow assemblages - be exposed during construction, the responsible Environmental

Control Officer should safeguard these, preferably *in situ*. The South African Heritage Resources Authority (SAHRA) should be alerted as soon as possible (Contact details: Mrs Colette Scheermeyer, P.O. Box 4637, Cape Town 8000, Tel: 021 462 4502, Email: cscheermeyer@sahra.org.za), so that appropriate action can be taken by a professional palaeontologist, at the developer's expense. Mitigation would normally involve the scientific recording and judicious sampling or collection of fossil material as well as associated geological data (e.g. stratigraphy, sedimentology, taphonomy) by a professional palaeontologist. The palaeontologist concerned with mitigation work will need a valid fossil collection permit from SAHRA and any material collected would have to be curated in an approved depository (e.g. museum or university collection). These recommendations should be included within the EMPr for the proposed solar energy facility development.

In this report the entire site (preferred and alternative layouts *plus* transmission lines) for the proposed Gemsbok Solar PV5 75 MW Solar PV Facility on Portion 8 of Gemsbok Bult Farm 120 has been assessed based on the worst case scenario. From a palaeontological heritage impact point of view, the applicant can select any 250 ha area within the surveyed area to construct the PV plant, provided that the recommended mitigation measures are implemented as applicable.

LIST OF ABBREVIATIONS

DEA	Department of Environmental Affairs
ECO	Environmental Control Officer
EMPr	Environmental Management Programme
EIA	Environmental Impact Assessment
HWC	Heritage Western Cape
PIA	Palaeontological Impact Assessment
PV	Photovoltaic
SAHRA	South African Heritage Resources Agency
Ma / mya	Million years ago

GLOSSARY

Definitions	
<i>Basement Rocks</i>	Ancient igneous and metamorphic rocks (usually unfossiliferous) underlying the sedimentary cover rocks in a given region
<i>Calcrete</i>	Pedogenic limestone (<i>i.e.</i> limestone generated by soil processes within soils and surface rock debris), generally associated with seasonally arid climates.
<i>Fossiliferous</i>	Containing fossil remains
<i>Igneous Rocks</i>	Rocks that have crystallised from a molten state (magma / lava); <i>e.g.</i> granite.
<i>Metamorphic</i>	Rocks that have recrystallized under conditions of altered (usually highly elevated) temperature and pressure; <i>e.g.</i> gneiss.
<i>Precambrian</i>	Older than 541 million years old (mya).
<i>Pleistocene Epoch</i>	Time period between c. 2.6 mya and 10 000 years ago (associated with a series of major glaciations in the northern hemisphere).

COMPLIANCE WITH THE APPENDIX 6 OF THE 2014 EIA REGULATIONS

Requirements of Appendix 6 – GN R982	Addressed in the Specialist Report
1. (1) A specialist report prepared in terms of these Regulations must contain-	
a) details of- <ul style="list-style-type: none"> i. the specialist who prepared the report; and ii. the expertise of that specialist to compile a specialist report including a curriculum vitae; 	Appendix A of this EIA Report
b) a declaration that the specialist is independent in a form as may be specified by the competent authority;	Appendix B of this EIA Report
c) an indication of the scope of, and the purpose for which, the report was prepared;	Section 1.1
d) the date and season of the site investigation and the relevance of the season to the outcome of the assessment;	Not Applicable
e) a description of the methodology adopted in preparing the report or carrying out the specialised process;	Section 1.1
f) the specific identified sensitivity of the site related to the activity and its associated structures and infrastructure;	Section 1.3
g) an identification of any areas to be avoided, including buffers;	Not Applicable
h) a map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	Section 1.3
i) a description of any assumptions made and any uncertainties or gaps in knowledge;	Section 1.1.4
j) a description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives on the environment;	Section 1.5, 1.6, 1.7 and 1.8
k) any mitigation measures for inclusion in the EMPr;	Section 1.7 and Section 1.8
l) any conditions for inclusion in the environmental authorisation;	Not Applicable
m) any monitoring requirements for inclusion in the EMPr or environmental authorisation;	Section 1.8
n) a reasoned opinion- <ul style="list-style-type: none"> i. as to whether the proposed activity or portions thereof should be authorised; and ii. if the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan; 	Section 1.9
o) a description of any consultation process that was undertaken during the course of preparing the specialist report;	Not Applicable
p) a summary and copies of any comments received during any consultation process and where applicable all responses thereto; and	Section 1.5.1
q) any other information requested by the competent authority.	Not applicable

TABLE OF CONTENTS

PALAEONTOLOGICAL HERITAGE	3
1.1. INTRODUCTION AND METHODOLOGY	3
1.1.1. SCOPE AND OBJECTIVES	3
1.1.2. TERMS OF REFERENCE	3
1.1.3. APPROACH AND METHODOLOGY	4
1.1.4. ASSUMPTIONS AND LIMITATIONS	5
1.1.5. SOURCES OF INFORMATION	6
1.2. DESCRIPTION OF PROJECT ASPECTS RELEVANT TO PALAEONTOLOGICAL HERITAGE IMPACTS	7
1.3. DESCRIPTION OF THE AFFECTED ENVIRONMENT	7
1.3.1. GEOLOGICAL CONTEXT	7
1.3.2. PALAEONTOLOGICAL HERITAGE	10
TABLE 1.1: FOSSIL HERITAGE RECORDED FROM THE MAJOR ROCK UNITS THAT ARE REPRESENTED WITHIN THE BROADER SCATEC SOLAR STUDY AREA NEAR KENHARDT	12
1.4. APPLICABLE LEGISLATION AND PERMIT REQUIREMENTS	12
1.5. IDENTIFICATION OF KEY ISSUES	14
1.5.1. KEY ISSUES IDENTIFIED DURING THE SCOPING PHASE	14
1.5.2. IDENTIFICATION OF POTENTIAL IMPACTS	14
1.5.3. CONSTRUCTION PHASE	14
1.5.4. OPERATIONAL PHASE	14
1.5.5. DECOMMISSIONING PHASE	14
1.5.6. CUMULATIVE IMPACTS	14
1.6. ASSESSMENT OF IMPACTS AND IDENTIFICATION OF MANAGEMENT ACTIONS	14
1.6.1. RESULTS OF THE FIELD STUDY	ERROR! BOOKMARK NOT DEFINED.
1.6.2. POTENTIAL IMPACTS (CONSTRUCTION PHASE)	15
1.6.3. POTENTIAL IMPACTS (OPERATIONAL AND DECOMMISSIONING PHASES)	16
1.6.4. CUMULATIVE IMPACTS	16
1.7. IMPACT ASSESSMENT SUMMARY	16

<u>1.8.</u>	<u>INPUT TO THE ENVIRONMENTAL MANAGEMENT PROGRAM</u>	<u>19</u>
<u>1.9.</u>	<u>CONCLUSION AND RECOMMENDATIONS</u>	<u>19</u>
<u>1.10.</u>	<u>REFERENCES</u>	<u>20</u>

TABLES AND FIGURES

Table 1.1: Fossil heritage recorded from the major rock units that are represented within the broader Nieuwehoop Solar Development study area near Kenhardt	12
Table 1.2 Impact assessment summary table for the Construction Phase	17
Table 1.3 Cumulative impact assessment summary table	18

PALAEONTOLOGICAL IMPACT ASSESSMENT

This chapter presents the Palaeontological Impact Assessment that was prepared by Dr. John Almond (of Natura Viva cc) as part of the Environmental Impact Assessment (EIA) for the proposed solar PV facility for the Phase 2 Nieuwehoop Solar Park near Kenhardt in the Northern Cape Province.

1.1. INTRODUCTION AND METHODOLOGY

1.1.1. *Scope and Objectives*

The proposed Gemsbok Solar PV5 75 MW Solar Photovoltaic (PV) Facility project area overlies potentially fossiliferous sedimentary rocks.

This report provides a desktop assessment of potential impacts on local palaeontological (*i.e.* fossil) heritage within the study area for the proposed Gemsbok Solar PV5 (75 MW) Solar PV Facility on Portion 8 of Gemsbok Bult Farm 120, situated c. 36 km north-east of Kenhardt, Northern Cape Province. The report contributes to the EIA for this alternative energy development and includes recommendations for inclusion in the EMPr (Part B of the EIA Report).

The overall objectives of the specialist study are to:

- Determine the current conditions in sufficient detail so that there is a baseline against which impacts can be identified and measured.
- Identify potential impacts that may occur during the construction, operational and decommissioning phases of the proposed development, as well as impacts associated with future environmental changes if the “no-go” option is implemented (both positive and negative).
- Assess the impacts in terms of direct, indirect and cumulative impacts.
- Provide recommendations with regards to potential monitoring programmes.
- Determine mitigation and/or management measures which could be implemented to as far as possible reduce the effect of negative impacts and enhance the effect of positive impacts.
- Incorporate and address all issues and concerns raised in relation to palaeontological impacts.

1.1.2. *Terms of Reference*

The Terms of Reference for the present study, as defined by the CSIR, are as follows:

1. Review detailed information relating to the project description and precisely define the environmental risks to palaeontological heritage, and consequences thereto.
2. Conduct a review of available information pertaining to the study area.
3. Draw on desktop information sources, the knowledge of local experts, information published in the scientific press and information derived from relevant EIAs and similar specialist studies previously conducted within the surrounding area.
4. Prepare and undertake a desktop study on the palaeontology and fossil heritage within the proposed project area, based on:
 - a review of all relevant palaeontological and geological literature, including geological maps and previous reports,
 - location and examination of fossil collections from the study area (*e.g.* museums), and

- data on the proposed development (e.g. location of footprint, depth and volume of bedrock excavation envisaged).
5. Describe the type and location of known fossil heritage sites in the study area, and characterize all items that may be affected by the proposed project.
 6. Describe the baseline environment and determine the *status quo* in terms of palaeontological heritage.
 7. Note fossils and associated sedimentological features of palaeontological relevance (photos, maps, aerial or satellite images, and stratigraphic columns).
 8. Analyse the stratigraphy, age and depositional setting of fossil-bearing units.
 9. Evaluate the potential for occurrence of palaeontological heritage features within the study area.
 10. Incorporate relevant information from other specialist reports/findings, if required.
 11. Identify and rank the highlights and sensitivities to development of fossil heritage within study area.
 12. Identify and rate potential direct, indirect and cumulative impacts of the proposed project on the palaeontology and fossil heritage during the construction, operational and decommissioning phases of the project. Study the cumulative impacts of the project by considering the impacts of existing industries / solar PV plants within the area (as well as those PV plants that are proposed), together with the impact of the proposed project.
 13. Provide recommendations and suggestions regarding fossil heritage management on site, including conservation measures, as well as promotion of local fossil heritage (e.g. for public education, schools) to ensure that the impacts are limited.
 14. Provide input to the EMPr, including mitigation and monitoring requirements to ensure that the impacts on the archaeological features and heritage features are limited.
 15. Provide specific recommendations for further palaeontological mitigation (if any).
 16. Compile an illustrated, fully-referenced review of palaeontological heritage within study area based on desktop study.

1.1.3. Approach and Methodology

In preparing a palaeontological desktop study the potentially fossiliferous rock units (groups, formations etc.) represented within the study area are determined from geological maps and satellite images. The known fossil heritage within each rock unit is inventoried from the published scientific literature, previous palaeontological impact studies in the same region, and the author's field experience and palaeontological database (consultation with professional colleagues as well as examination of institutional fossil collections may play a role here). This data is then used to assess the palaeontological sensitivity of each rock unit to development (provisional tabulations of palaeontological sensitivity of all formations in the Western, Eastern and Northern Cape have already been compiled by J. Almond and colleagues (e.g. Almond & Pether 2008). The likely impact of the proposed development on local fossil heritage is then determined on the basis of (1) the palaeontological sensitivity of the rock units concerned and (2) the nature and scale of the development itself, most significantly the extent of fresh bedrock excavation envisaged. When rock units of moderate to high palaeontological sensitivity are present within the development footprint, a Phase 1 field assessment study by a professional palaeontologist is usually warranted to identify any palaeontological hotspots and make specific recommendations for any mitigation required before or during the construction phase of the development. However, due to the low palaeontological sensitivity of the present study area a Phase 1 field assessment is not required and a desktop assessment is being undertaken instead (i.e. this study).

On the basis of the desktop and Phase 1 field assessment studies, the likely impact of the proposed development on local fossil heritage and any need for specialist mitigation are then

determined. Adverse palaeontological impacts normally occur during the construction rather than the operational or decommissioning phase. Phase 2 mitigation by a professional palaeontologist – normally involving the recording and sampling of fossil material and associated geological information (e.g. sedimentological data) may be required (a) in the pre-construction phase where important fossils are already exposed at or near the land surface and / or (b) during the construction phase when fresh fossiliferous bedrock has been exposed by excavations. To carry out mitigation, the palaeontologist involved will need to apply for a palaeontological collection permit from the relevant heritage management authorities for the Northern Cape, *i.e.* SAHRA (Contact details: Mrs Colette Scheermeyer, P.O. Box 4637, Cape Town 8000, Tel: 021 462 4502, Email: cscheermeyer@sahra.org.za). It should be emphasized that, providing appropriate mitigation is carried out, the majority of developments involving bedrock excavation can make a positive contribution to our understanding of local palaeontological heritage.

1.1.4. Assumptions and Limitations

The accuracy and reliability of palaeontological specialist studies as components of Heritage Impact Assessments are **generally** limited by the following constraints:

1. Inadequate database for fossil heritage for much of South Africa, given the large size of the country and the small number of professional palaeontologists carrying out fieldwork here. Most development study areas – including the overall Nieuwehoop Solar Development project area - have never been surveyed by a palaeontologist.
2. Variable accuracy of geological maps which underpin these desktop studies. For large areas of terrain these maps are largely based on aerial photographs alone, without ground-truthing. The maps generally depict only significant (“mappable”) bedrock units as well as major areas of superficial “drift” deposits (alluvium, colluvium) but for most regions give little or no idea of the level of bedrock outcrop, depth of superficial cover (soil etc.), degree of bedrock weathering or levels of small-scale tectonic deformation, such as cleavage. All of these factors may have a major influence on the impact significance of a given development on fossil heritage and can only be reliably assessed in the field.
3. Inadequate sheet explanations for geological maps, with little or no attention paid to palaeontological issues in many cases, including poor locality information.
4. The extensive relevant palaeontological “grey literature” - in the form of unpublished university theses, impact studies and other reports (e.g. of commercial mining companies) - that is not readily available for desktop studies.
5. Absence of a comprehensive computerized database of fossil collections in major South African institutions which can be consulted for impact studies. A Karoo fossil vertebrate database is now accessible for impact study work.

In the case of palaeontological desktop studies without supporting Phase 1 field assessments these limitations may variously lead to either:

(a) underestimation of the palaeontological significance of a given study area due to ignorance of significant recorded or unrecorded fossils preserved there, or

(b) overestimation of the palaeontological sensitivity of a study area, for example when originally rich fossil assemblages inferred from geological maps have in fact been destroyed by

tectonism or weathering, or are buried beneath a thick mantle of unfossiliferous “drift” (soil, alluvium *etc.*).

Since most areas of South Africa have not been studied palaeontologically, a palaeontological desktop study usually entails inferring the presence of buried fossil heritage within the study area from relevant fossil data collected from similar or the same rock units elsewhere, sometimes at localities far away. Where substantial exposures of bedrocks or potentially fossiliferous superficial sediments are present in the study area, the reliability of a palaeontological impact assessment may be significantly enhanced through field assessment by a professional palaeontologist.

In the case of the Nieuwehoop Solar Development project area near Kenhardt in the Northern Cape, bedrock exposure is limited due to extensive cover by superficial deposits (*e.g.* alluvium, soils, surface gravels), especially in areas of low relief, as well as by pervasive *bossieveld* vegetation. For this reason, as well as the low palaeontological sensitivity of the sedimentary rocks mapped in the project area, a desktop-level rather than field-based assessment was considered appropriate for this study. Despite the lack of palaeontological field data from the project area itself, confidence levels in the conclusions reached in the desktop study are moderately high because of the author’s field experience of the sedimentary rocks represented in the wider Bushmanland region (See reference list for previous palaeontological assessments in the area; *e.g.* Almond 2009, 2011, 2014a, 2014b, 2014c, 2014d). Recent palaeontological heritage assessments for several other alternative energy developments in the region have been taken into consideration (*e.g.* the Scatec Solar project area just to the west of the Mulilo Solar PV project area).

In terms of the impact assessment, the methodology adopted is outlined in Chapter 4 of the EIA Report, which also notes the developments within a 20 km radius that have been considered in order to assess cumulative impacts.

1.1.5. Sources of Information

The information used in this desktop study was based on the following sources:

1. A detailed project outline supplied by the CSIR - Environmental Management Services.
2. Previous desktop palaeontological assessment reports for study areas in the Kenhardt region by the author (Almond 2009, 2011, 2014a, 2014b, 2014c, 2014d).
3. A review of the relevant scientific literature, including published geological maps (*e.g.* 1: 250 000 scale geological map sheet 2920 Kenhardt published by the Council for Geoscience, Pretoria) and accompanying sheet explanations (*e.g.* Slabbert *et al.* 1999)
4. The author’s previous field experience with the formations concerned and their palaeontological heritage (*cf* Almond and Pether 2008; SAHRIS website).

1.1.6. Declaration of Independence of Specialists

Refer to Appendix A of this EIA Report for the Curriculum Vitae of Dr. John Almond, which highlights his experience and expertise. The declaration of independence by the specialist is provided in Box 13.1 below and included in Appendix B of this EIA Report.

BOX 13.1: DECLARATION OF INDEPENDENCE
--

I, John Almond, declare that I am an independent consultant and have no business, financial, personal or other interest in the proposed Gemsbok Solar PV5 Project, application or appeal in respect of which I was appointed, other than fair remuneration for work performed in connection with the activity, application or appeal. There are no circumstances that compromise the objectivity of my performing such work.



JOHN ALMOND

1.2. DESCRIPTION OF PROJECT ASPECTS RELEVANT TO PALAEOLOGICAL HERITAGE IMPACTS

As noted above, the Gemsbok Solar PV5 project area near Kenhardt is located in a region of Bushmanland that is underlain by potentially fossiliferous sedimentary rocks of Late Tertiary or Quaternary age as well as by unfossiliferous basement rocks (as discussed in Section 1.3 of this chapter). The construction phase of the proposed development will entail substantial excavations into the superficial sediment cover and locally into the underlying bedrock as well. These include, for example, surface clearance operations, excavations for the solar array footings, underground cables, access and internal gravel roads, 132 kV transmission line towers, on-site substation, laydown areas, stormwater channels, water pipelines (if required) and foundations for buildings (offices, operational control centre, warehouse/workshop). All these developments may adversely affect potential, legally-protected fossil heritage resources within the study area by destroying, disturbing or permanently sealing-in fossils at or beneath the surface of the ground that are then no longer available for scientific research or other public good.

The planning, operational and decommissioning phases of the proposed solar energy facility are very unlikely to involve additional adverse impacts on local palaeontological heritage, however.

A detailed description of the proposed project is included in Chapter 2 of the EIA Report.

1.3. DESCRIPTION OF THE AFFECTED ENVIRONMENT

In this section of the report an outline of the geology of the proposed Gemsbok Solar PV5 project area is first given, based on the relevant geological maps and scientific literature. This is followed by a brief review of fossil heritage that has previously been recorded from the sedimentary rock units that are represented within the project area.

1.3.1. Geological Context

The study area for the proposed Gemsbok Solar PV5 project, located on Portion 8 of Gemsbok Bult Farm 120, located some 36 km northeast of Kenhardt, Northern Cape, is situated in flat-lying terrain within the semi-arid Bushmanland region at elevations between c. 975 to 1120 m amsl. with a gentle slope to the southwest. It is drained by a dendritic network of shallow, southerly-flowing tributary streams of the Hartbeesrivier such as the Rugseersrivier system. The geology of the study area is shown on 1: 250 000 geology sheet 2920 Kenhardt (Council for Geoscience, Pretoria) (Figure 1.1). The entire area is underlain at depth by a variety of Precambrian basement rocks that are c. 2 billion years old and are assigned to the **Namaqua-**

Natal Province. These ancient igneous and high-grade metamorphic rocks - mainly granites and gneisses of the **Keimoes Suite** (granitoids) and **Korannaland Supergroup** (high grade metasediments) – are listed in the legend to Figure 1.1. The various basement rock units are described in the Kenhardt 1: 250 000 sheet explanation by Slabbert *et al.* (1999) and placed in the context of the Namaqua-Natal Province by Cornell *et al.* (2006). They crop out at surface as small patches and are entirely unfossiliferous and so will not be discussed further here. The Precambrian crustal rocks are transected by the NW-SE trending Boven Rugzeer Shear Zone that crosses the study area. The shear zone separates two major crustal blocks in Bushmanland known as the Kakamas Terrane and Areachap Terrane (Cornell *et al.* 2006, their fig. 18).

A large proportion of the basement rocks in the proposed project area are mantled by a range of superficial sediments of Late Caenozoic age, some of which are included within the **Kalahari Group**. These predominantly thin, unconsolidated deposits include small patches of calcretes (soil limestones), gravelly to sandy river alluvium, pan sediments along certain watercourses, surface gravels, colluvium (scree) as well as – especially – Quaternary to Recent aeolian (wind-blown) sands of the Gordonia Formation (Kalahari Group). The basement rocks in the Gersbok Solar PV5 study area are largely mantled by aeolian sands of the **Gordonia Formation** (“Kalahari sands”) as well as by Late Caenozoic alluvial deposits along shallow dendritic drainage lines.

The geology of the Late Cretaceous to Recent Kalahari Group is reviewed by Thomas (1981), Dingle *et al.* (1983), Thomas & Shaw (1991), Haddon (2000) and Partridge *et al.* (2006). The thickness of the unconsolidated Kalahari sands in the Bushmanland area is variable and often uncertain. The Gordonia Formation dune sands are considered to range in age from the Late Pliocene/Early Pleistocene to Recent, dated in part from enclosed Middle to Late Stone Age stone tools (Dingle *et al.*, 1983, p. 291). Note that the recent extension of the Pliocene - Pleistocene boundary from 1.8 Ma back to 2.588 Ma would place the older Gordonia Formation sands entirely within the Pleistocene Epoch. A number of older Kalahari formations underlie the young wind-blown surface sands in the main Kalahari depository to the north of the study area. However, at the latitude of the study area near Kenhardt (c. 29° S) Gordonia Formation sands less than 30 m thick are likely to be the main or perhaps only Kalahari sediments present (*cf* isopach map of the Kalahari Group, Figure 6 *in* Partridge *et al.*, 2006). These unconsolidated sands will be locally underlain by thin subsurface gravels along the buried palaeosurface and perhaps by calcretes of Pleistocene or younger age (*cf* Mokalanen Formation).




Figure 1.1. Extract from 1: 250 000 scale geological map sheet 2920 Kenhardt (Council for Geoscience, Pretoria) showing the geology of the Gembok Solar PV5 study area on the Portion 8 of Gembok Bult Farm 120, situated c. 36 km to the NE of Kenhardt, Northern Cape. The area – including the preferred and alternative sites as well as associated transmission lines - is *approximately* indicated by the pale blue polygon.

The main geological units represented within the Gembok Solar PV5 project area include:

PRECAMBRIAN BASEMENT ROCKS

KEIMOES SUITE

- Brown (Mge) = Gembokbult Granite
- Red (Mks) = Klipkoppies Granite
- Blue-green (Mf) = Friersdale Charnockite

KORANNALAND SUPERGROUP

- Dark blue (Mja) = Jacomynpan Group

LATE CAENOZOIC SUPERFICIAL SEDIMENTS

- Pale yellow with sparse red stipple (Qg) = aeolian sands of the Gordonia Formation (Kalahari Group)
- Pale yellow with dense black stipple = alluvial and pan sediments
- Dark yellow (Tec) = calcrete

1.3.2. Palaeontological Heritage

The Precambrian basement rocks represented within the study area are high grade metamorphic rocks that were last metamorphosed some 1 billion years ago and are entirely unfossiliferous. The sparse fossil record of Late Cenozoic superficial sediments in the Bushmanland region are briefly reviewed here (Refer also to Table 1.1). Note that, to the author's knowledge, there are no fossil records from the broader Nieuwehoop Solar Development project area itself and no palaeontological fieldwork has been undertaken here.

The diverse superficial deposits within the South African interior, including Bushmanland, have been comparatively neglected in palaeontological terms. However, sediments associated with ancient drainage systems, springs and pans may occasionally contain important fossil biotas, notably the bones, teeth and horn cores of mammals as well as remains of reptiles like tortoises (e.g. Skead 1980, Klein 1984, Brink 1987, Bousman *et al.* 1988, Bender & Brink 1992, Brink *et al.* 1995, MacRae 1999, Meadows & Watkeys 1999, Churchill *et al.* 2000, Partridge & Scott 2000, Brink & Rossouw 2000, Rossouw 2006, Almond *in* Macey *et al.* 2011). Other late Cenozoic fossil biotas that may occur within these superficial deposits include non-marine molluscs (bivalves, gastropods), ostrich egg shells, trace fossils (e.g. calcretised termitaria, coprolites, invertebrate burrows, rhizcretions), and plant material such as peats or palynomorphs (pollens) in organic-rich alluvial horizons (Scott 2000) and diatoms in pan sediments. In Quaternary deposits, fossil remains may be associated with human artefacts such as stone tools and are also of archaeological interest (e.g. Smith 1999 and references therein). Ancient solution hollows within extensive calcrete hardpans may have acted as animal traps in the past. As with coastal and interior limestones, they might occasionally contain mammalian bones and teeth (perhaps associated with hyaena dens) or invertebrate remains such as snail shells.

Diverse fossils associated with the ancient Tertiary drainage systems of the Karoo and Bushmanland region have been summarized by Almond *in* Macey *et al.* (2011) (See also articles by Cooke 1949, Wells 1964, Butzer *et al.* 1973, Helgren 1977, Klein 1984, Macrae 1999). They include remains of fish, reptiles, mammals, freshwater molluscs, petrified wood and trace fossils (e.g. De Wit 1990, 1993, De Wit & Bamford 1993, Bamford 2000, Bamford & De Wit 1993, Senut *et al.* 1996).

In the Brandvlei area to the southwest of Kenhardt lies the north-south trending Geelvloer Palaeo-valley, a Mid Tertiary palaeodrainage system that links up with the Commissioners Pan – Koa Valley system to the northwest. Here calcretised basal alluvial facies contain bones of hippopotamus-like artiodactyls called anthracotherids indicating a Miocene age (De Wit 1993, 1999, De Wit *et al.* 2000). Anthracotherids are an extinct group of amphibious mammalian herbivores only distantly related to true hippos that were widespread in the Miocene of Africa (Schneider & Marais 2004). Early to Mid-Miocene silicified woods from Brandvlei are referable to a number of extant tree families, including the Dipterocarpaceae that mainly inhabit tropical forests in Africa and Asia today. The fossil woods and associated sediments indicate that warm, tropical to subtropical climates prevailed in the Mid-Miocene and that perennial, low-sinuosity braided river systems supported lush riparian forests (De Wit & Bamford 1993, Bamford & De Wit 1993, Bamford 2000). Wet, weakly seasonal climates are suggested by the structure (indistinct growth rings) and dimensions (trunk diameters of over 50 cm) of the fossil woods (Bamford 2000).

Abraded Plio-Pleistocene fossil woods from relict alluvial terraces of the Sak River just north of Brandvlei include members of the Family Polygalaceae and also indicate humid growth conditions (Bamford & De Wit 1993). These terraces were formed by meandering rivers during intermittent pluvial (*i.e.* wetter), but still semi-arid, episodes following the onset of generally arid

conditions in the western portion of southern Africa towards the end of the Miocene. So far fossils have not been recorded from the Sakrivier system closer to Kenhardt.

Pan sediments in Bushmanland have also recently yielded interesting Pleistocene mammalian faunas in association with age-diagnostic archaeological material. Important fossil mammalian remains assigned to the Florisian Mammal Age (c. 300 000 – 12 000 BP; MacRae 1999) have recently been documented from stratigraphic units designated Group 4 to Group 6 (i.e. calcrete hardpan and below) at Bundu Pan, some 22 km northwest of Copperton (Kiberd 2006 and references therein). These are among very few Middle Pleistocene faunal records from stratified deposits in the southern Africa region (Klein 1980, 1984a, 1984b, 2000) and are therefore of high palaeontological significance. Characteristic extinct Pleistocene species recorded at Bundu Pan are the giant Cape Horse or Zebra (*Equus capensis*) and the Giant Hartebeest (*Megalotragus priscus*). Other extant to extinct taxa include species of warhog, blesbok, black wildebeest, springbok and baboon. There is additionally trace fossil evidence for hyaenids (tooth marks) as well as ostrich egg shell. Preliminary dating and the inferred ecology of the fossil taxa present suggests the presence of standing water within a grassy savanna setting during the 200 - 300 000 BP interval when the Bundu Pan faunal assemblage accumulated. A sequence of Earlier, Middle and Later Stone Age (ESA, MSA and LSA, respectively) artefact assemblages is also recorded from this site. Stratigraphic Groups 4 to 6 (i.e. calcrete hardpan and below) contain a Final Acheulian or transitional ESA/MSA artefact assemblage, while Groups 2 - 3 above the calcrete horizon contain a MSA artefact assemblage. Orton (2012) recorded a single fossil equid tooth associated with a rich MSA artefact assemblage from gravels overlying a calcrete hardpan on the farm Hoekplaas near Copperton. This horizon is probably equivalent to Group 3 of Kiberd's stratigraphy at Bundu Pan, and therefore somewhat younger than the Florisian mammal fauna reported there.

The fossil record of the Kalahari Group as a whole is generally sparse and low in diversity; no fossils are recorded here in the Kenhardt geology sheet explanation by Slabbert *et al.* (1999). The Gordonia Formation dune sands were mainly active during cold, drier intervals of the Pleistocene Epoch that were inimical to most forms of life, apart from hardy, desert-adapted species. Porous dune sands are not generally conducive to fossil preservation. However, mummification of soft tissues may play a role here and migrating lime-rich groundwaters derived from underlying lime-rich bedrocks may lead to the rapid calcretisation of organic structures such as burrows and root casts. Occasional terrestrial fossil remains that might be expected within this unit include calcretized rhizoliths (root casts) and termitaria (e.g. *Hodotermes*, the harvester termite), ostrich egg shells (*Struthio*), tortoise remains and shells of land snails (e.g. *Trigonephrus*) (Almond in Macey *et al.* 2011, Almond & Pether 2008). Other fossil groups such as freshwater bivalves and gastropods (e.g. *Corbula*, *Unio*), ostracods (seed shrimps), charophytes (stonewort algae), diatoms (microscopic algae within siliceous shells) and stromatolites (laminated microbial limestones) are associated with local watercourses and pans. Microfossils such as diatoms may be blown by wind into nearby dune sands (Du Toit 1954, Dingle *et al.*, 1983). These Kalahari fossils (or subfossils) can be expected to occur sporadically but widely, and the overall palaeontological sensitivity of the Gordonia Formation is therefore considered to be low. Underlying calcretes might also contain trace fossils such as rhizoliths, termite and other insect burrows, or even mammalian trackways. Mammalian bones, teeth and horn cores (also tortoise remains, and fish, amphibian or even crocodiles in wetter depositional settings) may be expected occasionally expected within Kalahari Group sediments and calcretes, notably those associated with ancient alluvial gravels (See Koa River Valley above). The younger (Pleistocene to Recent) fluvial and alluvial sands and gravels within the proposed development area are unlikely to contain many, if any, substantial fossil or subfossil remains.

Table 1.1: Fossil heritage recorded from the major rock units that are represented within the broader Nieuwehoop Solar Development study area near Kenhardt

GEOLOGICAL UNIT	ROCK TYPES AND AGE	FOSSIL HERITAGE	PALAEONTOLOGICAL SENSITIVITY
LATE CAENOZOIC SUPERFICIAL SEDIMENTS, especially ALLUVIAL AND PAN SEDIMENTS	fluvial, pan, lake and terrestrial sediments, including diatomite (diatom deposits), pedocretes (e.g. calcrete), colluvium (slope deposits such as scree), aeolian sands (Gordonia Formation, Kalahari Group) LATE TERTIARY, PLEISTOCENE TO RECENT	bones and teeth of wide range of mammals (e.g. mastodont proboscideans, rhinos, bovids, horses, micromammals), fish, reptiles (crocodiles, tortoises), ostrich egg shells, fish, freshwater and terrestrial molluscs (unionid bivalves, gastropods), crabs, trace fossils (e.g. calcretised termitaria, horizontal invertebrate burrows, stone artefacts), petrified wood, leaves, rhizoliths, stromatolites, diatom floras, peats and palynomorphs.	GENERALLY LOW BUT LOCALLY HIGH (e.g. Tertiary alluvium associated with old river courses)
Basement granites and gneisses NAMAQUA-NATAL PROVINCE	Highly-metamorphosed sediments, intrusive granites MID-PROTEROZOIC (c.1-2 billion years old)	None	ZERO

1.4. APPLICABLE LEGISLATION AND PERMIT REQUIREMENTS

All South African fossil heritage, including palaeontological sites and specimens, is protected by law (National Heritage Resources Act (Act 25 of 1999) and fossils cannot be collected, damaged, destroyed or disturbed without a permit from SAHRA or the relevant Provincial Heritage Resources Agency.

As mentioned previously, where palaeontological mitigation of a development project is required, the palaeontologist concerned with mitigation work would need a valid fossil collection permit from SAHRA and any material collected would have to be curated in an approved depository (e.g. museum or university collection). All palaeontological specialist work should conform to international best practice for palaeontological fieldwork and the study (e.g. data recording fossil collection and curation, final report) should adhere as far as possible to the minimum standards for Phase 2 palaeontological studies recently developed by SAHRA (2013).

The present palaeontological heritage assessment falls under Sections 35 and 38 (Heritage Resources Management) of the National Heritage Resources Act (Act 25 of 1999), and it will also inform the Environmental Management Programme for this project.

The various categories of heritage resources recognised as part of the National Estate in Section 3 of the National Heritage Resources Act include, among others:

- geological sites of scientific or cultural importance;
- palaeontological sites; and
- palaeontological objects and material, meteorites and rare geological specimens.

According to Section 35 of the National Heritage Resources Act (Act 25 of 1999), dealing with archaeology, palaeontology and meteorites:

- 1) The protection of archaeological and palaeontological sites and material and meteorites is the responsibility of a provincial heritage resources authority.
- 2) All archaeological objects, palaeontological material and meteorites are the property of the State.
- 3) Any person who discovers archaeological or palaeontological objects or material or a meteorite in the course of development or agricultural activity must immediately report the find to the responsible heritage resources authority, or to the nearest local authority offices or museum, which must immediately notify such heritage resources authority.
- 4) No person may, without a permit issued by the responsible heritage resources authority—
 - i. destroy, damage, excavate, alter, deface or otherwise disturb any archaeological or palaeontological site or any meteorite;
 - ii. destroy, damage, excavate, remove from its original position, collect or own any archaeological or palaeontological material or object or any meteorite;
 - iii. trade in, sell for private gain, export or attempt to export from the Republic any category of archaeological or palaeontological material or object, or any meteorite; or
 - iv. bring onto or use at an archaeological or palaeontological site any excavation equipment or any equipment which assist in the detection or recovery of metals or archaeological and palaeontological material or objects, or use such equipment for the recovery of meteorites.
- 5) When the responsible heritage resources authority has reasonable cause to believe that any activity or development which will destroy, damage or alter any archaeological or palaeontological site is under way, and where no application for a permit has been submitted and no heritage resources management procedure in terms of section 38 has been followed, it may—
 - a) serve on the owner or occupier of the site or on the person undertaking such development an order for the development to cease immediately for such period as is specified in the order;
 - b) carry out an investigation for the purpose of obtaining information on whether or not an archaeological or palaeontological site exists and whether mitigation is necessary;
 - c) if mitigation is deemed by the heritage resources authority to be necessary, assist the person on whom the order has been served under paragraph (a) to apply for a permit as required in subsection (4); and
 - d) recover the costs of such investigation from the owner or occupier of the land on which it is believed an archaeological or palaeontological site is located or from the person proposing to undertake the development if no application for a permit is received within two weeks of the order being served.

1.5. IDENTIFICATION OF KEY ISSUES

1.5.1. Key Issues Identified During the Scoping Phase

The only key issue identified by the specialist during the Scoping Phase is the potential loss of palaeontological heritage resources (fossils, fossil sites including their geological context) through surface clearance and excavations into sedimentary rocks during the construction phase of the project.

The Scoping Report was released for a 30-day comment period which extended from 23 October 2015 to 24 November 2015.

As noted above, based on the low palaeontological sensitivity of the area, this desktop Palaeontological Impact Assessment is being undertaken during the EIA Phase (*i.e.* prior to the commencement of construction of the Gemsbok Solar PV5 project (subject to the issuing of an Environmental Authorisation)). As mentioned above, this specialist assessment is conducted by Dr. John Almond in order to assess the significance of potential impacts of the proposed project on palaeontological resources (which is discussed in Section 1.6 of this chapter).

1.5.2. Identification of Potential Impacts

The potential impacts identified during the EIA Phase are:

1.5.3. Construction Phase

- Potential loss of palaeontological heritage resources through disturbance, damage or destruction of fossils and fossil sites (including associated geological contextual data) through surface clearance and excavation activities during the construction phase.

1.5.4. Operational Phase

No significant impacts on palaeontological heritage are anticipated during the operational phase of the development.

1.5.5. Decommissioning Phase

No significant impacts on palaeontological heritage are anticipated during the operational phase of the development.

1.5.6. Cumulative Impacts

- Potential cumulative loss of palaeontological heritage resources through disturbance, damage or destruction of fossils and fossil sites (including associated geological contextual data) through surface clearance and excavation activities during the construction phase of several alternative energy facilities within the broader Kenhardt region and other key electrical infrastructure developments within a 20 km radius of the proposed project site.

1.6. ASSESSMENT OF IMPACTS AND IDENTIFICATION OF MANAGEMENT ACTIONS

In this section of the report potential impacts of the construction, operational and decommissioning phases of the proposed PV solar facility development on palaeontological heritage are outlined and recommendations for any necessary monitoring or mitigation are

provided. Possible cumulative impacts in the light of other alternative energy development proposals in the Kenhardt region are also evaluated.

1.6.1. Potential Impact 1: Construction Phase

The construction phase of the proposed solar energy facility will entail substantial surface clearance and shallow excavations into the superficial sediment cover (aeolian sands, surface gravels, stream alluvium *etc.*), which may contain fossil remains, and in some cases also into the underlying unfossiliferous bedrock. These include, for example, surface clearance operations, excavations and foundations (which will likely be drilled and concreted into the ground) for the solar array footings, underground cables, access and internal gravel roads, 132 kV transmission line towers, on-site substation, laydown areas, stormwater channels, water pipelines and foundations for buildings (offices, operational control centre, warehouse/workshop). As a result, fossils at the ground surface or buried beneath it may be disturbed, damaged, destroyed or sealed-in while their scientifically informative sedimentary context will also be disturbed or destroyed.

Desktop analysis of the fossil records of the various rock units underlying the proposed project area indicates that the majority of these units are of zero to low palaeontological sensitivity (as discussed in Section 1.3.2 and Table 1.1 of this chapter). The basement rocks are entirely unfossiliferous while the overlying Late Caenozoic superficial sediments (wind-blown sands, alluvium, gravels *etc.*) are of low to very low palaeontological sensitivity. Construction of the solar panel arrays, overhead power lines, buildings and associated infrastructure is therefore unlikely to entail significant impacts on local fossil heritage resources.

The inferred impact of the proposed solar facility development on local fossil heritage is assessed in Table 1.2 below. This assessment applies only to the construction phase of the development since further impacts on fossil heritage during the operational and decommissioning phases of the solar energy facility are not anticipated.

The destruction, damage or disturbance out of context of fossils and fossil sites preserved at the ground surface or below ground represents a *direct negative* impact that is confined to the development footprint (*site specific*). Such impacts are made only during the construction period, and can usually be partially mitigated but cannot be fully rectified; *i.e.* they are *non-reversible* and of *permanent* duration. Since several of the sedimentary units represented within the study area do contain fossils of some sort, some level impact on fossil heritage is probable (*likely*). However, because of the generally very sparse occurrence of well-preserved, scientifically-valuable fossils within the superficial sediments, and because most of the fossils encountered are likely to be of widespread occurrence (low irreplaceability) the consequence of these impacts is rated as *slight*.

No previously recorded areas or sites of exceptional fossil heritage sensitivity or significance have been identified within the proposed project area as a whole. Due to the inferred scarcity of exceptional fossil remains within the study area, the overall impact significance of the construction phase of the proposed solar energy project is assessed as *VERY LOW* (without mitigation). Due to the paucity of palaeontological field studies within this part of Bushmanland, confidence levels for this desktop palaeontological heritage assessment are only moderate (medium).

Specialist palaeontological monitoring and mitigation for this project are not recommended, pending the discovery of new fossil sites during development, given its low impact significance. The Environmental Control Officer (ECO) responsible for the construction phase of the project should be aware of the necessity of conserving fossils and should monitor all substantial

excavations into sedimentary rocks for fossil remains. Proposed mitigation of chance fossil finds during the construction phase involves safeguarding of the fossils (preferably *in situ*) by the responsible ECO, reporting of finds to the SAHRA and, where appropriate, judicious sampling and recording of fossil material and associated geological data by a qualified palaeontologist (as discussed in Section 1.8 of this chapter). Should these recommended mitigation measures be fully implemented, the impact significance of the development would remain *VERY LOW* but small residual negative impacts (e.g. loss of undetected fossils) would remain. However, these negative impacts would be partially offset through the improved scientific understanding of local palaeontological heritage in a hitherto poorly-studied region of South Africa which would be considered as a significant *positive* outcome.

There are no fatal flaws in the proposed development proposal as far as fossil heritage is concerned.

1.6.2. Potential Impacts (Operational and Decommissioning Phases)

No significant impacts on fossil heritage resources are anticipated during the operational and decommissioning phases of the proposed solar energy facility.

1.6.3. Cumulative Impacts

The palaeontological heritage impact significance of all seven solar energy developments and associated electrical infrastructure proposed for Phase 2 of the Nieuwehoop Solar Development, as well as other proposed solar facilities and electrical infrastructure (discussed in Chapter 4 of the EIA Report) near Kenhardt (within a 20 km radius of the proposed project) are rated equally as very low. The potentially fossiliferous sedimentary rock units represented within the broader project area are of widespread occurrence and this is also likely to apply to most of the fossils they contain. It is concluded that the cumulative impact on fossil heritage resources posed by the proposed seven projects of the Nieuwehoop Solar Park Development and associated electrical infrastructure to the northeast of Kenhardt is of a low significance.

Given the generally low palaeontological sensitivity of the basement and overlying sedimentary rocks in the broader eastern Bushmanland region, significant cumulative impacts on fossil heritage are not anticipated here as a result of the various alternative energy and other infrastructure developments that have been proposed here (refer to the several recent palaeontological impact assessments undertaken by the author for projects near Kenhardt that are listed in the references).

1.7. IMPACT ASSESSMENT SUMMARY

The assessment of impacts on palaeontological heritage resources as well as recommended mitigation and monitoring measures, as discussed above, are collated in Tables 13.2 and 13.3 below.

The no-go option (no solar developments) will have a neutral impact on local palaeontological heritage resources.

Table 1.2 Impact assessment summary table for the Construction Phase

Construction Phase													
Direct Impacts													
Aspect/ Impact Pathway	Nature of Potential Impact/ Risk	Status	Spatial Extent	Duration	Consequence	Probability	Reversibility of Impact	Irreplac eability	Potential Mitigation Measures	Significance of Impact and Risk		Ranking of Residual Impact/ Risk	Confidence Level
										Without Mitigation/ Management	With Mitigation/ Management (Residual Impact/ Risk)		
Surface clearance and excavations into superficial sediments	Loss of fossil heritage at or beneath ground surface	Negative	Site	Permanent	Slight	Likely	Non- reversible	Low	<ul style="list-style-type: none"> Undertake monitoring of all substantial excavations into sedimentary rocks for fossil remains and safeguard any finds in situ. Appoint a professional palaeontologist to record and sample any chance fossil finds 	Very low	Very low	5	Medium

Table 1.3 Cumulative impact assessment summary table

Cumulative Impacts													
Aspect/ Impact Pathway	Nature of Potential Impact/ Risk	Status	Spatial Extent	Duration	Consequence	Probability	Reversibility of Impact	Irreplaceability	Potential Mitigation Measures	Significance of Impact and Risk		Ranking of Residual Impact/ Risk	Confidence Level
										Without Mitigation/ Management	With Mitigation/ Management (Residual Impact/ Risk)		
Surface clearance and excavation s into superficial sediments	Loss of fossil heritage at or beneath ground surface	Negative	Site	Permanent	Slight	Likely	Non- reversible	Low	<ul style="list-style-type: none"> Undertake monitoring of all substantial excavations into sedimentary rocks for fossil remains and safeguard any finds in situ. Appoint a professional palaeontologist to record and sample any chance fossil finds 	Very low	Very low	5	Medium

1.8. INPUT TO THE ENVIRONMENTAL MANAGEMENT PROGRAMME

Given the low palaeontological sensitivity of the proposed project area, as determined from desktop analysis, as well as the inferred very low impact significance of the alternative energy projects for fossil heritage conservation, no specialist palaeontological monitoring or mitigation is recommended here, pending the potential discovery of substantial new fossil remains during construction.

During the construction phase all substantial bedrock excavations should be monitored for fossil material by the responsible ECO. Should significant fossil remains - such as vertebrate bones and teeth, plant-rich fossil lenses, petrified wood or dense fossil burrow assemblages - be exposed during construction, the responsible ECO should safeguard these, preferably *in situ*. The SAHRA should be alerted as soon as possible (Contact details: Mrs Colette Scheermeyer, P.O. Box 4637, Cape Town 8000, Tel: 021 462 4502, Email: cscheermeyer@sahra.org.za), so that appropriate action can be taken by a professional palaeontologist, at the developer's expense. Mitigation would normally involve the scientific recording and judicious sampling or collection of fossil material as well as associated geological data (e.g. stratigraphy, sedimentology, taphonomy) by a professional palaeontologist.

The palaeontologist concerned with mitigation work will need a valid fossil collection permit from SAHRA and any material collected would have to be curated in an approved depository (e.g. museum or university collection). All palaeontological specialist work should conform to international best practice for palaeontological fieldwork and the study (e.g. data recording fossil collection and curation, final report) should adhere as far as possible to the minimum standards for Phase 2 palaeontological studies recently developed by SAHRA (2013).

No monitoring or mitigation is required during the operational and decommissioning phases of the development.

These mitigation recommendations (as summarised in Part B of the EIA Report) should be incorporated into the EMPr for each Solar PV energy facility proposed by Mulilo Renewable Project Developments.

1.9. CONCLUSION AND RECOMMENDATIONS

The study area for the Gemsbok Solar PV5 facility (including preferred and alternative sites as well as the associated 132 kV transmission line) is underlain at depth by Precambrian basement rocks (c. 1-2 billion years old) assigned to the Namaqua-Natal Province. These ancient igneous and high-grade metamorphic rocks - mainly granites and gneisses of the Keimoes Suite (granitoids) and Korannaland Supergroup (high grade metasediments) - crop out at surface in small areas and are entirely unfossiliferous. A large proportion of the basement rocks are mantled by a range of superficial sediments of Late Caenozoic age that may contain sparse fossil remains. These predominantly thin, unconsolidated deposits include small patches of calcretes, gravelly to sandy river alluvium, pan sediments, surface gravels, colluvium (scree) as well as Pleistocene to Recent wind-blown sands of the Gordonina Formation (Kalahari Group). Most of these younger rock units are of widespread occurrence and low palaeontological sensitivity. Scientifically important vertebrate fossil remains (e.g. Pleistocene mammalian bones and teeth) have been recorded within older stratified pan and river sediments elsewhere in the Bushmanland region where they are often associated with stone artefacts, while a limited range of trace fossils (e.g. plant root casts, termitaria and other invertebrate burrows) may be found within calcrete horizons.

No previously recorded areas or sites of exceptional fossil heritage sensitivity or significance have been identified within the Nieuwehoop Solar Park Development project area as a whole. Due to the inferred scarcity of scientifically important fossil remains within the Gemsbok Solar PV5 study area, the overall impact significance of the construction phase of the proposed solar energy project is assessed as VERY LOW (before and after mitigation). No significant impacts on fossil heritage are anticipated during the operational and decommissioning phases of the proposed solar energy facility. The potentially fossiliferous sedimentary rock units represented within the study area (e.g. Gordoniasands, calccrete) are of widespread occurrence and this is also likely to apply to most of the fossils they contain. It is concluded that the cumulative impacts on fossil heritage resources posed by the known alternative energy and other infrastructural developments in the region (as explained in Chapter 4 of the EIA Report) is of very low significance. There are no fatal flaws in the proposed solar facility development, nor are there objections to its authorisation as far as fossil heritage conservation is concerned, since significant impacts on scientifically valuable fossils or fossil sites are not anticipated here. The only proposed condition to accompany environmental authorisation is that the recommendations for monitoring and mitigation included in the EMPr are fully complied with. The no-go option (no solar developments) will have a neutral impact on local palaeontological heritage resources.

Given the low palaeontological sensitivity of the broader eastern Bushmanland region, as determined from desktop study, as well as the inferred very low impact significance of the Gemsbok Solar PV5 75 MW Solar PV Facility for fossil heritage conservation, no specialist palaeontological monitoring or mitigation is recommended here, pending the discovery of substantial new fossil remains during construction. Mitigation measures and monitoring recommendations for inclusion in the EMPr are discussed in Sections 1.6 and 1.8 of this report.

In this report the entire site (and transmission line) for the Gemsbok Solar PV3 75 MW Solar Photovoltaic (PV) Facility on Portion 3 of Gemsbok Bult Farm 120 has been assessed based on the worst case scenario. From a palaeontological heritage impact point of view, the applicant can select any 250 ha area within the surveyed area to build the PV plant, provided that the recommended mitigation measures are implemented as applicable.

1.10. REFERENCES

ALMOND, J.E. 2008a. Fossil record of the Loeriesfontein sheet area (1: 250 000 geological sheet 3018). Unpublished report for the Council for Geoscience, Pretoria, 32 pp.

ALMOND, J.E. 2009. Contributions to the palaeontology and stratigraphy of the Alexander Bay sheet area (1: 250 000 geological sheet 2816), 117 pp. Unpublished technical report prepared for the Council for Geoscience by Natura Viva cc, Cape Town.

ALMOND, J.E. 2011. Proposed Solar Cape Photovoltaic Electricity Generation Facility near Kenhardt, Northern Cape Province. Palaeontological impact assessment: desktop study, 18 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2014a. Proposed Exheredo CSP and PV solar energy facilities on the farm Styns Vley 280 near Kenhardt, Northern Cape Province. Palaeontological heritage assessment: desktop study, 28 pp. Natura Viva cc, Cape Town.

ALMOND, J.E. 2014b. Proposed Exheredo CSP and PV solar energy facilities near Kenhardt, Northern Cape Province. Palaeontological heritage assessment: combined desktop & field-based study, 61 pp. Natura Viva cc, Cape Town.

- ALMOND, J.E. 2014c. Proposed Gemsbok Solar PV1 Solar Energy Facility near Kenhardt, Northern Cape Province. Desktop study, 21 pp. Natura Viva cc, Cape Town.
- ALMOND, J.E. 2014d. Proposed Boven Solar PV1 Solar Energy Facility near Kenhardt, Northern Cape Province. Desktop study, 21 pp. Natura Viva cc, Cape Town.
- ALMOND, J.E. & PETHER, J. 2008. Palaeontological heritage of the Northern Cape. Interim SAHRA technical report, 124 pp. Natura Viva cc., Cape Town.
- BAMFORD, M.K. 2000. Cenozoic macro-plants. In: Partridge, T.C. & Maud, R.R. (Eds.) The Cenozoic of southern Africa, pp.351-356. Oxford University Press, Oxford.
- BAMFORD, M.K. & DE WIT, M.C.J. 1993. Taxonomic description of fossil wood from Cainozoic Sak River terraces, near Brandvlei, Bushmanland, South Africa. *Palaeontologia africana* 30: 71-80.
- BENDER, P.A. & BRINK, J.S. 1992. A preliminary report on new large mammal fossil finds from the Cornelia-Uitzoek site. *South African Journal of Science* 88: 512-515.
- BOUSMAN, C.B. et al. 1988. Palaeoenvironmental implications of Late Pleistocene and Holocene valley fills in Blydefontein Basin, Noupoot, C.P., South Africa. *Palaeoecology of Africa* 19: 43-67.
- BRINK, J.S. 1987. The archaeozoology of Florisbad, Orange Free State. *Memoirs van die Nasionale Museum* 24, 151 pp.
- BRINK, J.S. et al. 1995. A new find of *Megalotragus priscus* (Alcephalini, Bovidae) from the Central Karoo, South Africa. *Palaeontologia africana* 32: 17-22.
- BRINK, J.S. & ROSSOUW, L. 2000. New trial excavations at the Cornelia-Uitzoek type locality. *Navorsing van die Nasionale Museum Bloemfontein* 16, 141-156.
- BUTZER, K.W., HELGREN, D.M., FOCK, G. & STUCKENRATH, R. 1973. Alluvial terraces of the Lower Vaal River, South Africa: a re-appraisal and re-investigation. *Journal of geology* 81, 341-362.
- CHURCHILL, S.E. et al. 2000. Erfkroon: a new Florisian fossil locality from fluvial contexts in the western Free State, South Africa. *South African Journal of Science* 96: 161-163.
- COOKE, H.B.S. 1949. Fossil mammals of the Vaal River deposits. *Memoirs of the geological Survey of South Africa* 35, 1-117.
- CORNELL, D.H., THOMAS, R.J., MOEN, H.F.G., REID, D.L., MOORE, J.M. & GIBSON, R.L. 2006. The Namaqua-Natal Province. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) *The geology of South Africa*, pp. 461-499. Geological Society of South Africa, Marshalltown.
- DE WIT, M.C.J. 1990. Palaeoenvironmental interpretation of Tertiary sediments at Bosluispan, Namaqualand. *Palaeoecology of Africa and the surrounding islands* 21: 101-118.
- DE WIT, M.C.J. 1993. Cainozoic evolution of drainage systems in the north-western Cape. Unpublished PhD thesis, University of Cape Town, Cape Town, 371 pp.

- DE WIT, M.C.J. 1999. Post-Gondwana drainage and the development of diamond placers in western South Africa. *Economic Geology* 94: 721-740.
- DE WIT, M.C.J. & BAMFORD, M.K. 1993. Fossil wood from the Brandvlei area, Bushmanland as an indication of palaeoenvironmental changes during the Cainozoic. *Palaeontologia africana* 30: 81-89.
- DE WIT, M.C.J., MARSHALL, T.R. & PARTRIDGE, T.C. 2000. Fluvial deposits and drainage evolution. In: Partridge, T.C. & Maud, R.R. (Eds.) *The Cenozoic of southern Africa*, pp.55-72. Oxford University Press, Oxford.
- DINGLE, R.V., SIESSER, W.G. & NEWTON, A.R. 1983. *Mesozoic and Tertiary geology of southern Africa*. viii + 375 pp. Balkema, Rotterdam.
- DU TOIT, A. 1954. *The geology of South Africa*. xii + 611pp, 41 pls. Oliver & Boyd, Edinburgh.
- HADDON, I.G. 2000. Kalahari Group sediments. In: Partridge, T.C. & Maud, R.R. (Eds.) *The Cenozoic of southern Africa*, pp. 173-181. Oxford University Press, Oxford.
- HELGREN, D.M. 1977. Geological context of the Vaal River faunas. *South African Journal of Science* 73, 303-307.
- KIBERD, P. 2006. Bundu Farm: a report on archaeological and palaeoenvironmental assemblages from a pan site in Bushmanland, Northern Cape, South Africa. *South African Archaeological Bulletin* 61, 189-201.
- KLEIN, R. 1980. Environmental and ecological implications of large mammals from Upper Pleistocene and Hooocene sites in southern Africa. *Annals of the South African Museum* 81, 223-283.
- KLEIN, R.G. 1984. The large mammals of southern Africa: Late Pliocene to Recent. In: Klein, R.G. (Ed.) *Southern African prehistory and paleoenvironments*, pp 107-146. Balkema, Rotterdam.
- KLEIN, R. 2000. The Earlier Stone Age in southern Africa. *The South African Archaeological Bulletin* 40, 107-122.
- MACEY, P.H., SIEGFRIED, H.P., MINNAAR, H., ALMOND, J. AND BOTHA, P.M.W. 2011. The geology of the Loeriesfontein Area. Explanation to 1: 250 000 Geology Sheet 3018 Loeriesfontein, 139 pp. Council for Geoscience, Pretoria.
- MACRAE, C. 1999. *Life etched in stone. Fossils of South Africa*. 305 pp. The Geological Society of South Africa, Johannesburg.
- MEADOWS, M.E. & WATKEYS, M.K. 1999. Palaeoenvironments. In: Dean, W.R.J. & Milton, S.J. (Eds.) *The karoo. Ecological patterns and processes*, pp. 27-41. Cambridge University Press, Cambridge.
- ORTON, J. 2012. Heritage impact assessment for a proposed solar energy facility on the farm Hoekplaas near Copperton, Northern Cape, 32 pp. Archaeology Contracts Office, University of Cape Town, Cape Town.

- PARTRIDGE, T.C. & SCOTT, L. 2000. Lakes and Pans. In: Partridge, T.C. & Maud, R.R. (Eds.) *The Cenozoic of southern Africa*, pp.145-161. Oxford University Press, Oxford.
- PARTRIDGE, T.C., BOTHA, G.A. & HADDON, I.G. 2006. Cenozoic deposits of the interior. In: Johnson, M.R., Anhaeusser, C.R. & Thomas, R.J. (Eds.) *The geology of South Africa*, pp. 585-604. Geological Society of South Africa, Marshalltown.
- PICKFORD, M. & SENUT, B. 2002. The fossil record of Namibia. 39 pp. The Geological Survey of Namibia.
- ROSSOUW, L. 2006. Florisian mammal fossils from erosional gullies along the Modder River at Mitasrust Farm, Central Free State, South Africa. *Navorsing van die Nasionale Museum Bloemfontein* 22, 145-162.
- SAHRA 2013. Minimum standards: palaeontological component of heritage impact assessment reports, 15 pp. South African Heritage Resources Agency, Cape Town.
- SCHNEIDER, G. & MARAIS, C. 2004. Passage through time. The fossils of Namibia. 158 pp. Gamsberg MacMillan, Windhoek.
- SCOTT, L. 2000. Pollen. In: Partridge, T.C. & Maud, R.R. (Eds.) *The Cenozoic of southern Africa*, pp.339-35. Oxford University Press, Oxford.
- SENUT, B., PICKFORD, M., WARD, J., DE WIT, M., SPAGGIARI, R. & MORALES, J. 1996. Biochronology of the Cainozoic sediments at Bosluis Pan, Northern Cape Province, South Africa. *South African Journal of Science* 92: 249-251.
- SIEBRITS, L.B. 1989. Die geologie van die gebied Sakrivier. Explanation of 1: 250 000 geology sheet 3020, 19 pp. Council for Geoscience, Pretoria.
- SKEAD, C.J. 1980. Historical mammal incidence in the Cape Province. Volume 1: The Western and Northern Cape, 903pp. Department of Nature and Environmental Conservation, Cape Town.
- SLABBERT, M.J., MOEN, H.F.G. & BOELEMA, R. 1999. Die geologie van die gebied Kenhardt. Explanation to 1: 250 000 geology Sheet 2920 Kenhardt, 123 pp. Council for Geoscience, Pretoria.
- SMITH, A.B. 1999. Hunters and herders in the Karoo landscape. Chapter 15 in Dean, W.R.J. & Milton, S.J. (Eds.) *The Karoo; ecological patterns and processes*, pp. 243-256. Cambridge University Press, Cambridge.
- THOMAS, M.J. 1981. The geology of the Kalahari in the Northern Cape Province (Areas 2620 and 2720). Unpublished MSc thesis, University of the Orange Free State, Bloemfontein, 138 pp.
- THOMAS, R.J., THOMAS, M.A. & MALHERBE, S.J. 1988. The geology of the Nossob and Twee Rivieren areas. Explanation for 1: 250 000 geology sheets 2520-2620. 17pp. Council for Geoscience, Pretoria.
- THOMAS, D.S.G. & SHAW, P.A. 1991. *The Kalahari environment*, 284 pp. Cambridge University Press.

WELLS, L.H. 1964. The Vaal River 'Younger Gravels" faunal assemblage: a revised list. South African Journal of Science 60, 92-94.