

**Palaeontological Impact Assessment for  
the proposed development of the  
Harmony Moab PV Facility, southeast  
of Orkney,  
Free State Province**

**CTS22\_101**

**Desktop Study (Phase 1)**

**For**

**CTS Heritage**

**03 July 2022**

**Prof Marion Bamford**

Palaeobotanist

P Bag 652, WITS 2050

Johannesburg, South Africa  
[Marion.bamford@wits.ac.za](mailto:Marion.bamford@wits.ac.za)


## **Expertise of Specialist**

The Palaeontologist Consultant: Prof Marion Bamford  
Qualifications: PhD (Wits Univ, 1990); FRSSAf, mASSAf  
Experience: 33 years research and lecturing in Palaeontology  
25 years PIA studies and over 300 projects completed

## **Declaration of Independence**

This report has been compiled by Professor Marion Bamford, of the University of the Witwatersrand, sub-contracted by CTS Heritage, Cape Town, South Africa. The views expressed in this report are entirely those of the author and no other interest was displayed during the decision making process for the Project.

Specialist: Prof Marion Bamford

Signature: 

## **Executive Summary**

A Palaeontological Impact Assessment was requested for the proposed Moab PV facility for Harmony Mine, southeast of Orkney, near the town of Viljoenskroon, Free State.

To comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development.

The proposed site lies on the potentially fossiliferous Malmani Subgroup that could have trace fossils such as stromatolites below the land surface. The southernmost part is on moderately fossiliferous sands and alluvium of the Quaternary that overlies the Vryheid Formation. The area has been greatly disturbed by farming and mining activities and no potential rocky outcrops of dolomite or of traps for Quaternary fossils (pans) are visible from the satellite imagery. Nonetheless, a Fossil Chance Find Protocol should be added to the EMPr. Based on this information it is recommended that no further palaeontological impact assessment is required unless fossils are found by the contractor, environmental officer or other designated responsible person once excavations have commenced. Since the impact will be low, as far as the palaeontology is concerned, the project should be authorised.

## Table of Contents

Expertise of Specialist.....	1
Declaration of Independence.....	1
1. Background.....	4
2. Methods and Terms of Reference.....	7
3. Geology and Palaeontology.....	7
i. Project location and geological context.....	7
ii. Palaeontological context.....	9
4. Impact assessment.....	11
5. Assumptions and uncertainties.....	12
6. Recommendation.....	13
7. References.....	13
8. Chance Find Protocol.....	14
9. Appendix A - Examples of fossils .....	15
10. Appendix B - Details of specialist.....	17

Figure 1: Google Earth map of the general area to show the relative land marks.  
..... 6

Figure 2: Google Earth Map of the proposed development .....6

Figure 3: Geological map of the area around the project site.....7

Figure 4: SAHRIS palaeosensitivity map for the site .....10

## i. Background


Harmony Gold mines is proposing to develop photovoltaic farms (PVs) with grid connections on some of its properties in the Free State. This report is for the Moab PV Facility situated north and south of the R76 close to the town of Viljoenskroon in the Free State Province, southeast of Orkney. It falls within the jurisdiction of the Moqhaka Local Municipality in the Fezile Dabi District Municipality and is located within the existing Harmony Mine (Figures 1-2).

A Palaeontological Impact Assessment was requested for the Moab PV project. To comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development and is reported herein.


Table 1: National Environmental Management Act, 1998 (Act No. 107 of 1998) (NEMA) and Environmental Impact Assessment (EIA) Regulations, 2014 (as amended) - Requirements for Specialist Reports (Appendix 6).

	<b>A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:</b>	<b>Relevant section in report</b>
ai	Details of the specialist who prepared the report,	Appendix B
aii	The expertise of that person to compile a specialist report including a curriculum vitae	Appendix B
b	A declaration that the person is independent in a form as may be specified by the competent authority	Page 1
c	An indication of the scope of, and the purpose for which, the report was prepared	Section i.
ci	An indication of the quality and age of the base data used for the specialist report: SAHRIS palaeosensitivity map accessed - date of this report	Yes
cii	A description of existing impacts on the site, cumulative impacts of the proposed development and levels of acceptable change	Section 5
d	The date and season of the site investigation and the relevance of the season to the outcome of the assessment	N/A
e	A description of the methodology adopted in preparing the report or carrying out the specialised process	Section ii.
f	The specific identified sensitivity of the site related to the activity and its	Section 4

	<b>A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:</b>	<b>Relevant section in report</b>
	associated structures and infrastructure	
g	An identification of any areas to be avoided, including buffers	N/A
h	A map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	N/A
i	A description of any assumptions made and any uncertainties or gaps in knowledge;	Section vii.
j	A description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives, on the environment	Section vi.
k	Any mitigation measures for inclusion in the EMPr	Section 8, Appendix A
l	Any conditions for inclusion in the environmental authorisation	N/A
m	Any monitoring requirements for inclusion in the EMPr or environmental authorisation	Section 8, Appendix A
ni	A reasoned opinion as to whether the proposed activity or portions thereof should be authorised	Section 6
nii	If the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan	Sections 6, 8
o	A description of any consultation process that was undertaken during the course of carrying out the study	N/A
p	A summary and copies of any comments that were received during any consultation process	N/A
q	Any other information requested by the competent authority.	N/A
2	Where a government notice gazetted by the Minister provides for any protocol or minimum information requirement to be applied to a specialist report, the requirements as indicated in such notice will apply.	N/A



**Figure 1: Google Earth map of the general area to show the relative land marks. The Moab PV facility is shown by the labels.**



**Figure 2: Google Earth Map of the proposed development of the Moab PV facility with the sections shown by the pink and yellow outlines.**





**Figure 3: Geological map of the area around the proposed Moab PV facility. The location of the proposed project is indicated within the yellow rectangle. Abbreviations of the rock types are explained in Table 2. Map enlarged from the Geological Survey 1: 250 000 map 2626 West Rand.**

Table 2: Explanation of symbols for the geological map and approximate ages (Eriksson et al., 2006. Johnson et al., 2006; Partridge et al., 2006). SG = Supergroup; Fm = Formation; Ma = million years; grey shading = formations impacted by the project.

<b>Symbo l</b>	<b>Group/Formation</b>	<b>Lithology</b>	<b>Approximate Age</b>
Q	Quaternary	Alluvium, sand, soil	Quaternary, ca 1.0 Ma to present
Pv	Vryheid Fm, Eccca Group, Karoo SG	Shales, siltstone, mudstone, sandstone, cola	Early Permian Ca 290-280 Ma
Vdi	Diabase	Diabase (intrusive volcanic rocks)	Post Transvaal SG
Vh	Hekpoort Fm, Pretoria Group, Transvaal SG	Mafic lavas	Palaeoproterozoic Ca 2224 Ma
Vmd	Malmani Subgroup, Chuniespoort Group, Transvaal SG	Dolomite, limestone, chert	Palaeoproterozoic Ca 2500 Ma

The project lies in the north-central part of the main Karoo Basin where it unconformably overlies the southern margin of the Transvaal Basin that preserves the Transvaal Supergroup. Much of the area is covered by young sands and alluvium of Quaternary age (Figure 3).

In the Transvaal Basin the Transvaal Supergroup is divided into two Groups, the lower Chuniespoort Group and the upper Pretoria Group (with ten formations; Eriksson et al., 2006). The Chuniespoort Group is divided into the basal **Malmani Subgroup** that comprises dolomites and limestones and is divided into five formations based on chert content, stromatolitic morphology, intercalated shales and erosion surfaces. The top of the Chuniespoort Group has the Penge Formation and the Duitschland Formation.

Making up the lower Pretoria Group are the Timeball Hill Formation and the Boshhoek Formation. The Hekpoort, Dwaalheuwel, Strubenkop and Daspoort Formations form a sequence as the middle part of the Pretoria Group, Transvaal Supergroup, and represent rocks that are over 2060 million years old. The **Hekpoort Formation** is a massive lava deposit and is overlain by the Dwaalheuwel conglomerates, siltstone and sandstone (not present here). The Transvaal sequence has been interpreted as three major cycles of basin infill and tectonic activity with

the first deep basin sediments forming the Chuniespoort Group, the second cycle deposited the lower Pretoria Group, and the sediments in this area are from the interim lowstand that preceded the third cycle. These sediments were deposited in shallow lacustrine, alluvial fan and braided stream environments (Eriksson et al., 2012).

The Karoo Supergroup rocks cover a very large proportion of South Africa and extend from the northeast (east of Pretoria) to the southwest and across to almost the KwaZulu Natal south coast. It is bounded along the southern margin by the Cape Fold Belt and along the northern margin by the much older Transvaal Supergroup rocks. Representing some 120 million years (300 – 183Ma), the Karoo Supergroup rocks have preserved a diversity of fossil plants, insects, vertebrates and invertebrates.

During the Carboniferous Period South Africa was part of the huge continental landmass known as Gondwanaland and it was positioned over the South Pole. As a result, there were several ice sheets that formed and melted, and covered most of South Africa (Visser, 1986, 1989; Isbell et al., 2012). Gradual melting of the ice as the continental mass moved northwards and the earth warmed, formed fine-grained sediments in the large inland sea (Johnson et al., 2006).

Overlying the Dwyka Group rocks are rocks of the Ecca Group that are Early Permian in age. There are eleven formations recognised in this group but they do not all extend throughout the Karoo Basin. In the west and central part are the following formations, from base upwards: **Vryheid** and Volksrust Formations. All of these sediments have varying proportions of sandstones, mudstones, shales and siltstones and represent shallow to deep water settings, deltas, rivers, streams and overbank depositional environments.

#### v. Palaeontological context

The palaeontological sensitivity of the area under consideration is presented in Figure 4. The site for development is mostly in the Malmani Subgroup dolomites and cherts and the southernmost part is on Quaternary sands that overlie the Vryheid Formation. The Malmani Subgroup is considered as very highly sensitive for palaeontology as the dolomites sometimes contain trace fossils, stromatolites. The Vryheid Formation is also very highly sensitive because it could have fossil plants of the *Glossopteris* flora.


Stromatolites are the trace fossils that were formed by colonies of green algae and blue-green algae (Cyanobacteria) that grew in warm, shallow marine settings. These algae were responsible for releasing oxygen via the photosynthetic process where atmospheric carbon dioxide and water, using energy from the sun, are converted into carbon chains and

compounds that are the building blocks of all living organisms. The released carbon dioxide initially was taken up by the abundant reducing minerals to form oxides, e.g. iron oxide. Eventually free oxygen was released into the atmosphere and some was converted into ozone by the bombardment of cosmic rays. The ozone is critical for the filtering out of harmful ultraviolet rays.

Stromatolites are the layers upon layers of inorganic materials that were deposited during photosynthesis, namely calcium carbonate, magnesium carbonate, calcium sulphate and magnesium sulphate. These layers can be in the form of flat layers, domes or columns depending on the environment where they grew (Beukes, 1987). Some environments did not form stromatolites, just layers of limestone that later was converted to dolomite. The algae that formed the stromatolites are very rarely preserved, and they are microscopic so they can only be seen from thin sections studies under a petrographic microscope.

The Vryheid Formation is well known for the fossils of the *Glossopteris* flora that was widespread over Gondwanaland. The flora was dominated by *Glossopteris*, the leaves of glossopterid plants, as well as seeds, fructifications, roots and silicified wood. Other plants in this flora were lycopods, sphenophytes, ferns, cordaitaleans and early gymnosperms (Plumstead, 1969; Anderson and Anderson, 1985; Bamford, 2004). Coal seams were formed from buried peats of this flora that were later altered by heat and pressure. While no plants are discernible from coal itself, it is possible to find impressions of the plants in the shales and carbonaceous shales in lenses associated with the coals. In this part of the Free State the uppermost coal seam is between 25 and 50 m below the surface (Snyman, 1998; p 177).

Quaternary sands and alluvium do not preserve fossils because they are transported and porous. For preservation of fossils, a low energy deposit with sedimentation of fine grained silts or muds that exclude decomposing organisms such as bacteria, fungi and invertebrates is required to maintain a highly reducing environment (Cowan, 1995). Only if there are traps such as palaeo-pans or palaeo-springs that provide traps for water and fine sediments, would plants or bones be preserved and fossilised. No such features are visible in the satellite imagery in the project footprint.



**Figure 4: SAHRIS palaeosensitivity map for the site for the proposed Moab PV Facility shown within the lilac and pink outlines rectangle. Background colours indicate the following degrees of sensitivity: red = very highly sensitive; orange/yellow = high; green = moderate; blue = low; grey = insignificant/zero.**

From the SAHRIS map above the area is indicated as very highly sensitive (red) for the Malmani subgroup and moderately sensitive (green) for the Quaternary sands and alluvium that overlie the Vryheid Formation. Note the artificially horizontal contact line between the two strata that is based on the different interpretations of the adjacent geological maps. Both alternatives are on very highly sensitive rocks that are covered by soils and sands.

## vi. Impact assessment

An assessment of the potential impacts to possible palaeontological resources considers the criteria encapsulated in Table 3:

**Table 3a: Criteria for assessing impacts**

<b>PART A: DEFINITION AND CRITERIA</b>		
<b>Criteria for ranking of the SEVERITY/NATURE of environmental impacts</b>	<b>H</b>	Substantial deterioration (death, illness or injury). Recommended level will often be violated. Vigorous community action.
	<b>M</b>	Moderate/ measurable deterioration (discomfort). Recommended level will occasionally be violated. Widespread complaints.
	<b>L</b>	Minor deterioration (nuisance or minor deterioration). Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	<b>L+</b>	Minor improvement. Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	<b>M+</b>	Moderate improvement. Will be within or better than the recommended level. No observed reaction.
	<b>H+</b>	Substantial improvement. Will be within or better than the recommended level. Favourable publicity.
<b>Criteria for ranking the DURATION of impacts</b>	<b>L</b>	Quickly reversible. Less than the project life. Short term
	<b>M</b>	Reversible over time. Life of the project. Medium term
	<b>H</b>	Permanent. Beyond closure. Long term.
<b>Criteria for ranking the SPATIAL SCALE of impacts</b>	<b>L</b>	Localised - Within the site boundary.
	<b>M</b>	Fairly widespread - Beyond the site boundary. Local
	<b>H</b>	Widespread - Far beyond site boundary. Regional/ national
<b>PROBABILITY (of exposure to impacts)</b>	<b>H</b>	Definite/ Continuous
	<b>M</b>	Possible/ frequent
	<b>L</b>	Unlikely/ seldom

**Table 3b: Impact Assessment**

<b>PART B: Assessment</b>		
<b>SEVERITY/ NATURE</b>	<b>H</b>	-
	<b>M</b>	-
	<b>L</b>	Quaternary sands and soils do not preserve fossils; so far there are no records from the Quaternary or from the Malmani Subgroup of plant or animal fossils, or stromatolites, respectively, in this region so it is very unlikely that fossils occur on the site. The impact would be negligible
	<b>L+</b>	-
	<b>M+</b>	-

<b>PART B: Assessment</b>		
	<b>H</b> <b>+</b>	-
<b>DURATION</b>	<b>L</b>	-
	<b>M</b>	-
	<b>H</b>	Where manifest, the impact will be permanent.
<b>SPATIAL SCALE</b>	<b>L</b>	Since the only possible fossils within the area would be trace fossils in the dolomites of the Malmani subgroup, the spatial scale will be localised within the site boundary.
	<b>M</b>	-
	<b>H</b>	-
<b>PROBABILITY</b>	<b>H</b>	-
	<b>M</b>	-
	<b>L</b>	It is very unlikely that any fossils would be found in the loose soils and sands that cover the area or in the dolomites that might occur below them. Nonetheless, a Fossil Chance Find Protocol should be added to the eventual EMPr.

Based on the nature of the project, surface activities may impact upon the fossil heritage if preserved in the development footprint. The geological structures suggest that the rocks are the right age to contain fossils but are covered by soils. Furthermore, the material to be excavated are soils and this does not preserve fossils. Since there is a small chance that trace fossils from the Malmani Subgroup or plant fossils in the Vryheid Formation that might occur below the soils and may be disturbed a Fossil Chance Find Protocol has been added to this report. Taking account of the defined criteria, the potential impact to fossil heritage resources is low.

## vii. Assumptions and uncertainties

Based on the geology of the area and the palaeontological record as we know it, it can be assumed that the formation and layout of the dolomites, sandstones, shales and sands are typical for the country and might contain trace fossils or fossil plant, insect, invertebrate and vertebrate material. The sands of the Quaternary period would not preserve fossils. The area has been disturbed from farming and mining so no fossils would be present on the surface. It is unknown if the Malmani dolomite occurs below the surface or if it has stromatolites.

## viii. Recommendation

Based on experience and the lack of any previously recorded fossils from the area, it is extremely unlikely that any fossils would be preserved in the overlying sands and soils of the Quaternary. There is a very small chance that fossils may occur in the adjacent shales of the early Permian Vryheid Formation, or trace fossils in the Malmani Subgroup, so a Fossil Chance Find Protocol should be added to the EMPr. If fossils are found by the environmental officer, or other responsible person once excavations for foundations and amenities have commenced then they should be rescued and a palaeontologist called to assess and collect a representative sample. Since the impact on the palaeontological heritage would be low, as far as the palaeontology is concerned, the project should be authorised. There is no preferred alternative as far as the palaeontology is concerned.

## ix. References

Anderson, J.M., Anderson, H.M., 1985. Palaeoflora of Southern Africa: Prodrum of South African megaflores, Devonian to Lower Cretaceous. A.A. Balkema, Rotterdam. 423 pp.

Bamford, M.K. 2004. Diversity of woody vegetation of Gondwanan southern Africa. *Gondwana Research* 7, 153-164.

Beukes, N.J., 1987. Facies relations, depositional environments and diagenesis in a major early Proterozoic stromatolitic carbonate platform to basinal sequence, Campbellrand Subgroup, Transvaal Supergroup, southern Africa. *Sedimentary Geology* 54, 1-46.

Isbell, J.L., Henry, L.C., Gulbranson, E.L., Limarino, C.O., Fraiser, F.L., Koch, Z.J., Ciccio, P.I., Dineen, A.A., 2012. Glacial paradoxes during the late Paleozoic ice age: Evaluating the equilibrium line altitude as a control on glaciation. *Gondwana Research* 22, 1-19.

Johnson, M.R., van Vuuren, C.J., Visser, J.N.J., Cole, D.I., Wickens, H.deV., Christie, A.D.M., Roberts, D.L., Brandl, G., 2006. Sedimentary rocks of the Karoo Supergroup. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. Pp 461 - 499.

Plumstead, E.P., 1969. Three thousand million years of plant life in Africa. Geological Society of southern Africa, Annexure to Volume LXXII. 72pp + 25 plates.



Visser, J.N.J., 1986. Lateral lithofacies relationships in the glaciogene Dwyka Formation in the western and central parts of the Karoo Basin. Transactions of the Geological Society of South Africa 89, 373-383.


Visser, J.N.J., 1989. The Permo-Carboniferous Dwyka Formation of southern Africa: deposition by a predominantly subpolar marine icesheet. Palaeogeography, Palaeoclimatology, Palaeoecology 70, 377-391.

## x. Chance Find Protocol


### **Monitoring Programme for Palaeontology - to commence once the excavations / drilling activities begin.**

1. The following procedure is only required if fossils are seen on the surface and when excavations commence.
2. When excavations begin the rocks and must be given a cursory inspection by the environmental officer or designated person. Any fossiliferous material (stromatolites or plants) should be put aside in a suitably protected place. This way the project activities will not be interrupted.
3. Photographs of similar fossils must be provided to the developer to assist in recognizing the fossil plants, vertebrates, invertebrates or trace fossils in the shales and mudstones (for example see Figures 5-6). This information will be built into the EMP's training and awareness plan and procedures.
4. Photographs of the putative fossils can be sent to the palaeontologist for a preliminary assessment.
5. If there is any possible fossil material found by the developer/environmental officer then the qualified palaeontologist sub-contracted for this project, should visit the site to inspect the selected material and check the dumps where feasible.
6. Trace fossils, fossil plants or vertebrates that are considered to be of good quality or scientific interest by the palaeontologist must be removed, catalogued and housed in a suitable institution where they can be made available for further study. Before the fossils are removed from the site a SAHRA permit must be obtained. Annual reports must be submitted to SAHRA as required by the relevant permits.
7. If no good fossil material is recovered then no site inspections by the palaeontologist will be necessary. A final report by the palaeontologist must be sent to SAHRA once the project has been completed and only if there are fossils.
8. If no fossils are found and the excavations have finished then no further monitoring is required.

xi. Appendix A – Examples of fossils from the Malmani Subgroup



**Figure 5: Photographs of different types of stromatolites that occur in the Malmani dolomite in some areas.**



**Figure 6: Photographs of fossil plants of the *Glossopteris* flora that could occur in the shales of the Vryheid Formation.**

xii. Appendix B – Details of specialist

**Curriculum vitae (short) - Marion Bamford  
PhD**

**June 2022**

**I) Personal details**

Surname : **Bamford**  
First names : **Marion Kathleen**  
Present employment : Professor; Director of the Evolutionary  
Studies Institute.  
Member Management Committee of the NRF/DST  
Centre of  
Excellence Palaeosciences, University of the  
Witwatersrand,  
Johannesburg, South Africa  
Telephone : +27 11 717 6690  
Fax : +27 11 717 6694  
Cell : 082 555 6937  
E-mail : [marion.bamford@wits.ac.za](mailto:marion.bamford@wits.ac.za) ;  
[marionbamford12@gmail.com](mailto:marionbamford12@gmail.com)

**ii) Academic qualifications**

Tertiary Education: All at the University of the Witwatersrand:  
1980-1982: BSc, majors in Botany and Microbiology. Graduated April  
1983.  
1983: BSc Honours, Botany and Palaeobotany. Graduated April 1984.  
1984-1986: MSc in Palaeobotany. Graduated with Distinction, November  
1986.  
1986-1989: PhD in Palaeobotany. Graduated in June 1990.  
NRF Rating: C-2 (1999-2004); B-3 (2005-2015); B-2 (2016-2020); B-1  
(2021-2026)

**iii) Professional qualifications**

*Wood Anatomy Training (overseas as nothing was available in South  
Africa):*

1994 - Service d'Anatomie des Bois, Musée Royal de l'Afrique Centrale,  
Tervuren, Belgium, by Roger Dechamps  
1997 - Université Pierre et Marie Curie, Paris, France, by Dr Jean-Claude  
Koeniguer  
1997 - Université Claude Bernard, Lyon, France by Prof Georges Barale,  
Dr Jean-Pierre Gros, and Dr Marc Philippe

#### **iv) Membership of professional bodies/associations**

Palaeontological Society of Southern Africa  
Royal Society of Southern Africa - Fellow: 2006 onwards  
Academy of Sciences of South Africa - Member: Oct 2014 onwards  
International Association of Wood Anatomists - First enrolled: January 1991  
International Organization of Palaeobotany - 1993+  
Botanical Society of South Africa  
South African Committee on Stratigraphy - Biostratigraphy - 1997 - 2016  
SASQUA (South African Society for Quaternary Research) - 1997+  
PAGES - 2008 -onwards: South African representative  
ROCEEH / WAVE - 2008+  
INQUA - PALCOMM - 2011+onwards

#### **vii) Supervision of Higher Degrees**

All at Wits University

Degree	Graduated/ completed	Current
Honours	13	0
Masters	11	3
PhD	11	6
Postdoctoral fellows	15	1

#### **viii) Undergraduate teaching**

Geology II - Palaeobotany GEOL2008 - average 65 students per year  
Biology III - Palaeobotany APES3029 - average 45 students per year  
Honours - Evolution of Terrestrial Ecosystems; African Plio-Pleistocene Palaeoecology; Micropalaeontology - average 12-20 students per year.

#### **ix) Editing and reviewing**

Editor: Palaeontologia africana: 2003 to 2013; 2014 - Assistant editor  
Guest Editor: Quaternary International: 2005 volume  
Member of Board of Review: Review of Palaeobotany and Palynology: 2010 -  
Associate Editor Open Science UK: 2021 -  
Review of manuscripts for ISI-listed journals: 30 local and international journals  
Reviewing of funding applications for NRF, PAST, NWO, SIDA, National Geographic, Leakey Foundation

#### **x) Palaeontological Impact Assessments**

Selected from the past five years only - list not complete:

- Mala Mala 2017 for Henwood
- Modimolle 2017 for Green Vision
- Klippoortjie and Finaalspan 2017 for Delta BEC
- Ledjadja borrow pits 2018 for Digby Wells
- Lungile poultry farm 2018 for CTS
- Olienhout Dam 2018 for JP Celliers
- Isondlo and Kwasobabili 2018 for GCS

- Kanakies Gypsum 2018 for Cabanga
- Nababeep Copper mine 2018
- Glencore-Mbali pipeline 2018 for Digby Wells
- Remhoogte PR 2019 for A&HAS
- Bospoort Agriculture 2019 for Kudzala
- Overlooked Quarry 2019 for Cabanga
- Richards Bay Powerline 2019 for NGT
- Eilandia dam 2019 for ACO
- Eastlands Residential 2019 for HCAC
- Fairview MR 2019 for Cabanga
- Graspan project 2019 for HCAC
- Lieliefontein N&D 2019 for EnviroPro
- Skeerpoort Farm Mast 2020 for HCAC
- Vulindlela Eco village 2020 for 1World
- KwaZamakhule Township 2020 for Kudzala
- Sunset Copper 2020 for Digby Wells
- McCarthy-Salene 2020 for Prescali
- VLNR Lodge 2020 for HCAC
- Madadeni mixed use 2020 for EnviroPro
- Frankfort-Windfield Eskom Powerline 2020 for 1World
- Beaufort West PV Facility 2021 for ACO Associates
- Copper Sunset MR 2021 for Digby Wells
- Sannaspos PV facility 2021 for CTS Heritage
- Smithfield-Rouxville-Zastron PL 2021 for TheroServe

#### **xi) Research Output**

Publications by M K Bamford up to January 2022 peer-reviewed journals or scholarly books: over 160 articles published; 5 submitted/in press; 10 book chapters.

Scopus h-index = 30; Google scholar h-index = 35; -i10-index = 92

Conferences: numerous presentations at local and international conferences.