

**Palaeontological Impact Assessment for the
proposed new haul road and storm water
management for Palingpan Manganese Mine,
Northern Cape Province**

Desktop Study (Phase 1)

For

Lokisa Environmental Consulting, cc

16 February 2023

Prof Marion Bamford

Palaeobotanist

P Bag 652, WITS 2050

Johannesburg, South Africa

Marion.bamford@wits.ac.za

Expertise of Specialist

The Palaeontologist Consultant: Prof Marion Bamford
Qualifications: PhD (Wits Univ, 1990); FRSSAf, mASSAf
Experience: 34 years research and lecturing in Palaeontology
26 years PIA studies and over 350 projects completed

Declaration of Independence

This report has been compiled by Professor Marion Bamford, of the University of the Witwatersrand, sub-contracted by Lokisa Environmental Consulting cc, Groenkloof, South Africa. The views expressed in this report are entirely those of the author and no other interest was displayed during the decision making process for the Project.

Specialist: Prof Marion Bamford

Signature:

A handwritten signature in blue ink, appearing to read 'MKBamford', with a horizontal line underneath it.

Executive Summary

A Palaeontological Impact Assessment was requested for the proposed construction of a ±5km new gravel road that starts at the R325 (14.6 km from Postmasburg) and ends at the railway line.

To comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development.

The proposed road route lies on the potentially fossiliferous Ghaap Group dolomites, on what was previously known as the Lime Acres Member/Formation that might preserve trace fossils such as stromatolites. The very high palaeosensitivity indicated in the SAHRIS palaeosensitivity map is challenged and the area is more likely to be only moderately sensitive. Much of the area, however is covered by red sands and soils. Since there is a small chance that dolomites with stromatolites might occur below the soil and sands, and might be disturbed once excavations commence, a Fossil Chance Find Protocol should be added to the EMPr. Based on this information it is recommended that no further palaeontological impact assessment is required unless fossils are found by the contractor, environmental officer or other designated responsible person once excavations. Since the impact will be low, as far as the palaeontology is concerned, the project should be authorised.

Table of Contents

Expertise of Specialist	1
Declaration of Independence	1
1. Background	4
2. Methods and Terms of Reference	7
3. Geology and Palaeontology	7
i. Project location and geological context	7
ii. Palaeontological context	10
4. Impact assessment	11
5. Assumptions and uncertainties	13
6. Recommendation	13
7. References	14
8. Chance Find Protocol	15
9. Appendix A – Examples of fossils	16
10. Appendix B – Details of specialist	16
Figure 1: Google Earth map of the general area to show the relative land marks.	6
Figure 2: Google Earth Map of the proposed development	6
Figure 3: Geological map of the area around the project site.....	7
Figure 4: SAHRIS palaeosensitivity map for the site	11

1. Background

The Palingpan Manganese Mine requires the construction of a ±5km new gravel road that starts at the R325 (14.6 km from Postmasburg) and ends at the railway line (Figures 1-2).

A Palaeontological Impact Assessment was requested for the Palingpan Haul road project. To comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development and is reported herein.

Table 1: National Environmental Management Act, 1998 (Act No. 107 of 1998) (NEMA) and Environmental Impact Assessment (EIA) Regulations, 2014 (as amended) - Requirements for Specialist Reports (Appendix 6).

	A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:	Relevant section in report
ai	Details of the specialist who prepared the report,	Appendix B
a ii	The expertise of that person to compile a specialist report including a curriculum vitae	Appendix B
b	A declaration that the person is independent in a form as may be specified by the competent authority	Page 1
c	An indication of the scope of, and the purpose for which, the report was prepared	Section 1
ci	An indication of the quality and age of the base data used for the specialist report: SAHRIS palaeosensitivity map accessed – date of this report	Yes
cii	A description of existing impacts on the site, cumulative impacts of the proposed development and levels of acceptable change	Section 5
d	The date and season of the site investigation and the relevance of the season to the outcome of the assessment	N/A
e	A description of the methodology adopted in preparing the report or carrying out the specialised process	Section 2
f	The specific identified sensitivity of the site related to the activity and its associated structures and infrastructure	Section 4
g	An identification of any areas to be avoided, including buffers	N/A
h	A map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	N/A
i	A description of any assumptions made and any uncertainties or gaps in knowledge;	Section 5

	A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:	Relevant section in report
j	A description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives, on the environment	Section 4
k	Any mitigation measures for inclusion in the EMPr	Section 8, Appendix A
l	Any conditions for inclusion in the environmental authorisation	N/A
m	Any monitoring requirements for inclusion in the EMPr or environmental authorisation	Section 8, Appendix A
ni	A reasoned opinion as to whether the proposed activity or portions thereof should be authorised	Section 6
nii	If the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan	Sections 6, 8
o	A description of any consultation process that was undertaken during the course of carrying out the study	N/A
p	A summary and copies of any comments that were received during any consultation process	N/A
q	Any other information requested by the competent authority.	N/A
2	Where a government notice gazetted by the Minister provides for any protocol or minimum information requirement to be applied to a specialist report, the requirements as indicated in such notice will apply.	N/A



Figure 1: Google Earth map of the general area to show the relative landmarks. The Palingpan Haul Road project is shown by the white line.



Figure 2: Google Earth Map of the proposed development of a new haul road for Palingpan Manganese Mine with the proposed road route including stormwater infrastructure shown by the white line (option 1) and red line (option 2). Note - only the north western routes differ slightly. Map supplied by Lokisa.

2. Methods and Terms of Reference

The Terms of Reference (ToR) for this study were to undertake a PIA and provide feasible management measures to comply with the requirements of SAHRA.

The methods employed to address the ToR included:

1. Consultation of geological maps, literature, palaeontological databases, published and unpublished records to determine the likelihood of fossils occurring in the affected areas. Sources include records housed at the Evolutionary Studies Institute at the University of the Witwatersrand and SAHRA databases;
2. Where necessary, site visits by a qualified palaeontologist to locate any fossils and assess their importance (*not applicable to this assessment*);
3. Where appropriate, collection of unique or rare fossils with the necessary permits for storage and curation at an appropriate facility (*not applicable to this assessment*); and
4. Determination of fossils' representivity or scientific importance to decide if the fossils can be destroyed or a representative sample collected (*not applicable to this assessment*).

3. Geology and Palaeontology

i. Project location and geological context




Figure 3: Geological map of the area around the Palingpan Mine and haul road (yellow line). Abbreviations of the rock types are explained in Table 2. Map enlarged from the Geological Survey 1: 250 000 map 2822 Postmasburg.

Table 2: Explanation of symbols for the geological map and approximate ages (Eriksson et al., 2006; Johnson et al., 2006; Schier et al., 2018; Zeh et al., 2020). SG = Supergroup; Fm = Formation; Ma = million years; grey shading = formations impacted by the project.

Symbol	Group/Formation	Lithology	Approximate Age
Qs	Quaternary	Alluvium, sand, calcrete	Quaternary, ca 1.0 Ma to present
Vo	Ongeluk Fm, Postmasburg Group, Transvaal SG	Basaltic, andesitic lavas	Palaeoproterozoic Ca 2436 Ma
Vm	Makganyene Fm, Postmasburg Group, Transvaal SG	diamictite	Palaeoproterozoic Ca 2436 Ma
Vg	Gamagara Fm, Postmasburg Group, Transvaal SG	Shale, flagstone, quartzite, conglomeration	Palaeoproterozoic Ca 2436 Ma
Vad	Griquatown/Danielskuil Fm; Asbestos Hills Subgroup, Ghaap Group, Transvaal SG	Brown jaspilite, crocidolite, alternating shale and mudstones lenses	Palaeoproterozoic Ca 2489 Ma
Vak	Kuruman Fm, Asbestos Hills Subgroup, Ghaap Group, Transvaal SG	Banded ironstone, bands of amphibolite, lenses of conglomerate	Palaeoproterozoic Ca 2460 Ma
Vgl	Lime Acres Fm, Campbell Rand Subgroup, Ghaap Group, Transvaal SG	Dolomitic limestone and subordinate crystalline limestone; chert	Palaeoproterozoic Ca 2459 Ma

The project lies in the Griqualand West Basin of the Transvaal Supergroup.

The Late Archaean to early Proterozoic Transvaal Supergroup is preserved in three structural basins on the Kaapvaal Craton (Eriksson et al., 2006). In South Africa are the Transvaal and Griqualand West Basins, and the Kanye Basin is in southern Botswana. The Griqualand West Basin is divided into the Ghaap Plateau sub-basin and the Prieska sub-basin. Sediments in the lower parts of the basins are very similar but they differ somewhat higher up the sequences. Several tectonic events have greatly deformed the south western portion of the Griqualand West Basin between the two sub-basins

The Transvaal Supergroup comprises one of world's earliest carbonate platform successions (Beukes, 1987; Eriksson et al., 2006; Zeh et al., 2020). In some areas there are well preserved stromatolites that are evidence of the photosynthetic activity of blue green bacteria and green algae. These microbes formed colonies in warm, shallow seas.

The Transvaal Supergroup rocks in the Griqualand West Basin can be correlated with the rocks in the Transvaal Basin, closely according to Beukes and colleagues, or not so closely according to Moore and colleagues. Nonetheless, these rocks represent on a very large scale, a sequence of sediments filling the basins under conditions of lacustrine, fluvial, volcanic and glacial cycles in a tectonically active region. The predominantly carbonaceous sediments are evidence of the increase in the atmosphere of oxygen produced by algal colony photosynthesis, the so-called Great Oxygen Event (ca 2.40 – 2.32

Ga) and precursor to an environment where diverse life forms could evolve. The Neoproterozoic Transvaal Supergroup in South Africa contains the well-preserved stromatolitic Campbellrand-Malmani carbonate platform (Griqualand West Basin – Transvaal Basin respectively), which was deposited in shallow seawater shortly before the Great Oxidation Event (GOE).

Campbell Rand Subgroup, Ghaap Group

Based on the earlier works of Button (1976) and Beukes (1980, 1987) and updated after the application of sequence stratigraphy, Sumner and Beukes (2004) have described the 2650-2500 Ma Campbellrand and Malmani subgroups as being correlative and they form an extensive carbonate platform. Today there are preserved outcrops that cover 190,000 km² but these carbonate platforms probably originally covered the entire Kaapvaal Craton, >600,000 km² (ibid). The thick platform has predominantly peritidal facies in the north and east and deeper facies to the south and west. However, thinner basal sediments and platform slopes are preserved near Prieska. The ca 2.5 - 2.46 Ga Kuruman iron-formations conformably overlie the Campbellrand Subgroup and the Penge Formation overlies the Campbell Rand Subgroup. Both the Kuruman Iron Formation and the lower Penge Iron Formation consist of deep water, microbanded iron formation that formed on a stable marine shelf below wave base and then shallowed to sea level (Sumner and Beukes, 2006).

In other areas the **Campbell Rand Subgroup** has been divided into seven formations based on the different environmental settings that produces stromatolite, microbial mats, laminates, chert and carbonate platform. In this area there is only the uppermost member, the limestone-rich Lime Acres Member (old name used on the map) that contains economically important limestone, and completes this formation,

The **Asbestos Hills Subgroup** has three formations, the lower Kliphuis Formation, the Kuruman Formation and the Danielskuil Formation. They are all banded iron formations (BIF) and have vast economically important reserves and manganese deposits are closely associated with the BIF (Beukes et al., 2016).

Above the Asbestos Hills Subgroup is the **Postmasburg Group**. The Makganyene Formation has diamictites and shales from glacial conditions. Disconformably overlying these are the Ongeluk Formation basaltic andesitic lavas. According to Cornell et al. (1996) and Schroder et al. (2016) the Ongeluk Formation is equivalent to the lavas of the Hekpoort Formation in the Transvaal Basin.

QUATERNARY

There were two large basins dominating southern Africa during the Cenozoic, with the Kalahari Basin to the west and the Bushveld basin to the east. Both basins are bounded along their southern extent by the more or less west-east trending Griqualand-Transvaal Axis (Partridge et al., 2006). These sediments are not easy to date but recent attempts are gradually filling in the history of the sands, sand dunes and inter-dunes (Botha, 2021).

Quaternary Kalahari sands cover large parts of the rocks in this region, especially to the west. This is the largest and most extensive palaeo-erg in the world (Partridge et al., 2006) and is composed of extensive aeolian and fluvial sands, sand dunes, calcrete, scree

and colluvium. Periods of aridity have overprinted the sands, and calcrete and silcrete are common. Most geological maps indicate these sands simply descriptively (aeolian sand, gravelly sand, calcrete) or they are lumped together as the Gordonia Formation because the detailed regional lithostratigraphic work has not been done. Nonetheless, these sands have eroded from the interior and have been transported by wind or water to fill the basin. Reworking of the sands or stabilisation by vegetation has occurred. Probable ages of dune formation are around 100 kya (thousand years), 60 kya, 27-23 kya and 17-10 kya (in Botha, 2021).

ii. Palaeontological context

The palaeontological sensitivity of the area under consideration is presented in Figure 4. The site for development is in the Lime Acres Formation (Ghaap subgroup). The dolomites might preserve trace fossils such as stromatolites and then the formation is divided into various formations.

Stromatolites are the trace fossils that were formed by colonies of green algae and blue-green algae (Cyanobacteria) that grew in warm, shallow marine settings. These algae were responsible for releasing oxygen via the photosynthetic process where atmospheric carbon dioxide and water, using energy from the sun, are converted into carbon chains and compounds that are the building blocks of all living organisms. The released carbon dioxide initially was taken up by the abundant reducing minerals to form oxides, e.g. iron oxide. Eventually free oxygen was released into the atmosphere and some was converted into ozone by the bombardment of cosmic rays. The ozone is critical for the filtering out of harmful ultraviolet rays.

Stromatolites are the layers upon layers of inorganic materials that were deposited during photosynthesis, namely calcium carbonate, magnesium carbonate, calcium sulphate and magnesium sulphate. These layers can be in the form of flat layers, domes or columns depending on the environment where they grew (Beukes, 1987). Some environments did not form stromatolites, just layers of limestone that later were converted to dolomite. The algae that formed the stromatolites are very rarely preserved, and they are microscopic so they can only be seen from thin sections studies under a petrographic microscope.

KALAHARI GROUP

Aeolian sands and alluvium are fairly mobile and very porous so they not provide suitable conditions for preservation of organic matter (Cowan, 1995). Only in places where the sands have been waterlogged, such as palaeo-pans or palaeo-springs, is there any chance of fossilisation. For example, roots can be encased in calcium-rich or silica-rich sands and crusts, known as rhizoliths or rhizocretions, can form around the roots, invertebrates or bones around the margin of a pond, pan or spring (Klappa, 1980; Cramer and Hawkins, 2009; Peters et al., 2022).




Figure 4: SAHRIS palaeosensitivity map for the site for the proposed new haul road for Palingpan Manganese Mine shown with the yellow lines. Background colours indicate the following degrees of sensitivity: red = very highly sensitive; orange/yellow = high; green = moderate; blue = low; grey = insignificant/zero.

From the SAHRIS map above the area is indicated as very highly sensitive (red) so SAHRA would require a site visit based on the Palaeotechnical Report for the Northern Cape Province (Almond and Pether, 2009). The two palaeontologists, however coded the Ghaap Group as only moderately sensitive (green; page 41, 2009) based on the presence of stromatolites in some areas. I have carried out site visits on the adjacent properties to the north and south of Palingpan (Bamford, 2020 – Gloucester-Glosam; Bamford, 2021 McCarthy-Salene; respectively) and found only a few scattered pieces of dolomite with small stromatolites. I therefore support the recommendation of moderate sensitivity by Almond and Pether (2009) rather than the very highly sensitive coding provided in the SAHRIS palaeosensitivity map (Figure 4).

4. Impact assessment

An assessment of the potential impacts to possible palaeontological resources considers the criteria encapsulated in Table 3:

Table 3a: Criteria for assessing impacts

PART A: DEFINITION AND CRITERIA		
Criteria for ranking of the SEVERITY/NATURE of environmental impacts	H	Substantial deterioration (death, illness or injury). Recommended level will often be violated. Vigorous community action.
	M	Moderate/ measurable deterioration (discomfort). Recommended level will occasionally be violated. Widespread complaints.
	L	Minor deterioration (nuisance or minor deterioration). Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	L+	Minor improvement. Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	M+	Moderate improvement. Will be within or better than the recommended level. No observed reaction.
	H+	Substantial improvement. Will be within or better than the recommended level. Favourable publicity.
Criteria for ranking the DURATION of impacts	L	Quickly reversible. Less than the project life. Short term
	M	Reversible over time. Life of the project. Medium term
	H	Permanent. Beyond closure. Long term.
Criteria for ranking the SPATIAL SCALE of impacts	L	Localised - Within the site boundary.
	M	Fairly widespread – Beyond the site boundary. Local
	H	Widespread – Far beyond site boundary. Regional/ national
PROBABILITY (of exposure to impacts)	H	Definite/ Continuous
	M	Possible/ frequent
	L	Unlikely/ seldom

Table 3b: Impact Assessment

PART B: Assessment		
SEVERITY/NATURE	H	-
	M	-
	L	Soils and sand do not preserve fossils; so far there are no substantive records from the Lime Acres Fm of plant or animal fossils in this region, only rare scattered fragments so it is very unlikely that fossils occur on the site. The impact would be negligible
	L+	-
	M+	-
	H+	-
	DURATION	L
M		-
H		Where manifest, the impact will be permanent.

PART B: Assessment		
SPATIAL SCALE	L	Since the only possible fossils within the area would be trace fossils in the dolomites, the spatial scale will be localised within the site boundary.
	M	-
	H	-
PROBABILITY	H	-
	M	-
	L	It is extremely unlikely that any fossils would be found in the loose soils and sands that cover the area or in the dolomites that might occur below ground that will be disturbed. Nonetheless, a Fossil Chance Find Protocol should be added to the eventual EMPr.

Based on the nature of the project, surface activities may impact upon the fossil heritage if preserved in the development footprint. The geological structures suggest that the rocks are either much too old to contain body fossils but might contain trace fossils of ancient microbial life. Since there is a very small chance that fossils from the nearby Vryheid Formation may be disturbed a Fossil Chance Find Protocol has been added to this report. Taking account of the defined criteria, the potential impact to fossil heritage resources is very low.

5. Assumptions and uncertainties

Based on the geology of the area and the palaeontological record as we know it, it can be assumed that the formation and layout of the dolomites, sandstones, shales and sands are typical for the country and only some contain trace fossils such as stromatolites. The sands of the Quaternary period would not preserve fossils.

6. Recommendation

Based on experience and the lack of any previously recorded in situ fossils from the area, it is extremely unlikely that any fossils would be preserved in the overlying sands and soil of the Quaternary. There is a very small chance that trace fossils such as stromatolites may occur below ground in dolomites of the Lime Acres Formation so a Fossil Chance Find Protocol should be added to the EMPr. If fossils are found by the contractor, environmental officer, or other responsible person once excavations for the road and stormwater drainage have commenced then they should be rescued and a palaeontologist called to assess and collect a representative sample. The impact on the palaeontological heritage would be low, contrary to what is indicated on the SAHRIS palaeosensitivity map, so as far as the palaeontological heritage is concerned, the project should be authorised.

7. References

Almond, J.E., Pether, J. 2009. Palaeontological Heritage of the Northern Cape; Palaeotechnical Report for SAHRA. 115pp.

Bamford, M.K., 2020, November. Palaeontological Impact Assessment for the proposed Mining Rights application on Gloucester 674, Glosam Manganese, north of Postmasburg, Northern Cape Province. Unpublished report for AHSA.

Bamford, M.K., 2021. February. Palaeontological Impact Assessment for the proposed Salene Manganese MR on McCarthy 559, Kathu, Northern Cape Province. Unpublished report for Prescali.

Beukes, N.J., 1987. Facies relations, depositional environments and diagenesis in a major early Proterozoic stromatolitic carbonate platform to basinal sequence, Campbellrand Subgroup, Transvaal Supergroup, southern Africa. *Sedimentary Geology* 54, 1-46.

Beukes, N.J., Swindell, E.W.P., Wabo, H., 2016. Manganese Deposits of Africa. *Episodes*, 39(3), 1-33. DOI: 10.18814/epiiugs/2016/v39i2/95779

Button, A., 1976. Iron formation as an end member in carbonate sedimentary cycles in the Transvaal Supergroup, South Africa. *Economic Geology* 71, 193-201.

Cramer, M.D., Hawkins, H.-J., 2009. A physiological mechanism for the formation of root casts. *Palaeogeography, Palaeoclimatology, Palaeoecology* 274, 125-133.

Cowan, R., 1995. *History of Life*. 2nd Edition. Blackwell Scientific Publications, Boston. 462pp.

Eriksson, P.G., Altermann, W., Hartzler, F.J., 2006. The Transvaal Supergroup and its precursors. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. pp 237-260.

Johnson, M.R., van Vuuren, C.J., Visser, J.N.J., Cole, D.I., Wickens, H.deV., Christie, A.D.M., Roberts, D.L., Brandl, G., 2006. Sedimentary rocks of the Karoo Supergroup. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. Pp 461 – 499.

Klappa, C.F., 1980. Rhizoliths in terrestrial carbonates: classification, recognition, genesis and significance. *Sedimentology* 27, 613-629.

Peters, C.R., Bamford, M.K., Shields, J.P., 2022. Ch 33. Lower Bed II Olduvai Basin, Tanzania: Wetland Sedge Taphonomy, Seasonal Pasture, and Implications for Hominin Scavenging. In Reynolds, S.C., Bobe, R., (Eds). *African Paleoeology and Human Evolution*, Cambridge University Press & Assessment. 413-434.

Plumstead, E.P., 1969. Three thousand million years of plant life in Africa. Geological Society of southern Africa, Annexure to Volume LXXII. 72pp + 25 plates.

Schier, K., Bau, M., Münker, C., Beukes, N., Viehmann, S., 2018. Trace element and Nd isotope composition of shallow seawater prior to the Great Oxidation Event: Evidence from stromatolitic bioherms in the Paleoproterozoic Rooinekke and Nelani Formations, South Africa. *Precambrian Research* 315, 92-102.

Sumner, D.Y., Beukes, N.J., 2006. Sequence stratigraphic development of the Neoproterozoic Transvaal carbonate platform, Kaapvaal Craton, South Africa. *South African Journal of Geology* 109, 11–22.

Zeh, A., Wilson, A.H., Gerdes, A., 2020. Zircon U-Pb-Hf isotope systematics of Transvaal Supergroup – Constraints for the geodynamic evolution of the Kaapvaal Craton and its hinterland between 2.65 and 2.06 Ga. *Precambrian Research* 345, 105760.
<https://doi.org/10.1016/j.precamres.2020.105760>

8. Chance Find Protocol

Monitoring Programme for Palaeontology – to commence once the excavations and construction activities begin.

1. The following procedure is only required if fossils are seen on the surface and when excavations commence.
2. When excavations begin the rocks and discard must be given a cursory inspection by the environmental officer or designated person. Any fossiliferous material (trace fossils, plants, insects, bone or coal) should be put aside in a suitably protected place. This way the project activities will not be interrupted.
3. Photographs of similar fossils must be provided to the developer to assist in recognizing the trace fossils such as stromatolites in the dolomites or the Quaternary bones, rhizoliths, traces (for example see Figure 5). This information will be built into the EMP's training and awareness plan and procedures.
4. Photographs of the putative fossils can be sent to the palaeontologist for a preliminary assessment.
5. If there is any possible fossil material found by the contractor, developer or environmental officer then the qualified palaeontologist sub-contracted for this project, should visit the site to inspect the selected material and check the dumps where feasible.
6. Fossil plants or vertebrates that are considered to be of good quality or scientific interest by the palaeontologist must be removed, catalogued and housed in a suitable institution where they can be made available for further study. Before the fossils are removed from the site a SAHRA permit must be obtained. Annual reports must be submitted to SAHRA as required by the relevant permits.
7. If no good fossil material is recovered then no site inspections by the palaeontologist will be necessary. A final report by the palaeontologist must

be sent to SAHRA once the project has been completed and only if there are fossils.

8. If no fossils are found and the excavations have finished then no further monitoring is required.

9. Appendix A – Examples of fossils from the Malmani Subgroup




Figure 5: Photographs of various types of stromatolites in dolomite from the Malmani Subgroup which is the same age as the Ghaap Plateau Subgroup.

10. Appendix B – Details of specialist

Curriculum vitae (short) - Marion Bamford PhD January 2023

Present employment: Professor; Director of the Evolutionary Studies Institute.
Member Management Committee of the NRF/DSI Centre of Excellence Palaeosciences, University of the Witwatersrand,

Johannesburg, South Africa
 Telephone : +27 11 717 6690
 Cell : 082 555 6937
 E-mail : marion.bamford@wits.ac.za ;
marionbamford12@gmail.com

ii) Academic qualifications

Tertiary Education: All at the University of the Witwatersrand:
 1980-1982: BSc, majors in Botany and Microbiology. Graduated April 1983.
 1983: BSc Honours, Botany and Palaeobotany. Graduated April 1984.
 1984-1986: MSc in Palaeobotany. Graduated with Distinction, November 1986.
 1986-1989: PhD in Palaeobotany. Graduated in June 1990.

iii) Professional qualifications

Wood Anatomy Training (overseas as nothing was available in South Africa):
 1994 - Service d'Anatomie des Bois, Musée Royal de l'Afrique Centrale, Tervuren, Belgium, by Roger Dechamps
 1997 - Université Pierre et Marie Curie, Paris, France, by Dr Jean-Claude Koeniguer
 1997 - Université Claude Bernard, Lyon, France by Prof Georges Barale, Dr Jean-Pierre Gros, and Dr Marc Philippe

iv) Membership of professional bodies/associations

Palaeontological Society of Southern Africa
 Royal Society of Southern Africa - Fellow: 2006 onwards
 Academy of Sciences of South Africa - Member: Oct 2014 onwards
 International Association of Wood Anatomists - First enrolled: January 1991
 International Organization of Palaeobotany – 1993+
 Botanical Society of South Africa
 South African Committee on Stratigraphy – Biostratigraphy - 1997 - 2016
 SASQUA (South African Society for Quaternary Research) – 1997+
 PAGES - 2008 –onwards: South African representative
 ROCEEH / WAVE – 2008+
 INQUA – PALCOMM – 2011+onwards

v) Supervision of Higher Degrees

All at Wits University

Degree	Graduated/completed	Current
Honours	13	0
Masters	13	3
PhD	13	7
Postdoctoral fellows	14	4

vi) Undergraduate teaching

Geology II – Palaeobotany GEOL2008 – average 65 students per year
 Biology III – Palaeobotany APES3029 – average 25 students per year
 Honours – Evolution of Terrestrial Ecosystems; African Plio-Pleistocene Palaeoecology;
 Micropalaeontology – average 12 - 20 students per year.

vii) **Editing and reviewing**

Editor: *Palaeontologia africana*: 2003 to 2013; 2014 – Assistant editor

Guest Editor: *Quaternary International*: 2005 volume

Member of Board of Review: *Review of Palaeobotany and Palynology*: 2010 –

Associate Editor: *Cretaceous Research*: 2018-2020

Associate Editor: *Royal Society Open*: 2021 -

Review of manuscripts for ISI-listed journals: 30 local and international journals

viii) **Palaeontological Impact Assessments**

25 years' experience in PIA site and desktop projects

- Selected from recent projects only – list not complete:
- Skeerpoort Farm Mast 2020 for HCAC
- Vulindlela Eco village 2020 for 1World
- KwaZamakhule Township 2020 for Kudzala
- Sunset Copper 2020 for Digby Wells
- McCarthy-Salene 2020 for Prescali
- VLNR Lodge 2020 for HCAC
- Madadeni mixed use 2020 for Enviropro
- Frankfort-Windfield Eskom Powerline 2020 for 1World
- Beaufort West PV Facility 2021 for ACO Associates
- Copper Sunset MR 2021 for Digby Wells
- Sannaspos PV facility 2021 for CTS Heritage
- Smithfield-Rouxville-Zastron PL 2021 for TheroServe
- Glosam Mine 2022 for AHSA
- Wolf-Skilpad-Grassridge OHPL 2022 for Zutari
- Iziduli and Msenge WEFs 2022 for CTS Heritage
- Hendrina North and South WEFs & SEFs 2022 for Cabanga
- Dealesville-Springhaas SEFs 2022 for GIBB Environmental
- Vhuvhili and Mukondelei SEFs 2022 for CSIR
- Chemwes & Stilfontein SEFs 2022 for CTS Heritage
- Equestria Exts housing 2022 for Beyond Heritage
- Zeerust Salene boreholes 2022 for Prescali
- Tsakane Sewer upgrade 2022 for Tsimba
- Transnet MPP inland and coastal 2022 for ENVASS
- Ruighoek PRA 2022 for SLR Consulting (Africa)
- Namli MRA Steinkopf 2022 for Beyond Heritage

ix) **Research Output**

Publications by M K Bamford up to January 2022 peer-reviewed journals or scholarly books: over 170 articles published; 5 submitted/in press; 14 book chapters.

Scopus h-index = 30; Google Scholar h-index = 39; i10-index = 116 based on 6568 citations.

Conferences: numerous presentations at local and international conferences.