

**Palaeontological Impact Assessment for the  
proposed Palm Ridge X43 township  
development on Farm Rietspruit 152-IR,  
Midvaal, Gauteng Province**

**Desktop Study (Phase 1)**

**For**

**Beyond Heritage**

**18 November 2022**

**Prof Marion Bamford**

Palaeobotanist

P Bag 652, WITS 2050

Johannesburg, South Africa

[Marion.bamford@wits.ac.za](mailto:Marion.bamford@wits.ac.za)

## **Expertise of Specialist**

The Palaeontologist Consultant: Prof Marion Bamford  
Qualifications: PhD (Wits Univ, 1990); FRSSAf, mASSAf  
Experience: 33 years research and lecturing in Palaeontology  
25 years PIA studies and over 300 projects completed

## **Declaration of Independence**

This report has been compiled by Professor Marion Bamford, of the University of the Witwatersrand, sub-contracted by Beyond Heritage, Modimolle, South Africa. The views expressed in this report are entirely those of the author and no other interest was displayed during the decision making process for the Project.

Specialist: Prof Marion Bamford

A handwritten signature in blue ink, appearing to read 'MKBamford', written over a horizontal line.

Signature:

## **Executive Summary**

A Palaeontological Impact Assessment was requested for the proposed development of a residential township and amenities on Portion 34 of Farm Rietspruit 152-JR, Midvaal, alongside the K91 road, Ekurhuleni Metropolitan Municipality, Gauteng Province. The proposed site of approximately 8.58 hectares has a current zoning of "Agriculture" and will be known as Palm Ridge X43.

To comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development.

The proposed site lies on the potentially fossiliferous Black Reef Formation, basal Transvaal Supergroup that might preserve trace fossils but this is doubtful. Nonetheless, a Fossil Chance Find Protocol should be added to the EMP. Based on this information it is recommended that no further palaeontological impact assessment is required unless fossils are found by the contractor, environmental officer or other designated responsible person once excavations or drilling activities have commenced. Since the impact will be low, as far as the palaeontology is concerned, the project should be authorised.

# Table of Contents

Expertise of Specialist .....	1
Declaration of Independence .....	1
1. Background .....	4
2. Methods and Terms of Reference.....	7
3. Geology and Palaeontology.....	7
i. Project location and geological context .....	7
ii. Palaeontological context.....	8
4. Impact assessment.....	10
5. Assumptions and uncertainties.....	12
6. Recommendation.....	12
7. References .....	12
8. Chance Find Protocol .....	13
9. Appendix A – Examples of fossils .....	14
10. Appendix B – Details of specialist.....	15
Figure 1: Google Earth map of the general area to show the relative land marks. ....	6
Figure 2: Google Earth Map of the proposed development .....	6
Figure 3: Geological map of the area around the project site.....	7
Figure 4: SAHRIS palaeosensitivity map for the site .....	9

# 1. Background


Cosmopolitan Projects Johannesburg (Pty) Ltd proposes to establish a Residential Township, to be known as Palm Ridge X43, on Portion 34 of the Farm Rietspruit 152-IR. The proposed site of approximately 8.58 hectares has a current zoning of “Agriculture”. With an area of 3.03 hectares, the site consists of the proclaimed road K91 and divides the site in two separate areas. The proposed site is located 1 km from Sky City Mall (Figures 1-2). Midvaal is to the southeast. This project is within the jurisdiction of the Ekurhuleni Metropolitan Municipality, southern Gauteng Province.

A Palaeontological Impact Assessment was requested for the Palm Ridge X43 township project. To comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development and is reported herein.


Table 1: National Environmental Management Act, 1998 (Act No. 107 of 1998) (NEMA) and Environmental Impact Assessment (EIA) Regulations, 2014 (as amended) - Requirements for Specialist Reports (Appendix 6).

	<b>A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:</b>	<b>Relevant section in report</b>
ai	Details of the specialist who prepared the report,	Appendix B
aii	The expertise of that person to compile a specialist report including a curriculum vitae	Appendix B
b	A declaration that the person is independent in a form as may be specified by the competent authority	Page 1
c	An indication of the scope of, and the purpose for which, the report was prepared	Section 1
ci	An indication of the quality and age of the base data used for the specialist report: SAHRIS palaeosensitivity map accessed – date of this report	Yes
cii	A description of existing impacts on the site, cumulative impacts of the proposed development and levels of acceptable change	Section 5
d	The date and season of the site investigation and the relevance of the season to the outcome of the assessment	N/A
e	A description of the methodology adopted in preparing the report or carrying out the specialised process	Section 2
f	The specific identified sensitivity of the site related to the activity and its associated structures and infrastructure	Section 4
g	An identification of any areas to be avoided, including buffers	N/A

	<b>A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:</b>	<b>Relevant section in report</b>
h	A map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	N/A
i	A description of any assumptions made and any uncertainties or gaps in knowledge;	Section 5
j	A description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives, on the environment	Section 4
k	Any mitigation measures for inclusion in the EMPr	Section 8, Appendix A
l	Any conditions for inclusion in the environmental authorisation	N/A
m	Any monitoring requirements for inclusion in the EMPr or environmental authorisation	Section 8, Appendix A
ni	A reasoned opinion as to whether the proposed activity or portions thereof should be authorised	Section 6
nii	If the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan	Sections 6, 8
o	A description of any consultation process that was undertaken during the course of carrying out the study	N/A
p	A summary and copies of any comments that were received during any consultation process	N/A
q	Any other information requested by the competent authority.	N/A
2	Where a government notice gazetted by the Minister provides for any protocol or minimum information requirement to be applied to a specialist report, the requirements as indicated in such notice will apply.	N/A



**Figure 1: Google Earth map of the general area to show the relative land marks. The Palm Ridge township development on Portion 34 of Farm Rietspruit 152-IR shown by the red outline. Map from BID by iSquare.**



**Figure 2: Locality map of the proposed township development on Portion 34 of Farm Rietspruit 152-IR shown by the green rectangle. Map supplied by iSquare.**



## 2. Methods and Terms of Reference


The Terms of Reference (ToR) for this study were to undertake a PIA and provide feasible management measures to comply with the requirements of SAHRA.

The methods employed to address the ToR included:

1. Consultation of geological maps, literature, palaeontological databases, published and unpublished records to determine the likelihood of fossils occurring in the affected areas. Sources include records housed at the Evolutionary Studies Institute at the University of the Witwatersrand and SAHRA databases;
2. Where necessary, site visits by a qualified palaeontologist to locate any fossils and assess their importance (*not applicable to this assessment*);
3. Where appropriate, collection of unique or rare fossils with the necessary permits for storage and curation at an appropriate facility (*not applicable to this assessment*); and
4. Determination of fossils' representivity or scientific importance to decide if the fossils can be destroyed or a representative sample collected (*not applicable to this assessment*).

## 3. Geology and Palaeontology

### i. Project location and geological context



**Figure 3: Geological map of the area around the Palm Ridge township indicated within the yellow rectangle. Abbreviations of the rock types are explained in Table 2. Map enlarged from the Geological Survey 1: 250 000 map 2628 East Rand.**



Table 2: Explanation of symbols for the geological map and approximate ages (Eriksson et al., 2006. Zeh et al., 2020). SG = Supergroup; Fm = Formation; Ma = million years; grey shading = formations impacted by the project.

Symbol	Group/Formation	Lithology	Approximate Age
Q	Quaternary	Alluvium, sand, calcrete	Quaternary, ca 1.0 Ma to present
Vt	Timeball Hill Fm Pretoria Group, Transvaal SG	Quartzite	Palaeoproterozoic < 2420 Ma
Vbr	Black Reef Fm, Transvaal SG	Quartzite, conglomerate, shale, basalt	Ca 2650 – 2640 Ma
Vmd	Malmani Subgroup, Chuniespoort Group, Transvaal SG	Dolomite, chert	Ca 2750 – 2650 Ma
Rk	Klipriviersberg Group, Ventersdorp SG	Andesite, tuff	Ca 2714 Ma

The project lies in the central Transvaal Basin which is in the Kaapvaal Craton. The oldest rocks in the area are the predominantly volcanic rocks of the Ventersdorp Supergroup, namely the Klipriviersberg Group (Figure 3, Table 2). Overlying these are the strata of the Transvaal Supergroup. Much younger sands and alluvium have been deposited along the water courses.

Two of the basal members of the Transvaal Supergroup rocks are exposed in this area. At the base is the Black Reef Formation that is composed of diamictites, sandstone, conglomerate, shale, basalt that were deposited in a tectonically stable sag basin. Then a marine incursion began the formation of a carbonate platform (the Malmani Subgroup, Chuniespoort Group). To the west lies the Timeball Hill Formation of the Pretoria Group, Transvaal Supergroup.

The Malmani Subgroup is up to 2000m thick and has been divided into five formations based on the composition of cherts, stromatolites, limestones and shales. None of the five formations, however, has been recognised in this part of the basin.


## ii. Palaeontological context

The palaeontological sensitivity of the area under consideration is presented in Figure 4. The site for development is in the Black Reef Formation.

Since the three older gold-bearing deposits within the Witwatersrand Basin have been mined out in the upper layers and reserves are too deep for mining, the fourth gold-bearing stratum – the Black Reef Formation – is now being studied in detail (Nwaila et al., 2022). The Black Reef Formation occurs at the base of the 2.65–2.05 Ga Neoproterozoic Transvaal Supergroup (Transvaal Basin) and overlies an erosion surface developed on the older Witwatersrand and Ventersdorp strata. Three distinct Black Reef facies are found in the East Rand, namely the (1) channel facies, which may be locally absent and is variably gold-bearing, (2) blanket facies—a basal unit that is

widespread and poorly mineralized, and (3) pyritic facies—a pyrite-bearing unit that is typically well-mineralized (Nwaila et al., 2022). Such settings do not include any fossils or trace fossils in the geological literature.

According to the SAHRA Palaeotechnical Report for Gauteng (Groenewald et al., 2014) the Black Reef Formation may preserve stromatolites because it was considered as a possible equivalent of Black Reef Formation in the Northern Cape (Vryburg Formation) that contains stromatolitic carbonates. More recent work has shown that the Black Reef Formation is younger (Zeh et al., 2020).



**Figure 4: SAHRIS palaeosensitivity map for the site for the proposed Palm Ridge township shown within the yellow rectangle. Background colours indicate the following degrees of sensitivity: red = very highly sensitive; orange/yellow = high; green = moderate; blue = low; grey = insignificant/zero.**

From the SAHRIS map above the area is indicated as moderately sensitive (green) and this is for the Black Reef Formation. According to the geology this formation is predominantly arenaceous (sand, silica) and not carbonaceous (dolomite), therefore, it is unlikely to preserve stromatolites, but might have microbialites.

**Stromatolites** are the trace fossils that were formed by colonies of green algae and blue-green algae (Cyanobacteria) that grew in warm, shallow marine settings. These algae were responsible for releasing oxygen via the photosynthetic process where atmospheric carbon dioxide and water, using energy from the sun, are converted into carbon chains and compounds that are the building blocks of all living organisms. The released carbon

dioxide initially was taken up by the abundant reducing minerals to form oxides, e.g. iron oxide. Eventually free oxygen was released into the atmosphere and some was converted into ozone by the bombardment of cosmic rays. The ozone is critical for the filtering out of harmful ultraviolet rays. Stromatolites are the layers upon layers of inorganic materials that were deposited during photosynthesis, namely calcium carbonate, magnesium carbonate, calcium sulphate and magnesium sulphate. These layers can be in the form of flat layers, domes or columns depending on the environment where they grew (Beukes, 1987). Some environments did not form stromatolites, just layers of limestone that later was converted to dolomite. The algae that formed the stromatolites are very rarely preserved, and they are microscopic so they can only be seen from thin sections studies under a petrographic microscope.

**Microbialites** (sensu Burne and Moore, 1987) are organo-sedimentary deposits formed from interaction between benthic microbial communities (BMCs) and detrital or chemical sediments. In addition, microbialites contrast with other biological sediments in that they are generally not composed of skeletal remains. Archean carbonates mostly consist of stromatolites. These platforms could have been the site of early O<sub>2</sub> production on our planet. Stromatolites are the laminated, organo-sedimentary, non-skeletal products of microbial communities, which may have included cyanobacteria, the first photosynthetic organisms to produce oxygen. Another type of trace fossil has been termed Microbially-induced sedimentary structures (MISS sensu Noffke et al., 2001) or simply ‘fossil mats’ (sensu Tice et al., 2011). These include swirls, rip-ups, crinkled surfaces and wrinkles that were formed by the mucus extruded by littoral algae or microbes and bound together sand particles. Davies et al. (2016) caution against the assumption that all such structures are microbially induced unless there is additional evidence for microbes in the palaeoenvironment.

#### 4. Impact assessment

An assessment of the potential impacts to possible palaeontological resources considers the criteria encapsulated in Table 3:

**Table 3a: Criteria for assessing impacts**

<b>PART A: DEFINITION AND CRITERIA</b>		
<b>Criteria for ranking of the SEVERITY/NATURE of environmental impacts</b>	<b>H</b>	Substantial deterioration (death, illness or injury). Recommended level will often be violated. Vigorous community action.
	<b>M</b>	Moderate/ measurable deterioration (discomfort). Recommended level will occasionally be violated. Widespread complaints.
	<b>L</b>	Minor deterioration (nuisance or minor deterioration). Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	<b>L+</b>	Minor improvement. Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.

	<b>M+</b>	Moderate improvement. Will be within or better than the recommended level. No observed reaction.
	<b>H+</b>	Substantial improvement. Will be within or better than the recommended level. Favourable publicity.
<b>Criteria for ranking the DURATION of impacts</b>	<b>L</b>	Quickly reversible. Less than the project life. Short term
	<b>M</b>	Reversible over time. Life of the project. Medium term
	<b>H</b>	Permanent. Beyond closure. Long term.
<b>Criteria for ranking the SPATIAL SCALE of impacts</b>	<b>L</b>	Localised - Within the site boundary.
	<b>M</b>	Fairly widespread – Beyond the site boundary. Local
	<b>H</b>	Widespread – Far beyond site boundary. Regional/ national
<b>PROBABILITY (of exposure to impacts)</b>	<b>H</b>	Definite/ Continuous
	<b>M</b>	Possible/ frequent
	<b>L</b>	Unlikely/ seldom

**Table 3b: Impact Assessment**

<b>PART B: Assessment</b>		
<b>SEVERITY/NATURE</b>	<b>H</b>	-
	<b>M</b>	-
	<b>L</b>	Soils and sands do not preserve fossils; so far there are no records from the Black Reef Fm of trace fossils in this region so it is very unlikely that fossils occur on the site. The impact would be negligible
	<b>L+</b>	-
	<b>M+</b>	-
	<b>H+</b>	-
	<b>DURATION</b>	<b>L</b>
<b>M</b>		-
<b>H</b>		Where manifest, the impact will be permanent.
<b>SPATIAL SCALE</b>	<b>L</b>	Since the only possible fossils within the area would be trace fossils in the sandstones, the spatial scale will be localised within the site boundary.
	<b>M</b>	-
	<b>H</b>	-
<b>PROBABILITY</b>	<b>H</b>	-
	<b>M</b>	-
	<b>L</b>	It is extremely unlikely that any fossils would be found in the loose soils and sands that cover the area or in the quartzitic sandstones of the Black Reef Fm that will be excavated. Nonetheless, a Fossil Chance Find Protocol should be added to the eventual EMPr.

Based on the nature of the project, surface activities may impact upon the fossil heritage if preserved in the development footprint. The geological structures suggest that the

rocks are either much too old to contain fossils or the wrong type. Furthermore, the material to be excavated is the overlying soils and this does not preserve fossils. Since there is an extremely small chance that trace fossils from the Black Reef Formation may be disturbed a Fossil Chance Find Protocol has been added to this report. Taking account of the defined criteria, the potential impact to fossil heritage resources is extremely low.

## 5. Assumptions and uncertainties

Based on the geology of the area and the palaeontological record as we know it, it can be assumed that the formation and layout of the volcanic rocks, sandstones, shales and sands are typical for the country and only some might contain trace fossils such as stromatolites or microbialites. The overlying soils and sands of the Quaternary period would not preserve fossils. The category of moderately fossiliferous for the Black Reef Formation is based on an outdated correlation with the Vryburg Formation.

## 6. Recommendation

Based on experience and the lack of any previously recorded fossils from the area, it is extremely unlikely that any fossils would be preserved in the overlying soils and sands of the Quaternary. There is an extremely small chance that trace fossils may occur in the sandstones of the Palaeoproterozoic Black Reef Formation so a Fossil Chance Find Protocol should be added to the EMPr. If fossils are found by the contractor, environmental offic, or other responsible person once excavations for foundations and amenities have commenced then they should be rescued and a palaeontologist called to assess and collect a representative sample. The impact on the palaeontological heritage would be very low, so as far as the palaeontological heritage is concerned, the project should be authorised.

## 7. References

Beukes, N.J., 1987. Facies relations, depositional environments and diagenesis in a major early Proterozoic stromatolitic carbonate platform to basinal sequence, Campbellrand Subgroup, Transvaal Supergroup, southern Africa. *Sedimentary Geology* 54, 1-46.

Burne, R.V., Moore, L.S., 1987. Microbialites; organosedimentary deposits of benthic microbial communities. *Palaios* 2(3), 241-254.

Davies, N.S., Liu, A.G., Gibling, M.R., Miller, R.F., 2016. Resolving MISS conceptions and misconceptions: A geological approach to sedimentary surface textures generated by microbial and abiotic processes *Earth-Science Reviews* 154, 210–246.

Eriksson, P.G., Altermann, W., Hartzler, F.J., 2006. The Transvaal Supergroup and its precursors. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of*

South Africa. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. pp 237-260.

Groenewald, G., Groenewald, D., Groenewald, S., 2014. SAHRA Palaeotechnical Report. Palaeontological Heritage of Gauteng. 20 pages.

Homann. M., 2019. Earliest life on Earth: Evidence from the Barberton Greenstone Belt, South Africa. *Earth Science Reviews* 196, 102888.

Noffke, N., Gerdes, G., Klenke, T. and Krumbein, W.E. (2001). Microbially induced sedimentary structures – a new category within the classification of primary sedimentary structures. *Journal of Sedimentary Research*, A71, 649-656.

Nwaila, G.T., Manzi, M.S.D., Zhang, S.E., Bourdeau, J.E., Bam, L.C., Rose, D.H., Maselela, K., Reid, D.L., Ghorbani, Y., Durrheim, R. J., 2022. Constraints on the Geometry and Gold Distribution in the Black Reef Formation of South Africa Using 3D Reflection Seismic Data and Micro-X-ray Computed Tomography. *Natural Resources Research*, Vol. 31, No. 3, June 2022

Tice, M.M., Thornton, D.C.O., Pope, M.C., Olszewski, T.D., Gong, J., 2011. Archean Microbial Mat Communities. *Annual Review of Earth and Planetary Sciences* 39, 297–319.

Zeh, A., Wilson, A.H., Gerdes, A., 2020. Zircon U-Pb-Hf isotope systematics of Transvaal Supergroup – Constraints for the geodynamic evolution of the Kaapvaal Craton and its hinterland between 2.65 and 2.06 Ga. *Precambrian Research* 345, 105760.  
<https://doi.org/10.1016/j.precamres.2020.105760>

## 8. Chance Find Protocol

### **Monitoring Programme for Palaeontology – to commence once the excavation activities begin.**

1. The following procedure is only required if fossils are seen on the surface and when drilling/excavations commence.
2. When excavations begin the rocks and discard must be given a cursory inspection by the environmental officer or designated person. Any fossiliferous material (trace fossils such as stromatolites or microbialites) should be put aside in a suitably protected place. This way the project activities will not be interrupted.
3. Photographs of similar fossils must be provided to the developer to assist in recognizing the fossil plants, vertebrates, invertebrates or trace fossils in the shales and mudstones (for example see Figures 5-6). This information will be built into the EMP's training and awareness plan and procedures.
4. Photographs of the putative fossils can be sent to the palaeontologist for a preliminary assessment.




5. If there is any possible fossil material found by the contractor, developer or environmental officer then the qualified palaeontologist sub-contracted for this project, should visit the site to inspect the selected material and check the dumps where feasible.
6. Trace fossils, fossil plants or vertebrates that are considered to be of good quality or scientific interest by the palaeontologist must be removed, catalogued and housed in a suitable institution where they can be made available for further study. Before the fossils are removed from the site a SAHRA permit must be obtained. Annual reports must be submitted to SAHRA as required by the relevant permits.
7. If no good fossil material is recovered then no site inspections by the palaeontologist will be necessary. A final report by the palaeontologist must be sent to SAHRA once the project has been completed and only if there are fossils.
8. If no fossils are found and the excavations have finished then no further monitoring is required.

## Appendix A – Examples of fossils from the Malmani and Magaliesberg Formations



**Figure 5: Photographs of stromatolites (trace fossils of algal colonies) as seen in the field, from the Malmani Subgroup (Chuniespoort Group, Transvaal Supergroup).**



Magaliesberg Fm trace fossils, near Pretoria (all from Bosch & Eriksson, 2008): A – cracks,. B – sinuous structure, C – *Manchuriphycus*, D – circular structures. R1 coin for scale.

**Figure 6: Photographs of trace fossils from the Magaliesberg Formation (Pretoria Group, Transvaal Supergroup).**

## 9. Appendix B – Details of specialist

### **Curriculum vitae (short) - Marion Bamford PhD**

**July 2022**

#### **I) Personal details**

Surname : **Bamford**  
 First names : **Marion Kathleen**  
 Present employment: Professor; Director of the Evolutionary Studies Institute.  
 Member Management Committee of the NRF/DST Centre of  
 Excellence Palaeosciences, University of the Witwatersrand,  
 Johannesburg, South Africa  
 Telephone : +27 11 717 6690  
 Fax : +27 11 717 6694



Cell : 082 555 6937  
E-mail : [marion.bamford@wits.ac.za](mailto:marion.bamford@wits.ac.za) ;  
[marionbamford12@gmail.com](mailto:marionbamford12@gmail.com)

## ii) Academic qualifications

Tertiary Education: All at the University of the Witwatersrand:

1980-1982: BSc, majors in Botany and Microbiology. Graduated April 1983.

1983: BSc Honours, Botany and Palaeobotany. Graduated April 1984.

1984-1986: MSc in Palaeobotany. Graduated with Distinction, November 1986.

1986-1989: PhD in Palaeobotany. Graduated in June 1990.

NRF Rating: C-2 (1999-2004); B-3 (2005-2015); B-2 (2016-2020); B-1 (2021-2026)

## iii) Professional qualifications

*Wood Anatomy Training (overseas as nothing was available in South Africa):*

1994 - Service d'Anatomie des Bois, Musée Royal de l'Afrique Centrale, Tervuren, Belgium, by Roger Dechamps

1997 - Université Pierre et Marie Curie, Paris, France, by Dr Jean-Claude Koeniguer

1997 - Université Claude Bernard, Lyon, France by Prof Georges Barale, Dr Jean-Pierre Gros, and Dr Marc Philippe

## iv) Membership of professional bodies/associations

Palaeontological Society of Southern Africa

Royal Society of Southern Africa - Fellow: 2006 onwards

Academy of Sciences of South Africa - Member: Oct 2014 onwards

International Association of Wood Anatomists - First enrolled: January 1991

International Organization of Palaeobotany - 1993+

Botanical Society of South Africa

South African Committee on Stratigraphy - Biostratigraphy - 1997 - 2016

SASQUA (South African Society for Quaternary Research) - 1997+

PAGES - 2008 -onwards: South African representative

ROCEEH / WAVE - 2008+

INQUA - PALCOMM - 2011+onwards

## vii) Supervision of Higher Degrees

All at Wits University

Degree	Graduated/completed	Current
Honours	13	0
Masters	11	3
PhD	11	6
Postdoctoral fellows	15	1

## viii) Undergraduate teaching

Geology II - Palaeobotany GEOL2008 - average 65 students per year

Biology III - Palaeobotany APES3029 - average 45 students per year

Honours - Evolution of Terrestrial Ecosystems; African Plio-Pleistocene Palaeoecology;

Micropalaeontology - average 12-20 students per year.

## ix) Editing and reviewing

Editor: *Palaeontologia africana*: 2003 to 2013; 2014 – Assistant editor  
Guest Editor: *Quaternary International*: 2005 volume  
Member of Board of Review: *Review of Palaeobotany and Palynology*: 2010 –  
Associate Editor *Open Science UK*: 2021 -  
Review of manuscripts for ISI-listed journals: 30 local and international journals  
Reviewing of funding applications for NRF, PAST, NWO, SIDA, National Geographic,  
Leakey Foundation

## **x) Palaeontological Impact Assessments**

Selected from the past five years only – list not complete:

- Mala Mala 2017 for Henwood
- Modimolle 2017 for Green Vision
- Klipportjie and Finaalspan 2017 for Delta BEC
- Ledjadja borrow pits 2018 for Digby Wells
- Lungile poultry farm 2018 for CTS
- Olienhout Dam 2018 for JP Celliers
- Isondlo and Kwasobabili 2018 for GCS
- Kanakies Gypsum 2018 for Cabanga
- Nababeep Copper mine 2018
- Glencore-Mbali pipeline 2018 for Digby Wells
- Remhoogte PR 2019 for A&HAS
- Bospoort Agriculture 2019 for Kudzala
- Overlooked Quarry 2019 for Cabanga
- Richards Bay Powerline 2019 for NGT
- Eilandia dam 2019 for ACO
- Eastlands Residential 2019 for HCAC
- Fairview MR 2019 for Cabanga
- Graspan project 2019 for HCAC
- Lieliefontein N&D 2019 for EnviroPro
- Skeerpoort Farm Mast 2020 for HCAC
- Vulindlela Eco village 2020 for 1World
- KwaZamakhule Township 2020 for Kudzala
- Sunset Copper 2020 for Digby Wells
- McCarthy-Salene 2020 for Prescali
- VLNR Lodge 2020 for HCAC
- Madadeni mixed use 2020 for EnviroPro
- Frankfort-Windfield Eskom Powerline 2020 for 1World
- Beaufort West PV Facility 2021 for ACO Associates
- Copper Sunset MR 2021 for Digby Wells
- Sannaspos PV facility 2021 for CTS Heritage
- Smithfield-Rouxville-Zastron PL 2021 for TheroServe

## **xi) Research Output**

Publications by M K Bamford up to July 2022 peer-reviewed journals or scholarly books: over 165 articles published; 5 submitted/in press; 10 book chapters.  
Scopus h-index = 30; Google scholar h-index = 35; i10-index = 92

Conferences: numerous presentations at local and international conferences.