Palaeontological Impact Assessment for the proposed Su Casa Burial Estate, Witbank area, Mpumalanga Province

Desktop Study (Phase 1)

For

Eaglesage (Pty) Ltd & Amber Earth (Pty) Ltd

20 February 2022

Prof Marion Bamford

Palaeobotanist P Bag 652, WITS 2050 Johannesburg, South Africa Marion.bamford@wits.ac.za

Expertise of Specialist

The Palaeontologist Consultant: Prof Marion Bamford Qualifications: PhD (Wits Univ, 1990); FRSSAf, mASSAf

Experience: 33 years research and lecturing in Palaeontology

25 years PIA studies and over 300 projects completed

Declaration of Independence

This report has been compiled by Professor Marion Bamford, of the University of the Witwatersrand, sub-contracted by Amber Earth (Pty) Ltd who were appointed by Eaglesage (Pty) Ltd, Pretoria, South Africa. The views expressed in this report are entirely those of the author and no other interest was displayed during the decision making process for the Project.

Specialist: Prof Marion Bamford

MKBamfark

Executive Summary

A Palaeontological Impact Assessment was requested for the proposed private cemetery on Portion 10 of Farm Doornrug 302 JS, west of Witbank (Emalahleni), Mpumalanga Province.

To comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development.

The proposed site lies on the moderately fossiliferous rocks of the Dwyka Group (Karoo Supergroup) that could preserve plants of the early Glossopteris flora. Fossils, however have only been preserved from the mudstone facies, not the tillites and shales that occur in this region. Nonetheless, a Fossil Chance Find Protocol should be added to the EMPr. Based on this information it is recommended that no further palaeontological impact assessment is required unless fossils are found by the developer/ environmental officer/ other designated responsible person once excavations for infrastructure and graves have commenced. As far as the palaeontology is concerned, the project should be authorised.

Table of Contents

Ŀ	Expertise of Specialist 1
Γ	Declaration of Independence1
1.	Background4
2.	Methods and Terms of Reference7
3.	Geology and Palaeontology8
i.	Project location and geological context8
ii.	Palaeontological context9
4.	Impact assessment10
5.	Assumptions and uncertainties13
6.	Recommendation13
7.	References
8.	Chance Find Protocol14
9.	Appendix A – Examples of fossils16
10.	Appendix B – Details of specialist16
Figur	re 1: Google Earth map of the general area to show the relative land marks6
Figur	re 2: Google Earth Map of the proposed development6
Figur	re 3: Geological map of the area around the project site
Figur	re 4: SAHRIS palaeosensitivity map for the site

1. Background

There is a proposal to develop a private cemetery on a piece of vacant land on Portion 10 of Farm Doornrug 302 JS, south of the N4 and R104 between Balmoral and Witbank, and west of the south bound minor road (Figures 1, 2). The proposed activities entail but are not limited to the Su Casa Burial site, chapel, dining hall, ablution facilities, admin offices, cross landmark, fencing, landscaping, ash scattering garden, upgrade of the existing borehole, establishment of a new borehole, establishment of two ponds, wall of remembrance and upgrade of the existing road. The excavations of graves will be to a depth of about 2 m as and when required.

A Palaeontological Impact Assessment was requested for the Cemetery project. To comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development and is reported herein.

Table 1: National Environmental Management Act, 1998 (Act No. 107 of 1998) (NEMA) and Environmental Impact Assessment (EIA) Regulations, 2014 (as amended) - Requirements for Specialist Reports (Appendix 6).

	A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:	Relevant section in report
ai	Details of the specialist who prepared the report,	Appendix B
aii	The expertise of that person to compile a specialist report including a curriculum vitae	Appendix B
b	A declaration that the person is independent in a form as may be specified by the competent authority	Page 1
С	An indication of the scope of, and the purpose for which, the report was prepared	Section 1
ci	An indication of the quality and age of the base data used for the specialist report: SAHRIS palaeosensitivity map accessed – date of this report	Yes
cii	A description of existing impacts on the site, cumulative impacts of the proposed development and levels of acceptable change	Section 5
d	The date and season of the site investigation and the relevance of the season to the outcome of the assessment	N/A

	A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:	Relevant section in report
е	A description of the methodology adopted in preparing the report or carrying out the specialised process	Section 2
f	The specific identified sensitivity of the site related to the activity and its associated structures and infrastructure	Section 4
g	An identification of any areas to be avoided, including buffers	N/A
h	A map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	N/A
i	A description of any assumptions made and any uncertainties or gaps in knowledge;	Section 5
j	A description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives, on the environment	Section 4
k	Any mitigation measures for inclusion in the EMPr	Section 8, Appendix A
1	Any conditions for inclusion in the environmental authorisation	N/A
m	Any monitoring requirements for inclusion in the EMPr or environmental authorisation	Section 8, Appendix A
ni	A reasoned opinion as to whether the proposed activity or portions thereof should be authorised	Section 6
nii	If the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan	Sections 6, 8
0	A description of any consultation process that was undertaken during the course of carrying out the study	N/A
р	A summary and copies of any comments that were received during any consultation process	N/A
q	Any other information requested by the competent authority.	N/A
2	Where a government notice gazetted by the Minister provides for any protocol or minimum information requirement to be applied to a specialist report, the requirements as indicated in such notice will apply.	N/A



Figure 1: Google Earth map of the general area to show the relative land marks. The cemetery project site is shown by the red rectangle.



Figure 2: Google Earth Map of the proposed development of the Su Casa Burial Estate, west of Witbank with the section shown by the red outline. Map supplied by Amber Earth.

2. Methods and Terms of Reference

The Terms of Reference (ToR) for this study were to undertake a PIA and provide feasible management measures to comply with the requirements of SAHRA.

The methods employed to address the ToR included:

- 1. Consultation of geological maps, literature, palaeontological databases, published and unpublished records to determine the likelihood of fossils occurring in the affected areas. Sources included records housed at the Evolutionary Studies Institute at the University of the Witwatersrand and SAHRA databases;
- 2. Where necessary, site visits by a qualified palaeontologist to locate any fossils and assess their importance (*not applicable to this assessment*);
- 3. Where appropriate, collection of unique or rare fossils with the necessary permits for storage and curation at an appropriate facility (*not applicable to this assessment*); and
- 4. Determination of fossils' representivity or scientific importance to decide if the fossils can be destroyed or a representative sample collected (*not applicable to this assessment*).

3. Geology and Palaeontology

i. Project location and geological context




Figure 3: Geological map of the area around the proposed cemetery with the location indicated within the yellow rectangle. Abbreviations of the rock types are explained in Table 2. Map enlarged from the Geological Survey 1: 250 000 map 2528 Pretoria.

Table 2: Explanation of symbols for the geological map and approximate ages (Eriksson et al., 2006. Johnson et al., 2006). SG = Supergroup; Fm = Formation; Ma = million years; grey shading = formations impacted by the project.

Symbol	Group/Formation	Lithology	Approximate Age
Q	Quaternary	Alluvium, sand, calcrete	Neogene, ca 2.5 Ma to present
Jd	Jurassic dykes	Dolerite dykes, intrusive	Jurassic, approx. 180 Ma
Pe	Vryheid Fm, Ecca Group, Karoo SG	Shale, shaley sandstone, grit, sandstone, conglomerate, thin coal seams	Early Permian, ca 290-260 Ma
Pd	Dwyka Group, Karoo SG	Tillites, shale	Late Carboniferous to early Permian, ca 310-300 Ma
Mw	Wilgerivier Fm, Waterberg Group	Sandstone, quartzite, conglomerate	Ca 2050 – 2000 Ma
Mn	Nebo Granite, Bushveld Igneous Complex	Granite	Ca 2400 Ma
Di	Diabase	diabase	Post Transvaal SG

Symbol	Group/Formation	Lithology	Approximate Age
Vdr	Damwal Fm, Rooiberg	Volcanic rocks	Ca 2500 – 2400 Ma
	Group		
Vsi	Silverton Formation, Pretoria Group, Transvaal SG	Shale, carbonaceous in places, hornfels, chert	Ca 2500 – 2400 Ma

The project lies in the southeastern margin of the Transvaal Basin with the sediments of the Transvaal Supergroup, and the northern margin of the Karoo Basin with the lower Karoo Supergroup sediments. There are also outliers of the Waterberg Group. The project site is on shales and tillites of the Dwyka Group.

The Karoo Supergroup rocks cover a very large proportion of South Africa and extend from the northeast (east of Pretoria) to the southwest and across to almost the KwaZulu Natal south coast. It is bounded along the southern margin by the Cape Fold Belt and along the northern margin by the much older Transvaal Supergroup rocks. Representing some 120 million years (300 – 183Ma), the Karoo Supergroup rocks have preserved a diversity of fossil plants, insects, vertebrates and invertebrates.

During the Carboniferous Period South Africa was part of the huge continental landmass known as Gondwanaland and it was positioned over the South Pole. As a result, there were several ice sheets that formed and melted, and covered most of South Africa (Visser, 1986, 1989; Isbell et al., 2012). Gradual melting of the ice as the continental mass moved northwards and the earth warmed, formed fine-grained sediments in the large inland sea. These are the oldest rocks in the system and are exposed around the outer part of the ancient Karoo Basin, and are known as the Dwyka Group. They comprise tillites, diamictites, mudstones, siltstones and sandstones that were deposited as the basin filled. This group has been divided into two formations with Elandsvlei Formation occurring throughout the basin and the upper Mbizane Formation occurring only in the Free State and KwaZulu Natal (Johnson et al., 2006).

Overlying the Dwyka Group rocks are rocks of the Ecca Group that are Early Permian in age, then the Beaufort and Stormberg Groups. The whole Karoo sequence is capped by the Jurassic aged Drakensberg basalts. Associated with the latter are numerous intrusive dolerite dykes and sills that have cut through the Karoo sediments.

ii. Palaeontological context

The palaeontological sensitivity of the area under consideration is presented in Figure 4. The site for development is in the Dwyka Group that is indicated as having a moderate palaeosensitivity (green).

The Dwyka Group is made up of seven facies that were deposited in a marine basin under differing environmental settings of glacial formation and retreat (Visser, 1986, 1989; Johnson et al., 2006). In the north and east these are called the Mbizane Formation, and the Elandsvlei Formation in the south and west. Described below are the seven facies that occur in this group (Johnson et al., 2006 p. 463-465):

The <u>massive diamictite facies</u> comprises highly compacted diamictite that is clast-poor in the north. It was deposited in subaqueous or subglacial positions.

The <u>stratified diamictite</u> comprises alternating diamictite, mudrock, sandstone and conglomerate beds. They are interpreted as being rapidly deposited, sediment gravity flows but with some possible reworking of the subglacial diamictites.

The <u>massive carbonate-rich diamictite facies</u> is clast-poor and was formed by the rainout of debris, with the carbonate probably originating by crystallisation from interstitial waters.

The <u>conglomerate facies</u> ranges from single layer boulder beds to poorly sorted pebble and granule conglomerates. The boulder beds are interpreted as lodgement deposits whereas the poorly sorted conglomerates are a product of water-reworking of diamicton by high-density sediment gravity flows.

The <u>sandstone facies</u> were formed as turbidite deposits.

The <u>mudrock with stones facies</u> represents rainout deposits in the distal iceberg zone.

The <u>mudrock facies</u> consists of dark-coloured, commonly carbonaceous mudstone, shale or silty rhythmite that was formed when the mud or silt in suspension settled. This is the only fossiliferous facies of the Dwyka Group.

The Dwyka *Glossopteris* flora outcrops are very sporadic and rare. Of the seven facies that have been recognised in the Dwyka Group fossil plant fragments have only been recognised from the mudrock facies. They have been recorded from around Douglas only (Johnson et al., 2006; Anderson and McLachlan 1976) although the Dwyka Group

exposures are very extensive. Jurassic Dolerites do not contain fossils as they are igneous intrusives.




Figure 4: SAHRIS palaeosensitivity map for the site for the proposed cemeteryshown within the yellow rectangle. Background colours indicate the following degrees of sensitivity: red = very highly sensitive; orange/yellow = high; green = moderate; blue = low; grey = insignificant/zero.

4. Impact assessment

An assessment of the potential impacts to possible palaeontological resources considers the criteria encapsulated in Table 3:

Table 3a: Criteria for assessing impacts

PART A: DEFINITION AND CRITERIA			
Criteria for ranking of the SEVERITY/NATURE	Н	Substantial deterioration (death, illness or injury). Recommended level will often be violated. Vigorous community action.	
of environmental impacts	M	Moderate/ measurable deterioration (discomfort). Recommended level will occasionally be violated. Widespread complaints.	

	L	Minor deterioration (nuisance or minor deterioration). Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	L+	Minor improvement. Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	M+	Moderate improvement. Will be within or better than the recommended level. No observed reaction.
H		Substantial improvement. Will be within or better than the recommended level. Favourable publicity.
Criteria for ranking	L	Quickly reversible. Less than the project life. Short term
the DURATION of	M	Reversible over time. Life of the project. Medium term
impacts	Н	Permanent. Beyond closure. Long term.
Criteria for ranking	L	Localised - Within the site boundary.
the SPATIAL SCALE	M	Fairly widespread – Beyond the site boundary. Local
of impacts	Н	Widespread – Far beyond site boundary. Regional/ national
PROBABILITY	Н	Definite/ Continuous
(of exposure to	M	Possible/ frequent
impacts)	L	Unlikely/ seldom

Table 3b: Impact Assessment

PART B: Assessment				
	Н	-		
	M	-		
SEVERITY/NATURE	L	Soils do not preserve fossils; so far there are no records from the Dwyka Group of plant or animal fossils in this region so it is very unlikely that fossils occur on the site. The impact would be negligible		
	L+	-		
	M+	-		
	H+	-		
	L	-		
DURATION	M	-		
	Н	Where manifest, the impact will be permanent.		
SPATIAL SCALE	L	Since the only possible fossils within the area would be fossil plants in the mudstones of the Dwyka Group but only shales and tillites occur here, the spatial scale will be localised within the site boundary.		
	M	-		
	Н	-		

PART B: Assessment				
	Н	-		
	M	-		
PROBABILITY	L	It is extremely unlikely that any fossils would be found in the loose soils and sands that cover the area or in the shales and tillites that will be excavated. Nonetheless, a Fossil Chance Find Protocol should be added to the eventual EMPr.		

Based on the nature of the project, surface activities may impact upon the fossil heritage if preserved in the development footprint. The geological structures suggest that the rocks are either much too old to contain fossils or are the correct age but wrong lithology. Furthermore, the material to be excavated is soil and this does not preserve fossils. Since there is a very small chance that plant fossils from the Dwyka Group may be disturbed a Fossil Chance Find Protocol has been added to this report. Taking account of the defined criteria, the potential impact to fossil heritage resources is very low.

5. Assumptions and uncertainties

Based on the geology of the area and the palaeontological record as we know it, it can be assumed that the formation and layout of the dolorites, sandstones, shales and sands are typical for the country and most do not contain fossil plant, insect, invertebrate and vertebrate material. The sands of the Quaternary period would not preserve fossils. It is not known if fossils occur below ground but it is very unlikely.

6. Recommendation

Based on experience and the lack of any previously recorded fossils from the area, it is extremely unlikely that any fossils would be preserved in the Dwyka Group shales and tillites and not in the overlying soils and sands of the Quaternary. There is a very small chance that fossils may occur below ground in the adjacent shales and tillites of the Dwyka Group (Karoo Supergroup) so a Fossil Chance Find Protocol should be added to the EMPr. If fossils are found by the environmental officer, or other responsible person once excavations have commenced then they should be rescued and a palaeontologist called to assess and collect a representative sample. The impact on the palaeontological heritage would be low so as far as the palaeontology is concerned, the project should be authorised.

7. References

Anderson, J.M., Anderson, H.M., 1985. Palaeoflora of Southern Africa: Prodromus of South African megafloras, Devonian to Lower Cretaceous. A.A. Balkema, Rotterdam. 423 pp.

Eriksson, P.G., Altermann, W., Hartzer, F.J., 2006. The Transvaal Supergroup and its precursors. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). The Geology of South Africa. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. pp 237-260.

Isbell, J.L., Henry, L.C., Gulbranson, E.L., Limarino, C.O., Fraiser, F.L., Koch, Z.J., Ciccioli, P.l., Dineen, A.A., 2012. Glacial paradoxes during the late Paleozoic ice age: Evaluating the equilibrium line altitude as a control on glaciation. Gondwana Research 22, 1-19.

Johnson, M.R., van Vuuren, C.J., Visser, J.N.J., Cole, D.I., Wickens, H.deV., Christie, A.D.M., Roberts, D.L., Brandl, G., 2006. Sedimentary rocks of the Karoo Supergroup. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). The Geology of South Africa. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. Pp 461 – 499.

Plumstead, E.P., 1969. Three thousand million years of plant life in Africa. Geological Society of southern Africa, Annexure to Volume LXXII. 72pp + 25 plates.

Visser, J.N.J., 1986. Lateral lithofacies relationships in the glacigene Dwyka Formation in the western and central parts of the Karoo Basin. Transactions of the Geological Society of South Africa 89, 373-383.

Visser, J.N.J., 1989. The Permo-Carboniferous Dwyka Formation of southern Africa: deposition by a predominantly subpolar marine icesheet. Palaeogeography, Palaeoclimatology, Palaeoecology 70, 377-391.

8. Chance Find Protocol

Monitoring Programme for Palaeontology - to commence once the excavations begin.

- 1. The following procedure is only required if fossils are seen on the surface and when excavations commence.
- 2. When excavations begin the rocks and must be given a cursory inspection by the environmental officer or designated person. Any fossiliferous material (plants, insects, bone, coal) should be put aside in a suitably protected place. This way the project activities will not be interrupted.
- 3. Photographs of similar fossils must be provided to the developer to assist in recognizing the fossil plants, vertebrates, invertebrates or trace fossils in the

- shales and mudstones (for example see Figure 5). This information will be built into the EMP's training and awareness plan and procedures.
- 4. Photographs of the putative fossils can be sent to the palaeontologist for a preliminary assessment.
- 5. If there is any possible fossil material found by the developer/environmental officer then the qualified palaeontologist sub-contracted for this project, should visit the site to inspect the selected material and check the dumps where feasible.
- 6. Fossil plants or vertebrates that are considered to be of good quality or scientific interest by the palaeontologist must be removed, catalogued and housed in a suitable institution where they can be made available for further study. Before the fossils are removed from the site a SAHRA permit must be obtained. Annual reports must be submitted to SAHRA as required by the relevant permits.
- 7. If no good fossil material is recovered then no site inspections by the palaeontologist will be necessary. A final report by the palaeontologist must be sent to SAHRA once the project has been completed and only if there are fossils.
- 8. If no fossils are found and the excavations have finished then no further monitoring is required.

9. Appendix A – Examples of fossils from the Dwyka Group.



Figure 5: Photographs of fossil plants of the early Glossopteris flora that occur in the Dwyka Group sediments in north western South Africa.

10. Appendix B – Details of specialist

Curriculum vitae (short) - Marion Bamford PhD January 2022

Personal details

Surname : **Bamford**

First names : **Marion Kathleen**

Present employment: Professor; Director of the Evolutionary Studies Institute.

Member Management Committee of the NRF/DST Centre of Excellence Palaeosciences, University of the Witwatersrand,

Johannesburg, South Africa

Telephone : +27 11 717 6690 Fax : +27 11 717 6694 Cell : 082 555 6937

E-mail : marion.bamford@wits.ac.za;

marionbamford12@gmail.com

ii) Academic qualifications

Tertiary Education: All at the University of the Witwatersrand:

1980-1982: BSc, majors in Botany and Microbiology. Graduated April 1983.

1983: BSc Honours, Botany and Palaeobotany. Graduated April 1984.

1984-1986: MSc in Palaeobotany. Graduated with Distinction, November 1986.

1986-1989: PhD in Palaeobotany. Graduated in June 1990.

NRF Rating: C-2 (1999-2004); B-3 (2005-2015); B-2 (2016-2020); B-1 (2021-2026)

iii) Professional qualifications

Wood Anatomy Training (overseas as nothing was available in South Africa):

1994 - Service d'Anatomie des Bois, Musée Royal de l'Afrique Centrale, Tervuren, Belgium, by Roger Dechamps

1997 - Université Pierre et Marie Curie, Paris, France, by Dr Jean-Claude Koeniguer

1997 - Université Claude Bernard, Lyon, France by Prof Georges Barale, Dr Jean-Pierre

Gros, and Dr Marc Philippe

iv) Membership of professional bodies/associations

Palaeontological Society of Southern Africa

Royal Society of Southern Africa - Fellow: 2006 onwards

Academy of Sciences of South Africa - Member: Oct 2014 onwards

International Association of Wood Anatomists - First enrolled: January 1991

International Organization of Palaeobotany - 1993+

Botanical Society of South Africa

South African Committee on Stratigraphy - Biostratigraphy - 1997 - 2016

SASQUA (South African Society for Quaternary Research) - 1997+

PAGES - 2008 - onwards: South African representative

ROCEEH / WAVE - 2008+

INQUA - PALCOMM - 2011+onwards

vii) Supervision of Higher Degrees

All at Wits University

Degree	Graduated/completed	Current
Honours	13	0

Masters	11	3
PhD	11	6
Postdoctoral fellows	15	1

viii) Undergraduate teaching

Geology II – Palaeobotany GEOL2008 – average 65 students per year Biology III – Palaeobotany APES3029 – average 45 students per year Honours – Evolution of Terrestrial Ecosystems; African Plio-Pleistocene Palaeoecology; Micropalaeontology – average 12-20 students per year.

ix) Editing and reviewing

Editor: Palaeontologia africana: 2003 to 2013; 2014 - Assistant editor

Guest Editor: Quaternary International: 2005 volume

Member of Board of Review: Review of Palaeobotany and Palynology: 2010 -

Associate Editor Open Science UK: 2021 -

Review of manuscripts for ISI-listed journals: 30 local and international journals Reviewing of funding applications for NRF, PAST, NWO, SIDA, National Geographic, Leakey Foundation

x) Palaeontological Impact Assessments

Selected from the past five years only – list not complete:

- Mala Mala 2017 for Henwood
- Modimolle 2017 for Green Vision
- Klipoortjie and Finaalspan 2017 for Delta BEC
- Ledjadja borrow pits 2018 for Digby Wells
- Lungile poultry farm 2018 for CTS
- Olienhout Dam 2018 for JP Celliers
- Isondlo and Kwasobabili 2018 for GCS
- Kanakies Gypsum 2018 for Cabanga
- Nababeep Copper mine 2018
- Glencore-Mbali pipeline 2018 for Digby Wells
- Remhoogte PR 2019 for A&HAS
- Bospoort Agriculture 2019 for Kudzala
- Overlooked Quarry 2019 for Cabanga
- Richards Bay Powerline 2019 for NGT
- Eilandia dam 2019 for ACO
- Eastlands Residential 2019 for HCAC
- Fairview MR 2019 for Cabanga
- Graspan project 2019 for HCAC
- Lieliefontein N&D 2019 for EnviroPro
- Skeerpoort Farm Mast 2020 for HCAC
- Vulindlela Eco village 2020 for 1World
- KwaZamakhule Township 2020 for Kudzala
- Sunset Copper 2020 for Digby Wells
- McCarthy-Salene 2020 for Prescali
- VLNR Lodge 2020 for HCAC
- Madadeni mixed use 2020 for EnviroPro

- Frankfort-Windfield Eskom Powerline 2020 for 1World
- Beaufort West PV Facility 2021 for ACO Associates
- Copper Sunset MR 2021 for Digby Wells
- Sannaspos PV facility 2021 for CTS Heritage
- Smithfield-Rouxville-Zastron PL 2021 for TheroServe

xi) Research Output

Publications by M K Bamford up to January 2022 peer-reviewed journals or scholarly books: over 160 articles published; 5 submitted/in press; 10 book chapters. Scopus h-index = 30; Google scholar h-index = 35; -i10-index = 92 Conferences: numerous presentations at local and international conferences.