

Palaeontological Impact Assessment for the proposed upgrade and irrigation for Prieska Copper Mine, at Copperton, Northern Cape Province

Desktop Study (Phase 1)

For

Heritage Contracts and Archaeological Consultants

19 July 2020

Prof Marion Bamford

Palaeobotanist

P Bag 652, WITS 2050

Johannesburg, South Africa

Marion.bamford@wits.ac.za


Expertise of Specialist

The Palaeontologist Consultant: Prof Marion Bamford
Qualifications: PhD (Wits Univ, 1990); FRSSAf, ASSAf
Experience: 31 years research; 23 years PIA studies

Declaration of Independence

This report has been compiled by Professor Marion Bamford, of the University of the Witwatersrand, sub-contracted by Heritage Contracts and Archaeological Consultants, Modimolle, South Africa. The views expressed in this report are entirely those of the author and no other interest was displayed during the decision making process for the Project.

Specialist: Prof Marion Bamford

Signature: 

Executive Summary

A palaeontological Impact Assessment was requested for the upgrade of Priestka Copper Mine at Copperton, Northern Cape Province, and expansion of the irrigation system. To comply with the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development..

The proposed site lies on the Aeolian sands of the Gordonia Formation (Kalahari Group, Quaternary age), and not on the Dwyka Group. Windblown sands seldom preserve fossils and then only in such features as springs or palaeo-pans but no such features are evident. Nonetheless, a Fossil Chance Find Protocol should be added to the EMPr. Based on this information it is recommended that no palaeontological site visit is required unless the responsible person on site finds fossils and then a professional palaeontologist should make an assessment from photographs and collect fossils if deemed important.

Table of Contents

Expertise of Specialist.....	1
Declaration of Independence.....	1
1. Background.....	4
2. Methods and Terms of Reference	6
3i. Project location and geological context.....	7
3ii. Palaeontological context	8
4. Impact assessment	10
5. Assumptions and uncertainties.....	11
6. Recommendation.....	11
7. References.....	12
8. Chance Find Protocol	13
Appendix A (examples of fossils)	14
Appendix B (short CV of specialist)	15

1. Background

Vadocube has extended the Mining Rights boundary for Prieska Copper Mine at Copperton, Northern Cape Province (Figures 1 and 2) and have proposed some changes within the boundary. In this report the whole area is considered and not the activities within the area.

A Palaeontological Impact Assessment was requested for the amended project. To comply with the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development and is reported herein.

Table 1: Specialist report requirements in terms of Appendix 6 of the EIA Regulations (amended 2017)

	A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:	Relevant section in report
ai	Details of the specialist who prepared the report	Appendix B
aii	The expertise of that person to compile a specialist report including a curriculum vitae	Appendix B
b	A declaration that the person is independent in a form as may be specified by the competent authority	Page 1
c	An indication of the scope of, and the purpose for which, the report was prepared	Section 1
ci	An indication of the quality and age of the base data used for the specialist report: SAHRIS palaeosensitivity map accessed – date of this report	Yes
cii	A description of existing impacts on the site, cumulative impacts of the proposed development and levels of acceptable change	Section 5
d	The date and season of the site investigation and the relevance of the season to the outcome of the assessment	N/A
e	A description of the methodology adopted in preparing the report or carrying out the specialised process	Section 2
f	The specific identified sensitivity of the site related to the activity and its associated structures and infrastructure	Section 4
g	An identification of any areas to be avoided, including buffers	N/A
h	A map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	N/A
i	A description of any assumptions made and any uncertainties or gaps in knowledge;	Section 5
j	A description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives, on the environment	Section 4

k	Any mitigation measures for inclusion in the EMPr	Section 7, Appendix A
l	Any conditions for inclusion in the environmental authorisation	N/A
m	Any monitoring requirements for inclusion in the EMPr or environmental authorisation	Section 7, Appendix A
ni	A reasoned opinion as to whether the proposed activity or portions thereof should be authorised	N/A
nii	If the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan	N/A
o	A description of any consultation process that was undertaken during the course of carrying out the study	N/A
p	A summary and copies if any comments that were received during any consultation process	N/A
q	Any other information requested by the competent authority.	N/A



Figure 1: Google Earth map of the Prieska Copper Mine at Copperton, Northern Cape Province with the projects indicated within the thin black lines. The irrigation project stretches to the northeast and within 3km from the mine centre. Map supplied by HCAC.




Figure 2: Annotated Google Earth maps comparing the original proposed layout with the amended layout of the proposed development of Prieska Copper Mine at Copperton. Map supplied by HCAC.

2. Methods and Terms of Reference

The Terms of Reference (ToR) for this study were to undertake a PIA and provide feasible management measures to comply with the requirements of SAHRA.

The methods employed to address the ToR included:

1. Consultation of geological maps, literature, palaeontological databases, published and unpublished records to determine the likelihood of fossils occurring in the affected areas. Sources included records housed at the Evolutionary Studies Institute at the University of the Witwatersrand and SAHRA databases;
2. Where necessary, site visits by a qualified palaeontologist to locate any fossils and assess their importance (*not applicable to this assessment*);
3. Where appropriate, collection of unique or rare fossils with the necessary permits for storage and curation at an appropriate facility (*not applicable to this assessment*); and
4. Determination of fossils' representivity or scientific importance to decide if the fossils can be destroyed or a representative sample collected (*not applicable to this assessment*).

3. Geology and Palaeontology

i. Project location and geological context

The Prieska Copper Mine lies along the NNW-SSE trending Brakbosch Fault that separates the western Areachap Terrane from the Kaaie Terrane to the east (Cornell et al., 2006). This region was tectonically active about 2000 - 1000 million year ago when basal rocks were reworked, then juvenile pre-crustal and plutonic rocks formed during rifting, ocean-spreading and subduction, followed by collision and intense deformation and metamorphism. Finally large and small granitoids intruded through the terranes (ibid). These ancient rocks have been overlain by the much younger Karoo Supergroup rocks that were subsequently eroded and replaced by young sands of the Kalahari Group. The metamorphism has resulted in economically important minerals such as copper and zinc. First discovered in the 1890s, the Prieska ore body on farm Vogelstruis Bult 104 has been explored and mined by a number of different companies (Wilson, 1998).




Figure 2: Geological map of the area around Copperton in the Northern Cape Province with the Prieska Copper Mine footprint within the yellow rectangle. Abbreviations of the rock types are explained in Table 2. Map enlarged from the Geological Survey 1: 250 000 map 2922 Prieska.

Table 2: Explanation of symbols for the geological map and approximate ages (Cornell et al., 2006; Johnson et al., 2006; McCarthy et al., 2006). SG = Supergroup; Fm = Formation; Ma = million years; grey shading = formations impacted by the project.

Symbol	Group/Formation	Lithology	Approximate Age
Qg	Gordonia Fm, Kalahari Group	Aeolian sand, sand dunes	Quaternary, ca 2.5 Ma to present
C-Pd	Dwyka Group, Karoo SG	Shales, sandstone, coal	Lower Permian, Middle Ecca
Mu	Uitdraai Fm, Brulpan Group, Namaqua Metamorphic Province	Banded to massive quartzite	Ca 2000 Ma
Mhe	Hedley Plains Fm, Jacomyns Pan Group, Namaqua Metamorphic Province	Coal-silicate rocks, amphibolite	Ca 2000 Ma
Mv	Vogelstruisbult Fm, Jacomyns Pan Group, Namaqua Metamorphic Province	Schist and gneiss	Ca 2000 Ma
Ms	Spionkop Fm, Namaqua Metamorphic Province	Fine-grained quartzite, schist and gneiss	Ca 2000 Ma

ii. Palaeontological context

The palaeontological sensitivity of the area under consideration is presented in Figure 4. The Prieska ore body is underground with only a small area of surface exposure. Since the material to be mined is highly metamorphosed ancient extrusive rocks, it does not preserve any fossils. The rest of the surface rocks are the Aeolian sands of the Gordonia Formation (Kalahari Group, Quaternary age). The sands unconformably overlie much older rocks on the Farm Vogelstruis Bult, except to the far south of the farm where tillites, mudstones and shales of the Dwyka Group (Karoo Supergroup) (Figure 3, Table 2). According to von Brunn and Visser (1999), the formation present here is the Mbizane Formation but this is not indicated on the geological map.

The Karoo Supergroup preserves a range of plant and animal fossils from the basal Dwyka Group sediments that filled the basin when the glacial ice sheets melted, the Ecca shales with coal and plant fossils, the Beaufort Group with a series of amphibians, mammal-like reptiles (therapsids), gorgonopsians, paraisaurs and early mammals, and plants. Dinosaurs are present in the upper Stormberg Group.

Fossils are rare in the Dwyka Group because the environment was very cold but a few plants were present and were deposited in the marginal lake. The Dwyka Group is made up of seven facies that were deposited in a marine basin under differing environmental settings of glacial formation and retreat (Visser, 1986, 1989; Johnson et al., 2006). In the north these are called the Mbizane Formation, and the Elandsvlei Formation in the south. Described below are the seven facies (Johnson et al., 2006 p463-465):

The massive diamictite facies comprises highly compacted diamictite that is clast-poor in the north. It was deposited in subaqueous or subglacial positions.

The stratified diamictite comprises alternating diamictite, mudrock, sandstone and conglomerate beds. They are interpreted as being rapidly deposited, sediment gravity flows but with some possible reworking of the subglacial diamictites.

The massive carbonate-rich diamictite facies is clast-poor and was formed by the rainout of debris, with the carbonate probably originating by crystallisation from interstitial waters.

The conglomerate facies ranges from single layer boulder beds to poorly sorted pebble and granule conglomerates. The boulder beds are interpreted as lodgement deposits whereas the poorly sorted conglomerates are a product of water-reworking of diamicton by high-density sediment gravity flows.

The sandstone facies were formed as turbidite deposits.

The mudrock with stones facies represents rainout deposits in the distal iceberg zone.

The mudrock facies consists of dark-coloured, commonly carbonaceous mudstone, shale or silty rhythmite that was formed when the mud or silt in suspension settled. This is the only fossiliferous facies of the Dwyka Group.

The Dwyka *Glossopteris* flora outcrops are very sporadic and rare. Of the seven facies that have been recognised in the Dwyka Group fossil plant fragments have only been recognised from the mudrock facies. They have been recorded from around Douglas only (Johnson et al., 2006; Anderson and McLachlan 1976) although the Dwyka Group exposures are very extensive. Jurassic Dolerites do not contain fossils as they are igneous intrusives.



Figure 4: SAHRIS palaeosensitivity map for the site for the proposed upgrade and amendment to Prieska Copper Mine, Copperton shown within the yellow rectangle. Background colours indicate the following degrees of sensitivity: red = very highly sensitive; orange/yellow = high; green = moderate; blue = low; grey = insignificant/zero.

The Prieska Copper Mine footprint lies entirely on the Gordonia Formation Aeolian sands with a few outcrops of metamorphic rocks of the Namaqua Metamorphic Sequence. The latter is non-fossiliferous.

Although the Gordonia Formation is moderately sensitive as far as the palaeontology is concerned, fossils are extremely rare in Aeolian or windblown sands. Fossils of any size cannot be transported by wind but occasionally one can find tufas, spring sites or palaeo-pans beneath the sands and they are visible in relief on the satellite imagery. Such fossil traps may have preserved bones, silicified plant fragments (wood) or archaeological material.

From the SAHRIS map above the area is indicated as moderately sensitive (green) so a desktop study has been completed.

4. Impact assessment

An assessment of the potential impacts to possible palaeontological resources considers the criteria encapsulated in Table 3:

TABLE 3A: CRITERIA FOR ASSESSING IMPACTS

PART A: DEFINITION AND CRITERIA		
Criteria for ranking of the SEVERITY/NATURE of environmental impacts	H	Substantial deterioration (death, illness or injury). Recommended level will often be violated. Vigorous community action.
	M	Moderate/ measurable deterioration (discomfort). Recommended level will occasionally be violated. Widespread complaints.
	L	Minor deterioration (nuisance or minor deterioration). Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	L+	Minor improvement. Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	M+	Moderate improvement. Will be within or better than the recommended level. No observed reaction.
	H+	Substantial improvement. Will be within or better than the recommended level. Favourable publicity.
Criteria for ranking the DURATION of impacts	L	Quickly reversible. Less than the project life. Short term
	M	Reversible over time. Life of the project. Medium term
	H	Permanent. Beyond closure. Long term.
Criteria for ranking the SPATIAL SCALE of impacts	L	Localised - Within the site boundary.
	M	Fairly widespread – Beyond the site boundary. Local
	H	Widespread – Far beyond site boundary. Regional/ national
PROBABILITY (of exposure to impacts)	H	Definite/ Continuous
	M	Possible/ frequent
	L	Unlikely/ seldom

TABLE 3B: IMPACT ASSESSMENT

PART B: ASSESSMENT		
SEVERITY/NATURE	H	-
	M	-
	L	Wind-blown sands do not preserve plant fossils; so far there are no records from the Mbizane Fm of plant or animal fossils in this region so it is very unlikely that fossils occur on the site. The impact would be very unlikely.

PART B: ASSESSMENT		
	L+	-
	M+	-
	H+	-
DURATION	L	-
	M	-
	H	Where manifest, the impact will be permanent.
SPATIAL SCALE	L	Since only the possible fossils within the area would be fossil bone or plant fragments from the Gordonia Fm Aeolian sands, the spatial scale will be localised within the site boundary.
	M	-
	H	-
PROBABILITY	H	-
	M	-
	L	It is extremely unlikely that any fossils would be found in the loose or Aeolian sand that covers the region, and no features such as springs or palaeo-pans are visible in the satellite imagery. Nonetheless, a Fossil Chance Find protocol should be added to the eventual EMPr.

Based on the nature of the project, surface activities may impact upon the fossil heritage if preserved in the development footprint. The geological structures suggest that the rocks below the surface are much too old and metamorphosed to contain fossils. The Mbizane Formation (Dwyka Group, Karoo Supergroup) is south of the project site (carefully measured on the maps: the project extends 2.1km south of the railway line (fixed datum) while the Dwyka Group outcrops beyond 3km). The site lies entirely on the Gordonia Formation Aeolian sands and there is a small chance that there might be Quaternary fossils in the sands. Since there is an extremely small chance that fossils may be disturbed, a Fossil Chance Find Protocol has been added to this report. Taking account of the defined criteria, the potential impact to fossil heritage resources is extremely low.

5. Assumptions and uncertainties

Based on the geology of the area and the palaeontological record as we know it, it can be assumed that the formation and layout of the granitoids, schists and gneisses are typical for the country and do not contain fossil plant, insect, invertebrate and vertebrate material. The Aeolian sands of the Quaternary period would only preserve more robust fossils such as bones or silicified plants or wood in features such as spring sites or palaeo-pans, however no such feature is evident in the project footprint.

6. Recommendation

Based on experience and the lack of any previously recorded fossils from the area, it is extremely unlikely that any fossils would be preserved in the Aeolian sands of the Quaternary aged Gordonia Formation. There is a very small chance that fossil may occur in the adjacent mudstones of the Dwyka Group but this is not in the project footprint. A Fossil Chance Find

Protocol for Quaternary fossils should be added to the EMPr: if fossils are found once mining has commenced then they should be rescued and a palaeontologist called to assess and collect a representative sample.

7. References

Anderson, J.M., Anderson, H.M., 1985. Palaeoflora of Southern Africa: Prodrum of South African megaflores, Devonian to Lower Cretaceous. A.A. Balkema, Rotterdam. 423 pp.

Anderson, A.M., McLachlan, I.R., 1976. The plant record in the Dwyka and Ecca Series (Permian) of the south-western half of the great Karoo Basin, South Africa. *Palaeontologia africana* 19, 31-42.

Cornell, D.H., Thomas, R.J., Moen, H.F.G., Reid, D.L., Moore, J.M., Gibson, R.L., 2006. The Namaqua-Natal Province. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. Pp 325-379.

Johnson, M.R., van Vuuren, C.J., Visser, J.N.J., Cole, D.I., Wickens, H.deV., Christie, A.D.M., Roberts, D.L., Brandl, G., 2006. Sedimentary rocks of the Karoo Supergroup. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. Pp 461 – 499.

Plumstead, E.P., 1969. Three thousand million years of plant life in Africa. *Geological Society of southern Africa, Annexure to Volume LXXII*. 72pp + 25 plates.

Visser, J.N.J., 1986. Lateral lithofacies relationship in the glaciogenic Dwyka Formation in the western and central parts of the Karoo Basin. *Transactions of the Geological Society of South Africa* 89, 373-383.

Visser, J.N.J., 1989. The Permo-Carboniferous Dwyka Formation of southern Africa: deposition by a predominantly subpolar marine ice sheet. *Palaeogeography, Palaeoclimatology, Palaeoecology* 70, 377-391.

Von Brunn, V., Visser, J.N.J. 1999. Lithostratigraphy of the Mbizane Formation (Dwyka group). *South African Committee for Stratigraphy, Lithostratigraphic Series No. 32*, 10 pp. Council for Geoscience, Pretoria.

Wilson, M.G.C., 1998. Copper. In: Wilson, M.G.C. and Anhaeusser, C.R., 1998. *The mineral Resources of South Africa: Handbook*, Council for Geoscience, Pretoria. pp 209-227.

8. Chance Find Protocol

Monitoring Programme for Palaeontology – to commence once the excavations / drilling / mining activities begin.

1. The following procedure is only required if fossils are seen on the surface and when drilling/excavations/mining commence.
2. When excavations begin the rocks and must be given a cursory inspection by the environmental officer or designated person. Any fossiliferous material (plants, insects, bone) should be put aside in a suitably protected place. This way the project activities will not be interrupted.
3. Photographs of similar fossil plants must be provided to the developer to assist in recognizing the fossil plants in the shales and mudstones (for example see Figure 1.5). This information will be built into the EMP's training and awareness plan and procedures.
4. Photographs of the putative fossils can be sent to the palaeontologist for a preliminary assessment.
5. If there is any possible fossil material found by the developer/environmental officer/miners then the qualified palaeontologist sub-contracted for this project, should visit the site to inspect the selected material and check the dumps where feasible.
6. Fossil plants or vertebrates that are considered to be of good quality or scientific interest by the palaeontologist must be removed, catalogued and housed in a suitable institution where they can be made available for further study. Before the fossils are removed from the site a SAHRA permit must be obtained. Annual reports must be submitted to SAHRA as required by the relevant permits.
7. If no good fossil material is recovered then no site inspections by the palaeontologist will be necessary. A final report by the palaeontologist must be sent to SAHRA once the project has been completed and only if there are fossils.
8. If no fossils are found and the excavations have finished then no further monitoring is required.

Appendix A – Examples of fossils from the Quaternary sands.



Figure 5: Fragments of bone from a Quaternary paleo-pan.



Figure 6: Pieces of silicified wood from a Pleistocene alluvial deposit.

Appendix B – Details of specialist

Curriculum vitae (short) - Marion Bamford PhD April 2020

i) Personal details

Surname : **Bamford**
First names : **Marion Kathleen**
Present employment : Professor; Director of the Evolutionary Studies Institute.
Member Management Committee of the NRF/DST Centre of Excellence Palaeosciences, University of the Witwatersrand, Johannesburg, South Africa-
Telephone : +27 11 717 6690
Fax : +27 11 717 6694
Cell : 082 555 6937
E-mail : marion.bamford@wits.ac.za ; marionbamford12@gmail.com

ii) Academic qualifications

Tertiary Education: All at the University of the Witwatersrand:
1980-1982: BSc, majors in Botany and Microbiology. Graduated April 1983.
1983: BSc Honours, Botany and Palaeobotany. Graduated April 1984.
1984-1986: MSc in Palaeobotany. Graduated with Distinction, November 1986.
1986-1989: PhD in Palaeobotany. Graduated in June 1990.

iii) Professional qualifications

Wood Anatomy Training (overseas as nothing was available in South Africa):
1994 - Service d'Anatomie des Bois, Musée Royal de l'Afrique Centrale, Tervuren, Belgium, by Roger Dechamps
1997 - Université Pierre et Marie Curie, Paris, France, by Dr Jean-Claude Koeniguer
1997 - Université Claude Bernard, Lyon, France by Prof Georges Barale, Dr Jean-Pierre Gros, and Dr Marc Philippe

iv) Membership of professional bodies/associations

Palaeontological Society of Southern Africa
Royal Society of Southern Africa - Fellow: 2006 onwards
Academy of Sciences of South Africa - Member: Oct 2014 onwards
International Association of Wood Anatomists - First enrolled: January 1991
International Organization of Palaeobotany – 1993+

Botanical Society of South Africa
 South African Committee on Stratigraphy – Biostratigraphy - 1997 - 2016
 SASQUA (South African Society for Quaternary Research) – 1997+
 PAGES - 2008 –onwards: South African representative
 ROCEEH / WAVE – 2008+
 INQUA – PALCOMM – 2011+onwards

vii) Supervision of Higher Degrees

All at Wits University

Degree	Graduated/completed	Current
Honours	9	2
Masters	9	5
PhD	11	5
Postdoctoral fellows	10	4

viii) Undergraduate teaching

Geology II – Palaeobotany GEOL2008 – average 65 students per year
 Biology III – Palaeobotany APES3029 – average 25 students per year
 Honours – Evolution of Terrestrial Ecosystems; African Plio-Pleistocene Palaeoecology;
 Micropalaeontology – average 2-8 students per year.

ix) Editing and reviewing

Editor: Palaeontologia africana: 2003 to 2013; 2014 – Assistant editor
 Guest Editor: Quaternary International: 2005 volume
 Member of Board of Review: Review of Palaeobotany and Palynology: 2010 –
 Cretaceous Research: 2014 –
 Journal of African Earth Sciences: 2020 -

Review of manuscripts for ISI-listed journals: 25 local and international journals

x) Palaeontological Impact Assessments

Selected – list not complete:

- Thukela Biosphere Conservancy 1996; 2002 for DWAF
- Vioolsdrift 2007 for Xibula Exploration
- Rietfontein 2009 for Zitholele Consulting
- Bloeddrift-Baken 2010 for TransHex
- New Kleinfontein Gold Mine 2012 for Prime Resources (Pty) Ltd.
- Thabazimbi Iron Cave 2012 for Professional Grave Solutions (Pty) Ltd
- Delmas 2013 for Jones and Wagener
- Klipfontein 2013 for Jones and Wagener
- Platinum mine 2013 for Lonmin
- Syferfontein 2014 for Digby Wells
- Canyon Springs 2014 for Prime Resources

- Kimberley Eskom 2014 for Landscape Dynamics
- Yzermyne 2014 for Digby Wells
- Matimba 2015 for Royal HaskoningDV
- Commissiekraal 2015 for SLR
- Harmony PV 2015 for Savannah Environmental
- Glencore-Tweefontein 2015 for Digby Wells
- Umkomazi 2015 for JLB Consulting
- Ixia coal 2016 for Digby Wells
- Lambda Eskom for Digby Wells
- Alexander Scoping for SLR
- Perseus-Kronos-Aries Eskom 2016 for NGT
- Mala Mala 2017 for Henwood
- Modimolle 2017 for Green Vision
- Klipoortjie and Finaalspan 2017 for Delta BEC
- Ledjadja borrow pits 2018 for Digby Wells
- Lungile poultry farm 2018 for CTS
- Olienhout Dam 2018 for JP Celliers
- Isondlo and Kwasobabili 2018 for GCS
- Kanakies Gypsum 2018 for Cabanga
- Nababeep Copper mine 2018
- Glencore-Mbali pipeline 2018 for Digby Wells
- Remhoogte PR 2019 for A&HAS
- Bospoort Agriculture 2019 for Kudzala
- Overlooked Quarry 2019 for Cabanga
- Richards Bay Powerline 2019 for NGT
- Eilandia dam 2019 for ACO
- Eastlands Residential 2019 for HCAC
- Fairview MR 2019 for Cabanga
- Graspan project 2019 for HCAC
- Lielifontein N&D 2019 for Enviropro
- Skeerpoort Farm Mast 2020 for HCAC
- Vulindlela Eco village 2020 for 1World
- KwaZamakhule Township 2020 for Kudzala
- Sunset Copper 2020 for Digby Wells
-

xi) Research Output

Publications by M K Bamford up to December 2019 peer-reviewed journals or scholarly books: over 140 articles published; 5 submitted/in press; 8 book chapters.

Scopus h-index = 27; Google scholar h-index = 32; -i10-index = 80

Conferences: numerous presentations at local and international conferences.

xii) NRF Rating

NRF Rating: B-2 (2016-2020)

NRF Rating: B-3 (2010-2015)
NRF Rating: B-3 (2005-2009)
NRF Rating: C-2 (1999-2004)