

**Palaeontological Impact Assessment for the proposed  
King Cetshwayo Private Hospital and Training  
Hospital, near Ulundi,  
KwaZulu Natal Province**

**Desktop Study**

**For**

**Wagna Projects**

**02 December 2018**

**Prof Marion Bamford**

Palaeobotanist

P Bag 652, WITS 2050

Johannesburg, South Africa

[Marion.bamford@wits.ac.za](mailto:Marion.bamford@wits.ac.za)

## **Expertise of Specialist**

The Palaeontologist Consultant is: Prof Marion Bamford  
Qualifications: PhD (Wits Univ, 1990); FRSSAf, ASSAf  
Experience: 30 years research; 22 years PIA studies

## **Declaration of Independence**

This report has been compiled by Professor Marion Bamford, of the University of the Witwatersrand, sub-contracted by Wagna Projects, KwaZulu Natal, South Africa. The views expressed in this report are entirely those of the author and no other interest was displayed during the decision making process for the Project.

Specialist: Prof Marion Bamford

Signature: 

## **Executive Summary**

A palaeontological Impact Assessment was requested for the construction of the King Cetshwayo Private Hospital and Training Hospital just south of Ulundi D and north of Ulundi in northern KwaZulu Natal. To comply with the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development of a new hospital.

The proposed site lies on the tillites, shales, mudstones and sandstones of the Emakwezini Dwyka Group, Upper Carboniferous of the eastern part of the Main Karoo Basin. The mudrock facies of this group preserves impressions of fossils of the *Glossopteris* flora in other parts of the basin but none has been recorded from here. Nonetheless once excavations begin a monitoring programme and Chance Find Protocol should be added to the EMPr. Based on this information it is recommended that no palaeontological site visit is required until such time as excavations begin and if fossils are discovered.

## Table of Contents

Expertise of Specialist.....	1
Declaration of Independence.....	1
1. Background.....	4
2. Methods and Terms of Reference .....	6
i. Project location and geological context.....	6
ii. Palaeontological context .....	7
4. Impact assessment.....	9
5. Assumptions and uncertainties.....	10
6. Recommendation.....	10
7. References.....	11
8. Chance Find Protocol .....	12
Appendix A (examples of fossils) .....	13
Appendix B (short CV of specialist) .....	15

# 1. Background

A Palaeontological Impact Assessment was requested for the proposed construction of a private hospital and a training hospital, the King Cetshwayo Hospital between Ulundi D and Ulundi, northern KwaZulu Natal.

To comply with the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development of a new hospital and related infrastructure.

Table 1: Specialist report requirements in terms of Appendix 6 of the EIA Regulations (2014)

A specialist report prepared in terms of the Environmental Impact Regulations of 2014 must contain:	Relevant section in report
Details of the specialist who prepared the report	Appendix B
The expertise of that person to compile a specialist report including a curriculum vitae	Appendix B
A declaration that the person is independent in a form as may be specified by the competent authority	Page 1
An indication of the scope of, and the purpose for which, the report was prepared	Section 1
The date and season of the site investigation and the relevance of the season to the outcome of the assessment	N/A
A description of the methodology adopted in preparing the report or carrying out the specialised process	Section 2
The specific identified sensitivity of the site related to the activity and its associated structures and infrastructure	Section ii <b>Error! Reference source not found. 5</b>
An identification of any areas to be avoided, including buffers	N/A
A map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	N/A
A description of any assumptions made and any uncertainties or gaps in knowledge;	Section 5
A description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives, on the environment	Section 4
Any mitigation measures for inclusion in the EMPr	n/a
Any conditions for inclusion in the environmental authorisation	n/a
Any monitoring requirements for inclusion in the EMPr or environmental authorisation	Section 8

A reasoned opinion as to whether the proposed activity or portions thereof should be authorised	N/A
If the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMP, and where applicable, the closure plan	N/A
A description of any consultation process that was undertaken during the course of carrying out the study	N/A
A summary and copies if any comments that were received during any consultation process	N/A
Any other information requested by the competent authority.	N/A



Figure 1: Google Earth map of the proposed site for the King Cetshwayo Private Hospital and Training Hospital. The urban development to the north is Ulundi D. Map supplied by Wagna Projects.

## 2. Methods and Terms of Reference

The Terms of Reference (ToR) for this study were to undertake a PIA and provide feasible management measures to comply with the requirements of SAHRA.

The methods employed to address the ToR included:

1. Consultation of geological maps, literature, palaeontological databases, published and unpublished records to determine the likelihood of fossils occurring in the affected areas. Sources included records housed at the Evolutionary Studies Institute at the University of the Witwatersrand and SAHRA databases;
2. Where necessary, site visits by a qualified palaeontologist to locate any fossils and assess their importance (*not applicable to this assessment*);
3. Where appropriate, collection of unique or rare fossils with the necessary permits for storage and curation at an appropriate facility (*not applicable to this assessment*); and
4. Determination of fossils' representivity or scientific importance to decide if the fossils can be destroyed or a representative sample collected (*not applicable to this assessment*).

### 3. Geology and Palaeontology

#### i. Project location and geological context




Figure 2: Geological map of the area around Ulundi in northern KwaZulu Natal. The location of the proposed project is indicated with the arrow. Abbreviations of the rock types are explained in Table 2. Map enlarged from the Geological Survey 1: 1 000 000 map 1984.

Table 2: Explanation of symbols for the geological map and approximate ages (Johnson et al., 2006;; Marshall, 2006). SG = Supergroup; Fm = Formation.

Symbol	Group/Formation	Lithology	Approximate Age
Pvo	Volksrust Fm, Eccca Group	shale	Permian middle, Upper Eccca
Pv	Vryheid Fm, Eccca Group	Shales, sandstone, coal	Lower Permian, Middle Eccca
Pp	Pietermaritzburg Fm, Eccca Group	Shale	Lower Eccca, early Permian
C-Pd	Dwyka Group	Tillite, sandstone, mudstone, shale	Late Carboniferous to early Permian
O-S	Natal Group	Quartzitic sandstone, arkose, shale	Ordovician to Silurian
ZB	Basement complex	Potassic Granite, granodiorite	>3100 Ma

The oldest rocks in the area are the basement of Late Archean age and are part of the Kaap-Vaal Craton, comprising a number of granites and granodiorites. Unconformably overlying these rocks are the quartzitic sandstones and shales of the Natal Group. The overlying Dwyka tillites represent the deposits from the receding glaciers during the Upper Carboniferous. The next stratum is the Pietermaritzburg Formation shale that represent a major post-glacial transgression and a relatively shallow water setting (Johnson et al., 2006). It is the lowermost part of the Eccca Group of the Karoo Supergroup.

The Vryheid and Volksrust Formations are present in this area and the former has a number of coal lenses. In contrast the Volksrust Formation, comprising grey to black silty shales, represents a transgressive and possibly open shelf sequence made up mostly of muds that were deposited from suspension (Johnson et al., 2006).

## ii. Palaeontological context

The palaeontological sensitivity of the area under consideration is presented in Figure 3. The Site lies in the tillite, sandstone, mudstone and shales of the Dwyka Group that is Upper Carboniferous in age. These strata represent the sediments deposited by the retreating ice sheets that covered much of Gondwanaland, including South Africa.

The Dwyka Group is made up of seven facies that were deposited in a marine basin under differing environmental settings of glacial formation and retreat (Visser, 1986, 1989; Johnson et al., 2006). In the north these are called the Mbizane Formation, and the Elandsvlei Formation in the south. Described below are the seven facies (Johnson et al., 2006, p. 463-465):

The massive diamictite facies comprises highly compacted diamictite that is clast-poor in the north. It was deposited in subaqueous or subglacial positions.

The stratified diamictite comprises alternating diamictite, mudrock, sandstone and conglomerate beds. They are interpreted as being rapidly deposited, sediment gravity flows but with some possible reworking of the subglacial diamictites.

The massive carbonate-rich diamictite facies is clast-poor and was formed by the rainout of debris, with the carbonate probably originating by crystallisation from interstitial waters.



The conglomerate facies ranges from single layer boulder beds to poorly sorted pebble and granule conglomerates. The boulder beds are interpreted as lodgement deposits whereas the poorly sorted conglomerates are a product of water-reworking of diamicton by high-density sediment gravity flows.

The sandstone facies were formed as turbidite deposits.

The mudrock with stones facies represents rainout deposits in the distal iceberg zone.

The mudrock facies consists of dark-coloured, commonly carbonaceous mudstone, shale or silty rhythmite that was formed when the mud or silt in suspension settled. This is the only fossiliferous facies of the Dwyka Group.

The Dwyka *Glossopteris* flora outcrops are very sporadic and rare. Of the seven facies that have been recognised in the Dwyka Group fossil plant fragments have only been recognised from the mudrock facies. They have been recorded from around Douglas only (Johnson et al., 2006; Anderson and McLachlan 1976) although the Dwyka Group exposures are very extensive. Jurassic Dolerites do not contain fossils as they are igneous intrusives.



Figure 3: SAHRIS palaeosensitivity maps for the site for the proposed King Cetshwayo Private and Training Hospitals shown within the yellow rectangle. Colours indicate the following degrees of sensitivity: red = very highly sensitive; orange/yellow = high; green = moderate; blue = low; grey = insignificant/zero.

The Pietermaritzburg Formation overlies the Dwyka Group with a sharp contact and it is made up of dark silty mudrock and is heavily bioturbated especially in the sandy and silty beds. This formation represents a major post-glacial transgression with relatively shallow water on an unstable shelf (Johnson et al., 2006). No plant fossils have been recorded from this stratum and it predates terrestrial vertebrate fossils.

Vertebrate fossils do not occur in the Dwyka Group, but plant fossils of the *Glossopteris* flora, for example *Glossopteris* leaves, lycopods, sphenophytes, fern and wood fragments

have been reported from only the mudrock facies (Johnson et al., 2006; Anderson and Anderson, 1985). The fossils, however, are very sporadic and never abundant.

From the SAHRIS map above the area is indicated as moderately sensitive (green) so a desktop study is presented here. The area is highly disturbed from informal agriculture and pastoral practices so the ground surface and soils would not preserve any recognisable fossil plants. There might be fossils below the surface.

## 4. Impact assessment

An assessment of the potential impacts to possible palaeontological resources considers the criteria encapsulated in Table 3:

**TABLE 3A: CRITERIA FOR ASSESSING IMPACTS**

<b>PART A: DEFINITION AND CRITERIA</b>		
<b>Criteria for ranking of the SEVERITY/NATURE of environmental impacts</b>	<b>H</b>	Substantial deterioration (death, illness or injury). Recommended level will often be violated. Vigorous community action.
	<b>M</b>	Moderate/ measurable deterioration (discomfort). Recommended level will occasionally be violated. Widespread complaints.
	<b>L</b>	Minor deterioration (nuisance or minor deterioration). Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	<b>L+</b>	Minor improvement. Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	<b>M+</b>	Moderate improvement. Will be within or better than the recommended level. No observed reaction.
	<b>H+</b>	Substantial improvement. Will be within or better than the recommended level. Favourable publicity.
<b>Criteria for ranking the DURATION of impacts</b>	<b>L</b>	Quickly reversible. Less than the project life. Short term
	<b>M</b>	Reversible over time. Life of the project. Medium term
	<b>H</b>	Permanent. Beyond closure. Long term.
<b>Criteria for ranking the SPATIAL SCALE of impacts</b>	<b>L</b>	Localised - Within the site boundary.
	<b>M</b>	Fairly widespread – Beyond the site boundary. Local
	<b>H</b>	Widespread – Far beyond site boundary. Regional/ national
<b>PROBABILITY (of exposure to impacts)</b>	<b>H</b>	Definite/ Continuous
	<b>M</b>	Possible/ frequent
	<b>L</b>	Unlikely/ seldom

**TABLE 3B: IMPACT ASSESSMENT**

<b>PART B: ASSESSMENT</b>		
<b>SEVERITY/NATURE</b>	<b>H</b>	-
	<b>M</b>	.
	<b>L</b>	There is a small chance that fossil plants will occur in the mudrock facies of the Dwyka Group so the impact would be low
	<b>L+</b>	-
	<b>M+</b>	-
	<b>H+</b>	-
<b>DURATION</b>	<b>L</b>	-
	<b>M</b>	-
	<b>H</b>	Where manifest, the impact will be permanent.

PART B: ASSESSMENT		
SPATIAL SCALE	L	Since only the possible fossils within the area would be fossil plants from the <i>Glossopteris</i> flora in the mudrock, the spatial scale will be localised within the site boundary.
	M	-
	H	-
PROBABILITY	H	-
	M	-
	L	It is possible that fossil plants occur in the Dwyka group mudrock facies but they have not been recorded from here; where they do occur they are sporadic and not abundant. A Fossil Chance Find Protocol should be added to the EMPr.

Based on the nature of the project, surface activities are unlikely to impact on the fossil heritage given that the ground surface is already highly disturbed but they could be preserved below the round surface. A Fossil Chance Find Protocol has been added to this report. Taking account of the defined criteria, the potential impact to fossil heritage resources is low only once excavation and construction of foundations have penetrated below the surface.

## 5. Assumptions and uncertainties

Based on the geology of the area and the palaeontological record as we know it, it can be assumed that the formation and layout of the dolomites, sandstones, shales, coals and sands are typical for the country and do contain fossil plant and invertebrate material. Although not reported to date from the area around Ulundi it can be assumed that fossil plant impressions of the *Glossopteris* flora could occur in the mudstone facies of the Dwyka Group at this site because they have been reported from this formation to the west.

## 6. Recommendation

Based on experience and the lack of any previously recorded fossils from the area, it is nonetheless possible that fossil plant impressions could occur at the site but not on the surface, but possibly below the surface. It is recommended, therefore, that a Fossil Chance Find Protocol be added to the EMPr such that once excavations for the building foundations and infrastructure have commenced the geologist, environmentalist or other responsible person looks out for fossils and reports any occurrences to a professional palaeontologist for assessment of the scientific value, and to make a representative collection once an AMAFA permit has been obtained.

## 7. References

- Anderson, J.M., Anderson, H.M., 1985. Palaeoflora of Southern Africa: Prodrum of South African megaflores, Devonian to Lower Cretaceous. A.A. Balkema, Rotterdam. 423 pp.
- Anderson, A.M., 1981. The Umfolozia arthropod trackways in the Permian Dwyka and Ecca Series of South Africa. *Journal of Palaeontology* 55, 84-108.
- Anderson, A.M., McLachlan, I.R., 1976. The plant record in the Dwyka and Ecca Series (Permian) of the south-western half of the great Karoo Basin, South Africa. *Palaeontologia africana* 19, 31-42.
- Johnson, M.R., van Vuuren, C.J., Visser, J.N.J., Cole, D.I., Wickens, H.deV., Christie, A.D.M., Roberts, D.L., Brandl, G., 2006. Sedimentary rocks of the Karoo Supergroup. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. Pp 461 – 499.
- Marshall, G.G.A., 2006. The Natal Group. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. Pp 433-441.
- Plumstead, E.P., 1969. Three thousand million years of plant life in Africa. *Geological Society of southern Africa, Annexure to Volume LXXII*. 72pp + 25 plates.
- Visser, J.N.J., 1986. Lateral lithofacies relationship in the glaciogenic Dwyka Formation in the western and central parts of the Karoo Basin. *Transactions of the Geological Society of South Africa* 89, 373-383.
- Visser, J.N.J., 1989. The Permo-Carboniferous Dwyka Formation of southern Africa: deposition by a predominantly subpolar marine icesheet. *Palaeogeography, Palaeoclimatology, Palaeoecology* 70, 377-391.

## 8. Chance Find Protocol

### **Monitoring Programme for Palaeontology – to commence once the excavations for foundations and infrastructure have begun.**

1. The following procedure is only required if fossils are seen on the surface or and when excavations commence.
2. When excavations begin the rocks and must be given a cursory inspection by the environmental officer or designated person. Any fossiliferous material (plants, insects, bone, coal) should be put aside in a suitably protected place. This way the construction activities will not be interrupted.
3. Photographs of similar fossil plants must be provided to the developer to assist in recognizing the fossil plants in the shales and mudstones (for example see Figures 4, 5). This information will be built into the EMP's training and awareness plan and procedures.
4. Photographs of the putative fossils can be sent to the palaeontologist for a preliminary assessment.
5. If there is any possible fossil material found by the developer/environmental officer then the qualified palaeontologist sub-contracted for this project, should visit the site to inspect the selected material and check the dumps where feasible.
6. Fossil plants or invertebrates that are considered to be of good quality or scientific interest by the palaeontologist must be removed, catalogued and housed in a suitable institution where they can be made available for further study. Before the fossils are removed from the site a SAHRA or AMAFA permit must be obtained. Annual reports must be submitted to SAHRA as required by the relevant permits.
7. If no good fossil material is recovered then the site inspections by the palaeontologist will not be necessary. Annual reports by the palaeontologist must be sent to SAHRA.
8. If no fossils are found and the excavations have finished then no further monitoring is required.

## **Appendix A – Examples of Dwyka Fossil Plants**




Figure 4: Examples of fossil plants found in the Dwyka Group near Douglas (taken from Anderson and McLachlan 1976, plate 1)




Figure 5: More examples of fossil plants from the Dwyka Group near Douglas (taken from Anderson and McLachlan, 1976, plate 2).



## Appendix B – Details of specialist

### Curriculum vitae (short) - Marion Bamford PhD October 2018

#### i) Personal details

Surname : **Bamford**  
First names : **Marion Kathleen**  
Present employment : Professor; Director of the Evolutionary Studies Institute.  
Member Management Committee of the NRF/DST Centre of Excellence Palaeosciences, University of the Witwatersrand, Johannesburg, South Africa-  
Telephone : +27 11 717 6690  
Fax : +27 11 717 6694  
Cell : 082 555 6937  
E-mail : [marion.bamford@wits.ac.za](mailto:marion.bamford@wits.ac.za) ; [marionbamford12@gmail.com](mailto:marionbamford12@gmail.com)

#### ii) Academic qualifications

Tertiary Education: All at the University of the Witwatersrand:

1980-1982: BSc, majors in Botany and Microbiology. Graduated April 1983.

1983: BSc Honours, Botany and Palaeobotany. Graduated April 1984.

1984-1986: MSc in Palaeobotany. Graduated with Distinction, November 1986.

1986-1989: PhD in Palaeobotany. Graduated in June 1990.

#### iii) Professional qualifications

*Wood Anatomy Training (overseas as nothing was available in South Africa):*

1994 - Service d'Anatomie des Bois, Musée Royal de l'Afrique Centrale, Tervuren, Belgium, by Roger Dechamps

1997 - Université Pierre et Marie Curie, Paris, France, by Dr Jean-Claude Koeniguer

1997 - Université Claude Bernard, Lyon, France by Prof Georges Barale, Dr Jean-Pierre Gros, and Dr Marc Philippe

#### iv) Membership of professional bodies/associations

Palaeontological Society of Southern Africa

Royal Society of Southern Africa - Fellow: 2006 onwards

Academy of Sciences of South Africa - Member: Oct 2014 onwards

International Association of Wood Anatomists - First enrolled: January 1991

International Organization of Palaeobotany – 1993+

Botanical Society of South Africa

South African Committee on Stratigraphy – Biostratigraphy - 1997 - 2016

SASQUA (South African Society for Quaternary Research) – 1997+

PAGES - 2008 –onwards: South African representative

ROCEEH / WAVE – 2008+

INQUA – PALCOMM – 2011+onwards

## vii) Supervision of Higher Degrees

All at Wits University

Degree	Graduated/completed	Current
Honours	6	1
Masters	8	1
PhD	10	2
Postdoctoral fellows	9	3

## viii) Undergraduate teaching

Geology II – Palaeobotany GEOL2008 – average 65 students per year

Biology III – Palaeobotany APES3029 – average 25 students per year

Honours – Evolution of Terrestrial Ecosystems; African Plio-Pleistocene Palaeoecology;

Micropalaeontology – average 2-8 students per year.

## ix) Editing and reviewing

Editor: Palaeontologia africana: 2003 to 2013; 2014 – Assistant editor

Guest Editor: Quaternary International: 2005 volume

Member of Board of Review: Review of Palaeobotany and Palynology: 2010 –  
Cretaceous Research: 2014 -

Review of manuscripts for ISI-listed journals: 25 local and international journals

## x) Palaeontological Impact Assessments

Selected – list not complete:

- Thukela Biosphere Conservancy 1996; 2002 for DWAF
- Vioolsdrift 2007 for Xibula Exploration
- Rietfontein 2009 for Zitholele Consulting
- Bloeddrift-Baken 2010 for TransHex
- New Kleinfontein Gold Mine 2012 for Prime Resources (Pty) Ltd.
- Thabazimbi Iron Cave 2012 for Professional Grave Solutions (Pty) Ltd
- Delmas 2013 for Jones and Wagener
- Klipfontein 2013 for Jones and Wagener
- Platinum mine 2013 for Lonmin
- Syferfontein 2014 for Digby Wells
- Canyon Springs 2014 for Prime Resources
- Kimberley Eskom 2014 for Landscape Dynamics
- Yzermyne 2014 for Digby Wells
- Matimba 2015 for Royal HaskoningDV
- Commissiekraal 2015 for SLR
- Harmony PV 2015 for Savannah Environmental
- Glencore-Tweefontein 2015 for Digby Wells
- Umkomazi 2015 for JLB Consulting
- Ixia coal 2016 for Digby Wells

- Lambda Eskom for Digby Wells
- Alexander Scoping for SLR
- Perseus-Kronos-Aries Eskom 2016 for NGT
- Mala Mala 2017 for Henwood
- Modimolle 2017 for Green Vision
- Klipoortjie and Finaalspan 2017 for Delta BEC
- Ledjadja borrow pits 2018 for Digby Wells
- Lungile poultry farm 2018 for CTS
- Olienhout Dam 2018 for JP Celliers
- Isondlo and Kwasobabili 2018 for GCS
- Kanakies Gypsum 2018 for Cabanga
- Nababeep Copper mine 2018
- Glencore-Mbali pipeline 2018 for Digby Wells
- 

### **xi) Research Output**

Publications by M K Bamford up to June 2018 peer-reviewed journals or scholarly books: over 120 articles published; 5 submitted/in press; 8 book chapters.

Scopus h index = 26; Google scholar h index = 28;

Conferences: numerous presentations at local and international conferences.

### **xii) NRF Rating**

NRF Rating: B-2 (2016-2020)

NRF Rating: B-3 (2010-2015)

NRF Rating: B-3 (2005-2009)

NRF Rating: C-2 (1999-2004)