

**Palaeontological Impact Assessment for the proposed  
Dalpark Retail Development, near Carnival City,  
Boksburg, Gauteng Province**

**Site Visit (Phase 2) Report**

**For**

**Heritage Contracts and Archaeological Consulting**

**26 January 2021**

**Prof Marion Bamford**

Palaeobotanist

P Bag 652, WITS 2050

Johannesburg, South Africa

[Marion.bamford@wits.ac.za](mailto:Marion.bamford@wits.ac.za)

## **Expertise of Specialist**

The Palaeontologist Consultant is: Prof Marion Bamford  
Qualifications: PhD (Wits Univ, 1990); FRSSAf, ASSAf  
Experience: 32 years research; 24 years PIA studies

## **Declaration of Independence**

This report has been compiled by Professor Marion Bamford, of the University of the Witwatersrand, sub-contracted by Heritage Contracts and Archaeological Consulting, Modimolle, South Africa. The views expressed in this report are entirely those of the author and no other interest was displayed during the decision making process for the Project.

Specialist: Prof Marion Bamford

Signature:

A handwritten signature in blue ink, appearing to read 'MKBamford', with a horizontal line underneath it.

## **Executive Summary**

A palaeontological Impact Assessment was requested for the proposed retail development in Dalpark Ext 19, on Erf 3145, near Carnival City in Boksburg.

To comply with the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a site visit (Phase 2) Palaeontological Impact Assessment (PIA) was completed for the proposed development.

The proposed site lies mostly on moderately sensitive rocks of the Dwyka Group with a small section in the northwest on the shales of the Vryheid Formation (Ecca Group, Karoo Supergroup) that potentially have fossil plants of the Glossopteris flora. From the site visit observations, however, NO FOSSILS were found. The area is disturbed from urban activities and is covered by soil and vegetation. It is not possible to tell if there are any fossils below the surface until excavations for the buildings and amenities have commenced. Therefore, a Fossil Chance Find Protocol should be added to the EMPr. Based on this information it is recommended that no further palaeontological site visit is required unless fossils are revealed when the excavations commence. Then a professional palaeontologist should be called to rescue and collect a representative sample of fossils to be housed in a recognised institution.

## Table of Contents

Expertise of Specialist.....	1
Declaration of Independence.....	1
1. Background.....	4
2. Methods and Terms of Reference.....	6
3. Geology and Palaeontology.....	6
i. Project location and geological context.....	6
ii. Palaeontological context.....	8
4. Impact assessment.....	12
5. Assumptions and uncertainties.....	13
6. Recommendation.....	14
7. References.....	15
8. Chance Find Protocol.....	16

# 1. Background

The proposed retail development on a piece of vacant land near Carnival City in Dalpark Ext 19, Boksburg, Gauteng Province on Erf 3145 (Figure 1) has a small section on potentially very highly sensitive rocks according to the SAHRIS palaeosensitivity map. Most of the area is moderately sensitive.

Therefore, a Palaeontological Impact Assessment was requested for the Dalpark Retail Development. In order to comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a site visit (Phase 2) desktop Palaeontological Impact Assessment (PIA) was completed for the proposed development and is reported herein.

Table 1: Specialist report requirements in terms of Appendix 6 of the EIA Regulations (2017)

	<b>A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:</b>	<b>Relevant section in report</b>
ai	Details of the specialist who prepared the report	Appendix B
aii	The expertise of that person to compile a specialist report including a curriculum vitae	Appendix B
b	A declaration that the person is independent in a form as may be specified by the competent authority	Page 1
c	An indication of the scope of, and the purpose for which, the report was prepared	Section 1
ci	An indication of the quality and age of the base data used for the specialist report: SAHRIS palaeosensitivity map accessed – date of this report	Yes
cii	A description of existing impacts on the site, cumulative impacts of the proposed development and levels of acceptable change	Section 5
d	The date and season of the site investigation and the relevance of the season to the outcome of the assessment	N/A
e	A description of the methodology adopted in preparing the report or carrying out the specialised process	Section 2
f	The specific identified sensitivity of the site related to the activity and its associated structures and infrastructure	Section 4
g	An identification of any areas to be avoided, including buffers	N/A
h	A map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	N/A
i	A description of any assumptions made and any uncertainties or gaps in knowledge;	Section 5
j	A description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives, on the environment	Section 4

k	Any mitigation measures for inclusion in the EMPr	Appendix A
l	Any conditions for inclusion in the environmental authorisation	N/A
m	Any monitoring requirements for inclusion in the EMPr or environmental authorisation	Appendix A
ni	A reasoned opinion as to whether the proposed activity or portions thereof should be authorised	N/A
nii	If the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan	N/A
o	A description of any consultation process that was undertaken during the course of carrying out the study	N/A
p	A summary and copies if any comments that were received during any consultation process	N/A
q	Any other information requested by the competent authority.	N/A



Figure 1: Google Earth map of the proposed development of a Retail centre on Dalpark Ext 19, with the section shown by the yellow outline. Map supplied by HCAC.



Table 2: Explanation of symbols for the geological map and approximate ages (Eriksson et al., 2006. Johnson et al., 2006; McCarthy et al., 2006; Robb et al., 2006; van der Westhuizen et al., 2006). SG = Supergroup; Fm = Formation; Ma = million years; grey shading = formations impacted by the project.

Symbol	Group/Formation	Lithology	Approximate Age
Q	Quaternary	Alluvium, sand, calcrete	Neogene, ca 2.5 Ma to present
Jd	Jurassic dykes	Dolerite dykes, intrusive	Jurassic, approx. 180 Ma
Pv	Vryheid Fm, Ecca Group, Karoo SG	Shales, sandstone, coal	Early Permian, Middle Ecca
C-Pd	Dwyka Group, Karoo SG	Tillites, diamictites, mudstone, sandstone	Late Carboniferous to early Permian
Vm	Malmani Subgroup, Chuniespoort Group, Transvaal SG	Dolomite, chert	Ca 2750 – 2650 Ma
Vbr	Black Reef Fm, Ventersdorp SG	Quartzite, conglomerate, shale, basalt	Ca 2650 – 2640 Ma
Rt	Turfontein Subgroup, Central Rand Group, Witwatersrand SG	Conglomerate, quartzite	Ca 2750 Ma
Rjo	Johannesburg Subgroup, Central Rand Group, Witwatersrand SG	Quartzite, conglomerate, shale	

The site is in the Transvaal Basin and has rocks of three major time frames exposed in close proximity (Figure 2). Oldest rocks in the area are from the Witwatersrand Supergroup (the Turfontein and Johannesburg Subgroups). Unconformably overlying these rocks are the uppermost Black Reef Formation and the Malmani Subgroup. Much younger rocks of the basal Karoo Supergroup overlie this very uneven basement of different rocks. The oldest Karoo rocks are called the Dwyka Group.

During the Carboniferous Period South Africa was part of the huge continental landmass known as Gondwanaland and it was positioned over the South Pole. As a result there were several ice sheets that formed and melted, and covered most of South Africa (Visser, 1986, 1989; Isbell et al., 2012). Gradual melting of the ice as the continental mass moved northwards and the earth warmed, formed fine-grained sediments in the large inland sea. These are the oldest rocks in the system and are exposed around the outer part of the ancient Karoo Basin, and are known as the Dwyka Group. They comprise tillites, diamictites, mudstones, siltstones and sandstones that were deposited as the basin filled. This group has been divided into two formations with Elandsvlei Formation occurring throughout the basin and the upper Mbizane Formation occurring only in the Free State and KwaZulu Natal (Johnson et al., 2006).

Overlying the Dwyka Group rocks are rocks of the Ecca Group that are Early Permian in age. There are eleven formations recognised in this group but they do not all extend throughout the Karoo Basin. In the west and central part are the following formations, from base upwards: Prince Albert Formation, Whitehill Formation, Collingham Formation, Laingsburg /



Ripon Formations, Tierberg / Fort Brown Formations, and Waterford Formation. In the Free State and KwaZulu Natal, from the base upwards are the Pietermaritzburg Formation, Vryheid Formation and the Volksrust Formation. All of these sediments have varying proportions of sandstones, mudstones, shales and siltstones and represent shallow to deep water settings, deltas, rivers, streams and overbank depositional environments.

ii. Palaeontological context

The palaeontological sensitivity of the area under consideration is presented in Figure 3. The site for development is in the Dwyka Group tillites, diamictites, mudstones and sandstones, with only the narrow western section occurring on the Vryheid Formation.




Figure 3: SAHRIS palaeosensitivity map for the site for the proposed Dalpark Ext 19 Retail development on Erf 3147 shown within the yellow rectangle. Background colours indicate the following degrees of sensitivity: red = very highly sensitive; orange/yellow = high; green = moderate; blue = low; grey = insignificant/zero.

The site for the project is in the Dwyka Group tillites, sandstone, mudstone and shales, and these potentially could preserve fossils. Around 300-290 Ma the climate in southern Africa was still relatively cool, but there were well developed Carboniferous floras in the northern hemisphere. In South Africa, however, much of the land surface was covered by ice sheets. As they melted they dropped the moraine trapped in the ice, together with limited plant matter from the vegetation that gradually recovered and colonised the land surface. Terrestrial vertebrates had not evolved at this time. The late Carboniferous flora comprised *Glossopteris* leaves and seeds, wood, and other plants such as lycopods, sphenophytes and ferns.

The Dwyka Group is made up of seven facies that were deposited in a marine basin under differing environmental settings of glacial formation and retreat (Visser, 1986, 1989; Johnson et al., 2006). In the north and east these are called the Mbizane Formation, and the Elandsvlei Formation in the south and west. Described below are the seven facies (Johnson et al., 2006 p. 463-465):

The massive diamictite facies comprises highly compacted diamictite that is clast-poor in the north. It was deposited in subaqueous or subglacial positions.

The stratified diamictite comprises alternating diamictite, mudrock, sandstone and conglomerate beds. They are interpreted as being rapidly deposited, sediment gravity flows but with some possible reworking of the subglacial diamictites.

The massive carbonate-rich diamictite facies is clast-poor and was formed by the rainout of debris, with the carbonate probably originating by crystallisation from interstitial waters.

The conglomerate facies ranges from single layer boulder beds to poorly sorted pebble and granule conglomerates. The boulder beds are interpreted as lodgement deposits whereas the poorly sorted conglomerates are a product of water-reworking of diamicton by high-density sediment gravity flows.

The sandstone facies were formed as turbidite deposits.

The mudrock with stones facies represents rainout deposits in the distal iceberg zone.

The mudrock facies consists of dark-coloured, commonly carbonaceous mudstone, shale or silty rhythmite that was formed when the mud or silt in suspension settled. This is the only fossiliferous facies of the Dwyka Group.

The Dwyka *Glossopteris* flora outcrops are very sporadic and rare. Of the seven facies that have been recognised in the Dwyka Group, fossil plant fragments have only been recognised from the mudrock facies. They have been recorded from around Douglas only (Johnson et al., 2006; Anderson and McLachlan 1976) although the Dwyka Group exposures are very extensive. Jurassic Dolerites do not contain fossils as they are igneous intrusives.

The overlying Vryheid Formation has a good record of fossil plants but they are scattered and unpredictable in their distribution. The rich vegetation occupying the deltas, river margins and floodplains formed peats in some areas. Buried peats were altered by heat and pressure over time and were transformed into coal seams.

Although no plant structure is visible in colas, often the lenses between the coals preserve impressions of the plants. In this formation they are known as the Glossopteris flora, named after the dominant plant group that is now extinct. Other plants in the flora include lycopods, sphenophytes, ferns and early gymnosperms (Plumstead, 1969, Anderson and Anderson, 1985).

From the SAHRIS map above (Figure 3) the majority of the area is indicated as moderately sensitive (green) for the Dwyka Group, and very highly sensitive (red) for the Vryheid Formation. Therefore, a site visit was undertaken.

### **lil Site visit observations**

As part of the Archaeological Heritage survey Dr Jaco van der Walt also looked for fossils in the project footprint. The photographs presented below were taken by him.

The area is already disturbed by previous urban or peri-urban activities, and the land is covered with soils and vegetation, secondary grassland and introduced trees. There were no rocky outcrops or exposures of shales, and NO FOSSILS were seen.



Figure 4: General view of the area covered with soils and vegetation.



Figure 5: Another general view with small shrubs growing in the grassland



Figure 6: Gravel road near the eastern side of the project area where roads and paths expose the underlying soils and gravel.



Figure 7: View along the western margin. This is the section underlain by the potentially fossiliferous Vryheid Formation. No rocks, shales or fossils are visible on the surface.

## 4. Impact assessment

An assessment of the potential impacts to possible palaeontological resources considers the criteria encapsulated in Table 3:

**TABLE 3A: CRITERIA FOR ASSESSING IMPACTS**

<b>PART A: DEFINITION AND CRITERIA</b>		
<b>Criteria for ranking of the SEVERITY/NATURE of environmental impacts</b>	<b>H</b>	Substantial deterioration (death, illness or injury). Recommended level will often be violated. Vigorous community action.
	<b>M</b>	Moderate/ measurable deterioration (discomfort). Recommended level will occasionally be violated. Widespread complaints.
	<b>L</b>	Minor deterioration (nuisance or minor deterioration). Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	<b>L+</b>	Minor improvement. Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.
	<b>M+</b>	Moderate improvement. Will be within or better than the recommended level. No observed reaction.
	<b>H+</b>	Substantial improvement. Will be within or better than the recommended level. Favourable publicity.
<b>Criteria for ranking the DURATION of impacts</b>	<b>L</b>	Quickly reversible. Less than the project life. Short term
	<b>M</b>	Reversible over time. Life of the project. Medium term
	<b>H</b>	Permanent. Beyond closure. Long term.
<b>Criteria for ranking the SPATIAL SCALE of impacts</b>	<b>L</b>	Localised - Within the site boundary.
	<b>M</b>	Fairly widespread – Beyond the site boundary. Local
	<b>H</b>	Widespread – Far beyond site boundary. Regional/ national

PROBABILITY (of exposure to impacts)	H	Definite/ Continuous
	M	Possible/ frequent
	L	Unlikely/ seldom

**TABLE 3B: IMPACT ASSESSMENT**

PART B: ASSESSMENT		
SEVERITY/NATURE	H	-
	M	-
	L	Soils and sands do not preserve plant fossils; so far there are no records from the Dwyka Group or Vryheid formation of plant or animal fossils in this region so it is very unlikely that fossils occur on the site. The impact would be very unlikely.
	L+	-
	M+	-
	H+	-
DURATION	L	-
	M	-
	H	Where manifest, the impact will be permanent.
SPATIAL SCALE	L	Since only the possible fossils within the area would be fossil plants from the <i>Glossopteris</i> flora in the shales, the spatial scale will be localised within the site boundary.
	M	-
	H	-
PROBABILITY	H	-
	M	-
	L	It is extremely unlikely that any fossils would be found in the loose soils that cover the area. It is unknown if there are fossils below ground, therefore, a Fossil Chance Find Protocol should be added to the eventual EMPr.

Based on the nature of the project, surface activities may impact upon the fossil heritage if preserved in the development footprint. The geological structures suggest that the rocks are either much too old to contain fossils or in the footprint they are the correct age and type to preserve fossils. **The site visit confirmed that there are NO fossils on the surface.** Since there is a small chance that fossils may occur below ground the Vryheid Formation, and may be disturbed a Fossil Chance Find Protocol has been added to this report. Taking account of the defined criteria, the potential impact to fossil heritage resources is low.

## 5. Assumptions and uncertainties

Based on the geology of the area and the palaeontological record as we know it, it can be assumed that the formation and layout of the dolomites, sandstones, shales and sands are typical for the country and could contain fossil plant, insect, invertebrate and vertebrate material. The sands of the Quaternary period would not preserve fossils. No fossils were seen on the surface during the site visit. It is not known if there are fossiliferous shales of the Vryheid Formation below the ground.

## 6. Recommendation

Based on experience and the lack of any previously recorded fossils from the area, it is unlikely that any fossils would be preserved in the shales below the ground because such fossils are patchy in their distribution and unpredictable. There is a small chance that fossils may occur below ground in the shales of the early Permian Vryheid Formation so a Fossil Chance Find Protocol should be added to the EMPr. If fossils are found once excavations for foundations and amenities have commenced then they should be rescued and a palaeontologist called to assess them and collect a representative sample. Any fossils collected must be done with a SAHRA permit and the fossils housed in a recognised institution.

## 7. References

- Anderson, A.M., McLachlan, I.R., 1976. The plant record in the Dwyka and Ecca Series (Permian) of the south-western half of the great Karoo Basin, South Africa. *Palaeontologia africana* 19, 31-42.
- Anderson, J.M., Anderson, H.M., 1985. *Palaeoflora of Southern Africa: Prodrum of South African megaflores, Devonian to Lower Cretaceous*. A.A. Balkema, Rotterdam. 423 pp.
- Isbell, J.L., Henry, L.C., Gulbranson, E.L., Limarino, C.O., Fraiser, F.L., Koch, Z.J., Ciccioli, P.I., Dineen, A.A., 2012. Glacial paradoxes during the late Paleozoic ice age: Evaluating the equilibrium line altitude as a control on glaciation. *Gondwana Research* 22, 1-19.
- Johnson, M.R., van Vuuren, C.J., Visser, J.N.J., Cole, D.I., Wickens, H.deV., Christie, A.D.M., Roberts, D.L., Brandl, G., 2006. Sedimentary rocks of the Karoo Supergroup. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. Pp 461 – 499.
- McCarthy, T.S., 2006. The Witwatersrand Supergroup. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. Pp 155-186.
- Plumstead, E.P., 1969. Three thousand million years of plant life in Africa. Geological Society of southern Africa, Annexure to Volume LXXII. 72pp + 25 plates.
- Robb, L.J., Brandl, G., Anhaeusser, C.R., Poujol, M., 2006. Archaean Granitoid Intrusions. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. Pp 57-94.
- Van der Westhuizen, W.A., de Bruijn, H., Meintjes, P.G., 2006. The Ventersdorp Supergroup. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). *The Geology of South Africa*. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. pp 187-208.
- Visser, J.N.J., 1986. Lateral lithofacies relationships in the glaciogene Dwyka Formation in the western and central parts of the Karoo Basin. *Transactions of the Geological Society of South Africa* 89, 373-383.
- Visser, J.N.J., 1989. The Permo-Carboniferous Dwyka Formation of southern Africa: deposition by a predominantly subpolar marine icesheet. *Palaeogeography, Palaeoclimatology, Palaeoecology* 70, 377-391.



## 8. Chance Find Protocol

**Monitoring Programme for Palaeontology – to commence once the excavations / drilling activities begin.**

1. The following procedure is only required if fossils are seen on the surface and when excavations commence.
2. When excavations begin the rocks and must be given a cursory inspection by the environmental officer or designated person. Any fossiliferous material (plants, insects, bone, coal) should be put aside in a suitably protected place. This way the excavation activities will not be interrupted.
3. Photographs of similar fossil plants must be provided to the developer to assist in recognizing the fossil plants in the shales and mudstones (for example see Figure 8). This information will be built into the EMP's training and awareness plan and procedures.
4. Photographs of the putative fossils can be sent to the palaeontologist for a preliminary assessment.
5. If there is any possible fossil material found by the developer/environmental officer/miners then the qualified palaeontologist sub-contracted for this project, should visit the site to inspect the selected material and check the dumps where feasible.
6. Fossil plants or vertebrates that are considered to be of good quality or scientific interest by the palaeontologist must be removed, catalogued and housed in a suitable institution where they can be made available for further study. Before the fossils are removed from the site a SAHRA permit must be obtained. Annual reports must be submitted to SAHRA as required by the relevant permits.
7. If no good fossil material is recovered then no site inspections by the palaeontologist will be necessary. A final report by the palaeontologist must be sent to SAHRA once the project has been completed and only if there are fossils.
8. If no fossils are found and the excavations have finished then no further monitoring is required.

# Appendix A – Examples of fossils plants from the Vryheid Formation



Figure 8: Examples of fossil plants as impressions in the rocks, from the Vryheid Formation.

## Appendix B – Details of specialist

### Curriculum vitae (short) - Marion Bamford PhD January 2021

#### i) Personal details

Surname : **Bamford**  
First names : **Marion Kathleen**  
Present employment : Professor; Director of the Evolutionary Studies Institute.  
Member Management Committee of the NRF/DST Centre of Excellence Palaeosciences, University of the Witwatersrand, Johannesburg, South Africa-  
Telephone : +27 11 717 6690  
Fax : +27 11 717 6694  
Cell : 082 555 6937  
E-mail : [marion.bamford@wits.ac.za](mailto:marion.bamford@wits.ac.za) ; [marionbamford12@gmail.com](mailto:marionbamford12@gmail.com)

#### ii) Academic qualifications

Tertiary Education: All at the University of the Witwatersrand:

1980-1982: BSc, majors in Botany and Microbiology. Graduated April 1983.

1983: BSc Honours, Botany and Palaeobotany. Graduated April 1984.

1984-1986: MSc in Palaeobotany. Graduated with Distinction, November 1986.

1986-1989: PhD in Palaeobotany. Graduated in June 1990.

#### iii) Professional qualifications

*Wood Anatomy Training (overseas as nothing was available in South Africa):*

1994 - Service d'Anatomie des Bois, Musée Royal de l'Afrique Centrale, Tervuren, Belgium, by Roger Dechamps

1997 - Université Pierre et Marie Curie, Paris, France, by Dr Jean-Claude Koeniguer

1997 - Université Claude Bernard, Lyon, France by Prof Georges Barale, Dr Jean-Pierre Gros, and Dr Marc Philippe

#### iv) Membership of professional bodies/associations

Palaeontological Society of Southern Africa

Royal Society of Southern Africa - Fellow: 2006 onwards

Academy of Sciences of South Africa - Member: Oct 2014 onwards

International Association of Wood Anatomists - First enrolled: January 1991

International Organization of Palaeobotany – 1993+

Botanical Society of South Africa

South African Committee on Stratigraphy – Biostratigraphy - 1997 - 2016  
 SASQUA (South African Society for Quaternary Research) – 1997+  
 PAGES - 2008 –onwards: South African representative  
 ROCEEH / WAVE – 2008+  
 INQUA – PALCOMM – 2011+onwards

### **vii) Supervision of Higher Degrees**

All at Wits University

Degree	Graduated/completed	Current
Honours	11	2
Masters	10	5
PhD	11	4
Postdoctoral fellows	10	4

### **viii) Undergraduate teaching**

Geology II – Palaeobotany GEOL2008 – average 65 students per year  
 Biology III – Palaeobotany APES3029 – average 25 students per year  
 Honours – Evolution of Terrestrial Ecosystems; African Plio-Pleistocene Palaeoecology;  
 Micropalaeontology – average 2-8 students per year.

### **ix) Editing and reviewing**

Editor: *Palaeontologia africana*: 2003 to 2013; 2014 – Assistant editor  
 Guest Editor: *Quaternary International*: 2005 volume  
 Member of Board of Review: *Review of Palaeobotany and Palynology*: 2010 –  
*Cretaceous Research*: 2014 –  
*Journal of African Earth Sciences*: 2020 –

Review of manuscripts for ISI-listed journals: 25 local and international journals

## **x) Palaeontological Impact Assessments**

Selected – list not complete:

- Thukela Biosphere Conservancy 1996; 2002 for DWAF
- Vioolsdrift 2007 for Xibula Exploration
- Rietfontein 2009 for Zitholele Consulting
- Bloeddrift-Baken 2010 for TransHex
- New Kleinfontein Gold Mine 2012 for Prime Resources (Pty) Ltd.
- Thabazimbi Iron Cave 2012 for Professional Grave Solutions (Pty) Ltd
- Delmas 2013 for Jones and Wagener
- Klipfontein 2013 for Jones and Wagener
- Platinum mine 2013 for Lonmin
- Syferfontein 2014 for Digby Wells
- Canyon Springs 2014 for Prime Resources
- Kimberley Eskom 2014 for Landscape Dynamics

- Yzermyne 2014 for Digby Wells
- Matimba 2015 for Royal HaskoningDV
- Commissiekraal 2015 for SLR
- Harmony PV 2015 for Savannah Environmental
- Glencore-Tweefontein 2015 for Digby Wells
- Umkomazi 2015 for JLB Consulting
- Ixia coal 2016 for Digby Wells
- Lambda Eskom for Digby Wells
- Alexander Scoping for SLR
- Perseus-Kronos-Aries Eskom 2016 for NGT
- Mala Mala 2017 for Henwood
- Modimolle 2017 for Green Vision
- Klipoortjie and Finaalspan 2017 for Delta BEC
- Ledjadja borrow pits 2018 for Digby Wells
- Lungile poultry farm 2018 for CTS
- Olienhout Dam 2018 for JP Celliers
- Isondlo and Kwasobabili 2018 for GCS
- Kanakies Gypsum 2018 for Cabanga
- Nababeep Copper mine 2018
- Glencore-Mbali pipeline 2018 for Digby Wells
- Remhoogte PR 2019 for A&HAS
- Bospoort Agriculture 2019 for Kudzala
- Overlooked Quarry 2019 for Cabanga
- Richards Bay Powerline 2019 for NGT
- Eilandia dam 2019 for ACO
- Eastlands Residential 2019 for HCAC
- Fairview MR 2019 for Cabanga
- Graspan project 2019 for HCAC
- Lieliefontein N&D 2019 for Enviropro
- Skeerpoort Farm Mast 2020 for HCAC
- Vulindlela Eco village 2020 for 1World
- KwaZamakhule Township 2020 for Kudzala
- Sunset Copper 2020 for Digby Wells
- McCarthy-Salene 2020 for Prescali
- VLNR Lodge 2020 for HCAC
- Madadeni mixed use 2020 for Enviropro

## **xi) Research Output**

Publications by M K Bamford up to December 2019 peer-reviewed journals or scholarly books: over 150 articles published; 5 submitted/in press; 8 book chapters.

Scopus h index = 29; Google scholar h index = 36;

Conferences: numerous presentations at local and international conferences.

**xii) NRF Rating**

NRF Rating: B-2 (2016-2020)

NRF Rating: B-3 (2010-2015)

NRF Rating: B-3 (2005-2009)

NRF Rating: C-2 (1999-2004)