Palaeontological Impact Assessment for the proposed Mopane Solar Energy Facilities near Carltonville, Gauteng and North West Provinces

Mopane Solar PV 5

Site Visit Report (Phase 2)

For

Beyond Heritage

08 January 2023

Prof Marion Bamford Palaeobotanist P Bag 652, WITS 2050 Johannesburg, South Africa Marion.bamford@wits.ac.za

Expertise of Specialist

The Palaeontologist Consultant: Prof Marion Bamford Qualifications: PhD (Wits Univ, 1990); FRSSAf, ASSAf Experience: 34 years research; 26 years PIA studies

Declaration of Independence

This report has been compiled by Professor Marion Bamford, of the University of the Witwatersrand, sub-contracted by Beyond Heritage, Modimolle, South Africa. The views expressed in this report are entirely those of the author and no other interest was displayed during the decision making process for the Project.

Specialist: Prof Marion Bamford

Millamfus

Signature:

Executive Summary

A Palaeontological Impact Assessment was requested for the proposed development of the Mopane Solar PV 5, a part of the cluster of Energy Facilities (SEFs), west northwest of Carltonville, along the Gauteng - North West Province border. Mopane Solar PV 5 will be located on Portion 2 of Rooidraai 85 IQ, JB Marks Local Municipality, North West Province.

To comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a site visit (Phase 2) Palaeontological Impact Assessment (PIA) was completed for the proposed development.

The proposed site lies on the potentially fossiliferous Malmani Subgroup (Transvaal Supergroup) that could preserve trace fossils such as stromatolites or microbialites in the dolomites. The site visit and walk through in December 2022 (summer) by the palaeontologist confirmed that dolomites were very uncommon in the area for the proposed solar collectors and infrastructure. NO FOSSILS of any kind were seen in this area although stromatolites have been reported from other outcrops of this lithotype. Nonetheless, a Fossil Chance Find Protocol should be added to the EMPr. Based on this information it is recommended that no further palaeontological impact assessment is required unless fossils are found by the contractor, developer, environmental officer or other designated responsible person once excavations for pole foundations for solar collectors and infrastructure have commenced. Since the impact will be low, as far as the palaeontology is concerned, the project should be authorised.

Table of Contents

Exper	tise of Specialist	2
De	eclaration of Independence	2
1.	Background	5
2.	Methods and Terms of Reference	9
3.	Geology and Palaeontology	9
i.	Project location and geological context	9
ii.	Palaeontological context	.11
iii.	Site visit observations	12
4.	Impact assessment	.14
5.	Assumptions and uncertainties	.16
6.	Recommendation	.16
7.	References	.16
8.	Chance Find Protocol	.17
9.	Appendix A – Examples of fossils	.18
10.	Appendix B – Details of specialist	.19
Figure	e 1: Google Earth map of the project area	8
Figure	es 2: Google Earth map of the project footprint	8
Figure	e 3: Geological map of the area around the project site	9
Figure	e 4: SAHRIS palaeosensitivity map for the site for the project	.11
Figure	e 5-6: Site visit photographs 13-	14

1. Background

In view of the growing electricity demand and in an effort to use renewable energy resources, Ursa Energy (Pty) Ltd, Fornax Energy (Pty) Ltd and Volans Energy (Pty) Ltd is assessing the feasibility of energy generation facilities, consisting of the construction, operation and maintenance of Photovoltaic (PV) Power Plants with a maximum generation capacity up to 100 MW, each, at the point of connection.

The Mopane Solar PV 5 project site is on Portion 2 of the farm Rooidraai 85 IQ, located within the JB Marks Local Municipality, Dr Kenneth Kaunda District Municipality, North West Province (Figures 1-2).

The project site is located ±7 km northwest of Welverdiend along the border between Gauteng and the North West Province. The Eskom Carmel Main Transmission Substation (MTS) is located 16.4 km southeast of project sites.

The development is located 5.5km north of R501 with access from both the D859 (Preferred) and R501 (alternative).

The developed area (footprint) required for the proposed project will be up to 366 hectares,. The final size and location of the project footprint will be assessed following the outcomes of the Public Participation Process and of the recommendations and conclusions of the Specialist Studies to be conducted during the Environmental Impact Assessment (EIA) process.

The proposed development (the Photovoltaic (PV) Power Plants and connection infrastructure) consists of the installation of the following equipment:

- Photovoltaic modules (mono-crystalline, poly-crystalline, or bi-facial modules)
- Mounting systems for the PV arrays (single-axis horizontal trackers or fixed structures) and related foundations
- Internal cabling and string boxes
- DC/AC inverters
- Medium voltage stations, hosting LV/MV power transformers
- Medium voltage receiving station(s)
- Workshops & warehouses
- One on-site high-voltage substation and one high-voltage busbar with metering and protection devices
- One on-site high-voltage substation with high-voltage power transformers, stepping up the voltage to 400kV/132kV and one high-voltage busbar with metering and protection devices
- One on-site switching station, with one high-voltage busbar with metering and protection devices
- Battery Energy Storage Systems (BESS), with a Maximum Export Capacity up to 100 MW and a 5-hour storage capacity up to 1250 MWh, with a footprint up to 10 ha, next to the on-site high-voltage substation, within the PV plant footprint / fenced areas
- Electrical system and UPS (Uninterruptible Power Supply) devices
- Lighting system

- Grounding system
- Internal roads
- Fencing of the site and alarm and video-surveillance system
- Water access point, water supply pipelines, water treatment facilities
- Small scale patented wastewater treatment system.

During the construction phase, the site may be provided with additional activities that will be removed at the end of construction:

- Water access point, water supply pipelines, water treatment facilities
- Prefabricated buildings
- Workshops and warehouses

The proposed SEF site and the grid connection are on very highly sensitive rocks of the Malmani Subgroup that might preserve trace fossils such as stromatolites, so a site visit is required by SAHRA as part of the EIA process (https://sahris.sahra.org/map.palaeo.

A Palaeontological Impact Assessment was requested for the Mopane Solar PV 5 project. To comply with the regulations of the South African Heritage Resources Agency (SAHRA) in terms of Section 38(8) of the National Heritage Resources Act, 1999 (Act No. 25 of 1999) (NHRA), a site visit and walkthrough (Phase 2) Palaeontological Impact Assessment (PIA) was completed for the proposed development and is reported herein.

Table 1: National Environmental Management Act, 1998 (Act No. 107 of 1998) (NEMA) and Environmental Impact Assessment (EIA) Regulations, 2014 (as amended) - Requirements for Specialist Reports (Appendix 6).

	A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:	Relevant section in report
ai	Details of the specialist who prepared the report,	Appendix B
aii	The expertise of that person to compile a specialist report including a curriculum vitae	Appendix B
b	A declaration that the person is independent in a form as may be specified by the competent authority	Page 2
С	An indication of the scope of, and the purpose for which, the report was prepared	Section 1
ci	An indication of the quality and age of the base data used for the specialist report: SAHRIS palaeosensitivity map accessed – date of this report	Yes
cii	A description of existing impacts on the site, cumulative impacts of the proposed development and levels of acceptable change	Section 5
d	The date and season of the site investigation and the relevance of the season to the outcome of the assessment	N/A
е	A description of the methodology adopted in preparing the report or carrying out the specialised process	Section 2

	A specialist report prepared in terms of the Environmental Impact Regulations of 2017 must contain:	Relevant section in report
f	The specific identified sensitivity of the site related to the activity and its associated structures and infrastructure	Section 4
g	An identification of any areas to be avoided, including buffers	N/A
h	A map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;	N/A
i	A description of any assumptions made and any uncertainties or gaps in knowledge;	Section 5
j	A description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives, on the environment	Section 4
k	Any mitigation measures for inclusion in the EMPr	Section 8, Appendix A
l	Any conditions for inclusion in the environmental authorisation	N/A
m	Any monitoring requirements for inclusion in the EMPr or environmental authorisation	Section 8, Appendix A
ni	A reasoned opinion as to whether the proposed activity or portions thereof should be authorised	Section 6
nii	If the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan	Sections 6, 8
0	A description of any consultation process that was undertaken during the course of carrying out the study	N/A
р	A summary and copies of any comments that were received during any consultation process	N/A
q	Any other information requested by the competent authority.	N/A
2	Where a government notice gazetted by the Minister provides for any protocol or minimum information requirement to be applied to a specialist report, the requirements as indicated in such notice will apply.	




Figure 1: Google Earth map of the proposed development showing the Mopane SEF cluster area (yellow), and the OHPL routes (turquoise and blue), and the relevant landmarks.




Figure 2: Google Earth map for the proposed Mopane PVs 4 & 5 (orange polygon) on Portion 2 of Farm Rooidraai 85 IQ.

2. Methods and Terms of Reference

The Terms of Reference (ToR) for this study were to undertake a PIA and provide feasible management measures to comply with the requirements of SAHRA.

The methods employed to address the ToR included:

- 1. Consultation of geological maps, literature, palaeontological databases, published and unpublished records to determine the likelihood of fossils occurring in the affected areas. Sources included records housed at the Evolutionary Studies Institute at the University of the Witwatersrand and SAHRA databases;
- 2. Where necessary, site visits by a qualified palaeontologist to locate any fossils and assess their importance, as is the case here;
- 3. Where appropriate, collection of unique or rare fossils with the necessary permits for storage and curation at an appropriate facility (*not applicable to this assessment*); and
- 4. Determination of fossils' representivity or scientific importance to decide if the fossils can be destroyed or a representative sample collected (*not applicable to this assessment*).

3. Geology and Palaeontology

i. Project location and geological context




Figure 3: Geological map of the area around the Farm Rooidraai for the Mopane Solar PV 5 (orange rectangle). Abbreviations of the rock types are explained in Table 2. Map enlarged from the Geological Survey 1: 250 000 map 2626 West Rand.

Table 2: Explanation of symbols for the geological map and approximate ages (Johnson et al., 2006; Partridge et al., 2006; Zeh et al., 2020). SG = Supergroup; Fm = Formation; Ma = million years; grey shading = formations impacted by the project.

Symbol	Group/Formation	Lithology	Approximate Age
Qs	Quaternary	Alluvium, sand, calcrete	Neogene, ca 2.5 Ma to present
Vdi	diabase	Diabase	Post-Transvaal SG
Vh	Hekpoort Fm, Pretoria Group, Transvaal SG	Andesite, agglomerate, tuff	
Vt	Timeball Hill Fm Pretoria Group, Transvaal SG	Quartzite	< 2420 Ma
Vmd	Malmani Subgroup, Chuniespoort Group, Transvaal SG	Dolomite, chert	Ca 2750 – 2650 Ma
Vbr	Black Reef Fm, Transvaal SG	Quartzite, conglomerate, shale, basalt	Ca 2650 – 2640 Ma
R-Vr	Rietgat Fm, Platberg Group, Ventersdorp SG	Amygdaloidal lava, agglomerate, tuff	Mesoarchaean Ca 2720 Ma
Rg	Government Subgroup, West Rand Group, Witwatersrand SG	Quartzite, shale, greywacke, conglomerate	Mesoarchaean Ca 2890 Ma

The project lies in the south western part of the Transvaal Basin where the lower rocks of the Transvaal Supergroup are exposed, in particular the dolomites of the Malmani Subgroup (Chuniespoort Group, Transvaal Supergroup; ca 2585-2480 Ma), (Figure 3).

The Late Archaean to early Proterozoic Transvaal Supergroup is preserved in three structural basins on the Kaapvaal Craton (Eriksson et al., 2006). In South Africa are the Transvaal and Griqualand West Basins, and the Kanye Basin is in southern Botswana. The Griqualand West Basin is divided into the Ghaap Plateau sub-basin and the Prieska sub-basin. Sediments in the lower parts of the basins are very similar but they differ somewhat higher up the sequences. Several tectonic events have greatly deformed the south western portion of the Griqualand West Basin between the two sub-basins

In the Transvaal Basin the Transvaal Supergroup is divided into two Groups, the lower Chuniespoort Group and the upper Pretoria Group (with ten formations; Eriksson et al., 2006). The Chuniespoort Group is divided into the basal Malmani Subgroup that comprises dolomites and limestones and is divided into five formations based on chert content, stromatolitic morphology, intercalated shales and erosion surfaces. The top of the Chuniespoort Group has the Penge Formation and the Duitschland Formation.

Making up the lower Pretoria Group are the Timeball Hill Formation and the Boshoek Formation. The Hekpoort, Dwaalheuwel, Strubenkop and Daspoort Formations form a sequence as the middle part of the Pretoria Group, Transvaal Supergroup, and represent rocks that are over 2060 million years old. The Hekpoort Formation is a massive lava deposit and is overlain by the rest of the Transvaal Supergroup.

The Transvaal sequence has been interpreted as three major cycles of basin infill and tectonic activity with the first deep basin sediments forming the Chuniespoort Group, the second cycle deposited the lower Pretoria Group, and the sediments in this area are from the interim lowstand that preceded the third cycle. These sediments were deposited in shallow lacustrine, alluvial fan and braided stream environments (Eriksson et al., 2012).

ii. Palaeontological context

The palaeontological sensitivity of the area under consideration is presented in Figure 4. The site for development is in the very highly sensitive Malmani Subgroup (Transvaal Supergroup).




Figure 4: SAHRIS palaeosensitivity map for the site for the proposed Mopane PV 5 on Portion 2 of Rooidraai 85 IQ (yellow polygon). Background colours indicate the following degrees of sensitivity: red = very highly sensitive; orange/yellow = high; green = moderate; blue = low; grey = insignificant/zero.

The Transvaal Supergroup rocks represent on a very large scale, a sequence of sediments filling the basins under conditions of lacustrine, fluvial, volcanic and glacial cycles in a tectonically active region. The predominantly carbonaceous sediments are evidence of the increase in the atmosphere of oxygen produced by algal colony photosynthesis, the so-called Great Oxygen Event (ca 2.40 – 2.32 Ga) and precursor to an environment where diverse life forms could evolve. The Neoarchean-Paleoproterozoic Transvaal Supergroup in South Africa contains the well-preserved stromatolitic Campbellrand -Malmani

carbonate platform (Griqualand West Basin – Transvaal Basin respectively), which was deposited in shallow seawater shortly before the Great Oxidation Event (GOE).

The Transvaal Supergroup comprises one of world's earliest carbonate platform successions (Beukes, 1987; Eriksson et al., 2006; Zeh et al., 2020). In some areas there are well preserved stromatolites that are evidence of the photosynthetic activity of blue green bacteria and green algae. These microbes formed colonies in warm, shallow seas and deposited layer upon layer of minerals, often in domes or columns. The minerals are predominantly calcium carbonate, calcium sulphate, magnesium carbonate and magnesium sulphate. Only very rarely are the bacteria and algae preserved but the stromatolites are traces of their activity, hence called trace fossils. These fossils are protected by legislation, therefore the Malmani Subgroup palaeosensitivity is very high (red; Figure 4, SAHRIS).

iii. Site visit observations

The site was walked through for both proposed Solar PV 5 solar collectors and infrastructure. The land had been cleared of vegetation in the past for grazing or agriculture and so there were very few bushes, mostly low secondary grassland remains. There were very few rocky outcrops and these were examined for stromatolites. Most were quartzite, sandstone or chert and not dolomite. The rare dolomite outcrops did not have any stromatolites.

Rocks were also rare in the Mopane Solar PV 5 footprint and only quartzite and sandstone were see (Figures 5-6). No dolomite and no stromatolites were seen.



Figure 5: Photographs of site visit for the Mopane Solar PV 5 project. A-B – general views showing fairly flat topography, low vegetation. C - rare outcrop of chert and quartzite. D – farm track exposes the red sandy soil, and low vegetation.



Figure 6: Photographs of site visit for the Mopane Solar PV 5 project. A – Farm road along flat topography towards a powerline. B – bushes associated with an outcrop of chert. No dolomites and no stromatolites.

4. Impact assessment

An assessment of the potential impacts to possible palaeontological resources considers the criteria encapsulated in Table :

Table 4a: Criteria for assessing impacts

PART A: DEFINITION AND CRITERIA			
	Н	Substantial deterioration (death, illness or injury). Recommended level will often be violated. Vigorous community action.	
Criteria for ranking of the	M	Moderate/ measurable deterioration (discomfort). Recommended level will occasionally be violated. Widespread complaints.	
SEVERITY/NATURE of environmental impacts	L	Minor deterioration (nuisance or minor deterioration). Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.	
	L+	Minor improvement. Change not measurable/ will remain in the current range. Recommended level will never be violated. Sporadic complaints.	

	M+	Moderate improvement. Will be within or better than the recommended level. No observed reaction.	
	Н+	Substantial improvement. Will be within or better than the recommended level. Favourable publicity.	
Criteria for ranking	L	Quickly reversible. Less than the project life. Short term	
the DURATION of	M	Reversible over time. Life of the project. Medium term	
impacts	Н	Permanent. Beyond closure. Long term.	
Criteria for ranking	L	Localised - Within the site boundary.	
the SPATIAL SCALE	M	Fairly widespread – Beyond the site boundary. Local	
of impacts	Н	Widespread - Far beyond site boundary. Regional/ national	
PROBABILITY	Н	Definite/ Continuous	
(of exposure to	M	Possible/ frequent	
impacts)	L	Unlikely/ seldom	

Table 4b: Impact Assessment

PART B: Assessment			
	Н	-	
	M	-	
SEVERITY/NATURE	L	Soils do not preserve plant fossils; so far there are no records from the Malmani Subgroup of trace fossils of straomatolites in this region so it is very unlikely that fossils occur on the site. The impact would be very unlikely.	
	L+	-	
	M+	-	
	H+	-	
	L	-	
DURATION	M	-	
	Н	Where manifest, the impact will be permanent.	
SPATIAL SCALE	L	Since the only possible fossils within the area would be trace fossils such as stromatolites in the dolomites, the spatial scale will be localised within the site boundary.	
	M	-	
	Н	-	
	Н	-	
	M	-	
PROBABILITY	L	It is extremely unlikely that any fossils would be found in the loose sand that will be excavated but there might be stromatolites in the dolomites. Therefore, a Fossil Chance Find Protocol should be added to the eventual EMPr.	

Based on the nature of the project, surface activities may impact upon the fossil heritage if preserved in the development footprint. The geological structures suggest that the rocks are the correct age and type to preserve fossils. The site visit and walk through

confirmed that there were NO FOSSILS in the project footprint. There were very few dolomite exposures. Since there is an extremely small chance that trace fossils from the Malmani Subgroup may be disturbed a Fossil Chance Find Protocol has been added to this report. Taking account of the defined criteria, the potential impact to fossil heritage resources is extremely low.

5. Assumptions and uncertainties

Based on the geology of the area and the palaeontological record as we know it, it can be assumed that the formation and layout of the dolomites, sandstones, shales and sands are typical for the country and only some do contain trace fossils, fossil plant, insect, invertebrate and vertebrate material. The site visit and walk through in mid December 2022 by the palaeontologist confirmed that there are NO FOSSILS in the proposed area for the solar collectors or in the adjacent grid infrastructure sites. The overlying sands and soils of the Quaternary period would not preserve fossils.

6. Recommendation

Based on the fossil record but confirmed by the site visit and walk through there are NO FOSSILS of the project footprint. Rocky outcrops were rare and mostly not dolomitic in composition. This apparent lack of any kind of dolomite is probably the reason why the area has been classified as Malmani Subgroup, only, without the more detailed differences for the formations, such as, stromatolites, oolitic and algal dolomites that are used elsewhere to distinguish the formations. It is extremely unlikely that any fossils would be preserved in the overlying soils and sands of the Quaternary.

There is a very small chance that trace fossils may occur in below the ground surface in the dolomites of the Malmani Subgroup so a Fossil Chance Find Protocol should be added to the EMPr. If fossils are found by the environmental officer or other responsible person once excavations and drilling have commenced, then they should be rescued and a palaeontologist called to assess and collect a representative sample.

Since there is an extremely small chance of fossils being impacted by this project, as far as the palaeontology is concerned, the project should be authorised

7. References

Beukes, N.J., 1987. Facies relations, depositional environments and diagenesis in a major early Proterozoic stromatolitic carbonate platform to basinal sequence, Campbellrand Subgroup, Transvaal Supergroup, southern Africa. Sedimentary Geology 54, 1-46.

Eriksson, P.G., Altermann, W., Hartzer, F.J., 2006. The Transvaal Supergroup and its precursors. In: Johnson, M.R., Anhaeusser, C.R. and Thomas, R.J., (Eds). The Geology of South Africa. Geological Society of South Africa, Johannesburg / Council for Geoscience, Pretoria. pp 237-260.

Plumstead, E.P., 1969. Three thousand million years of plant life in Africa. Geological Society of southern Africa, Annexure to Volume LXXII. 72pp + 25 plates.

Zeh, A., Wilson, A.H., Gerdes, A., 2020. Zircon U-Pb-Hf isotope systematics of Transvaal Supergroup – Constraints for the geodynamic evolution of the Kaapvaal Craton and its hinterland between 2.65 and 2.06 Ga. Precambrian Research 345, 105760. https://doi.org/10.1016/j.precamres.2020.105760

8. Chance Find Protocol

Monitoring Programme for Palaeontology – to commence once the excavations / drilling activities begin.

- 1. The following procedure is only required if fossils are seen on the surface and when drilling/excavations commence.
- 2. When excavations begin the rocks and discard must be given a cursory inspection by the environmental officer or designated person. Any fossiliferous material (trace fossils, fossils of plants, insects, bone or coalified material) should be put aside in a suitably protected place. This way the project activities will not be interrupted.
- 3. Photographs of similar fossils must be provided to the developer to assist in recognizing the fossil plants, vertebrates, invertebrates or trace fossils in the shales and mudstones (for example see Figures 9-10). This information will be built into the EMP's training and awareness plan and procedures.
- 4. Photographs of the putative fossils can be sent to the palaeontologist for a preliminary assessment.
- 5. If there is any possible fossil material found by the developer/environmental officer then the qualified palaeontologist sub-contracted for this project, should visit the site to inspect the selected material and check the dumps where feasible.
- 6. Trace fossils, fossil plants or vertebrates that are considered to be of good quality or scientific interest by the palaeontologist must be removed, catalogued and housed in a suitable institution where they can be made available for further study. Before the fossils are removed from the site a SAHRA permit must be obtained. Annual reports must be submitted to SAHRA as required by the relevant permits.
- 7. If no good fossil material is recovered then no site inspections by the palaeontologist will be necessary. A final report by the palaeontologist must be sent to SAHRA once the project has been completed and only if there are fossils.
- 8. If no fossils are found and the excavations have finished then no further monitoring is required.

Appendix A – Examples of fossils from the Malmani Subgroup



Figure 9: Photographs of dolomite and stromatolites as seen in the field.



Figure 10: Field photographs of more types of trace fossils in dolomite.

9. Appendix B – Details of specialists

Curriculum vitae (short) - Marion Bamford PhD January 2023

Present employment: Professor; Director of the Evolutionary Studies Institute.

Member Management Committee of the NRF/DSI Centre of Excellence Palaeosciences, University of the Witwatersrand,

Johannesburg, South Africa

Telephone : +27 11 717 6690 Cell : 082 555 6937

E-mail : <u>marion.bamford@wits.ac.za</u>;

marionbamford12@gmail.com

ii) Academic qualifications

Tertiary Education: All at the University of the Witwatersrand:

1980-1982: BSc, majors in Botany and Microbiology. Graduated April 1983.

1983: BSc Honours, Botany and Palaeobotany. Graduated April 1984.

1984-1986: MSc in Palaeobotany. Graduated with Distinction, November 1986.

1986-1989: PhD in Palaeobotany. Graduated in June 1990.

iii) Professional qualifications

Wood Anatomy Training (overseas as nothing was available in South Africa):

1994 - Service d'Anatomie des Bois, Musée Royal de l'Afrique Centrale, Tervuren, Belgium, by Roger Dechamps

1997 - Université Pierre et Marie Curie, Paris, France, by Dr Jean-Claude Koeniguer

1997 - Université Claude Bernard, Lyon, France by Prof Georges Barale, Dr Jean-Pierre

Gros, and Dr Marc Philippe

iv) Membership of professional bodies/associations

Palaeontological Society of Southern Africa

Royal Society of Southern Africa - Fellow: 2006 onwards

Academy of Sciences of South Africa - Member: Oct 2014 onwards

International Association of Wood Anatomists - First enrolled: January 1991

International Organization of Palaeobotany - 1993+

Botanical Society of South Africa

South African Committee on Stratigraphy - Biostratigraphy - 1997 - 2016

SASQUA (South African Society for Quaternary Research) - 1997+

PAGES - 2008 - onwards: South African representative

ROCEEH / WAVE - 2008+

INQUA - PALCOMM - 2011+onwards

v) Supervision of Higher Degrees

All at Wits University

Degree	Graduated/completed	Current
Honours	13	0
Masters	13	3
PhD	13	7
Postdoctoral fellows	14	4

vi) Undergraduate teaching

Geology II - Palaeobotany GEOL2008 - average 65 students per year

Biology III - Palaeobotany APES3029 - average 25 students per year

Honours - Evolution of Terrestrial Ecosystems; African Plio-Pleistocene Palaeoecology;

Micropalaeontology – average 12 - 20 students per year.

vii) Editing and reviewing

Editor: Palaeontologia africana: 2003 to 2013; 2014 - Assistant editor

Guest Editor: Quaternary International: 2005 volume

Member of Board of Review: Review of Palaeobotany and Palynology: 2010 -

Associate Editor: Cretaceous Research: 2018-2020

Associate Editor: Royal Society Open: 2021 -

Review of manuscripts for ISI-listed journals: 30 local and international journals

viii) Palaeontological Impact Assessments

25 years' experience in PIA site and desktop projects

- Selected from recent projects only list not complete:
- Skeerpoort Farm Mast 2020 for HCAC
- Vulindlela Eco village 2020 for 1World
- KwaZamakhule Township 2020 for Kudzala
- Sunset Copper 2020 for Digby Wells
- McCarthy-Salene 2020 for Prescali
- VLNR Lodge 2020 for HCAC
- Madadeni mixed use 2020 for EnviroPro
- Frankfort-Windfield Eskom Powerline 2020 for 1World
- Beaufort West PV Facility 2021 for ACO Associates
- Copper Sunset MR 2021 for Digby Wells
- Sannaspos PV facility 2021 for CTS Heritage
- Smithfield-Rouxville-Zastron PL 2021 for TheroServe
- Glosam Mine 2022 for AHSA
- Wolf-Skilpad-Grassridge OHPL 2022 for Zutari
- Iziduli and Msenge WEFs 2022 for CTS Heritage
- Hendrina North and South WEFs & SEFs 2022 for Cabanga
- Dealesville-Springhaas SEFs 2022 for GIBB Environmental
- Vhuvhili and Mukondeleli SEFs 2022 for CSIR
- Chemwes & Stilfontein SEFs 2022 for CTS Heritage
- Equestria Exts housing 2022 for Beyond Heritage
- Zeerust Salene boreholes 2022 for Prescali
- Tsakane Sewer upgrade 2022 for Tsimba
- Transnet MPP inland and coastal 2022 for ENVASS
- Ruighoek PRA 2022 for SLR Consulting (Africa)
- Namli MRA Steinkopf 2022 for Beyond Heritage

ix) Research Output

Publications by M K Bamford up to January 2022 peer-reviewed journals or scholarly books: over 170 articles published; 5 submitted/in press; 14 book chapters. Scopus h-index = 30; Google Scholar h-index = 39; -i10-index = 116 based on 6568 citations.

Conferences: numerous presentations at local and international conferences.