PALAEONTOLOGICAL IMPACT SCREENING REPORT

PROPOSED CLOCOLAN SOLID WASTE DISPOSAL FACILITY

On the Farm: Ferndale 349 of the Setsoto Municipality within the Thabo Mofutsanyane District Municipality in the Free State Province of South Africa

Developer: Setsoto Municipality



22 March 2012



Metsi Metseng Geological & Environmental Services

Suite 91 Private Bag X62 Bethlehem 9700 info@mmges.co.za Fax 086 743 6864 +27 82 070 0735 +27 82 829 4978 www.mmges.co.za VAT 4260183498 Reg 1999/30444/23

EXECUTIVE SUMMARY

The development of a Solid Waste Disposal Facility near Clocolan in the Eastern Free State is an initiative of Setsoto Municipality. The purpose of this Palaeontological Impact Assessment is to identify exposed and potential palaeontological heritage on the site of the proposed development, to assess the impact the development may have on this resource, and to make recommendations as to how this impact might be mitigated.

The Setsoto Municipality plans to develop a solid waste disposal facility located to the west of Clocolan and Hlohlowane on the farm Ferndale 349. The area is next to the R703 tar road leading from Clocolan to Excelsior. The facility footprint is approximately 3 ha.

A basic assessment of the topography and geology of the area was made by using appropriate geological (1:250 000) maps in conjunction with Google Earth. A review of the literature on the geological formations exposed at surface in the development site and the fossils that have been associated with these geological strata was undertaken.

The proposed Clocolan solid waste disposal site is underlain by the Triassic Molteno Formation that consists of coarse-grained grey sandstone and dark grey mudstone. Soils are derived from the underlying rock and are generally deep and relatively high in fertility.

The Molteno Formation has a high palaeontological sensitivity rating. Through adequate monitoring and mitigation measures during excavations within the bedrock, the impact severity can be lowered to beneficial. The exposure and subsequent reporting of fossils (that would otherwise have remained undiscovered) will be a beneficial palaeontological impact.

It is recommended that the resident ECO be trained by a professional palaeontologist in the recognition of fossil material. If fossil material is later discovered it must be appropriately protected and the discovery reported to a palaeontologist for the removal thereof.

That all earth-moving activities within the bedrock with a potential impact on the Adelaide Subgroup be monitored by a palaeontologist. That a monitoring report be submitted to SAHRA after the completion of the earth works phase.

	SIGNIFICANCE RATING						
	Tomporal		Degree of	Impact Severity		Overall Significance	
Rock Unit	Temporal Scale	Spatial Scale	Confidence	With mitigation	Without mitigation	With mitigation	Without mitigation
Molteno Formation	permanent	international	unsure	beneficial	severe	beneficial	Negative

TABLE OF CONTENT

1.	INTR	ODUCTION	1
1.	1.	Legal Requirements	1
2.	AIMS	S AND METHODS	1
3.	PRO	POSED DEVELOPMENT DESCRIPTION	1
4.	GEO	LOGY OF THE AREA	2
4.	1.	The Molteno Formation	2
5.	PALA	AEONTOLOGY OF THE AREA	3
5.	1.	The Molteno Formation	3
6.	PALA	AEONTOLOGICAL SIGNIFICANCE AND RATING	4
7.	PALA	AEONTOLOGICAL IMPACT AND MITIGATION	4
8.	CON	CLUSION	5
9.	REFE	ERENCES	6
10.	Q	UALIFICATIONS AND EXPERIENCE OF THE AUTHOR	7
11.	AI	PPENDIX A - METHODOLOGY FOR ASSESSING THE SIGNIFICANCE OF IMPACTS	8

LIST OF FIGURES

Figure 3.1	Locality Map	.2
Figure 4.1	The Geology (Geo Map 2826- Windburg) of the Clocolan Waste Site Development	.3
Figure 7.1	Palaeontological Impact of the Proposed Clocolan Waste Disposal Facility	.5

LIST OF TABLES

Table 6.1	Palaeontological Significance of Geological Units on Site	4
Table 6.3	Significance Rating Table as Per CES Template	4
Table 7.1	Site Specific Mitigation Measures	5

1. INTRODUCTION

The development of a Solid Waste Disposal Facility near Clocolan in the Eastern Free State is an initiative of Setsoto Municipality. The purpose of this Palaeontological Impact Assessment is to identify exposed and potential palaeontological heritage on the site of the proposed development, to assess the impact the development may have on this resource, and to make recommendations as to how this impact might be mitigated.

1.1. Legal Requirements

This report forms part of the Scoping and Environmental Impact Assessment for the CLOCOLAN WASTE DISPOSAL SITE and complies with the requirements for the South African National Heritage Resource Act No 25 of 1999. In accordance with Section 38 (Heritage Resources Management), a Palaeontological Impact Assessment is required to assess any potential impacts to palaeontological heritage within the development footprint of the Clocolan Solid Waste Disposal site.

Categories of heritage resources recognised as part of the National Estate in Section 3 of the Heritage Resources Act, and which therefore fall under its protection, include:

- geological sites of scientific or cultural importance;
- objects recovered from the soil or waters of South Africa, including archaeological and palaeontological objects and material, meteorites and rare geological specimens; and
- objects with the potential to yield information that will contribute to an understanding of South Africa's natural or cultural heritage.

2. AIMS AND METHODS

After discussions with LHL Engineers a request for a Palaeontological Impact Screening was received. Following the *"SAHRA APM Guidelines: Minimum Standards for the Archaeological & Palaeontological Components of Impact Assessment Reports"* the aims of the Palaeontological Impact Screening were:

- identifying exposed and subsurface rock formations that are considered to be palaeontologically significant;
- assessing the level of palaeontological significance of these formations;
- commenting on the impact of the development on these exposed and/or potential fossil resources;
- making recommendations as to how the developer should conserve or mitigate damage to these resources.

A basic assessment of the topography and geology of the area was made by using appropriate geological (1:250 000) maps in conjunction with Google Earth. The only limitation on this methodology is the scale of mapping, which restricts comparison of the geology to the 1:250 000 scale. This restriction only applies in areas where major changes in the geological character of the area occur over very short distances or on the geological transformation zones.

A review of the literature on the geological formations exposed at surface in the development site and the fossils that have been associated with these geological strata was undertaken.

3. PROPOSED DEVELOPMENT DESCRIPTION

The Setsoto Municipality plans to develop a solid waste disposal facility located to the west of Clocolan and Hlohlowane on the farm Ferndale 349. The area is next to the R703 tar road leading

from Clocolan to Excelsior. The proposed facility general GPS co-ordinates are 28°54'30.15" S and 27°32'20.57" E (Figure 2.1). The facility footprint is approximately 3 ha.

The solid waste disposal facility will be constructed with accompanying infrastructure to supply in the demand for waste disposal for the next 10 years. These waste cells will be approximately 35m wide, 180m long and 2m deep.

Waste will be dumped in the cells and compacted until the waste reaches ground level when it will be covered. When the cells are full the actual land building will start. The waste will be dumped, compacted and covered in 2m layers until an eventual height of 13 m is reached.




Figure 3.1 Locality Map

4. GEOLOGY OF THE AREA

The geology around development area is underlain by the Molteno Formation (Trm) of the Karoo Supergroup as illustrated in Figure 4.1.

4.1. The Molteno Formation

The proposed development site is underlain by the Triassic Molteno Formation that consists of coarse-grained grey sandstone and dark grey mudstone. Soils are derived from the underlying rock and are generally deep and relatively high in fertility.




Figure 4.1 The Geology (Geo Map 2826- Windburg) of the Clocolan Waste Site Development

5. PALAEONTOLOGY OF THE AREA

5.1. The Molteno Formation

The Triassic Molteno Formation can have a moderate to high potential for plant fossils from the *Dicroidium* assemblage (Johnson et al, 2006). The *Dicroidium* assemblage is an extinct genus of fork-leaved seed ferns that were distributed over Gondwanaland during the Triassic Period.

Invertebrate fossils are restricted to trace fossils.

6. PALAEONTOLOGICAL SIGNIFICANCE AND RATING

The predicted palaeontological impact of the development is based on the initial mapping assessment and literature reviews as well as information gathered during the field investigation.

The palaeontological significance and rating is summarised in Table 7.1 and 7.2. For the methodology and definitions of impact rating and significance see Appendix A (CES 2011).

Geological Unit	Rock Type and Age	Fossil Heritage	Vertebrate Biozone	Palaeontological Sensitivity
Molteno Formation	Fluvial, braided river sandstone and mudstone TRIASSIC	Plant fossils such as <i>Dicroidium</i> assemblages Invertebrate fossils are restricted to trace fossils.	None	High sensitivity

 Table 6.1
 Palaeontological Significance of Geological Units on Site

Table 6.2	Significance Rating Table as Per CES Template
-----------	---

Rock Unit	Temporal Scale	Spatial Scale (area in which	Degree of confidence (confidence with which	(severity of ne or how bene	severity gative impacts, ficial positive would be)	Overall Sig (The combinat other criteria signific	ion of all the as an overall
NOCK ONIC	(duration of impact)	impact will have an effect)	one has predicted the significance of an impact)	With mitigation	Without mitigation	With mitigation	Without mitigation
Molteno Formation	permanent	international	unsure	beneficial	severe	beneficial	Negative

There is a possibility that tree fossils could be encountered during excavation of bedrock within the development footprint and these fossils would be of low significance. If effective mitigation measures are in place at the time of exposure, and the fossils are successfully excavated for study, this would represent a beneficial palaeontological impact.

7. PALAEONTOLOGICAL IMPACT AND MITIGATION

The predicted palaeontological impact of the development is based on the initial mapping assessment and literature reviews.

The Molteno Formation consists mainly of coarse-grained sandstone with thin layers of interbedded mudstone. The excavation of the underlying bedrock will have the potential to uncover the mud rock and sandstone of the Molteno Formation. Therefore, monitoring and mitigation in terms of the palaeontological heritage are required if the bedrock is exposed.

The following colour coding method was developed to classify a development area's palaeontological impact as illustrated in Figure 7.1:

- Red colouration indicates a very high possibility of finding fossils of a specific assemblage zone. Fossils will most probably be present in all outcrops on the site/route and the chances of finding fossils during the construction phase are very high.
- Orange colouration indicates a possibility of finding fossils of a specific assemblage zone either in outcrops or in bedrock on the site/route.
- Green colouration indicates that there is limit/no possibility of finding fossils in that section of the site/route development.




Figure 7.1 Palaeontological Impact of the Proposed Clocolan Waste Disposal Facility

The proposed development involves the excavation of waste disposal cells and infrastructure such as roads and buildings. The construction phase will require excavation of bedrock and has the potential to impact directly on fossil heritage if the Molteno mudstone and sandstone are exposed. From Figure 7.1 the following mitigation measures are recommended:

Colour Coding (Figure 7.1)	Mitigation Recommended	
	The resident ECO must be trained by a professional palaeontologist in	
	the recognition of fossils. If fossil material is later discovered it must	
	be appropriately protected and the discovery reported to a	
Orange Site	palaeontologist for the removal thereof as per SAHRA legislation. All	
	earth-moving activities within the bedrock are to be monitored by the	
	ECO under guidance of a palaeontologist. A monitoring report should	
	be submitted to SAHRA after completion of the earth-moving activity.	

Table 7.1	Site Specific Mitigation Measures
-----------	-----------------------------------

8. CONCLUSION

The proposed development site is underlain by the Triassic Molteno Formation that consists of coarse-grained grey sandstone and dark grey mudstone. Soils are derived from the underlying rock and are generally deep and relatively high in fertility.

The development site for the Clocolan Solid Waste Disposal Facility is underlain by the Triassic Molteno Formation. Deep soils are expected over the entire development site. There is a moderate potential for fossil material in the underlying mudstones that may be uncovered during excavations.

Through adequate monitoring and mitigation measures during excavations in the underlying bedrock the impact severity can be lowered to beneficial. The exposure and subsequent reporting of fossils (that would otherwise have remained undiscovered) to a qualified palaeontologist for excavation will have a beneficial palaeontological impact.

It is recommended that:

- The resident ECO must also be trained by a professional palaeontologist in the recognition of fossils. If fossil material is later discovered it must be appropriately protected and the discovery reported to a palaeontologist for the removal thereof as per SAHRA legislation.
- All earth-moving activities within the underlying bedrock with potential impact are to be monitored by a palaeontologist. A monitoring report should be submitted to SAHRA after completion of the earth-moving activities.

9. REFERENCES

Coastal & Environmental Services, 2011. Proposed Thomas River Energy Facility, Eastern Cape Province of South Africa, East London. Final Scoping Report, East London, South Africa.

Groenewald, G.H., 1996. Stratigraphy of the Tarkastad Subgroup, Karoo Supergroup, South Africa: Unpublished Ph.D. Thesis, University of Port Elizabeth, South Africa, 145 p.

Johnson MR , Anhaeusser CR and Thomas RJ (Eds), 2006. The Geology of South Africa. GSSA, Council for Geoscience, Pretoria, 691pp.

McCarthy, T. and Rubidge, B.S. 2005. The Story of Earth and Life. Struik Publishers, Cape T

10. QUALIFICATIONS AND EXPERIENCE OF THE AUTHOR

Dr Gideon Groenewald has a PhD in Geology from the Nelson Mandela Metropolitan University (1996) and the National Diploma in Nature Conservation from the University of South Africa (1990). He specialises in research on South African Permian and Triassic sedimentology and macrofossils with an interest in biostratigraphy, and palaeoecological aspects. He has extensive experience in the locating of fossil material in the Karoo Supergroup and has more than 20 years of experience in locating, collecting and curating fossils, including exploration field trips in search of new localities in the southern, western, eastern and north-eastern parts of the country. His publication record includes multiple articles in internationally recognized journals. Dr Groenewald is accredited by the Palaeontological Society of Southern Africa (society member for 25 years).

Declaration of Independence

I, Gideon Groenewald, declare that I am an independent specialist consultant and have no financial, personal or other interest in the proposed development, nor the developers or any of their subsidiaries, apart from fair remuneration for work performed in the delivery of palaeontological heritage assessment services. There are no circumstances that compromise the objectivity of my performing such work.

idea Grenewale G

Dr Gideon Groenewald Geologist

11. APPENDIX A - METHODOLOGY FOR ASSESSING THE SIGNIFICANCE OF IMPACTS

Although specialists will be given relatively free rein on how they conduct their research and obtain information, they will be required to provide their reports to the EAP in a specific layout and structure, so that a uniform specialist report volume can be produced.

To ensure a direct comparison between various specialist studies, a standard rating scale has been defined and will be used to assess and quantify the identified impacts. This is necessary since impacts have a number of parameters that need to be assessed. Four factors need to be considered when assessing the significance of impacts, namely:

- 1. Relationship of the impact to **temporal** scales the temporal scale defines the significance of the impact at various time scales, as an indication of the duration of the impact.
- 2. Relationship of the impact to **spatial** scales the spatial scale defines the physical extent of the impact.
- 3. The severity of the impact the **severity/beneficial** scale is used in order to scientifically evaluate how severe negative impacts would be, or how beneficial positive impacts would be on a particular affected system (for ecological impacts) or a particular affected party.

The severity of impacts can be evaluated with and without mitigation in order to demonstrate how serious the impact is when nothing is done about it. The word 'mitigation' means not just 'compensation', but also the ideas of containment and remedy. For beneficial impacts, optimization means anything that can enhance the benefits. However, mitigation or optimization must be practical, technically feasible and economically viable.

4. The **likelihood** of the impact occurs - the likelihood of impacts taking place as a result of project actions differs between potential impacts. There is no doubt that some impacts would occur (e.g. loss of vegetation), but other impacts are not as likely to occur (e.g. vehicle accident), and may or may not result from the proposed development. Although some impacts may have a severe effect, the likelihood of them occurring may affect their overall significance.

The *environmental significance* scale is an attempt to evaluate the importance of a particular impact. This evaluation needs to be undertaken in the relevant context, as an impact can either be ecological or social, or both. The evaluation of the significance of an impact relies heavily on the values of the person making the judgment. For this reason, impacts of especially a social nature need to reflect the values of the affected society.

Negative impacts that are ranked as being of "VERY HIGH" and "HIGH" significance will be investigated further to determine how the impact can be minimised or what alternative activities or mitigation measures can be implemented. These impacts may also assist decision makers i.e. lots of HIGH negative impacts may bring about a negative decision.

For impacts identified as having a negative impact of "**MODERATE**" significance, it is standard practice to investigate alternate activities and/or mitigation measures. The most effective and practical mitigations measures will then be proposed.

For impacts ranked as "LOW" significance, no investigations or alternatives will be considered. Possible management measures will be investigated to ensure that the impacts remain of low significance.

Table 9-1: Criterion used to rate the significance of an impact

Significance Rating Table				
Temporal Scale (The duration of the impact)				
Short term	Less than 5 years (Many construction phase impacts are of a short duration)			
Medium term	Between 5 and 20 years			
Long term	Between 20 and 40 years (From a human perspective almost permanent).			
Permanent	Over 40 years or resulting in a permanent and lasting change that will always be there			
	Spatial Scale (The area in which any impact will have an affect)			
Individual	Impacts affect an individual.			
Localised	Impacts affect a small area, often only a portion of the project area.			
Project Level	Impacts affect the entire project area.			
Surrounding Areas	Impacts that affect the area surrounding the development			
Municipal	Impacts affect either the Local Municipality, or any towns within them.			
Regional	Impacts affect the wider district municipality or the province as a whole.			
National	Impacts affect the entire country.			
International/Global	Impacts affect other countries or have a global influence.			
Will definitely occur	Impacts will definitely occur.			
Degree of Conf	idence or Certainty (The confidence to predicted the significance of an impact)			
Definite	More than 90% sure of a particular fact. Should have substantial supportive data.			
Probable	Over 70% sure of a particular fact, or of the likelihood of that impact occurring.			
Possible	Only over 40% sure of a particular fact or of the likelihood of an impact occurring.			
Unsure	Less than 40% sure of a particular fact or of the likelihood of an impact occurring.			

Table 9-2: The severity rating scale

Impact severity			
(The severity of negative impacts, or how beneficial positive	e impacts would be on a particular affected system or party)		
Very severe	Very beneficial		
An irreversible and permanent change to the affected	A permanent and very substantial benefit to the		
system(s) or party(ies) which cannot be mitigated. For	affected system(s) or party(ies), with no real		
example the permanent loss of land.	alternative to achieving this benefit. For example the		
	vast improvement of sewage effluent quality.		
Severe	Beneficial		
Long term impacts on the affected system(s) or	A long term impact and substantial benefit to the		
party(ies) that could be mitigated. However, this	affected system(s) or party(ies). Alternative ways of		
mitigation would be difficult, expensive or time	achieving this benefit would be difficult, expensive or		
consuming, or some combination of these. For	time consuming, or some combination of these. For		
example, the clearing of forest vegetation.	example an increase in the local economy.		
Moderately severe	Moderately beneficial		
Medium to long term impacts on the affected	A medium to long term impact of real benefit to the		
system(s) or party (ies), which could be mitigated.	affected system(s) or party(ies). Other ways of		
For example constructing the sewage treatment	optimising the beneficial effects are equally difficult,		
facility where there was vegetation with a low	expensive and time consuming (or some combination		
conservation value.	of these), as achieving them in this way. For example		
	a 'slight' improvement in sewage effluent quality.		
Slight	Slightly beneficial		
Medium or short term impacts on the affected	A short to medium term impact and negligible benefit		
system(s) or party(ies). Mitigation is very easy, cheap,	to the affected system(s) or party(ies). Other ways of		
less time consuming or not necessary. For example a	optimising the beneficial effects are easier, cheaper		
temporary fluctuation in the water table due to water	and quicker, or some combination of these.		
abstraction.			
No effect	Don't know/Can't know		
The system(s) or party(ies) is not affected by the	In certain cases it may not be possible to determine		
proposed development.	the severity of an impact		

Table 3: Overall significance appraisal

Querall Significance (The combination of all	the above criteria as an everall significance)			
	the above criteria as an overall significance)			
VERY HIGH NEGATIVE	VERY BENEFICIAL			
These impacts would be considered by society as constituting a major and usually permanent change to the (natural and/or social) environment, and usually result in severe or very severe effects, or				
	usually result in severe or very severe effects, or			
beneficial or very beneficial effects.				
	ed by informed society as being of VERY HIGH			
significance.				
	infrastructure in a rural area, which previously had			
	ted parties as resulting in benefits with VERY HIGH			
significance.				
	BENEFICIAL			
	ffects on the social and/or natural environment.			
	by society as constituting an important and usually			
	environment. Society would probably view these			
impacts in a serious light.				
	which is fairly common elsewhere, would have a			
significance rating of HIGH over the long term, as t				
	ct the natural system, and the impact on affected			
parties (such as people growing crops in the soil) w				
	SOME BENEFITS			
	long term effects on the social and/or natural			
	need to be considered by society as constituting a			
	to the (natural and/or social) environment. These			
impacts are real but not substantial.				
	on type of low diversity may be regarded as			
MODERATELY significant.				
	FEW BENEFITS			
	short term effects on the social and/or natural			
	be considered by the public and/or the specialist as			
	nort term change to the (natural and/or social)			
environment. These impacts are not substantial ar	•			
	e of a wetland habitat, as these systems is adapted			
to fluctuating water levels.				
	ple employed as a result of a development would			
only result in benefits of LOW significance to peop				
	IFICANCE			
There are no primary or secondary effects at all th				
Example: A change to the geology of a particular formation may be regarded as severe from a				
geological perspective, but is of NO significance in the overall context.				
DON'T KNOW In certain cases it may not be possible to determine the significance of an impact. For example, the				
	s on the social or natural environment given the			
available information.				
	nt on people's psychological perspective of the			
environment.				