

PALAEONTOLOGICAL IMPACT ASSESSMENT REPORT

PROPOSED REITZ SOLID WASTE FACILITY

Reitz, Free State Province of South Africa

***Farm: Townlands in the Nketoana Local Municipality within the
Thabo Mofutsanyane District Municipality***

Developer: Nketoana local Municipality

Consultant:



16 March 2012



Metsi Metseng Geological & Environmental Services

Suite 91
Private Bag X62
Bethlehem 9700
info@mmges.co.za
Fax 086 743 6864
+27 82 070 0735
+27 82 829 4978
www.mmges.co.za
VAT 4260183498
Reg 1999/30444/23

EXECUTIVE SUMMARY

The development of a Solid Waste Disposal Facility near Reitz in the Eastern Free State is an initiative of Nketoana Local Municipality. The purpose of this Palaeontological Impact Assessment is to identify exposed and potential palaeontological heritage on the site of the proposed development, to assess the impact the development may have on this resource, and to make recommendations as to how this impact might be mitigated.

The Nketoana Local Municipality plans to develop a solid waste disposal facility on the Townlands approximately 3 km north-east of Reitz Town in the Eastern Free State Province. The installation's footprint is approximately 11.5ha.

A basic assessment of the topography and geology of the area was made by using appropriate geological (1:250 000, 2728 Frankfort) map in conjunction with Google Earth. A review of the literature on the geological formations underlying the development site and the fossils that have been associated with these geological strata was undertaken.

The Reitz Waste Disposal Site development is underlain by the Late Permian to Early Triassic Normandien Formation of the Adelaide Subgroup that consists of grey mudstone, dark-grey shale, siltstone and sandstone. Soils are derived from the underlying rock and are generally deep and high in fertility.

The Adelaide Subgroup has a high palaeontological sensitivity rating. Through adequate monitoring and mitigation measures during excavations within the bedrock, the high impact severity can be lowered to beneficial. The exposure and subsequent reporting of fossils (that would otherwise have remained undiscovered) will be a beneficial palaeontological impact.

It is recommended that the resident ECO be trained by a professional palaeontologist in the recognition of fossil material. If fossil material is later discovered it must be appropriately protected and the discovery reported to a palaeontologist for the removal thereof.

That all earth-moving activities within the bedrock with a potential impact on the Adelaide Subgroup be monitored by a palaeontologist. That a monitoring report be submitted to SAHRA after the completion of the earth works phase.

SIGNIFICANCE RATING							
Rock Unit	Temporal Scale	Spatial Scale	Degree of Confidence	Impact Severity		Overall Significance	
				With mitigation	Without mitigation	With mitigation	Without mitigation
Adelaide Subgroup	permanent	international	possible	beneficial	very severe	beneficial	High negative

TABLE OF CONTENT

1. INTRODUCTION	1
1.1. Legal Requirements	1
2. PROPOSED DEVELOPMENT DESCRIPTION	1
3. AIMS AND METHODS	1
4. GEOLOGY OF THE AREA	2
4.1. The Adelaide Subgroup	2
4.1.1. The Normandien Formation.....	2
5. PALAEONTOLOGY OF THE AREA	4
5.1. The Adelaide Subgroup	4
5.1.1. The Normandien Formation.....	4
6. PALAEONTOLOGICAL SIGNIFICANCE AND RATING	4
7. PALAEONTOLOGICAL IMPACT AND MITIGATION	5
8. CONCLUSION.....	6
9. REFERENCES.....	7
10. QUALIFICATIONS AND EXPERIENCE OF THE AUTHOR	8
11. APPENDIX A - METHODOLOGY FOR ASSESSING THE SIGNIFICANCE OF IMPACTS	9

LIST OF FIGURES

Figure 2-1 Locality map of proposed development	2
Figure 4-1 The Geology (Geo Map 2728- Frankfort) of the Reitz Waste Site Development.....	3
Figure 7-1 Palaeontological Impact of the Proposed Reitz Waste Disposal Facility	5

LIST OF TABLES

Table 6.1 Palaeontological Significance of Geological Units on Site	4
Table 6.2 Significance Rating Table as Per CES Template.....	4
Table 7.1 Site Specific Mitigation Measures	6

1. INTRODUCTION

The development of a Solid Waste Disposal Facility near Reitz in the Eastern Free State is an initiative of the Nketoana Local Municipality. The purpose of this Palaeontological Impact Assessment is to identify exposed and potential palaeontological heritage on the site of the proposed development, to assess the impact the development may have on this resource, and to make recommendations as to how this impact might be mitigated.

1.1. Legal Requirements

This report forms part of the Scoping and Environmental Impact Assessment for the REITZ WASTE DISPOSAL SITE and complies with the requirements for the South African National Heritage Resource Act No 25 of 1999. In accordance with Section 38 (Heritage Resources Management), a Palaeontological Impact Assessment is required to assess any potential impacts to palaeontological heritage within the development footprint of the Senekal Solid Waste Disposal site.

Categories of heritage resources recognised as part of the National Estate in Section 3 of the Heritage Resources Act, and which therefore fall under its protection, include:

- geological sites of scientific or cultural importance;
- objects recovered from the soil or waters of South Africa, including archaeological and palaeontological objects and material, meteorites and rare geological specimens; and
- objects with the potential to yield information that will contribute to an understanding of South Africa's natural or cultural heritage.

2. PROPOSED DEVELOPMENT DESCRIPTION

The Nketoana Local Municipality plans to develop a solid waste disposal facility on the Townlands approximately 3 km north-east of Reitz Town in the Eastern Free State Province (See Locality Map Figure 2-1). The disposal site's footprint area is approximately 11.5ha. Initially the development will consist of 3 new waste cells which will be 30m wide, 140m long and 2m deep. Provision is further made for 2 more waste cells in the future.

3. AIMS AND METHODS

After discussions with LHL Engineers a request for a Screening Palaeontological Impact Assessment (PIA) was received. Following the *"SAHRA APM Guidelines: Minimum Standards for the Archaeological & Palaeontological Components of Impact Assessment Reports"* the aims of the PIA were:

- identifying exposed and subsurface rock formations that are considered to be palaeontologically significant;
- assessing the level of palaeontological significance of these formations;
- commenting on the impact of the development on these exposed and/or potential fossil resources;
- making recommendations as to how the developer should conserve or mitigate damage to these resources.

A basic assessment of the topography and geology of the area was made by using appropriate geological (1:250 000, 2728 Frankfort) maps in conjunction with Google Earth. The only limitation on this methodology is the scale of mapping, which restricts comparison of the geology to the 1:250 000 scale. This restriction only applies in areas where major changes in the geological character of the area occur over very short distances or on the geological transformation zones.

A review of the literature on the geological formations underlying the development site and the fossils that have been associated with these geological strata was undertaken.




Figure 3-1 Locality map of proposed development

4. GEOLOGY OF THE AREA

The entire development and surrounding area is underlain by the Normandien (Pne) Formation of the Adelaide (Pa) Subgroup of the Beaufort Group of the Karoo Supergroup. Quaternary (Yellow) sediments occur in the valley floors as illustrated in Figure 4.1.

4.1. The Adelaide Subgroup

The development site is underlain by the Late Permian to Early Triassic Adelaide Subgroup that consists of grey mudstone, dark-grey shale, siltstone and sandstone. Soils are derived from the underlying rock and are generally deep and relatively high in fertility.

4.1.1. The Normandien Formation

The Late Permian to Early Triassic Normandien Formation comprises a brightly coloured mudstone that underlies the prominent sandstone of the Katberg Formation of the Tarkastad Subgroup.





Figure 4-1 The Geology (Geo Map 2728- Frankfort) of the Reitz Waste Site Development

5. PALAEOLOGY OF THE AREA

5.1. The Adelaide Subgroup

The late Permian to Triassic Adelaide Subgroup can have a moderate to high potential for fossils from the *Dicynodon* and *Lystrosaurus* Assemblage Zones (Rubidge et al, 1995; Johnson et al, 2006). Plant fossils expected from these rocks include examples of *Glossopteris* assemblages and examples of other genera include *Cyclodendron*, *Phyllothea* and *Noeggerathiopsis*. Invertebrate fossils are restricted to trace fossils, including casts of some vertebrate burrows (Groenewald, 1996)

5.1.1. The Normandien Formation

The Late Permian to Early Triassic Normandien Formation of the Adelaide Subgroup is very productive and is palaeontologically known to contain fossils of the *Dicynodon* and *Lystrosaurus* Assemblage zones, including casts of vertebrate burrows (Groenewald, 1996).

6. PALAEOLOGICAL SIGNIFICANCE AND RATING

The predicted palaeontological impact of the development is based on the initial mapping assessment and literature reviews.

The palaeontological significance and rating is summarised in Table 7.1 and 7.2. For the methodology and definitions of impact rating and significance see Appendix A (CES 2011).

Table 6.1 Palaeontological Significance of Geological Units on Site

Geological Unit	Rock Type and Age	Fossil Heritage	Vertebrate Biozone	Palaeontological Sensitivity
Adelaide Subgroup and the Normandien Formation	Fluvial and lacustrine mudstones and sandstones. LATE PERMIAN TO EARLY TRIASSIC	Vertebrate fossils from the <i>Dicynodon</i> and <i>Lystrosaurus</i> assemblage zones can be expected. Plant fossils such as <i>Glossopteris</i> assemblages and other genera including <i>Cyclodendron</i> , <i>Phyllothea</i> and <i>Noeggerathiopsis</i> . Invertebrate fossils are restricted to trace fossils, including casts of some vertebrate burrows	<i>Dicynodon</i> and <i>Lystrosaurus</i> Assemblage Zones	High sensitivity

Table 6.2 Significance Rating Table as Per CES Template

Rock Unit	Temporal Scale (duration of impact)	Spatial Scale (area in which impact will have an effect)	Degree of confidence (confidence with which one has predicted the significance of an impact)	Impact severity (severity of negative impacts, or how beneficial positive impacts would be)		Overall Significance (The combination of all the other criteria as an overall significance)	
				With mitigation	Without mitigation	With mitigation	Without mitigation
Adelaide Subgroup	permanent	international	possible	beneficial	very severe	beneficial	High negative

There is a possibility that vertebrate fossils could be encountered during excavation of bedrock within the development footprint and these fossils would be of international significance. If effective mitigation measures are in place at the time of exposure, and the fossils are successfully excavated for study, this would represent a beneficial palaeontological impact.

Unfortunately within the Adelaide Subgroup, there is no way of assessing the likelihood of encountering vertebrate fossils during excavation. As evidenced in other similar areas with exposures, fossils were apparently absent or very scarce over large areas but locally dense accumulations were found.

Therefore, vertebrate fossils within the development site could be characterised as rare but highly significant. The damage and/or loss of these fossils due to inadequate mitigation would be a highly negative palaeontological impact. However, the exposure and subsequent reporting of fossils (that would otherwise have remained undiscovered) to a qualified palaeontologist for excavation will be a beneficial palaeontological impact.

7. PALAEOLOGICAL IMPACT AND MITIGATION

The predicted palaeontological impact of the development is based on the initial mapping assessment and literature reviews.

The Adelaide Subgroup is interbedded mud- and siltstone that do have potential to yield fossils. The excavation of the different cells on the slopes will have the potential to uncover the mud rock and sandstone of the Adelaide Subgroup. However, a geotechnical survey indicated soils deeper than 2m and therefore a small possibility exists that underlying bedrock may be uncovered. If underlying bedrock is uncovered then monitoring and mitigation in terms of the palaeontological heritage are required.




Figure 7-1 Palaeontological Impact of the Proposed Reitz Waste Disposal Facility

The following colour coding method was developed to classify a development area's palaeontological impact as illustrated in Figure 7-1:

- Red colouration indicates a very high possibility of finding fossils of a specific assemblage zone. Fossils will most probably be present in all outcrops on the site/route and the chances of finding fossils during the construction phase are very high.
- Orange colouration indicates a possibility of finding fossils of a specific assemblage zone either in outcrops or in bedrock on the site/route.
- Green colouration indicates that there is no possibility of finding fossils in that section of the site/route development.

The proposed development involves the excavation of waste disposal cells and infrastructure such as roads and buildings. The construction phase will require excavation of very deep soils and possibly bedrock and has the potential to impact directly on fossil heritage if the Adelaide Subgroup mudstone is exposed. From Figure 7.1 the following mitigation measures are recommended:

Table 7.1 Site Specific Mitigation Measures

Colour Coding (Figures. 7-1)	Mitigation Recommended
Orange Sites	The resident ECO must be trained by a professional palaeontologist in the recognition of fossils. If fossil material is later discovered it must be appropriately protected and the discovery reported to a palaeontologist for the removal thereof as per SAHRA legislation. All earth-moving activities within bedrock are to be monitored by a palaeontologist. A monitoring report should be submitted to SAHRA after completion of the earth-moving activity.

8. CONCLUSION

The development site for the Reitz Solid Waste Disposal Facility is underlain by the Late Permian Normandien Formation of the Adelaide Subgroup. Deep soils occur over the entire development site. There is a moderate potential for fossil material in the underlying mudstones that may be uncovered during excavations.

Through adequate monitoring and mitigation measures during excavations in the underlying bedrock the high impact severity can be lowered to beneficial. The exposure and subsequent reporting of fossils (that would otherwise have remained undiscovered) to a qualified palaeontologist for excavation will have a beneficial palaeontological impact.

It is recommended that:

- The resident ECO must also be trained by a professional palaeontologist in the recognition of fossils. If fossil material is later discovered it must be appropriately protected and the discovery reported to a palaeontologist for the removal thereof as per SAHRA legislation.
- All earth-moving activities within the underlying bedrock with potential impact are to be monitored by a palaeontologist. A monitoring report should be submitted to SAHRA after completion of the earth-moving activities.

9. REFERENCES

Coastal & Environmental Services, 2011. Proposed Thomas River Energy Facility, Eastern Cape Province of South Africa, East London. Final Scoping Report, East London, South Africa.

Groenewald, G.H., 1996. Stratigraphy of the Tarkastad Subgroup, Karoo Supergroup, South Africa: Unpublished Ph.D. Thesis, University of Port Elizabeth, South Africa, 145 p.

Johnson MR , Anhaeusser CR and Thomas RJ (Eds), 2006. The Geology of South Africa. GSSA, Council for Geoscience, Pretoria, 691pp.

McCarthy, T. and Rubidge, B.S. 2005. The Story of Earth and Life. Struik Publishers, Cape T

Rubidge, B.S. (Ed.). 1995. Biostratigraphy of the Beaufort Group (Karoo Supergroup). SACS Biostratigraphic Series, vol. 1.

10. QUALIFICATIONS AND EXPERIENCE OF THE AUTHOR

Dr Gideon Groenewald has a PhD in Geology from the Nelson Mandela Metropolitan University (1996) and the National Diploma in Nature Conservation from the University of South Africa (1990). He specialises in research on South African Permian and Triassic sedimentology and macrofossils with an interest in biostratigraphy, and palaeoecological aspects. He has extensive experience in the locating of fossil material in the Karoo Supergroup and has more than 20 years of experience in locating, collecting and curating fossils, including exploration field trips in search of new localities in the southern, western, eastern and north-eastern parts of the country. His publication record includes multiple articles in internationally recognized journals. Dr Groenewald is accredited by the Palaeontological Society of Southern Africa (society member for 25 years).

Declaration of Independence

I, Gideon Groenewald, declare that I am an independent specialist consultant and have no financial, personal or other interest in the proposed development, nor the developers or any of their subsidiaries, apart from fair remuneration for work performed in the delivery of palaeontological heritage assessment services. There are no circumstances that compromise the objectivity of my performing such work.

A handwritten signature in black ink, reading "Gideon Groenewald", with a horizontal line underneath it.

Dr Gideon Groenewald
Geologist

11. APPENDIX A - METHODOLOGY FOR ASSESSING THE SIGNIFICANCE OF IMPACTS

Although specialists will be given relatively free rein on how they conduct their research and obtain information, they will be required to provide their reports to the EAP in a specific layout and structure, so that a uniform specialist report volume can be produced.

To ensure a direct comparison between various specialist studies, a standard rating scale has been defined and will be used to assess and quantify the identified impacts. This is necessary since impacts have a number of parameters that need to be assessed. Four factors need to be considered when assessing the significance of impacts, namely:

1. Relationship of the impact to **temporal** scales - the temporal scale defines the significance of the impact at various time scales, as an indication of the duration of the impact.
2. Relationship of the impact to **spatial** scales - the spatial scale defines the physical extent of the impact.
3. The severity of the impact - the **severity/beneficial** scale is used in order to scientifically evaluate how severe negative impacts would be, or how beneficial positive impacts would be on a particular affected system (for ecological impacts) or a particular affected party.

The severity of impacts can be evaluated with and without mitigation in order to demonstrate how serious the impact is when nothing is done about it. The word 'mitigation' means not just 'compensation', but also the ideas of containment and remedy. For beneficial impacts, optimization means anything that can enhance the benefits. However, mitigation or optimization must be practical, technically feasible and economically viable.

4. The **likelihood** of the impact occurs - the likelihood of impacts taking place as a result of project actions differs between potential impacts. There is no doubt that some impacts would occur (e.g. loss of vegetation), but other impacts are not as likely to occur (e.g. vehicle accident), and may or may not result from the proposed development. Although some impacts may have a severe effect, the likelihood of them occurring may affect their overall significance.

The **environmental significance** scale is an attempt to evaluate the importance of a particular impact. This evaluation needs to be undertaken in the relevant context, as an impact can either be ecological or social, or both. The evaluation of the significance of an impact relies heavily on the values of the person making the judgment. For this reason, impacts of especially a social nature need to reflect the values of the affected society.

Negative impacts that are ranked as being of "**VERY HIGH**" and "**HIGH**" significance will be investigated further to determine how the impact can be minimised or what alternative activities or mitigation measures can be implemented. These impacts may also assist decision makers i.e. lots of **HIGH** negative impacts may bring about a negative decision.

For impacts identified as having a negative impact of "**MODERATE**" significance, it is standard practice to investigate alternate activities and/or mitigation measures. The most effective and practical mitigations measures will then be proposed.

For impacts ranked as "**LOW**" significance, no investigations or alternatives will be considered. Possible management measures will be investigated to ensure that the impacts remain of low significance.

Table 9-1: Criterion used to rate the significance of an impact

Significance Rating Table	
Temporal Scale (The duration of the impact)	
Short term	Less than 5 years (Many construction phase impacts are of a short duration)
Medium term	Between 5 and 20 years
Long term	Between 20 and 40 years (From a human perspective almost permanent).
Permanent	Over 40 years or resulting in a permanent and lasting change that will always be there
Spatial Scale (The area in which any impact will have an affect)	
Individual	Impacts affect an individual.
Localised	Impacts affect a small area, often only a portion of the project area.
Project Level	Impacts affect the entire project area.
Surrounding Areas	Impacts that affect the area surrounding the development
Municipal	Impacts affect either the Local Municipality, or any towns within them.
Regional	Impacts affect the wider district municipality or the province as a whole.
National	Impacts affect the entire country.
International/Global	Impacts affect other countries or have a global influence.
Will definitely occur	Impacts will definitely occur.
Degree of Confidence or Certainty (The confidence to predicted the significance of an impact)	
Definite	More than 90% sure of a particular fact. Should have substantial supportive data.
Probable	Over 70% sure of a particular fact, or of the likelihood of that impact occurring.
Possible	Only over 40% sure of a particular fact or of the likelihood of an impact occurring.
Unsure	Less than 40% sure of a particular fact or of the likelihood of an impact occurring.

Table 9-2: The severity rating scale

Impact severity	
(The severity of negative impacts, or how beneficial positive impacts would be on a particular affected system or party)	
Very severe	Very beneficial
An irreversible and permanent change to the affected system(s) or party(ies) which cannot be mitigated. For example the permanent loss of land.	A permanent and very substantial benefit to the affected system(s) or party(ies), with no real alternative to achieving this benefit. For example the vast improvement of sewage effluent quality.
Severe	Beneficial
Long term impacts on the affected system(s) or party(ies) that could be mitigated. However, this mitigation would be difficult, expensive or time consuming, or some combination of these. For example, the clearing of forest vegetation.	A long term impact and substantial benefit to the affected system(s) or party(ies). Alternative ways of achieving this benefit would be difficult, expensive or time consuming, or some combination of these. For example an increase in the local economy.
Moderately severe	Moderately beneficial
Medium to long term impacts on the affected system(s) or party (ies), which could be mitigated. For example constructing the sewage treatment facility where there was vegetation with a low conservation value.	A medium to long term impact of real benefit to the affected system(s) or party(ies). Other ways of optimising the beneficial effects are equally difficult, expensive and time consuming (or some combination of these), as achieving them in this way. For example a 'slight' improvement in sewage effluent quality.
Slight	Slightly beneficial
Medium or short term impacts on the affected system(s) or party(ies). Mitigation is very easy, cheap, less time consuming or not necessary. For example a temporary fluctuation in the water table due to water abstraction.	A short to medium term impact and negligible benefit to the affected system(s) or party(ies). Other ways of optimising the beneficial effects are easier, cheaper and quicker, or some combination of these.
No effect	Don't know/Can't know
The system(s) or party(ies) is not affected by the proposed development.	In certain cases it may not be possible to determine the severity of an impact

Table 3: Overall significance appraisal

Overall Significance (The combination of all the above criteria as an overall significance)	
VERY HIGH NEGATIVE	VERY BENEFICIAL
<p>These impacts would be considered by society as constituting a major and usually permanent change to the (natural and/or social) environment, and usually result in severe or very severe effects, or beneficial or very beneficial effects.</p> <p>Example: The loss of a species would be viewed by informed society as being of VERY HIGH significance.</p> <p>Example: The establishment of a large amount of infrastructure in a rural area, which previously had very few services, would be regarded by the affected parties as resulting in benefits with VERY HIGH significance.</p>	
HIGH NEGATIVE	BENEFICIAL
<p>These impacts will usually result in long term effects on the social and/or natural environment. Impacts rated as HIGH will need to be considered by society as constituting an important and usually long term change to the (natural and/or social) environment. Society would probably view these impacts in a serious light.</p> <p>Example: The loss of a diverse vegetation type, which is fairly common elsewhere, would have a significance rating of HIGH over the long term, as the area could be rehabilitated.</p> <p>Example: The change to soil conditions will impact the natural system, and the impact on affected parties (such as people growing crops in the soil) would be HIGH.</p>	
MODERATE NEGATIVE	SOME BENEFITS
<p>These impacts will usually result in medium to long term effects on the social and/or natural environment. Impacts rated as MODERATE will need to be considered by society as constituting a fairly important and usually medium term change to the (natural and/or social) environment. These impacts are real but not substantial.</p> <p>Example: The loss of a sparse, open vegetation type of low diversity may be regarded as MODERATELY significant.</p>	
LOW NEGATIVE	FEW BENEFITS
<p>These impacts will usually result in medium to short term effects on the social and/or natural environment. Impacts rated as LOW will need to be considered by the public and/or the specialist as constituting a fairly unimportant and usually short term change to the (natural and/or social) environment. These impacts are not substantial and are likely to have little real effect.</p> <p>Example: The temporary change in the water table of a wetland habitat, as these systems is adapted to fluctuating water levels.</p> <p>Example: The increased earning potential of people employed as a result of a development would only result in benefits of LOW significance to people who live some distance away.</p>	
NO SIGNIFICANCE	
<p>There are no primary or secondary effects at all that are important to scientists or the public.</p> <p>Example: A change to the geology of a particular formation may be regarded as severe from a geological perspective, but is of NO significance in the overall context.</p>	
DON'T KNOW	
<p>In certain cases it may not be possible to determine the significance of an impact. For example, the significance of the primary or secondary impacts on the social or natural environment given the available information.</p> <p>Example: The effect of a particular development on people's psychological perspective of the environment.</p>	